

2015

# Spatiotemporal Correlations between Water Footprint and Agricultural Inputs: A Case Study of Maize Production in Northeast China

Peili Duan

Lijie Qin

*See next page for additional authors*

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Follow this and additional works at: [https://digitalcommons.uri.edu/nrs\\_facpubs](https://digitalcommons.uri.edu/nrs_facpubs)

---

## Citation/Publisher Attribution

Duan, P., Qin, L., Wang, Y., & He, H. (2015). Spatiotemporal Correlations between Water Footprint and Agricultural Inputs: A Case Study of Maize Production in Northeast China. *Water*, 7(8), 4026-4040. doi: 10.3390/w7084026  
Available at: <http://dx.doi.org/10.3390/w7084026>

This Article is brought to you for free and open access by the Natural Resources Science at DigitalCommons@URI. It has been accepted for inclusion in Natural Resources Science Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact [digitalcommons@etal.uri.edu](mailto:digitalcommons@etal.uri.edu).

---

**Authors**

Peili Duan, Lijie Qin, Yeqiao Wang, and Hongshi He

Article

## Spatiotemporal Correlations between Water Footprint and Agricultural Inputs: A Case Study of Maize Production in Northeast China

Peili Duan <sup>1</sup>, Lijie Qin <sup>1,\*</sup>, Yeqiao Wang <sup>2</sup> and Hongshi He <sup>1,3</sup>

<sup>1</sup> School of Geographical Sciences, Northeast Normal University, 5268 Renmin Street, Changchun 130024, China; E-Mails: duanpl069@nenu.edu.cn (P.D.); heh@missouri.edu (H.H.)

<sup>2</sup> Department of Natural Resources science, University of Rhode Island, Kingston, RI 02881, USA; E-Mail: yqwang@uri.edu

<sup>3</sup> School of Natural Resources, University of Missouri, Columbia, MO 65211, USA

\* Author to whom correspondence should be addressed; E-Mail: qinlj953@nenu.edu.cn; Tel.: +86-0431-8509-9123.

Academic Editor: Markus Disse

Received: 21 April 2015 / Accepted: 15 July 2015 / Published: 23 July 2015

---

**Abstract:** To effectively manage water resources in agricultural production, it is necessary to understand the spatiotemporal variation of the water footprint (WF) and the influences of agricultural inputs. Employing spatial autocorrelation analysis and a geographically weighted regression (GWR) model, we explored the spatial variations of the WF and their relationships with agricultural inputs from 1998 to 2012 in Northeast China. The results indicated that: (1) the spatial distribution of WFs for the 36 major maize production prefectures was heterogeneous in Northeast China; (2) a cluster of high WFs was found in southeast Liaoning Province, while a cluster of low WFs was found in central Jilin Province, and (3) spatial and temporal differentiation in the correlations between the WF of maize production and agricultural inputs existed according to the GWR model. These correlations increased over time. Our results suggested that localized strategies for reducing the WF should be formulated based on specific relationships between the WF and agricultural inputs.

**Keywords:** spatiotemporal differences; agricultural inputs; geographically weighted regression; water footprint; maize; Northeast China

---

## 1. Introduction

Freshwater has an uneven temporal and spatial distribution from the perspective of availability and quantity. A growing population and increased demand for agricultural products have led to increasing competition for freshwater resources over the past several decades [1–3]. As a sector of intense water usage, agriculture is the largest user of freshwater, accounting for roughly 70% of total water usage [4–6]. Water scarcity is a major constraint in some agricultural production activities. Water usage for producing crops is expected to increase by 0.7% per year from its estimated level of 6400 billion  $\text{m}^3 \cdot \text{year}^{-1}$  in 2000 to 9060 billion  $\text{m}^3 \cdot \text{year}^{-1}$  by 2050 as a result of the need to feed a projected global population of 9.2 billion [7,8]. As a result, it is important to evaluate water resource utilization in agricultural production to meet future freshwater supply challenges [9,10].

The water footprint (WF) concept provides a new way of evaluating water utilization in agricultural production [11]. For a given crop, the WF refers to the total amount of freshwater that is consumed and contaminated in the crop production process. It contains three components: the amount of rainwater consumed, the amount of irrigation water in the form of surface or ground water consumed, and the amount of freshwater required to assimilate the load of pollutants [12]. These components are defined as green WF, blue WF and grey WF, respectively [13,14]. The WF has led to a better understanding of the water requirement of crop production, which is attracting increased attention and has motivated a large number of studies [15–19]. Thus far, several studies have shed light on the spatial differences in the WFs of a variety of types of crops production in China. Sun *et al.* assessed the spatiotemporal variability of the WF for grain in an irrigation district of China during 1960–2008 [20]. Tian *et al.* analysed the temporal and spatial variation of the WF of China's five major food crops from 1978 to 2010 [21]. Wang *et al.* calculated and compared the WF of each grain crop and the integrated WFs of grain products with actual and virtual crop patterns in various regions of China for 2010 [22]. These previous studies suggested that the WF exhibits variability in different regions at the national level. However, to the best of our knowledge, few studies have tried to explore the spatial pattern characteristics of the WF at the prefecture level. Studies regarding the WF prefecture levels are more important because they can directly relate water usage to crop production. Besides, the underlying factors influencing WFs were mostly investigated from the perspective of climate. In fact, the WF of crop production primarily depends on agricultural inputs rather than the regional climate and its variation [23]. Therefore, it is essential to investigate the relationships between the WF and agricultural inputs.

To better understand the correlations between the WF of crop production and agricultural inputs, an integrated approach is much needed. The geographically weighted regression (GWR) model is an expansion of the classical multivariate linear regression model, which considers not only the spatial non-stationary of relationships but also, in part, the spatial dependence, as it relies on local rather than global parameters [24]. Additionally, the GWR model has been applied in social economics, urban geography, meteorology, forestry and many other disciplines [25–27]. Few studies have been conducted on the analysis of correlations between the WF and agricultural inputs over space with the GWR method.


We selected Northeast China, which provides one-third of China's food production, as our study region. In this region, maize is one of the primary crops. The major objectives of this paper were: (1) to determine the spatial distribution of the WF of maize production at the prefecture level; (2) to identify the spatial patterns and clusters of the WF using global and local spatial autocorrelation analyses; and

(3) to analyse the spatial relationship between the WF and the three agricultural inputs based on the GWR model. This study could provide valuable information for designing and implementing effective measures for reducing the WF of maize production in Northeast China.


## 2. Material and Methods

### 2.1. Study Area

Northeast China is located between 118°53'–135°05' E and 38°43'–53°33' N, and consists of Liaoning, Jilin, and Heilongjiang Provinces (Figure 1). The region covers an area of  $7.92 \times 10^5$  km<sup>2</sup>, of which farmland occupies  $2.64 \times 10^5$  ha, representing 16.5% of the total arable land in China [28]. The maize sowing area and yield at the prefecture level in 2012 are shown in Figure 2. The climate in the region is cold temperate and humid in the north and semi-humid in the west. The mean annual air temperature ranged from −4.2 °C to 10.9 °C, decreasing from south to north. The precipitation is generally concentrated in the summer and autumn (from May to October) [29], which coincides with the growing period of maize in Northeast China. The annual precipitation decreased from the southeast (1070 mm) to the northwest (450 mm). Maize, as the major grain crop in Northeast China, is highly seasonal with a single crop each year. The differences in the physical geography significantly affect the growth periods and crop coefficients of maize in each province [30] (Table 1).



**Figure 1.** Location of Northeast China and the names of its 36 prefectures in three provinces.



**Figure 2.** The maize sowing area and yield at the prefecture level in 2012.

**Table 1.** Growth periods and literature values of crop coefficient  $K_c$  for maize in Liaoning, Jilin and Heilongjiang Provinces.

Province	Growth Periods			Crop Coefficients ( $K_c$ )		
	Sowing Date	Harvesting Date	Total Days	Initial Season ( $K_{c\ ini}$ )	Mid-Season ( $K_{c\ mid}$ )	Late-Season ( $K_{c\ late}$ )
Liaoning	25-April	20-September	149	0.38	1.22	0.60
Jilin	1-May	24-September	147	0.40	1.22	0.58
Heilongjiang	16-May	30-September	137	0.36	1.20	0.55


Note: Source: China Meteorological Data Sharing Service System and Huang *et al.* (2011) [30].

### 2.2. Data Sources

The climate data of 36 weather stations, one location per prefecture, were taken from the China Meteorological Data Sharing Service System [31] for the period of 1998 to 2012, including monthly average maximum and minimum temperature, relative humidity, wind speed, sunshine hours and precipitation. The agricultural data, including maize yield, maize sowing area, agricultural machinery power, effective irrigation area and consumption of chemical fertilizers, were obtained from the statistical yearbooks 1998–2012 of the three provinces.

### 2.3. Methods

First, the WFs of maize production were accounted for in Northeast China. Next, the spatial patterns of the WF were investigated. Finally, the relationships between the WF and agricultural inputs were explored. The overall calculation process flow chart is diagrammed in Figure 3.



**Figure 3.** Diagram for WF assessment and the correlations between WF and agricultural inputs.

2.3.1. WF Calculation Methodology

The WF of crop production was estimated following the calculation framework developed by Hoekstra *et al.* [14], which is the total water consumption during crop growth process, including blue WF, green WF and grey WF [32], namely:

$$WF_{total} = WF_{green} + WF_{blue} + WF_{grey} \tag{1}$$

where  $WF_{total}$  is the WF of crop production ( $m^3 \cdot ton^{-1}$ );  $WF_{green}$  is the green WF ( $m^3 \cdot ton^{-1}$ );  $WF_{blue}$  is the blue WF ( $m^3 \cdot ton^{-1}$ ); and  $WF_{grey}$  is the grey WF ( $m^3 \cdot ton^{-1}$ ).

The  $WF_{green}$  and  $WF_{blue}$  are calculated as crop water use divided by crop yield. The green and blue water use is calculated by accumulation of daily evapotranspiration over the growing period.

$$WF_{green} = \frac{CWU_{green}}{Y} = 10 \times \frac{ET_{green}}{Y} \tag{2}$$

$$WF_{blue} = \frac{CWU_{blue}}{Y} = 10 \times \frac{ET_{blue}}{Y} \tag{3}$$

where  $CWU_{green}$  and  $CWU_{blue}$  are green and blue water usage (mm);  $ET_{green}$  and  $ET_{blue}$  are green and blue water evapotranspiration (mm);  $Y$  is the crop yield ( $ton \cdot ha^{-1}$ ); and the factor 10 converts water depths in millimetres into water volumes per land surface in  $m^3 \cdot ha^{-1}$ .

The  $WF_{\text{grey}}$  is calculated as the chemical application rate to the field per hectare ( $AR$ ,  $\text{kg}\cdot\text{ha}^{-1}$ ) times the leaching-run-off fraction ( $\alpha$ ) divided by the maximum acceptable concentration ( $C_{\text{max}}$ ,  $\text{kg}\cdot\text{m}^{-3}$ ) minus the natural concentration for the pollutant considered ( $C_{\text{nat}}$ ,  $\text{kg}\cdot\text{m}^{-3}$ ) and divided by the crop yield ( $Y$ ,  $\text{ton}\cdot\text{ha}^{-1}$ ).

$$WF_{\text{grey}} = \frac{(\alpha \times AR) / (C_{\text{max}} - C_{\text{nat}})}{Y} \quad (4)$$

Considering the availability of local data, in this study, we quantified the grey WF related to nitrogen use only. We have further taken  $\alpha$  as 10% following Chapagain *et al.* [12],  $C_{\text{nat}}$  as zero, and  $C_{\text{max}}$  as  $12 \text{ mg}\cdot\text{L}^{-1}$  following China water quality standards.

### 2.3.2. ET Calculations Based on the CROPWAT Model

The green and blue water evapotranspiration are estimated based on crop water requirement by employing the CROPWAT 8.0 model [33,34], which are calculated as:

$$ET_{\text{green}} = \min(ET_c, P_{\text{eff}}) \quad (5)$$

$$ET_{\text{blue}} = \max(0, ET_c - P_{\text{eff}}) \quad (6)$$

where  $ET_c$  is the crop evapotranspiration and  $P_{\text{eff}}$  is the effective precipitation over the crop growing period (mm).

$ET_c$  is calculated by employing the CROPWAT model based on the FAO Penman-Monteith method [35].

$$ET_c = K_c ET_0 \quad (7)$$

where  $K_c$  is the crop coefficient;  $ET_0$  is the reference evapotranspiration, calculated as follows:

$$ET_0 = \frac{0.408\Delta(R_n - G) + \gamma \frac{900}{T + 273} U_2 (e_s - e_a)}{\Delta + \gamma(1 + 0.34U_2)} \quad (8)$$

where  $R_n$  is the net radiation ( $\text{MJ}\cdot\text{m}^{-2}\cdot\text{d}^{-1}$ );  $G$  the soil heat flux ( $\text{MJ}\cdot\text{m}^{-2}\cdot\text{d}^{-1}$ );  $T$  the average air temperature ( $^{\circ}\text{C}$ );  $U_2$  the wind speed at 2 m height ( $\text{m}\cdot\text{s}^{-1}$ );  $e_s$  the saturation vapor pressure (kPa);  $e_a$  the actual vapor pressure (kPa);  $\Delta$  the slope of saturation vapor pressure *versus* air temperature curve ( $\text{kPa}\cdot^{\circ}\text{C}^{-1}$ ); the hygrometer constant ( $\text{kPa}\cdot^{\circ}\text{C}^{-1}$ ).

$P_{\text{eff}}$  is calculated according to a formula developed by the USDA Soil Conservation Service [34,36]:

$$P_{\text{eff}} = \begin{cases} P/125(125 - 0.2P) & P \leq 250 \\ 125 + 0.1P & P > 250 \end{cases} \quad (9)$$

where  $P$  is the precipitation at a month step (mm).

### 2.3.3. Spatial Autocorrelation

Both global and local spatial autocorrelation test results are used to confirm the presence of strong spatial dependence and heterogeneity in the distribution of the WF, supporting the use of the GWR technique.



Global Moran's  $I$  is applied to detect the overall degree of spatial autocorrelation among the regions. The equation is given as:

$$I = \frac{n \sum_{i=1}^n \sum_{j=1}^n W_{ij} (X_i - \bar{X})(X_j - \bar{X})}{\left( \sum_{i=1}^n \sum_{j=1}^n W_{ij} \right) \cdot \sum_{i=1}^n (X_i - \bar{X})^2} \quad (10)$$

where  $n$  is the number of spatial unit;  $X_i$  and  $X_j$  are the values of WF of spatial units  $i$  and  $j$ , respectively;  $\bar{X}$  is the mean of  $X$ ;  $W_{ij}$  is a matrix of spatial weight applied to the comparison between spatial units  $i$  and  $j$ . Generally,  $W$  is defined using an adjacency standard. When the units  $i$  are adjacent to units  $j$ ,  $W_{ij}$  is equal to 1, otherwise  $W_{ij}$  is equal to 0. The value of Moran's  $I$  varies between  $-1$  and  $1$ . The significance of the spatial autocorrelation can be tested by  $Z$  statistics, which is given as:

$$Z = \frac{1 - E(I)}{\sqrt{\text{Var}(I)}} \quad (11)$$

where  $E(I)$  and  $\text{Var}(I)$  denote the expectation value and the variance of Moran's  $I$ , respectively [37]. If  $Z$ -value is greater than 0 and statistically significant, an obvious positive autocorrelation exists in the spatial distribution [38].

The local indicator of spatial autocorrelation index (LISA) takes each spatial unit as the inspection object and analyses its properties of spatial autocorrelation [39]. The equation is given as:

$$I_i = Z_i \sum W_{ij} Z_j \quad (12)$$

where  $I_i$  is the local autocorrelation index of WF and  $Z_i$  and  $Z_j$  are the standardization of values of WF of spatial units  $i$  and  $j$ .

#### 2.3.4. Geographically Weighted Regression (GWR) Model

The GWR model extends the traditional regression framework by embedding the spatial location of the data into the regression parameter [40,41]. The local estimation of the parameters with GWR is expressed as follows [42]:

$$y_i = \beta_0(u_i, v_i) + \sum_k \beta_k(u_i, v_i) x_{ik} + \varepsilon_i \quad i = 1, 2, \dots, n \quad (13)$$

where  $y_i$  is the dependent variable in the  $i$ th location;  $x_{ik}$  is the  $i$ th value of the  $k$ th independent variable;  $k$  represents the number of independent variables;  $i$  represents the number of samplings;  $(u_i, v_i)$  denotes the coordinates;  $\beta_0(u_i, v_i)$  represents the intercept value, and  $\beta_k(u_i, v_i)$  is the continued function;  $\varepsilon_i$  represents the model residual.

To calculate the spatial distribution of the weightings, the optimal bandwidth is required. The Cross-Validation (CV) or Akaike Information Criterion (AIC) is usually used for calculation. In this paper the AIC was chosen, which was fixed by the maximum likelihood principle to determine the optimal bandwidth [43].

GWR model analysis was performed using the software ArcGIS10.1. In this study, the WF of maize production was treated as the dependent variable, and three agricultural inputs, namely, agricultural


machinery power (AMP), effective irrigation area (EIA) and chemical fertilizers consumption (CFC) were chosen as the independent variables of the models.

### 3. Results

#### 3.1. Spatial Distribution of WF of Maize Production

From 1998 to 2012, the average annual WF of maize production was  $1029 \text{ m}^3 \cdot \text{ton}^{-1}$  in Northeast China. Green WF was the biggest part of the total WF of maize production with the proportion at 51%, followed by grey and blue WF, which accounted for 28% and 21%, respectively. This suggests that the WF of maize production in this region largely relies on green water resources, *i.e.*, consumptive use of rainwater, which had a non-uniform seasonal distribution.

The WF of maize production exhibited significant differences on the prefecture level. As can be seen from 1998, the highest values of WF were observed in Baicheng, where it was  $1608 \text{ m}^3 \cdot \text{ton}^{-1}$  (Figure 4). This could be explained by the highest evapotranspiration. Dandong took second place with  $1347 \text{ m}^3 \cdot \text{ton}^{-1}$ . The lowest WFs were calculated for Siping with  $691 \text{ m}^3 \cdot \text{ton}^{-1}$ , and Tieling with  $756 \text{ m}^3 \cdot \text{ton}^{-1}$ , which could be explained by the higher maize yields in these two prefectures. Compared to 1998, the WFs of all prefectures showed a decreasing trend in 2012. Dandong had replaced Baicheng with the highest WF, which was  $1313 \text{ m}^3 \cdot \text{ton}^{-1}$ , followed by Baicheng with  $1291 \text{ m}^3 \cdot \text{ton}^{-1}$ . Tieling had replaced Siping as the prefecture with the lowest WF, which was  $667 \text{ m}^3 \cdot \text{ton}^{-1}$ . Siping ranked second, with a value of  $672 \text{ m}^3 \cdot \text{ton}^{-1}$ .



**Figure 4.** Spatial distribution of WF of maize production in 1998 and 2012.

#### 3.2. Spatial Pattern Characteristics of the WF of Maize Production

Both the global and local spatial autocorrelation test results confirm the presence of strong spatial dependence and heterogeneity in the distribution of the WF, supporting the use of the GWR technique.


Global Moran's  $I$  of WFs in 1998 and 2012 (Table 2) were 0.5512 and 0.5530, respectively, and all  $Z$ -values were significant ( $p$ -value = 0.01, with 999 permutations), which demonstrated a positive spatial

correlation of prefecture-level WFs in the entire study region. In other words, prefecture units with higher and lower levels of WFs were spatially clustered. Meanwhile, the spatial autocorrelation of WFs in 2012 was slightly stronger than that in 1998.

**Table 2.** Global Moran's  $I$  of water footprints (WFs) of maize production in 1998 and 2012.

Year	Moran's $I$	Z-Value	$p$ -Value
1998	0.5512	2.6366	0.01
2012	0.5530	2.4741	0.01

As for the local spatial autocorrelation, LISA statistics can be classified into four categories and mapped in a LISA cluster map (Figure 5). For instance, prefectures that have high WFs surrounded by high WFs (HH) and those with low WFs surrounded by low WFs (LL) represent potential spatial clusters. Prefectures with high WFs surrounded by low WFs (HL) and low WFs surrounded by high WFs (LH) suggest potential spatial outliers. The cluster of high WF values (HH) was located in southeast Liaoning in 1998 (composed of the three prefectures of Dandong, Anshan and Yingkou), but it increased in 2012 (composed of the five prefectures of Dandong, Liaoyang, Anshan, Yingkou and Dalian), indicating that the WF disparity in this region decreased. The cluster of low WF values (LL) included the four prefectures of Changchun, Siping, Liaoyuan and Tieling, which were primarily located in central Jilin Province, and the number of prefectures in this cluster remained relatively stable. Other forms of local spatial autocorrelation included one HL-type prefecture (Yanbian) in both years.



**Figure 5.** LISA cluster map for WFs of maize production in 1998 and 2012.

### 3.3. Spatial Variations in Correlations between Agricultural Inputs and the WF of Maize Production

Table 3 summarizes the results of the GWR between the WF and agricultural inputs. Results of the GWR models demonstrated that the  $R^2$  and adjusted  $R^2$  values in 1998 and 2012 were approximately equal, both representing over 63% of the total variance of the WF. The local parameters are described by six-number summaries consisting of the minimum, maximum, mean, lower quartile, median and upper


quartile values, which presented large differences. In 1998, the correlation between the EIA and the WF was still positive; however, the AMP and CFC were either positive or negative. In 2012, the correlations between the AMP, EIA and the WF were positive and the CFC was negative.

**Table 3.** Statistical description of the geographically weighted regression (GWR) model parameter for 1998 and 2012.

Descriptive Index	1998			2012		
	AMP	EIA	CFC	AMP	EIA	CFC
Minimum value	-2.283	0.464	-1.624	-3.746	1.653	-2.624
Maximum value	0.933	2.033	0.369	-0.517	2.313	-0.347
Mean	-0.591	1.315	-0.657	-1.550	1.977	-1.598
Upper quartile	-1.610	1.091	-1.113	-2.262	1.814	-1.902
Median	-0.350	1.373	-0.648	-1.039	1.961	-1.547
Lower quartile	0.460	1.539	-0.205	-0.748	2.174	-1.368
AIC	481.242			487.305		
R <sup>2</sup>	0.635			0.631		
Adjusted R <sup>2</sup>	0.581			0.593		

Notes: AMP: Agricultural machinery power; EIA: Effective irrigation area; CFC: Chemical fertilizers consumption.

The relationships between the WF and agricultural inputs were spatially non-stationary (Figures 6 and 7). In 1998, most regression coefficients of the AMP were found to be negative, except for south-central Liaoning Province. The CFC had a positive correlation with the WF in most areas, other than south Liaoning Province. The EIA had a positive correlation with the WF in the entire region. Compared to 1998, the regression coefficients of all agricultural inputs increased in 2012, suggesting the correlations were becoming stronger. Overall the correlations of the AMP and CFC were changing from positive to negative. The AMP and CFC had negative correlation with the WF in the entire region, while the EIA still had a positive correlation with the WF. The absolute value of regression coefficient of the AMP, EIA and CFC all exhibited a decreasing trend from the centre to the south and north.



**Figure 6.** Spatial distribution of regression coefficients of agricultural inputs in 1998.




Figure 7. Spatial distribution of regression coefficients of agricultural inputs in 2012.

#### 4. Discussion

In this study, we mapped the distribution of the WF of maize production at the prefecture level in Northeast China. A combination of the climatic condition, growing period and fertilizer consumption lead to the diversity of the WFs among 36 prefectures. The distribution of the WF in Northeast China was clustered, indicated by the significantly global Moran's  $I$  values. The LISA method detected the regions with high or low concentrations of WF. The spatial clustered regions with high WF of maize production were distributed in southeast Liaoning Province. The soils in these regions are poor and essentially unsuitable for crop production. The spatial clustered regions with low WF of maize production were distributed in central Jilin Province, which is located on the Songnen Plain and considered the Golden-Maize-Belt in China. It can be attributed to such factors as favourable climate condition, soil quality, proper irrigation facilities and relatively higher maize yield. These findings can help policy-makers prioritize maize production area and optimize maize production distribution.

The factors impacting the WF are miscellaneous. To explore the WF and its influence mechanisms, scholars began to study the relationships between WF and individual factors, such as climatic factors and agricultural input [23,44,45]. The methodology employed in this study only included correlation analysis. However, in reality, the relationships between the WF and influential factors are not always consistent in different areas. In particular, we employed the GWR technique to examine the spatial and temporal differences in the correlation between three agricultural inputs and the WF of maize production, including the AMP, EIA and CFC. The regression coefficients of the GWR also displayed a high variability in space, which could better explain the spatial differences of the correlation between the agricultural inputs and the WF. The spatial variations in the relationships are affected by the level of agricultural development.

In addition, the GWR analysis can help decision-makers clearly see the results of past relationships between the WF and agricultural inputs on a local scale and make appropriate decisions for future planning to reduce the WF of maize production. According to the results of the GWR model, the correlation between the AMP and WF was mainly positive in the study area. Attention should be paid to improving the mechanical technologies. The AMP in south-central Liaoning shifted from a positive to a

negative correlation with the WF, so the amount of machinery should be adjusted appropriately. The correlation between the EIA and WF was positive in the study area. Thus, irrigation water should be reduced in this region. Water-saving irrigation techniques, such as drip or sprinkler irrigation, should be employed appropriately to inhibit the growing WF. The CFC had a negative correlation with the WF, which indicated that properly managed fertilizer use contributed to reducing the WF. Meanwhile, the CFC shifted from a positive to a negative correlation with the WF in south Liaoning. In these areas, increasing fertilizer usage or adjusting the fertilizer proportions and increasing the usage of organic fertilizers should be considered.

To the best of our knowledge, this is the first study focusing on the spatial pattern of the WF and the relationships between the WF and agricultural inputs. Nevertheless, there are still limitations in this study. There are some uncertainties in the quantification of the WF of maize production. On the one hand, the agricultural data were primarily derived from statistical data, and there are discrepancies between these values and reality. On the other hand, the grey WF was estimated based on a simplified approach, which left out local factors that influence the precise leaching rates. In addition, our spatial analysis was based on the prefecture scale, but the distribution of the WF was also not homogeneous at the prefecture level. Therefore, this study cannot identify the spatial heterogeneity within each unit.

## 5. Conclusions

The spatial distribution and cluster areas of the WFs of maize production in Northeast China were investigated and the relationships between the WF and agricultural inputs were explored. The results demonstrated that the spatial differences of the WFs of maize production were obvious in Northeast China. Two cluster regions were detected. The prefectures with higher WFs were formed in southeast Liaoning Province while the prefectures with lower WFs were formed in central Jilin Province. From the perspective of the GWR results, the WF and agricultural inputs relationships were significantly spatially non-stationary. Finally, the combination of the GWR along with the GIS approaches constitutes a powerful tool for better displaying spatiotemporally varying relationships between the WF and agricultural inputs. Extension of this study could focus on adding other driving forces, such as population, plantation structure and policy in the spatiotemporal analysis of the WF in the studied areas.

## Acknowledgments

This work was funded by the Major Science and Technology Program for Water Pollution Control and Treatment (2014ZX07201-011), the Natural Science Foundation of China (41171038), the Humanity and Social Science Foundation of Ministry of Education (10YJA840032), and the Fundamental Research Funds for the Central Universities (12SSXT150).

## Author Contributions

Peili Duan carried out the calculation, result analysis and drafted the manuscript, which was revised by all authors. All authors read and approved the final manuscript.

## Conflicts of Interest

The authors declare no conflict of interest.

## References

1. Falkenmark, M.; Rockstrom, J.; Karlberg, L. Present and future water requirements for feeding humanity. *Food Secur.* **2009**, *1*, 59–69.
2. De Fraiture, C.; Wichelns, D. Satisfying future water demands for agriculture. *Agric Water Manag.* **2010**, *97*, 502–511.
3. Erkin, A.E.; Hoekstra, A.Y. Water footprint scenarios for 2050: A global analysis. *Environ. Int.* **2014**, *64*, 71–82.
4. Molden, D.; Oweis, T.; Pasquale, S.; Kijne, J.; Hanjra, M.A.; Bindraban, P.; Bouman, B.; Cook, S.; Erenstein, O.; Farahani, H. Water for food, water for life, a comprehensive assessment of water management in agriculture. In *Pathways for Increasing Agricultural Water Productivity*; Earthscan and International Water Management Institute: London, UK; Colombo, Sri Lanka, 2007; pp. 279–310.
5. Xiong, W.; Holman, I.; Lin, E.D.; Conway, D.; Jiang, J.H.; Xu, Y.L.; Li, Y. Climate change, water availability and future cereal production in China. *Agric. Ecosyst. Environ.* **2010**, *135*, 58–69.
6. Fader, M.; Gerten, D.; Thamm, M.; Heinke, J.; Lotze-Campen, H.; Lucht, W.; Cramer, W. Internal and external green-blue agricultural water footprints of nations, and related water and land savings through trade. *Hydrol. Earth Syst. Sci.* **2011**, *15*, 1641–1660.
7. Rosegrant, M.W.; Ringler, C.; Zhu, T. Water for agriculture: Maintaining food security under growing scarcity. *Annu. Rev. Environ. Resour.* **2009**, *34*, 205–222.
8. Mekonnen, M.M.; Hoekstra, A.Y. Water footprint benchmarks for crop production: A first global assessment. *Ecol. Indic.* **2014**, *46*, 214–223.
9. Wu, P.; Jin, J.; Zhao, X. Impact of climate change and irrigation technology advancement on agricultural water use in China. *Clim. Chang.* **2010**, *100*, 797–805.
10. Cao, X.C.; Wu, P.T.; Wang, Y.B.; Zhao, X.N. Assessing blue and green water utilisation in wheat production of China from the perspectives of water footprint and total water use. *Hydrol. Earth Syst. Sci.* **2014**, *18*, 3165–3178.
11. Hoekstra, A.Y.; Chapagain, A.K. *Globalization of Water: Sharing the Planet's Freshwater Resources*; Blackwell Publishing: Oxford, UK, 2008.
12. Chapagain, A.K.; Hoekstra, A.Y.; Savenije, H.H.G.; Gautam, R. The water footprint of cotton consumption: An assessment of the impact of worldwide consumption of cotton products on the water resources in the cotton producing countries. *Ecol. Econ.* **2006**, *60*, 186–203.
13. Chapagain, A.K.; Hoekstra, A.Y. The blue, green and grey water footprint of rice from production and consumption perspectives. *Ecol. Econ.* **2011**, *70*, 749–758.
14. Hoekstra, A.Y.; Chapagain, A.K.; Aldaya, M.M.; Mekonnen, M.M. *Water Footprint Assessment Manual: Setting the Global Standard*; Earthscan: London, UK, 2011.
15. Mekonnen, M.M.; Hoekstra, A.Y. A global and high-resolution assessment of the green, blue and grey water footprint of wheat. *Hydrol. Earth Syst. Sci.* **2010**, *14*, 1259–1276.

16. Mekonnen, M.M.; Hoekstra, A.Y. The green, blue and grey water footprint of crops and derived crop products. *Hydrol. Earth Syst. Sci.* **2011**, *15*, 1577–1600.
17. Bocchiola, D.; Nana, E.; Soncini, A. Impact of climate change scenarios on crop yield and water footprint of maize in the Po valley of Italy. *Agric. Water Manag.* **2013**, *116*, 50–61.
18. Ababaei, B.; Etedali, H. Estimation of water footprint components of Iran's wheat production: Comparison of global and national scale estimates. *Environ. Process.* **2014**, *1*, 193–205.
19. Gheewala, S.; Silalertruksa, T.; Nilsalab, P.; Mungkung, R.; Perret, S.; Chaiyawannakarn, N. Water footprint and impact of water consumption for food, feed, fuel crops production in Thailand. *Water-Sui* **2014**, *6*, 1698–1718.
20. Sun, S.K.; Wu, P.T.; Wang, Y.B.; Zhao, X.N.; Liu, J.; Zhang, X.H. The temporal and spatial variability of water footprint of grain: A case study of an irrigation district in China from 1960 to 2008. *J. Food Agric. Environ.* **2012**, *10*, 1246–1251.
21. Tian, T.H.; Zhu, D.J.; Wang, H.M.; Zang, M.D. Water footprint calculation of China's main food crops: 1987–2010. *China Popul. Resour. Environ.* **2013**, *23*, 122–128.
22. Wang, Y.B.; Wu, P.T.; Engel, B.A.; Sun, S.K. Application of water footprint combined with a unified virtual crop pattern to evaluate crop water productivity in grain production in China. *Sci. Total Environ.* **2014**, *497*, 1–9.
23. Sun, S.K.; Wu, P.T.; Wang, Y.B.; Zhao, X.N.; Liu, J.; Zhang, X.H. The impacts of interannual climate variability and agricultural inputs on water footprint of crop production in an irrigation district of China. *Sci. Total Environ.* **2013**, *444*, 498–507.
24. Zhang, H.T.; Guo, L.; Chen, J.Y.; Fu, P.H.; Gu, J.L.; Liao, G.Y. Modeling of spatial distributions of farmland density and its temporal change using geographically weighted regression model. *Chin. Geogr. Sci.* **2014**, *24*, 191–204.
25. Lesage, J.P. A spatial econometric examination of China's economic growth. *Geogr. Inf. Sci.* **1999**, *5*, 143–153.
26. Kamarianakis, Y.; Feidas, H.; Kokolatos, G.; Chrysoulakis, N.; Karatzias, V. Evaluating remotely sensed rainfall estimates using nonlinear mixed models and geographically weighted regression. *Environ. Model. Softw.* **2008**, *23*, 1438–1447.
27. Tang, Q.Y.; Xu, W.; Ai, F.L. A GWR-based study on spatial pattern and structural determinants of Shanghai's housing price. *Econ. Geogr.* **2012**, *32*, 52–58.
28. NBSC. *China Statistical Yearbook*; China Statistics Press: Beijing, China, 2011.
29. Tan, J.; Yang, P.; Liu, Z.; Wu, W.; Zhang, L.; Li, Z.; You, L.; Tang, H.; Li, Z. Spatio-temporal dynamics of maize cropping system in Northeast China between 1980 and 2010 by using spatial production allocation model. *J. Geogr. Sci.* **2014**, *24*, 397–410.
30. Huang, J.; Gao, Y.; Li, S.C. Temporal variation of virtual water of selected agricultural products in Northeast China. *Acta Sci. Nat. Univ. Pekin.* **2011**, *47*, 505–512.
31. China Meteorological Administration (CMA). *China Meteorological Data Sharing Service System*; CMA: Beijing, China, 2013.
32. Duan, P.L.; Qin, L.J. Temporal and spatial changes of water footprint of maize production in Jilin Province. *Water Sci. Technol. Water Supply* **2014**, *14*, 1067–1075.



33. Allen, R.G.; Pereira, L.S.; Raes, D.; Smith, M. *Crop Evapotranspiration-Guidelines for Computing Crop Water Requirements-Fao Irrigation and Drainage Paper 56*; Food and Agriculture Organization of the United Nations (FAO): Rome, Italy, 1998; Volume 300, p. 6541.
34. Food and Agriculture Organization of the United Nations (FAO). *Cropwat Model*; FAO: Rome, Italy, 2010.
35. Allen, R.G.; Pruitt, W.O.; Wright, J.L.; Howell, T.A.; Ventura, F.; Snyder, R.; Itenfisu, D.; Steduto, P.; Berengena, J.; Yrisarry, J.B.; *et al.* A recommendation on standardized surface resistance for hourly calculation of reference ET<sub>0</sub> by the FAO56 Penman-Monteith method. *Agric. Water Manag.* **2006**, *81*, 1–22.
36. Liu, J.; Wu, P.; Wang, Y.; Zhao, X.; Sun, S.; Cao, X. Impacts of changing cropping pattern on virtual water flows related to crops transfer: A case study for the hetao irrigation district, China. *J. Sci. Food Agric.* **2014**, *94*, 2992–3000.
37. Cheng, Y.; Wang, Y.; Wang, Z.; Luo, X. Changing rural development inequality in Jilin Province, Northeast China. *Chin. Geogr. Sci.* **2013**, *23*, 620–633.
38. Cheng, Y.Q.; Wang, Z.Y.; Ye, X.Y.; Wei, Y.D. Spatiotemporal dynamics of carbon intensity from energy consumption in China. *J. Geogr. Sci.* **2014**, *24*, 631–650.
39. Anselin, L. Local indicators of spatial association—Lisa. *Geogr. Anal.* **1995**, *27*, 93–115.
40. Fotheringham, A.S.; Charlton, M.E.; Brunson, C. Geographically weighted regression: A natural evolution of the expansion method for spatial data analysis. *Environ. Plan. A* **1998**, *30*, 1905–1927.
41. Fotheringham, A.S.; Brunson, C.; Charlton, M.E. *Geographically Weighted Regression: The Analysis of Spatially Varying Relationships*; Wiley: Chichester, UK, 2002.
42. Yang, L.X.; Wang, G.S.; Yao, S.M.; Yuan, S.F. A study on the spatial heterogeneity of grain yield per hectare and driving factors based on Esda-Gwr. *Econ. Geogr.* **2012**, *32*, 120–127.
43. Shi, S.Q.; Cao, Q.W.; Yao, Y.M.; Tang, H.J.; Yang, P.; Wu, W.B.; Xu, H.Z.; Liu, J.; Li, Z.G. Influence of climate and socio-economic factors on the spatio-temporal variability of soil organic matter: A case study of central Heilongjiang Province, China. *J. Integr. Agric.* **2014**, *13*, 1486–1500.
44. Sun, S.K.; Wu, P.T.; Wang, Y.B.; Zhao, X.N. Impacts of climate change on water footprint of spring wheat production: The case of an irrigation district in China. *Span. J. Agric. Res.* **2012**, *10*, 1176–1187.
45. Sun, S.K.; Wu, P.T.; Wang, Y.B.; Zhao, X.N. Temporal variability of water footprint for maize production: The case of Beijing from 1978 to 2008. *Water Resour. Manag.* **2013**, *27*, 2447–2463.