

# Journal Pre-proof



Mutations in NBEAL2 do not impact Weibel-Palade body biogenesis and Von Willebrand factor secretion in Gray Platelet Syndrome Endothelial Colony Forming Cells

Marije Kat, PhD, Iris van Moort, PhD, Petra E. Bürgisser, MSc, Taco W. Kuijpers, MD PhD, Menno Hofman, MSc, Marie Favier, MD, Rémi Favier, MD, Coert Margadant, PhD, Jan Voorberg, PhD, Ruben Bierings, PhD

PII: S2475-0379(23)00055-9

DOI: <https://doi.org/10.1016/j.rpth.2023.100086>

Reference: RPTH 100086

To appear in: *Research and Practice in Thrombosis and Haemostasis*

Received Date: 9 June 2022

Revised Date: 13 December 2022

Accepted Date: 25 January 2023

Please cite this article as: Kat M, van Moort I, Bürgisser PE, Kuijpers TW, Hofman M, Favier M, Favier R, Margadant C, Voorberg J, Bierings R, Mutations in NBEAL2 do not impact Weibel-Palade body biogenesis and Von Willebrand factor secretion in Gray Platelet Syndrome Endothelial Colony Forming Cells, *Research and Practice in Thrombosis and Haemostasis* (2023), doi: <https://doi.org/10.1016/j.rpth.2023.100086>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 The Author(s). Published by Elsevier Inc. on behalf of International Society on Thrombosis and Haemostasis.

**Mutations in NBEAL2 do not impact Weibel-Palade body biogenesis and Von Willebrand factor secretion in Gray Platelet Syndrome Endothelial Colony Forming Cells**

Marije Kat PhD<sup>1,\*</sup>, Iris van Moort PhD<sup>2,\*</sup>, Petra E. Bürgisser MSc<sup>2</sup>, Taco W. Kuijpers MD PhD<sup>1,3</sup>, Menno Hofman MSc<sup>1</sup>, Marie Favier MD<sup>4</sup>, Rémi Favier MD<sup>4,5</sup>, Coert Margadant PhD<sup>6</sup>, Jan Voorberg PhD<sup>1,7</sup> and Ruben Bierings PhD<sup>2</sup>

<sup>1</sup>Molecular Hematology, Sanquin Research and Landsteiner Laboratory, Amsterdam University Medical Center, University of Amsterdam, The Netherlands. <sup>2</sup>Hematology, Erasmus MC University Medical Center, Rotterdam, The Netherlands. <sup>3</sup>Pediatric Immunology, Rheumatology and Infectious Diseases, Emma Children's Hospital, Amsterdam University Medical Center, location AMC, Amsterdam, The Netherlands. <sup>4</sup>Assistance Publique-Hôpitaux de Paris, French Reference Center for Platelet Disorders, Armand Trousseau Children's Hospital, Paris, France. <sup>5</sup>INSERM, UMR 1287, Gustave Roussy Cancer Campus, Paris-Saclay University, Villejuif, France. <sup>6</sup>Angiogenesis Laboratory, Cancer Center Amsterdam, Amsterdam University Medical Center, location VUmc, The Netherlands. <sup>7</sup>Experimental Vascular Medicine, Amsterdam University Medical Center, University of Amsterdam, The Netherlands.

Running title: Weibel-Palade bodies in NBEAL2 deficient ECFCs

Corresponding author: Dr. Ruben Bierings, Department of Hematology, Erasmus MC University Medical Center, Rotterdam, The Netherlands. e-mail: r.bierings@erasmusmc.nl

\*: these authors contributed equally to this work.

Word count abstract:	241	Figure count:	3
Word count:	1883	Number of references:	38

### Essentials

- Biallelic mutations in NBEAL2 lead to Gray Platelet Syndrome (GPS), which affects  $\alpha$ -granules.
- NBEAL2 interactor SEC22B controls formation of  $\alpha$ -granules and Weibel-Palade bodies (WPBs).
- GPS patient-derived endothelial cells with NBEAL2 mutations were characterized *ex vivo*.
- NBEAL2 is not essential for WPB formation, maturation, and VWF secretion from endothelial cells.

**Abstract****Background**

Gray Platelet Syndrome (GPS) patients with Neurobeachin-like 2 (NBEAL2) deficiency produce platelets lacking alpha-granules (AGs) and present with lifelong bleeding symptoms. AGs are lysosome-related organelles (LROs) and store the hemostatic protein Von Willebrand factor (VWF) and the transmembrane protein P-selectin. Weibel-Palade bodies (WPBs) are LROs of endothelial cells and also store VWF and P-selectin. In megakaryocytes, NBEAL2 links P-selectin on AGs to the SNARE protein SEC22B on the endoplasmic reticulum (ER), thereby preventing premature release of cargo from AG precursors. In endothelial cells, SEC22B drives VWF trafficking from ER to Golgi and promotes the formation of elongated WPBs, but it is unclear if this requires NBEAL2.

**Objectives**

To investigate a potential role for NBEAL2 in WPB biogenesis and VWF secretion using NBEAL2 deficient endothelial cells.

**Methods**

Interaction of SEC22B with NBEAL2 in endothelial cells was investigated by interactomic mass spectrometry and pull down analysis. Endothelial Colony Forming Cells (ECFCs) were isolated from healthy controls and 3 unrelated GPS patients with mutations in *NBEAL2*.

**Results**

We show that SEC22B binds to NBEAL2 in ECs. GPS patient-derived ECFCs are deficient of NBEAL2, but reveal normal formation and maturation of WPBs and normal WPB cargo recruitment. Neither basal nor histamine-induced VWF secretion are altered in the absence of NBEAL2.

**Conclusions**

While NBEAL2 deficiency causes absence of AGs in GPS patients, it has no impact on WPB functionality in ECs. Our data highlight the difference in regulatory mechanisms between these two hemostatic storage compartments.



## Introduction

Gray platelet syndrome (GPS) is a rare platelet storage pool disorder (SPD) that is characterized by thrombocytopenia, enlarged platelets and lack of platelet alpha-granules (AGs), causing increased bleeding tendency, myelofibrosis and splenomegaly.<sup>1</sup> AGs store bioactive compounds, including Von Willebrand Factor (VWF), a crucial protein for hemostasis and wound healing.<sup>2</sup> Weibel-Palade bodies (WPBs) are the endothelial cell (EC) equivalent of AGs and contain VWF as their main cargo.<sup>3</sup> Both AGs and WPBs are lysosome-related organelles (LROs), a family of cell-type specific (secretory) organelles that acquire cargo and membrane components from the endolysosomal system and which have synergistic functions in immunity, hemostasis, inflammation and angiogenesis.<sup>4,5</sup> LRO biogenesis and degranulation are in many respects regulated by universal mechanisms that are shared between cell types that contain LROs.<sup>4</sup> Genetic defects causing SPDs often affect LRO biogenesis and release in multiple cell types simultaneously, including endothelial WPBs.<sup>5</sup> Vascular injury triggers VWF secretion from WPBs, producing adhesive strings that mediate platelet adherence to the damaged vessel wall.<sup>3</sup> Platelets subsequently become activated and release AG content to stabilize the platelet plug.<sup>5</sup> Evidently, in GPS AG degranulation is absent, resulting in platelet aggregation defects and bleeding complications in patients.

GPS is caused by autosomal recessive mutations in neurobeachin-like 2 (NBEAL2),<sup>6-8</sup> while autosomal dominant mutations in the transcription factor Growth Factor Independent 1B (GFI1B) underlie a related 'GPS-like' syndrome.<sup>9</sup> NBEAL2 is required for AG biogenesis, potentially through retention of cargo proteins in AG precursors during maturation or by retention of AGs flowing their formation in megakaryocytes.<sup>10,11</sup> GPS neutrophils also exhibit defects in a subset of their LROs, the specific granules (SGs), which include loss of SG cargo due to premature release of SGs.<sup>12</sup> Taken together this suggests that NBEAL2 potentially serves a general role in regulation of LROs.<sup>12</sup> The SNARE protein SEC22B is a central player in transport between the endoplasmic reticulum (ER) and the Golgi<sup>13</sup> that has recently been implicated in AG and WPB biogenesis. We have previously uncovered that in ECs, SEC22B

controls the formation and morphology of WPBs.<sup>14,15</sup> In megakaryocytes, NBEAL2 links SEC22B to the leukocyte receptor P-selectin present on AGs in a tripartite complex which acts to prevent the premature release of AG cargo from AG precursors through a mechanism that needs further clarification.<sup>16</sup> Given the prominent role of SEC22B in WPB biogenesis and because P-selectin is also present on WPBs<sup>17</sup>, we investigated whether NBEAL2 plays a similar role in the endothelial secretory pathway.

## **Materials and Methods**

### ***Cell culture and isolation of endothelial cells***

Pooled, cryopreserved human umbilical vein endothelial cells (HUVECs) were purchased from Promocell. Endothelial colony forming cells (ECFCs) were isolated from heparinized venous blood as described.<sup>18</sup> Control ECFCs were isolated from healthy, pseudonymized volunteers participating in Sanquin's internal blood donor system. GPS ECFCs were isolated from three unrelated GPS patients, coded C (6 clones), D (2 clones) and E (2 clones), that were previously described<sup>12</sup>, and correspond to patient C, D and E from that study. The study was performed according to national regulations regarding the use of human materials, informed consent and the Declaration of Helsinki and was approved by the Medical Ethical Committee of the Academic Medical Center in Amsterdam. HUVECs and ECFCs were cultured as described.<sup>19</sup> Expression of interleukin-8 (IL-8) was upregulated by incubation with 10 ng/ml interleukin-1 $\beta$  (IL-1 $\beta$ ) as described previously.<sup>20</sup> Immunocytochemistry, immunoblotting and VWF secretion assays were performed essentially as described.<sup>19,21</sup>

Antibodies and dilutions are described in the Supplemental Materials and Methods.

## Results and Discussion

### ***NBEAL2 interacts with SEC22B in primary ECs***

NBEAL2 was previously shown to be part of a tripartite complex linking SEC22B and P-selectin in megakaryocytes.<sup>16</sup> We have recently performed an interactomic analysis of SEC22B in ECs, which identified the SNARE protein STX5 as a novel regulator of WPB size as well as VWF trafficking and secretion.<sup>22</sup> We revisited that dataset, which is publically accessible via the PRIDE repository with dataset identifier PXD027516, and found that NBEAL2 was also identified as a specific interactor of SEC22B in ECs (**Figure 1A**). To confirm this we used mEGFP-SEC22B as bait in a pulldown assay in HUVECs followed by immunoblot analysis. Because ECs also contain P-selectin and, like megakaryocytes, express it on the limiting membrane of their LROs (i.e. WPBs), we also investigated whether the tripartite SEC22B/NBEAL2/P-selectin complex is present in ECs. We indeed confirmed a specific interaction with endogenous NBEAL2, but found no evidence of such a tripartite complex in ECs as P-selectin coprecipitation was not observed (**Figure 1B**).

### ***NBEAL2 deficient ECFCs produce WPBs***

A previous study showed that ECs in tissue sections of patients lacking platelet AGs have WPBs containing VWF and P-selectin.<sup>23</sup> However, it has since been shown that defects in NBEAL2 and GFI1B both lead to AG loss,<sup>6,7,24</sup> and it was unclear whether these patients were NBEAL2 deficient. To investigate the role of NBEAL2 in WPB regulation, we isolated ECFCs from three GPS patients with different mutations in *NBEAL2* (**Figure 1C,D**). These patients were previously shown to have platelets with no AGs, diminished AG cargo and elevated steady-state cell surface P-selectin<sup>12,25</sup>, and also neutrophils with reduced SG cargo (including lactoferrin), elevated steady-state surface expression of the SG protein CD66b and defective NETosis<sup>12</sup>, which is thought to be the result of impaired AG and SG retention in platelets and neutrophils respectively. We first confirmed that GPS patient-derived ECFCs are NBEAL2-

deficient using immunoblot analysis of ECFC lysates (**Figure 1E, Supplemental Figure 1**), which showed that NBEAL2 was undetectable in ECFCs of patient C and E and all but lost in ECFCs of patient D. Next we assessed the effect of *NBEAL2* mutations on WPB formation in cultured GPS ECFCs. Immunofluorescent staining of VWF revealed rod-shaped WPBs in all clones, with no apparent changes in number or morphology compared to those in healthy donor ECFCs (**Figure 2A**). Furthermore, in CTRL and GPS ECFCs SEC22B staining was observed on the ER and Golgi, while P-selectin was present on WPBs of CTRL as well as GPS ECFCs, suggesting that their intracellular localization is independent of NBEAL2 (**Figure 2B**). The lack of overlap between SEC22B and P-selectin further highlights that the presence of a tripartite SEC22B/NBEAL2/P-selectin complex in ECs is unlikely because of significant spatial separation. Immunolocalization of NBEAL2 resulted in a dispersed intracellular staining pattern with no overlap with VWF or resemblance to any other intracellular structures (**Supplemental Figure 2**).

Since a previous report suggested that endogenously synthesized (VWF) as well as endocytosed (fibrinogen) AG cargo proteins were not retained in AG precursors during AG biogenesis in NBEAL2 deficient megakaryocytes<sup>10</sup>, we next tested whether the composition of WPBs was altered in GPS ECFCs. For this we examined the presence of 3 WPB cargo proteins: Angiopoietin 2 (Ang-2), which is endogenously synthesized and included into newly forming WPBs at the level of the *trans*-Golgi network (TGN), CD63, a tetraspanin that recycles from the plasma membrane and is delivered to WPBs via the endosomal compartment, and interleukin-8 (IL-8), a chemokine that is included in WPBs at the TGN after upregulation of its synthesis by interleukin-1 $\beta$ .<sup>17,20,26–28</sup> IF stainings in CTRL and GPS ECFC clones revealed no change in Ang-2, CD63 and IL-8 localization to WPBs (**Figure 3A, Supplemental Figure 3A, Supplemental Figure 4**), which shows Golgi and post-Golgi WPB cargo recruitment and retention are unaffected by loss of NBEAL2. In addition, we observed normal recruitment of the small GTPase Rab27A, which is regarded as a maturation marker for WPBs and recruits effectors that act as positive and negative regulators of WPB exocytosis<sup>29</sup>, indicating that this

proximal event in recruitment of the WPB exocytotic machinery is not impaired by the absence of NBEAL2 (**Figure 3B, Supplemental Figure 3B**). Taken together our data suggest that NBEAL2 is not crucial for WPB biogenesis, cargo recruitment or maturation, hinting towards different regulatory mechanisms with regard to storage compartment biogenesis and (cargo) retention in comparison with megakaryocytes and platelets.<sup>16</sup>

### ***NBEAL2 is dispensable for basal and regulated VWF secretion***

Since GPS is associated with a bleeding tendency<sup>1</sup>, we examined a possible contribution of disrupted endothelial VWF secretion. Accumulating evidence suggests that deficiency of platelet AGs and neutrophil SGs in GPS is underpinned by premature release of these granules or their cargo from megakaryocytes and neutrophils, respectively, due to defective retention.<sup>10,12,25</sup> After biosynthesis WPBs are stored in ECs until exocytosis in response to activation, while at steady-state WPBs will stochastically undergo basal secretion.<sup>3</sup> To test whether NBEAL2 deficiency impacts WPB retention at steady state or during stimulated secretion, we examined VWF storage and secretion during basal and histamine-induced release. No difference was seen in composition of intracellular or secreted VWF, with regard to proteolytic processing of ProVWF to mature VWF (**Figure 3C**) or to multimeric composition of secreted VWF (**Figure 3D**), indicating normal regulation of synthesis and secretion. Indeed, both basal (unstimulated) VWF secretion during 24 hours and 30-minute histamine-stimulated VWF secretion remained unchanged (**Figure 3E,F**). As VWF storage and secretion from NBEAL2 deficient ECs remains functional, this further points towards platelet defects as the prime cause for bleeding symptoms in GPS.<sup>1</sup> Our findings that basal VWF release, which accounts for the bulk of circulating VWF, and histamine-stimulated VWF release are unaffected in GPS ECFCs are also consistent with the observations that GPS patients and *Nbeal2*<sup>-/-</sup> mice have normal plasma levels of VWF<sup>30,31</sup> and that DDAVP infusion increases plasma VWF in GPS patients.<sup>32</sup>

Our results show that NBEAL2 is not required for WPB biogenesis nor WPB cargo retention, indicating that, contrary to its role in LRO cargo retention in megakaryocytes and neutrophils, in the endothelial secretory pathway it is nonessential. Tubular condensation of sufficient amounts of VWF in the lumen of the TGN is a key requirement for WPB biogenesis, and this shapes emerging WPBs into their characteristic elongated form.<sup>15</sup> The essential role for VWF in this process is highlighted by the absence of WPBs in engineered and patient-derived VWF deficient endothelial cells<sup>26,33</sup> and the appearance of pseudo-WPBs after ectopic expression of VWF in non-endothelial cells.<sup>34,35</sup> This dependence of WPB biogenesis on condensation of VWF is in contrast to the formation of platelet AGs, which are still present in platelets of patients with type 3 Von Willebrand Disease caused by mutations that abrogate VWF synthesis.<sup>36,37</sup> We speculate that the difference between AG and WPB biogenesis in terms of NBEAL2 dependence may lie in the fact that WPBs are in many respects a nearly finished product after emergence from the TGN, while the cargo destined for AGs first proceeds through a number of endosomal compartments following its exit from the Golgi. During their synthesis in megakaryocytes, AGs are preceded by an intermediate stage that display the hallmarks of multi vesicular bodies (MVBs).<sup>38</sup> These MVBs are late endosomal compartments that carry VWF originating from the TGN and will eventually progress to AG precursors, the site where NBEAL2 is thought to associate with P-selectin in megakaryocytes to retain AG cargo.<sup>10</sup> Possibly, the molecular composition of the NBEAL2-dependent mechanism of LRO cargo retention differs between cell types, as is underscored by the notion that while neutrophils retain SGs and their cargo in a NBEAL2-dependent manner, they lack P-selectin.<sup>12</sup> From a functional point of view, NBEAL2-dependent mechanisms that retain LROs or LRO cargo may be more crucial for cells such as neutrophils and megakaryocyte-derived platelets, which are unable to replenish secretory organelles after their initial synthesis, than for ECs that can continuously generate new WPBs.

### **Acknowledgements**

Work in our laboratory was funded by grants from the Landsteiner Stichting voor Bloedtransfusie Research (LSBR-1707 and LSBR-2005), the Dutch Thrombosis Foundation (TSN 2017-01) and the Netherlands Organization for Scientific Research (NWO NWA.1160.18.038). We thank the patients for their contribution and participation in our study.

### **Authorship Contributions**

MK, IVM, PEB and MH performed research and analyzed data; TK, RF and MF provided vital reagents and expertise; MK, IVM, CM, JV, and RB designed the research and wrote the paper.

### **Disclosure of Conflict of Interest**

The authors report no conflicts of interest.

## References

1. Nurden AT, Nurden P. The gray platelet syndrome: Clinical spectrum of the disease. *Blood Rev.* **2007**;21(1):21–36.
2. Blair P, Flaumenhaft R. Platelet  $\alpha$ -granules: Basic biology and clinical correlates. *Blood Rev.* **2009**;23(4):177–189.
3. Schillemans M, Karampini E, Kat M, Bierings R. Exocytosis of Weibel–Palade bodies: how to unpack a vascular emergency kit. *J. Thromb. Haemost.* **2019**;17(1):6–18.
4. Delevoye C, Marks MS, Raposo G. Lysosome-related organelles as functional adaptations of the endolysosomal system. *Curr. Opin. Cell Biol.* **2019**;59:147–158.
5. Karampini E, Bierings R, Voorberg J. Orchestration of Primary Hemostasis by Platelet and Endothelial Lysosome-Related Organelles. *Arterioscler. Thromb. Vasc. Biol.* **2020**;40(6):1441–1453.
6. Kahr WHA, Hinckley J, Li L, et al. Mutations in NBEAL2, encoding a BEACH protein, cause gray platelet syndrome. *Nat. Genet.* **2011**;43(8):738–740.
7. Gunay-Aygun M, Falik-Zaccari TC, Vilboux T, et al. NBEAL2 is mutated in gray platelet syndrome and is required for biogenesis of platelet  $\alpha$ -granules. *Nat. Genet.* **2011**;43(8):732–4.
8. Albers CA, Cvejic A, Favier R, et al. Exome sequencing identifies NBEAL2 as the causative gene for gray platelet syndrome. *Nat. Genet.* **2011**;43(8):735–7.
9. Monteferrario D, Bolar NA, Marneth AE, et al. A dominant-negative GFI1B mutation in the gray platelet syndrome. *N. Engl. J. Med.* **2014**;370(3):245–53.
10. Lo RW, Li L, Leung R, Pluthero FG, Kahr WHA. NBEAL2 (Neurobeachin-Like 2) Is Required for Retention of Cargo Proteins by  $\alpha$ -Granules During Their Production by Megakaryocytes. *Arterioscler. Thromb. Vasc. Biol.* **2018**;38(10):2435–2447.



11. Guerrero JA, Bennett C, van der Weyden L, et al. Gray platelet syndrome: proinflammatory megakaryocytes and  $\alpha$ -granule loss cause myelofibrosis and confer metastasis resistance in mice. *Blood*. **2014**;124(24):3624–3635.
12. Aarts CEM, Downes K, Hoogendijk AJ, et al. Neutrophil specific granule and NETosis defects in gray platelet syndrome. *Blood Adv*. **2021**;5(2):549–564.
13. Xu D, Joglekar AP, Williams AL, Hay JC. Subunit structure of a mammalian ER/Golgi SNARE complex. *J. Biol. Chem*. **2000**;275(50):39631–39639.
14. Karampini E, Bürgisser PE, Olins J, et al. Sec22b determines Weibel-Palade body length by controlling anterograde ER-Golgi transport. *Haematologica*. **2021**;106(4):1138–1147.
15. Kat M, Margadant C, Voorberg J, Bierings R. Dispatch and delivery at the ER–Golgi interface: how endothelial cells tune their hemostatic response. *FEBS J*. **2022**;289(22):6863–6870.
16. Lo RW, Li L, Pluthero FG, Leung R, Eto K, Kahr WHA. The endoplasmic reticulum protein SEC22B interacts with NBEAL2 and is required for megakaryocyte  $\alpha$ -granule biogenesis. *Blood*. **2020**;136(6):715–725.
17. Harrison-Lavoie KJ, Michaux G, Hewlett L, et al. P-selectin and CD63 use different mechanisms for delivery to Weibel-Palade bodies. *Traffic*. **2006**;7(6):647–62.
18. Martin-Ramirez J, Kok MGM, Hofman M, et al. Individual with Subclinical Atherosclerosis Have Impaired Proliferation of Blood Outgrowth Endothelial Cells, Which Can Be Restored by Statin Therapy. *PLoS One*. **2014**;9(6):e99890.
19. Kat M, Bürgisser PE, Janssen H, et al. GDP/GTP exchange factor MADD drives activation and recruitment of secretory Rab GTPases to Weibel-Palade bodies. *Blood Adv*. **2021**;5(23):5116–5127.
20. Bierings R, van den Biggelaar M, Kragt A, Mertens K, Voorberg J, van Mourik JA.

- Efficiency of von Willebrand factor-mediated targeting of interleukin-8 into Weibel-Palade bodies. *J. Thromb. Haemost.* **2007**;5(12):2512–2519.
21. van Hooren KWEM, van Breevoort D, Fernandez-Borja M, et al. Phosphatidylinositol-3,4,5-triphosphate-dependent Rac exchange factor 1 regulates epinephrine-induced exocytosis of Weibel-Palade bodies. *J. Thromb. Haemost.* **2014**;12(2):273–281.
  22. Kat M, Karampini E, Hoogendijk AJ, et al. Syntaxin 5 determines Weibel-Palade body size and Von Willebrand factor secretion by controlling Golgi architecture. *Haematologica.* **2022**;2021.12.10.472050.
  23. Gebrane-Younes J, Cramer EM, Orcel L, Caen JP. Gray platelet syndrome. Dissociation between abnormal sorting in megakaryocyte ??-granules and normal sorting in Weibel-Palade bodies of endothelial cells. *J. Clin. Invest.* **1993**;92(6):3023–3028.
  24. Ward CM, Morel-Kopp M-C, Chen Q, et al. GFI1B Mutation Causes a Platelet Function Defect With Reduced Alpha-Granule Content and Abnormal Aggregation Response. *Blood.* **2013**;122(21):566–566.
  25. Sims MC, Mayer L, Collins JH, et al. Novel manifestations of immune dysregulation and granule defects in gray platelet syndrome. *Blood.* **2020**;136(17):1956–1967.
  26. Schillemans M, Kat M, Westeneng J, et al. Alternative trafficking of Weibel-Palade body proteins in CRISPR/Cas9-engineered von Willebrand factor-deficient blood outgrowth endothelial cells. *Res. Pract. Thromb. Haemost.* **2019**;3(4):718–732.
  27. Karampini E, Schillemans M, Hofman M, et al. Defective AP-3-dependent VAMP8 trafficking impairs Weibel-Palade body exocytosis in Hermansky-Pudlak Syndrome type 2 blood outgrowth endothelial cells. *Haematologica.* **2019**;104(10):2091–2099.
  28. Knipe L, Meli A, Hewlett L, et al. A revised model for the secretion of tPA and cytokines from cultured endothelial cells. *Blood.* **2010**;116(12):2183–2191.
  29. Bierings R, Hellen N, Kiskin N, et al. The interplay between the Rab27A effectors Slp4-

- a and MyRIP controls hormone-evoked Weibel-Palade body exocytosis. *Blood*. **2012**;120(13):2757–2767.
30. Gunay-Aygun M, Zivony-Elboun Y, Gumruk F, et al. Gray platelet syndrome: natural history of a large patient cohort and locus assignment to chromosome 3p. *Blood*. **2010**;116(23):4990–5001.
31. Deppermann C, Cherpokova D, Nurden P, et al. Gray platelet syndrome and defective thrombo-inflammation in Nbeal2-deficient mice. *J. Clin. Invest.* **2013**;123(8):3331–3342.
32. Pfueller SL, Howard MA, White JG, Menon C, Berry EW. Shortening of bleeding time by 1-deamino-8-arginine vasopressin (DDAVP) in the absence of platelet von Willebrand factor in Gray platelet syndrome. *Thromb. Haemost.* **1987**;58(4):1060–3.
33. Groeneveld DJ, van Bekkum T, Dirven RJ, et al. Angiogenic characteristics of blood outgrowth endothelial cells from patients with von Willebrand disease. *J. Thromb. Haemost.* **2015**;13(10):1854–1866.
34. Voorberg J, Fontijn R, Calafat J, Janssen H, van Mourik JA, Pannekoek H. Biogenesis of von Willebrand factor-containing organelles in heterologous transfected CV-1 cells. *EMBO J.* **1993**;12(2):749–758.
35. Michaux G, Abbitt KB, Collinson LM, Haberichter SL, Norman KE, Cutler DF. The physiological function of von Willebrand's factor depends on its tubular storage in endothelial Weibel-Palade bodies. *Dev. Cell.* **2006**;10(2):223–232.
36. Nurden P, Nurden AT, La Marca S, Punzo M, Baronciani L, Federici AB. Platelet morphological changes in 2 patients with von Willebrand disease type 3 caused by large homozygous deletions of the von Willebrand factor gene. *Haematologica*. **2009**;94(11):1627–1629.
37. Swinkels M, Atiq F, Bürgisser PE, et al. Quantitative 3D microscopy highlights altered

von Willebrand factor  $\alpha$ -granule storage in patients with von Willebrand disease with distinct pathogenic mechanisms. *Res. Pract. Thromb. Haemost.* **2021**;5(6):e12595.

38. Heijnen HFG, Debili N, Vainchencker W, Breton-Gorius J, Geuze HJ, Sixma JJ. Multivesicular Bodies Are an Intermediate Stage in the Formation of Platelet  $\alpha$ -Granules. *Blood.* **1998**;91(7):2313–2325.

Journal Pre-proof

## Figure Legends

**Figure 1. Mutations in the SEC22B interactor NBEAL2 are associated with loss of NBEAL2 expression in GPS patient-derived endothelial cells.** (A) Label-free quantification (LFQ) of SEC22B and NBEAL2 protein levels in GFP pull down samples of mEGFP or mEGFP-SEC22B baits expressed in HUVECs determined by mass spectrometry analysis.<sup>22</sup> (B) Pull down analysis of mEGFP or mEGFP-SEC22B bait expressed in HUVECs using GFP or control (ctrl) beads and Western blot analysis of GFP, NBEAL2, and P-selectin. Molecular weights of protein ladder marker bands are indicated in kDa on the left. (C) Domain structure of full length (2754 a.a.) NBEAL2 showing the position of the mutations present in patients C, D and E. (D) Table showing *NBEAL2* gene mutations, changes on protein level and type of mutation in GPS patients included in this study (m: male, f: female). Gene transcript identifier: NM\_015175 / Protein identifier: NP\_055990. (E) Western blot of NBEAL2 expression in ECFCs from 3 GPS patients (C, D and E; 2 clones were analyzed for patient D) compared to ECFCs isolated from healthy control donors (CTRL 1-3).  $\alpha$ -tubulin is shown as a loading control. Molecular weights of protein ladder marker bands are indicated in kDa on the left.

**Figure 2. GPS patient-derived endothelial cells generate WPBs despite NBEAL2 deficiency.** Representative maximal projections of confocal immunofluorescent analysis of (A) VWF (red) and nuclei (Hoechst, blue) and (B) SEC22B (red), P-selectin (green) and nuclei (blue) in CTRL and GPS patient-derived ECFCs. Grayscale images for SEC22B and P-selectin channels are shown below. Scale bars represent 20  $\mu$ m.

**Figure 3. NBEAL2 is dispensable for WPB cargo sorting and maturation and for basal and secretagogue-induced VWF secretion.** Representative maximal projections of confocal immunofluorescent analysis of (A) VWF (red), CD63 (green) and Angiopoietin 2 (Ang-2, blue) or (B) VWF (red), Rab27A (green), and nuclei (Hoechst, blue) in CTRL and GPS ECFCs.

Boxed areas are shown below in grayscale. Yellow arrowheads indicate example WPBs. Scale bars represent 20  $\mu\text{m}$ . (C) Western blot analysis of VWF content in lysates and media from CTRL and GPS patient-derived ECFCs. (D) VWF multimer analysis of 24 hour basal media from CTRL and and GPS patient-derived ECFCs. UL: ultra-large, HMW: high molecular weight, LMW: low molecular weight, D: dimer. (E) Percentage of stored and secreted VWF in 24 hours under basal conditions (n=7, mean $\pm$ SEM, two-way ANOVA, not significant). (F) Secreted VWF in 30 minutes under unstimulated conditions (vehicle) or in the presence of 100  $\mu\text{M}$  histamine expressed as a percentage of the total intracellular VWF levels in unstimulated cells (n=4, mean $\pm$ SEM, two-way ANOVA, not significant).

Figure 1







Figure 2

A



B








Figure 3



**Rebuttal Figure 1**

**Rebuttal Figure 1:** Representative images of post-confluent monolayers of CTRL and GPS ECFCs used for VWF secretion assays, stained for VWF (green) and nuclei (Hoechst). Separate channels are shown in grayscale on the right. Scale bar represents 50  $\mu\text{m}$ .