

BMJ Open EffectiveNess of a multimodal preHAbilitation program in patieNts with bladder canCER undergoing radical cystectomy: protocol of the ENHANCE multicentre randomised controlled trial

Emine Akdemir ^{1,2}, Maike G Sweegers,^{1,2} Alina Vrieling ³, Helene Rundqvist,⁴ Richard P Meijer,⁵ Annemarie M Leliveld-Kors,⁶ Antoine G van der Heijden,⁷ Vera C Rutten ⁸, Evert L Koldewijn,⁹ Siebe D Bos,¹⁰ Carl J Wijburg,¹¹ Tom A T Marcelissen,¹² Bart C Bongers,^{13,14} Valesca P Retèl ^{1,15}, Wim H van Harten,^{1,16} Anne M May,¹⁷ Wim G Groen,^{18,19} Martijn M Stuiver^{1,20}

To cite: Akdemir E, Sweegers MG, Vrieling A, *et al*. EffectiveNess of a multimodal preHAbilitation program in patieNts with bladder canCER undergoing radical cystectomy: protocol of the ENHANCE multicentre randomised controlled trial. *BMJ Open* 2023;**13**:e071304. doi:10.1136/bmjopen-2022-071304

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-071304>).

WGG and MMS are joint senior authors.

Received 21 December 2022
Accepted 16 February 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to
Dr Martijn M Stuiver;
m.stuiver@nki.nl

ABSTRACT

Introduction Radical cystectomy (RC) is the standard treatment for patients with non-metastatic muscle-invasive bladder cancer, as well as for patients with therapy refractory high-risk non-muscle invasive bladder cancer. However, 50–65% of patients undergoing RC experience perioperative complications. The risk, severity and impact of these complications is associated with a patient's preoperative cardiorespiratory fitness, nutritional and smoking status and presence of anxiety and depression. There is emerging evidence supporting multimodal prehabilitation as a strategy to reduce the risk of complications and improve functional recovery after major cancer surgery. However, for bladder cancer the evidence is still limited. The aim of this study is to investigate the superiority of a multimodal prehabilitation programme versus standard-of-care in terms of reducing perioperative complications in patients with bladder cancer undergoing RC.

Methods and analysis This multicentre, open label, prospective, randomised controlled trial, will include 154 patients with bladder cancer undergoing RC. Patients are recruited from eight hospitals in The Netherlands and will be randomly (1:1) allocated to the intervention group receiving a structured multimodal prehabilitation programme of approximately 3–6 weeks, or to the control group receiving standard-of-care. The primary outcome is the proportion of patients who develop one or more grade ≥ 2 complications (according to the Clavien-Dindo classification) within 90 days of surgery. Secondary outcomes include cardiorespiratory fitness, length of hospital stay, health-related quality of life, tumour tissue biomarkers of hypoxia, immune cell infiltration and cost-effectiveness. Data collection will take place at baseline, before surgery and 4 and 12 weeks after surgery.

Ethics and dissemination Ethical approval for this study was granted by the Medical Ethics Committee NedMec (Amsterdam, The Netherlands) under reference number 22–595/NL78792.031.22. Results of the study will be published in international peer-reviewed journals.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This is a multicentre randomised controlled trial investigating the effects of multimodal prehabilitation in an understudied group of patients with bladder cancer undergoing radical cystectomy.
- ⇒ This data collected in this study enables exploration of the effects of prehabilitation on tumour hypoxia and immune cell infiltration.
- ⇒ The intervention accounts for heterogeneity in cancer treatment by offering additional support during neoadjuvant treatment when appropriate.
- ⇒ This study includes a cost-effectiveness analysis from a societal perspective.
- ⇒ Due to the two-group design and the multimodal intervention, it will not be possible to disentangle the independent effects of physical exercise training, nutritional support, psychological counselling and smoking cessation.

Trial registration number NCT05480735.

INTRODUCTION

Bladder cancer is the 10th most common diagnosed cancer worldwide, with over 573 000 new patients and 213 000 deaths each year.¹ Radical cystectomy (RC) is the standard treatment for patients with non-metastatic muscle-invasive bladder cancer,^{2,3} as well as for patients with therapy refractory high-risk non-muscle invasive bladder cancer.⁴ RC is a challenging and costly surgical procedure with high morbidity and mortality rates⁵: 50–65% of the patients experience perioperative complications, of which 10–20% are high-grade.^{6–8} Low cardiorespiratory fitness,^{9,10} poor nutritional status,¹¹ the

presence of anxiety and depression¹² and smoking¹³ increase the risk of perioperative complications, length of hospital stay and the associated medical costs.^{9 10} In daily clinical practice, 25% of patients with muscle-invasive bladder cancer receive neoadjuvant treatment, which might further impair preoperative cardiorespiratory fitness and nutritional status.^{14–17}

Emerging evidence has identified the preoperative period as a window of opportunity to address lifestyle. Multimodal preoperative interventions, including physical exercise training, nutritional support, psychological counselling and smoking cessation, that aim to increase a patient's tolerance to surgery, reduce the incidence, severity and impact of complications, accelerate and improve the quality of recovery and improve quality of life have been termed *prehabilitation*.

To date, no adequately powered trial has been performed to establish the effectiveness of a multimodal prehabilitation programme for reducing the incidence, severity and impact of complications and costs associated with RC for bladder cancer. A few studies investigated the impact of a single modal^{18 19} or multimodal^{20 21} prehabilitation programme on functional recovery following RC. A phase I/II trial with 54 patients showed improvement in patient-reported quality of life after 4 weeks of a supervised physical exercise training programme before surgery.¹⁹ A feasibility randomised controlled trial with 60 patients provided evidence that a 3–6 weeks supervised vigorous aerobic exercise training programme before surgery led to improvements in cardiorespiratory fitness parameters and possibly fewer surgical complications.¹⁸ A randomised controlled trial with 107 patients demonstrated improvement in walking distance during 7 days after surgery²⁰ and improved muscle power.²² Finally, a randomised controlled trial with 70 patients showed significantly better functional capacity in the intervention group that followed a multimodal programme compared with the control group, at 4 weeks after surgery.²¹ However, the physical exercise intervention in these studies including patients with bladder cancer had limitations with regard to therapeutic validity²³ due to a short duration of the intervention,^{20 22} relatively low exercise intensity^{20–22} or only patient-reported compliance with the exercise intervention.^{20–22}

Preclinical studies indicate that exercise may directly affect tumour characteristics such as normalisation of tumour vasculature and immune cell infiltration, thereby enhancing tumour perfusion and reducing hypoxia.^{24 25} These factors are associated with treatment efficacy and survival.^{26–28} The current study provides a unique opportunity to explore the effects of prehabilitation on tumour inflammation and hypoxia markers, which has not yet been studied in human patients.

Study objectives

The ENHANCE study is designed to investigate the effectiveness of a multimodal prehabilitation programme versus standard-of-care in patients approaching RC. The

primary aim is to investigate the superiority of a multimodal prehabilitation programme in terms of reducing one or more grade ≥ 2 perioperative complications within 90 days in patients with bladder cancer undergoing RC. Secondary outcomes include changes in preoperative cardiorespiratory fitness, length of hospital stay, health-related quality of life, tumour tissue biomarkers of hypoxia, immune cell infiltration and cost-effectiveness.

METHODS AND ANALYSIS

The ENHANCE study is a multicentre, open label, two-arm randomised controlled trial. Patient inclusion and data collection started in August 2022. Ethical approval, extending to all participating centres, was granted by the Medical Ethical Committee NedMec (Amsterdam, The Netherlands), under reference number 22–595/NL78792.031.22. The trial is prospectively registered in Clinical Trials on 29 July 2022.

Study population

The study aims to include 154 patients who meet the following inclusion criteria: aged ≥ 18 years, histologically confirmed, primary, bladder cancer (cTa-4N0-3M0) and planned to undergo RC. Surgery will not be delayed in favour of prehabilitation. Hence, patients who are scheduled for surgery within 3 weeks are not eligible for the trial. Patients who express the intention to follow a similar exercise training programme regardless of randomisation outcome, patients with severe cognitive or psychiatric disorders, patients with a contraindication to perform physical exercise training or a cardiopulmonary exercise test (CPET) and patients unable to read or understand the Dutch language will also be excluded.

Recruitment and randomisation

Patients are recruited from eight academic or teaching hospitals across different regions in The Netherlands (Catharina Hospital, Erasmus University Medical Center, Maastricht University Medical Center+, Noordwest Hospital Group, Radboud University Medical Center, Rijnstate Hospital, University Medical Center Groningen and University Medical Center Utrecht). The Netherlands Cancer Institute is the coordinating centre. After establishment of diagnosis and indication for surgery, the urologist or nurse specialist invites patients to participate in the study and provides the patient information letter. The study coordinator contacts these patients to provide further oral information and answer potential questions about the study. When a patient agrees to participate, written informed consent is obtained. After collection of baseline data, patients are randomised in a 1:1 ratio, using a minimisation algorithm²⁹ aimed to achieve optimal balance between the two study arms with regard to the recruiting hospital, neoadjuvant treatment (yes/no), nodal status (N0/N1-3) and type of surgery (open/robot-assisted RC). Minimisation is done using the Minirand package in R V.4.0.4.^{30 31} The algorithm

includes a random component to ensure blinding of treatment allocation. Due to the nature of the intervention, blinding of participants and research investigators is not possible. After randomisation, patients in both the intervention and control groups receive a leaflet with recommendations on physical activity, diet and smoking cessation, according to the latest guidelines for patients with cancer.^{32 33} These recommendations are not further individualised or actively supported.

Patients who do not wish to participate in the study are asked to participate in a one-time questionnaire as described in [table 1](#). A participant flow diagram is shown in [figure 1](#).

Control group—standard-of-care

Patients randomised to the standard-of-care arm receive care as usual, which does not include a comprehensive multimodal prehabilitation programme. In all participating centres, enhanced recovery after surgery protocols are used to optimise medical conditions to enhance recovery.³⁴ These protocols include advice on smoking cessation when a patient is smoking and a referral to a dietician when malnutrition is detected. Preoperatively, the urologist determines whether the patient is physically fit for surgery. Patients in the control group are not prohibited to be physically active or seek counselling for nutritional advice, psychological support or smoking cessation.

Intervention group—prehabilitation

Patients randomised to the intervention arm will participate in a multimodal prehabilitation programme including supervised physical exercise training, nutritional support and—when relevant—psychological counselling and professional support for smoking-cessation, to enhance their health status. The intervention starts as soon as possible after baseline measurements and randomisation, approximately 3–6 weeks before surgery and is continued until surgery. Patients who are included before undergoing neoadjuvant treatment (including chemotherapy and immunotherapy) will participate in a physical exercise training programme from inclusion until completion of neoadjuvant treatment to prevent the often observed decline in physical fitness throughout neoadjuvant therapy. Subsequently, these patients will participate in the full multimodal prehabilitation programme until the date of surgery. Adverse events related to the intervention are monitored.

Physical exercise training programme

The physical exercise training programme consists of three training sessions per week under the supervision of a physical therapist. An overview of the training programme and training progression is described in [table 2](#). The programme will be delivered at a physical therapy practice near a patient's home to minimise travel time, preferably supervised by a physical therapist affiliated with Onconet, a nationwide network of physical

therapists with additional competencies in cancer care. The study coordinator provides the physical therapists instructions via a video call covering the specifics of the training programme for the current study.

Training sessions last 1 hour and consist of individualised aerobic interval training (three times a week), resistance training (two times a week) and relaxation exercises (one time a week). Results of the CPET will be used to establish the individual training intensity for aerobic interval training. For the resistance training, six large muscle groups will be targeted, in two sets of a defined maximum number of repetitions. The aim of these training sessions is to improve cardiorespiratory fitness, and muscle strength and mass. Relaxation exercises consist of guided breathing exercises and progressive muscle relaxation.³⁵ The aim of these exercises is to reduce possible anxiety and stress. In addition to the supervised sessions, patients are encouraged to be moderately active on at least two additional days per week for 30 min.

The supervised sessions during neoadjuvant treatment consist of twice-weekly moderate-intensity aerobic and resistance training. Patients receive the physical exercise training programme as described above for the remaining 3–6 weeks before surgery.

Nutritional support

Patients receive tailored advice from a registered dietician at the participating hospitals aiming at a total protein intake of 1.9–2.3 g/per kg of fat-free mass as estimated with bioelectrical impedance analysis, to promote an anabolic state. Dietary advice will emphasise the benefit of spreading protein consumption over three meals, with a goal of 25–30 g protein per meal and includes advice for optimal energy intake. To achieve this, and to increase adaptive responses of the skeletal muscle, participants will receive high-quality protein supplements containing 30 g of whey protein, 20 µg of vitamin D and 250 mg of calcium³⁶ in standardised supplements produced for the purpose of the study (FrieslandCampina, The Netherlands). These supplements are prescribed after each physical exercise training session and daily at least 1 hour before sleep or in the morning (depending on the patient's preference). The dietician provides intake consultation and one or two follow-up sessions to evaluate nutritional intake. Protein intake will be restricted according to the severity of renal impairment. Depending on the patient's protein intake, patients will be advised to either reduce or drop the intake of protein supplements. If necessary, additional dietary advice will be provided to reduce nutritional protein intake. Nutritional support starts at 3–6 weeks before surgery for all patients.

Psychological counselling

Patients are screened for anxiety and depression using the Hospital Anxiety and Depression Scale questionnaire at baseline.³⁷ If a patient's score falls between 11 and 18, the option of psychological counselling is discussed and referral is arranged for patients indicating a need for such

Table 1 Outcome measures in the intervention and control group

Description of outcome	Assessment	Description of measure	T0a	T0b	T1	T2	T3
Primary outcome							
Proportion of patients who develop one or more grade ≥ 2 perioperative complications within 90 days	Perioperative complications, grade ≥ 2 according to Clavien-Dindo classification	Number of grade ≥ 2 complication is abstracted from medical records				X	X
Secondary outcomes							
Proportion of patients who develop one or more high-grade (≥ 3) complications, number of complications, length of hospital stay and readmissions within 90 days	Perioperative complications, grade ≥ 3 according to Clavien-Dindo classification, length of hospital stay, readmissions	Number of grade ≥ 3 complications, number of complications, length of hospital stay and readmissions abstracted from medical records				X	X
Cardiorespiratory fitness	Cardiopulmonary exercise test on a cycle ergometer using a ramp protocol ⁴⁰	Peak oxygen uptake (VO_{2peak} in mL/kg/min)	X		X		
Physical functioning	Short physical performance battery ⁸⁴	Five items organised into three subscales of balance, walking speed and lower extremity muscle strength	X		X		X
Upper extremity muscle strength	Handgrip strength using a handheld dynamometer ⁸⁵	The maximum score of three attempts of both hands in kilogram (kg)	X		X		X
Lower extremity functional muscle strength	30-s sit-to-stand test ⁸⁶	The number of sit to stands within 30s	X		X		X
Nutritional intake	24-hour recall	Protein (total gram) and caloric intake (total kcal)	X		X		
Nutritional status	Patient-generated subjective global assessment short form ⁸⁷	Four items assessing weight (status), nutritional intake, symptoms and physical functioning	X	X	X	X	X
	Nil per mouth consumption during hospitalisation	The number of days after RC abstracted from medical records				X	X
Body composition	Bioelectrical impedance analysis	Body mass (kg), body height (cm), (subcutaneous) fat mass (%) and (upper limb) muscle mass (kg) are measured, and BMI (kg/m ²)	X		X		X
Health-related quality of life	EORTC quality of life questionnaire core 30 ⁸⁸	Thirty items, organised into five functional scales (physical, role, emotional, cognitive and social), three symptom scales (pain, fatigue and emesis), six items (dyspnoea, sleep disturbance, appetite loss, constipation, diarrhoea and financial impact), and an overall quality of life scale	X	X	X	X	X
Bladder cancer-related quality of life	EORTC muscle-invasive bladder cancer specific module ⁸⁹	Thirty items, assessing urinary symptoms, bowel symptoms, sexual functioning, urostomy problems, difficulties associated with the use of a catheter and body image	X	X	X	X	X

Continued

Table 1 Continued

Description of outcome	Assessment	Description of measure	T0a	T0b	T1	T2	T3
Anxiety and depression	Hospital Anxiety and Depression Scale ³⁷	Seven items assessing anxiety and seven items assessing depression	X	X	X	X	X
Fatigue	Multidimensional fatigue inventory ⁹⁰	Twenty items, categorised into five scales: general fatigue, physical fatigue, reduced activity, reduced motivation and mental fatigue	X	X	X	X	X
Physical activity	Short questionnaire to assess health-enhancing physical activity ⁹¹	Eleven items, organised into four different physical activities measuring frequency, duration and intensity (physical activity to and from home, household activities, activities at work and physical activities performed during leisure time)	X	X	X	X	X
Tumour hypoxia and immune cell infiltration	Two tumour biopsies: from routine diagnostic investigation and from surgical tumour excision	Immunohistochemistry analysis will be performed for hypoxia and immune cell infiltration biomarkers using the laboratory of the Karolinska Institutet (Sweden) and their according protocols	X		X		
Cost-effectiveness	EuroQol 5-dimension-5 levels ⁴⁴	Five items (dimensions) multiattribute utility questionnaire that measures mobility, self-care, usual activities, pain/discomfort and anxiety/depression in five levels	X	X	X	X	X
	iMTA Medical Consumption Questionnaire ⁴² and Productivity Costs Questionnaire ⁴³	Patient-reported productivity losses and medical consumption		X	X	X	X
	Healthcare costs	Medical activities abstracted from the management systems of the hospitals					X
Other outcomes							
Socio-demographic and clinical data	Socio-demographic data, disease and treatment characteristics will be abstracted from medical records or reported by the patient	Patient reported: place of birth, sex, marital status, living and work situation, education, lifestyle variables and the self-administrated comorbidity questionnaire ⁹²	X				
		Medical records: birth month and year, date of diagnosis, date and type of treatment, type of urinary diversion, tumour characteristics, ASA score, WHO score					
		Patient reported: smoking status	X	X	X	X	X
Coping mechanism	Sense of coherence questionnaire ⁹³	Thirteen items, categorised into three scales: comprehensibility, manageability and meaningfulness	X				
Compliance to the intervention	Adherence rates	Patient reported: self-composed (activity) diary. Physical therapist: adherence of the physical exercise training intervention on standardised training session forms	X	X	X		
Patient evaluation	Self-composed questionnaire	Patients in the intervention group: satisfaction with the programme and willingness to participate in focus group Patients in the control group: evaluate contamination Both groups: evaluation of possible post-surgical intervention					X

Continued

Table 1 Continued

Description of outcome	Assessment	Description of measure	T0a	T0b	T1	T2	T3
Non-participation	Self-composed questionnaire	Patient-reported outcomes: socio-demographic, health-related and bladder cancer-related quality of life, anxiety and depression, fatigue, physical activity, coping mechanism and reason(s) for not participating. Medical records: birth month and year, date of diagnosis, date and type of treatment, tumour characteristics, ASA score, WHO score	X				

ASA, American Society of Anesthesiologists; BMI, body mass index; EORTC, European Organisation for Research and Treatment of Cancer; RC, radical cystectomy; VO₂peak, oxygen uptake at peak exercise.

counselling. Patients who score ≥ 19 are directly referred to psychological counselling. Referred patients receive an initial counselling session of 1.5 hours, and additional sessions at the discretion of the psychological counsellor and the patient.

Smoking cessation

Intensive counselling and nicotine replacement therapy is offered to all smoking patients in the 3–6 weeks before surgery. Counselling includes at least one in-person session and one or more telephone or in-person follow-up sessions by trained counsellors. If the patient indicates to be smoking and is willing to quit, the treating physician will refer the patient.

Study outcomes

Data are collected before randomisation (T0a), before surgery (T1), and 4 (T2), and 12 (T3) weeks after surgery. If applicable, additional data will be collected after neoadjuvant treatment (T0b). At T0a, clinical data (eg, treatment, clinical stage) will be abstracted from the medical records and socio-demographic characteristics (eg, age, sex and education) will be obtained via a questionnaire. In addition, coping mechanisms³⁸ will be assessed via a questionnaire as this can help understand participation, attendance and dropouts. Data is collected in Castor EDC and access is restricted to the investigator team. Due to the low risk of the intervention a data safety monitoring board is not instated.


**Figure 1** Study flow chart.

Table 2 Content of the supervised physical exercise training programme in the intervention group

Frequency	Intensity	Time	Type
Three times a week	Aerobic interval training, consisting of four intervals of alternating effort performed on a cycle ergometer: <ul style="list-style-type: none"> ▶ 4 min of low-intensity exercise, defined as 30% of the workload achieved at VO_2peak. ▶ 2–3 min of high-intensity exercise, defined as 90% of the workload achieved at VO_2peak.* 	24–28 min	Aerobic interval training
Two times a week	During neoadjuvant treatment: <ul style="list-style-type: none"> ▶ Week 1–3: consisting of 20 min on a cycle ergometer and 10 min on a different aerobic machine, corresponding to 30% of the heart rate reserve. ▶ Week 4–6: consisting of 20 min on a cycle ergometer and 10 min on a different aerobic machine, corresponding to 40% of the heart rate reserve. ▶ Week 7–9: consisting of 20 min on a cycle ergometer and 10 min on a different aerobic machine, corresponding to 50% of the heart rate reserve. ▶ Week 10–12 (and further): consisting of 20 min on a cycle ergometer and 10 min on a different aerobic machine, corresponding to 60% of the heart rate reserve. 	30 min	Aerobic training
Two times a week	Resistance training, consisting of training six large muscle groups (leg press, bench press or chest press, abdominal crunch, pull over, low row and step up)† in two sets: <ul style="list-style-type: none"> ▶ Week 1: maximal 15 repetitions at ~65% of 1RM per set. ▶ Week 2: maximal 12 repetitions at ~70% of 1RM per set (weight week 1+10%). ▶ Week 3: maximal 10 repetitions at ~75% of 1RM per set (weight week 2+10%). ▶ From week 4 maximal 10 repetitions per set.‡ 	~20 min	Resistance training
One time a week	Progressive muscle relaxation techniques. ³⁵	~20 min	Relaxation exercises

*Further tailoring is done by the physical therapists: if a patient is not able to complete the high-intensity interval, the intensity will be reduced by 10%. The intensity can be reduced further in steps of 10% until the patient can complete all four high-intensity intervals. If a patient is able to complete all high-intensity intervals, moderate and high intensity will be increased by 10%.⁹⁴ During neoadjuvant treatment⁹⁵: if a patient scores below a Borg score of 12 intensity is increased, if a patient scores above a Borg score of 15 intensity is decreased.

†Physical therapists can offer alternative resistance exercises targeting the same muscle group to accommodate a patient's abilities and preferences.

‡If the patient is able to do two repetitions more than planned, the load will be increased by 10%. The load will be decreased by 10% if the number of repetitions the patient achieves is two less than the planned number of repetitions.

1RM, one-repetition maximum; CPET, cardiopulmonary exercise test; VO_2 peak, oxygen uptake at peak exercise.

Primary outcome

The primary outcome is the proportion of patients who develop one or more grade ≥ 2 perioperative complications within 90 days after surgery. Complications are graded according to the Clavien-Dindo classification system³⁹ as described in [table 3](#).

Secondary outcomes

Secondary outcomes are the proportion of patients who develop one or more high-grade (grade ≥ 3) complications, total number of complications, length of hospital stay, number of readmissions, disease status (progression/recurrence) and additional treatment within 90 days. Secondary outcomes also include the intermediate outcomes of the intervention: cardiorespiratory fitness (measured using the standardised CPET)⁴⁰, physical functioning, upper and lower extremity (functional) muscle strength, nutritional intake, nutritional status, body composition, health-related quality of life, bladder cancer-related quality of life, anxiety and depression, fatigue, physical activity and adherence to the intervention. Direct and indirect costs are collected for the cost-effectiveness analysis. Tumour hypoxia, immune cell

infiltration and pathological response are assessed as explorative outcomes. Participants in the control group will be asked whether they participated in any (structured) lifestyle programme during the preoperative period, to monitor contamination. In both groups, participants will be asked whether they received any postoperative intervention. The timing and type of outcome measures are presented in [table 1](#).

Costs

For direct costs in both groups, medical records will be used to gather data on treatment, complications, length of hospital stay and number of readmissions, follow-up and diagnostics. For the intervention group, the costs for the prehabilitation programme, including physical exercise training, nutritional support, psychological counselling and smoking cessation, will be determined by means of the activity-based costing⁴¹ method. Costs for neoadjuvant treatment are expected to be equal for both arms and will therefore not be included in the calculation. For indirect costs, the iMTA Medical Consumption Questionnaire (iMCQ)⁴² will be used to gather data on medical consumption outside the hospital. The iMCQ includes

Table 3 The Clavien-Dindo classification of surgical complications

Grade	Definition
Grade I	Any deviation from the normal postoperative course without the need for pharmacological treatment or surgical, endoscopic and radiological interventions. Allowed therapeutic regimens are: drugs as antiemetics, antipyretics, analgetics, diuretics and electrolytes and physical therapy. This grade also includes wound infections opened at the bedside.
Grade II	Requiring pharmacological treatment with drugs other than such allowed for grade I complications. Blood transfusions and total parenteral nutrition are also included.
Grade III	Requiring surgical, endoscopic or radiological intervention.
Grade-IIIa	Intervention not under general anaesthesia.
Grade-IIIb	Intervention under general anaesthesia.
Grade IV	Life-threatening complication (including CNS complications)* requiring IC/ICU-management.
Grade-IVa	Single organ dysfunction (including dialysis).
Grade-IVb	Multiorgan dysfunction.
Grade V	Death of a patient.

Adapted from the study by Dindo *et al.*³⁹
*Brain haemorrhage, ischaemic stroke, subarachnoidal bleeding, but excluding transient ischaemic attacks.
CNS, central nervous system; IC, intermediate care; ICU, intensive care unit.

questions related to frequently occurring contacts with healthcare providers. For patients who currently have remunerative employment, productivity losses will be obtained through the iMTA Productivity Costs Questionnaire.⁴³ To calculate quality-adjusted life years, utilities will be derived from the EuroQol 5-dimension-5 levels (EQ-5D-5L) questionnaire.⁴⁴

Compliance

To monitor therapeutic validity, compliance with the physical exercise training programme is assessed by attendance rates and compliance to all parts of the training programme as scored by the physical therapist on standardised training session forms, as well as by patient self-report via an activity diary. Compliance to nutritional supplement intake is assessed via a diary. Whether smoking cessation has been successful is assessed through questionnaires.

Satisfaction

Patients in the intervention group are asked to complete a short questionnaire at the end of the study about the

perceived effectiveness of and satisfaction with the programme, whether they would suggest any changes and whether they would recommend it to other patients with bladder cancer. In addition, patients will be asked if they are willing to be contacted for participation in a focus group where the intervention programme and changes that may positively affect implementation will be further discussed.

Sample size calculation

Previous studies in diverse types of cancer reported a reduction of postoperative complications in the intervention group compared with the control group, with ORs ranging from 0.11 to 0.88.^{45–51} This study aims to reduce the number of patients with any grade ≥ 2 perioperative complication within 90 days from 60%^{52 53} to 35% (relative risk 0.58, OR 0.36).^{54 55} Assuming a two-sided Fisher's exact test with a power of 80% and an alpha of 0.05, in total 140 patients will be needed. To account for 10% dropout, including dropout due to cancelling of the planned surgery, 154 patients will be included. Approximately 380 patients with bladder cancer undergo RC in the eight participating hospitals annually. This implies that it is feasible to complete inclusion within two and a half years, if recruitment rate is at least 17%.

Statistical analysis

Primary outcome

All analyses will be performed on an intention-to-treat basis. Descriptive statistics will be calculated to describe and evaluate the comparability of the two groups at baseline on socio-demographic and clinical variables, and to assess the adequacy of the randomisation. Patients who do not receive the planned surgery or have an open-closed procedure, independent of group allocation, will be excluded from the primary analysis. The proportion of patients who develop any grade ≥ 2 complications will be compared in the two study arms by using Fisher's exact test and a Poisson regression model with a log link, adjusted for the stratification factors and relevant baseline imbalances. The relative risk will be reported with a 95% CI based on robust SEs.⁵⁶

Secondary study outcomes

Between-group differences over time will be evaluated in measures of physical functioning and patient-reported outcomes using linear mixed effects regression analysis. For high-grade complications, length of hospital stay and number of readmissions, Poisson regression models with an appropriate link function will be used. For continuous outcomes, differences in mean change scores between the two study arms will be accompanied by effect sizes. Standardised effect sizes will be calculated by subtracting the mean change scores of the control group from those of the intervention group, and subsequently dividing this by the pooled SD. Effect sizes of 0.2 are considered small, 0.5 moderate and 0.8 large.⁵⁷ A p value < 0.05 will be considered statistically significant.

Intervention fidelity

Descriptive statistics will be used to summarise compliance rates of the supervised exercise sessions as well as home-based physical activity, supplement consumption and smoking cessation. Compliance rates are based on number of completed training sessions, supplementation consumption and number of patients who stop smoking in the study intervention. Whether the level of compliance is associated with changes over time in primary and secondary study outcomes will be evaluated using generalised linear mixed effects models.

Non-participants

Baseline data of participants will be compared with those of non-participants using χ^2 statistics for categorical variables and analysis of variance for continuous variables.

Exploratory analysis

Exploratory analyses will be performed to explore the moderating effect of intervention duration, and differences in effectiveness of the physical exercise training programme between those who received neoadjuvant chemotherapy and those who did not. This will be done by adding interaction terms to the model and by performing stratified analyses if the interaction term is statistically significant at $p < 0.10$. Exploratory analyses will also be executed to study the relationship of post-intervention/preoperative physical fitness parameters (ie, oxygen uptake at peak exercise) and nutritional status, with perioperative outcomes (grade ≥ 2 complications yes/no and number of days in the hospital). For this analysis, univariable and multivariable Poisson regression analyses will be used.

Economic evaluation

A trial-based and model-based economic evaluation will be performed, based on the intention-to-treat analysis. The model-based evaluation will use literature for the potential long-term consequences and parametric survival methods to extrapolate the trial data beyond the included follow-up. The analysis will be approached from a societal perspective of The Netherlands and a lifelong time horizon. A Markov decision model will be built, with relevant health states derived from the EQ-5D-5L questionnaire. Outcomes are (1) the incremental costs per reduced proportion of patients who develop one or more grade ≥ 2 perioperative complications within 90 days (trial-based), and (2) incremental costs, incremental quality-adjusted life years and the incremental cost-effectiveness ratio (model-based). An estimation of the degree of uncertainty around each input parameter will be included with the use of probabilistic sensitivity analyses. Parameter values will be drawn randomly from the assigned distributions, using Monte Carlo simulations.⁵⁸ To capture necessary support regarding adoption and further research, value of information analyses will be performed.⁵⁹ Where appropriate, Dutch guidelines for costing studies will be used in applying tariffs to units of resource use.⁶⁰ Finally,

a budget impact analysis will be performed according to the International Society for Pharmacoeconomics and Outcomes Research guidelines.⁶¹

Patient and public involvement

Patients or the public were not involved in the development of the study design. A patient representative is currently involved in the study and input will be obtained whenever relevant during the trial. Annual consortium meetings with the urologists are organised.

DISCUSSION

A considerable proportion of patients with bladder cancer scheduled for RC has a poor cardiorespiratory fitness,^{9 10} is malnourished^{11 62} and/or is an active smoker at diagnosis.⁶³ This implies that there is a substantial potential for improving cardiorespiratory fitness and nutritional status in this patient population. Here, the rationale and design of a multimodal prehabilitation programme for patients with bladder cancer who are scheduled for RC is presented. It is hypothesised that the programme will be effective in reducing the number and severity of perioperative complications.

This study has several strengths, including its multi-centre, randomised design, the use of an intention-to-treat basis for the data analysis and a minimisation technique to ensure blinded treatment allocation and comparable groups. Most importantly, the study intervention consists of a tailored programme for physical exercise training and nutritional support following current best-practice for prehabilitation. Moreover, a cost-effectiveness analysis will be performed to anticipate smooth implementation and reimbursement, and tumour hypoxia and immune cell infiltration analysis will be performed exploratively. Intervention fidelity will be monitored in detail, as recommended previously,⁶⁴ as will adverse events related to the intervention. Another notable strength is the additional analysis in non-participants. Selective non-participation is a serious risk for the generalisability of physical exercise training studies. Previous physical exercise training studies in other cancer populations have shown relevant differences between those who participate and those who do not.^{65 66} It has previously been described that patients who were eligible for prehabilitation programmes for colon cancer surgery expressed several reservations.⁶⁷ It is vital to understand the characteristics of non-participants and reasons for non-participation in bladder cancer prehabilitation. This will not only help judge the generalisability of the results but will also support implementation in a way that will maximise the potential value of the prehabilitation programme and achieve equitable health outcomes. Finally, the exploratory subgroup analysis in patients who receive neoadjuvant treatment might be relevant to inform future studies on risk stratification.



To limit barriers to participation and adherence, the exercise programme will be delivered as near to the patients' homes as possible, to minimise travel time. This is a very important factor for patients with cancer according to previous studies.^{55 68–70} The availability of and close collaboration with the nationwide Onconet network of physical therapists, who are specifically educated to supervise patients with cancer, is an important advantage. An additional benefit of this approach is that it will facilitate implementation if the intervention proves to be effective.

This study also has some limitations. Patients with bladder cancer will be followed for a period of 90 days after surgery, meaning that longer-term evaluation of outcomes will not be possible. Because up to 60% of patients report complications within 90 days after surgery,⁷¹ it is expected that the time frame will be adequate for our primary outcome. A possible limitation is the risk of contamination in the control group. An evaluation questionnaire will be used in the control group to determine whether patients were physically active or received a (structured) lifestyle intervention preoperatively and postoperatively. Although objective measurements of habitual physical activity may provide a more detailed insight into physical activity levels not prone to recall bias, the collection of physical activity levels will be restricted to using questionnaires for feasibility reasons. This programme is designed to maximally improve a patient's health status by including physical exercise training, nutritional support, psychological counselling and smoking cessation. It is not likely that patients who are randomised to the control group would initiate a programme consisting of all these components on their own. The multimodal approach prohibits disentangling of the individual effects of each lifestyle component in the prehabilitation programme. Considering the number of intervention components and the prevalence of bladder cancer, a larger study using a full factorial design is unlikely to be feasible in this population. Moreover, the current best-practice for other types of cancer supports the use of multimodal interventions over unimodal approaches.^{72 73} Higher levels of physical exercise training have been demonstrated to be beneficial for both cancer prevention and, in some solid tumours, progression of disease and cancer-related mortality.^{74 75} However, the underlying biological mechanism has yet to be demonstrated. It is expected that the prehabilitation programme has positive effects on the tumour microenvironment. The hypoxic tumour microenvironment is a common characteristic of a solid tumour when oxygen levels become low, as a result of the rapid proliferation of tumour cells⁷⁶ and is linked to poor prognosis in bladder cancer.⁷⁷ Physical exercise training has regulatory effects on the angiogenesis of skeletal muscles, which has raised interest in whether these effects might translate to solid tumours.⁷⁸ Preclinical research has shown that training may acutely reduce tumour hypoxia through vascular normalisation and thereby improve the perfusion of tumour tissue.^{79–82} In addition, exercise training has

been suggested to alter immune cell infiltration in solid tumours and thereby contribute to enhanced immune surveillance and improved vascular function.²⁵ However, current evidence is inconsistent and inconclusive.⁸³ Clinical trials are very limited and this preoperative setting provides an excellent opportunity to investigate the potential role of prehabilitation on tumour hypoxia and immune cell infiltration.

To summarise, this study will provide empirical evidence on the benefits of multimodal prehabilitation for patients with bladder cancer planned for RC who are at high risk of perioperative complications and a long recovery period. When proven (cost-)effective, the study results will support implementation of a multimodal prehabilitation programme for patients with bladder cancer in daily clinical practice.

Author affiliations

¹Division of Psychosocial Research and Epidemiology, The Netherlands Cancer Institute, Amsterdam, The Netherlands

²Center for Quality of Life, The Netherlands Cancer Institute, Amsterdam, The Netherlands

³Department for Health Evidence, Radboud University Medical Center, Nijmegen, The Netherlands

⁴Department of Laboratory Medicine, Karolinska Institutet, Stockholm, Sweden

⁵Department of Oncological Urology, University Medical Center Utrecht, Utrecht, The Netherlands

⁶Department of Urology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

⁷Department of Urology, Radboud University Medical Center, Nijmegen, The Netherlands

⁸Department of Urology, Erasmus MC Cancer Institute, Erasmus University Medical Center, Rotterdam, The Netherlands

⁹Department of Urology, Catharina Hospital, Eindhoven, The Netherlands

¹⁰Department of Urology, Noordwest Hospital Group, Alkmaar, The Netherlands

¹¹Department of Urology, Rijnstate Hospital, Arnhem, The Netherlands

¹²Department of Urology, Maastricht University Medical Center+, Maastricht, The Netherlands

¹³Department of Nutrition and Movement Sciences, Nutrition and Translational Research in Metabolism (NUTRIM), Faculty of Health, Medicine and Life Sciences, Maastricht University, Maastricht, The Netherlands

¹⁴Department of Surgery, Nutrition and Translational Research in Metabolism (NUTRIM), Faculty of Health, Medicine and Life Sciences, Maastricht University, Maastricht, The Netherlands

¹⁵Erasmus School of Health Policy and Management, Erasmus University Rotterdam, Rotterdam, The Netherlands

¹⁶Department of Health Technology and Services Research, University of Twente, Enschede, The Netherlands

¹⁷Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, The Netherlands

¹⁸Department of Medicine for Older People, Amsterdam University Medical Center Locatie VUmc, Amsterdam, The Netherlands

¹⁹Amsterdam Public Health, Aging & Later Life, Amsterdam, The Netherlands

²⁰Center of Expertise Urban Vitality, Faculty of Health, Amsterdam University of Applied Sciences, Amsterdam, The Netherlands

Acknowledgements Research at the Netherlands Cancer Institute is supported by institutional grants of the Dutch Cancer Society and of the Dutch Ministry of Health, Welfare and Sport. The protein supplements in the clinical study will be kindly provided by FrieslandCampina.

Contributors AV, AML-K, AGH, ELK, VPR, WHVH, AMM, WGG and MMS conceived the study. AV, WHVH, AMM, WGG, MMS, MGS and EA contributed to the design of the study protocol. RPM, AML-K, AGH, VCR, ELK, SDB, CJW, TATM and BCB reviewed the study protocol with a clinical perspective. VPR wrote the statistical approach of the economic evaluation. HR composed the protocol of tumour hypoxia and immune

cell infiltration analysis. AV, WHVH, AMM, WGG, MMS, MGS, and EA wrote the manuscript. All authors read, commented on and approved the manuscript.

Funding This work was supported by the Dutch Cancer Society (KWF, Delflandlaan 17, 1062 EA, Amsterdam, The Netherlands), grant number NKI 2020-1/12877. The funder has no role in the design, data collection and analysis of the study.

Competing interests CJW received a research grant from ZonMw and gives courses on robot-assisted radical cystectomy during conferences entitled with EAU. TATM received consulting fees as member of the advisory board prostate cancer of Janssen-Cilag B.V. BCB participates on a Data Safety Monitoring Board of an international prehabilitation study, is board member of Netherlands Society of Human Movement sciences, member of Exercise is Medicine of Vereniging voor Sportgeneeskunde and member of the Scientific Committee of Fit4Surgery. RPM received research grants from Janssen, Roche and Astellas, an educational grant from Merck, received consulting fees from Merck, MSD, Janssen, Bristol-Myers Squibb Company, received support for attending the 2022 Global Congress on Bladder Cancer, and is EAU Panel member of the Muscle Invasive Bladder Cancer Guideline. MMS is member of the Scientific Committee of Fit4Surgery and board member of Onconet.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Emine Akdemir <http://orcid.org/0000-0003-4220-5242>

Alina Vrieling <http://orcid.org/0000-0002-8178-2125>

Vera C Rutten <http://orcid.org/0000-0003-2452-5357>

Valesca P Retèl <http://orcid.org/0000-0002-5624-3234>

REFERENCES

- Sung H, Ferlay J, Siegel RL, *et al*. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2021;71:209–49.
- Stein JP, Skinner DG. Radical cystectomy for invasive bladder cancer: long-term results of a standard procedure. *World J Urol* 2006;24:296–304.
- Alfred Witjes J, Lebre T, Compérat EM, *et al*. Updated 2016 EAU guidelines on muscle-invasive and metastatic bladder cancer. *Eur Urol* 2017;71:462–75.
- Babjuk M, Burger M, Capoun O, *et al*. European association of urology guidelines on non-muscle-invasive bladder cancer (TA, T1, and carcinoma in situ). *Eur Urol* 2022;81:75–94.
- Maibom SL, Joensen UN, Poulsen AM, *et al*. Short-Term morbidity and mortality following radical cystectomy: a systematic review. *BMJ Open* 2021;11:e043266.
- Shabsigh A, Korets R, Vora KC, *et al*. Defining early morbidity of radical cystectomy for patients with bladder cancer using a standardized reporting methodology. *Eur Urol* 2009;55:164–74.
- Novara G, De Marco V, Aragona M, *et al*. Complications and mortality after radical cystectomy for bladder transitional cell cancer. *J Urol* 2009;182:914–21.
- Bochner BH, Sjoberg DD, Laudone VP, *et al*. A randomized trial of robot-assisted laparoscopic radical cystectomy. *N Engl J Med* 2014;371:389–90.
- Prentis JM, Trenell MI, Vasdev N, *et al*. Impaired cardiopulmonary reserve in an elderly population is related to postoperative morbidity and length of hospital stay after radical cystectomy. *BJU Int* 2013;112:E13–9.
- Tolchard S, Angell J, Pyke M, *et al*. Cardiopulmonary reserve as determined by cardiopulmonary exercise testing correlates with length of stay and predicts complications after radical cystectomy. *BJU Int* 2015;115:554–61.
- Arora K, Hanson KT, Habermann EB, *et al*. Early complications and mortality following radical cystectomy: associations with malnutrition and obesity. *Bladder Cancer* 2018;4:377–88.
- Mavros MN, Athanasiou S, Gkegkes ID, *et al*. Do psychological variables affect early surgical recovery? *PLoS One* 2011;6:e20306.
- Sathianathen NJ, Weight CJ, Jarosek SL, *et al*. Increased surgical complications in smokers undergoing radical cystectomy. *Bladder Cancer* 2018;4:403–9.
- Jack S, West MA, Raw D, *et al*. The effect of neoadjuvant chemotherapy on physical fitness and survival in patients undergoing oesophagogastric cancer surgery. *Eur J Surg Oncol* 2014;40:1313–20.
- West MA, Parry MG, Lythgoe D, *et al*. Cardiopulmonary exercise testing for the prediction of morbidity risk after rectal cancer surgery. *Br J Surg* 2014;101:1166–72.
- Sinclair R, Navidi M, Griffin SM, *et al*. The impact of neoadjuvant chemotherapy on cardiopulmonary physical fitness in gastro-oesophageal adenocarcinoma. *Ann R Coll Surg Engl* 2016;98:396–400.
- Thomson IG, Wallen MP, Hall A, *et al*. Neoadjuvant therapy reduces cardiopulmonary function in patients undergoing oesophagectomy. *Int J Surg* 2018;53:86–92.
- Banerjee S, Manley K, Shaw B, *et al*. Vigorous intensity aerobic interval exercise in bladder cancer patients prior to radical cystectomy: a feasibility randomised controlled trial. *Support Care Cancer* 2018;26:1515–23.
- Kaye DR, Schafer C, Thelen-Perry S, *et al*. The feasibility and impact of a presurgical exercise intervention program (prehabilitation) for patients undergoing cystectomy for bladder cancer. *Urology* 2020;145:106–12.
- Jensen BT, Petersen AK, Jensen JB, *et al*. Efficacy of a multiprofessional rehabilitation programme in radical cystectomy pathways: a prospective randomized controlled trial. *Scand J Urol* 2015;49:133–41.
- Minnella EM, Awasthi R, Bousquet-Dion G, *et al*. Multimodal prehabilitation to enhance functional capacity following radical cystectomy: a randomized controlled trial. *Eur Urol Focus* 2021;7:132–8.
- Jensen BT, Laustsen S, Jensen JB, *et al*. Exercise-based prehabilitation is feasible and effective in radical cystectomy pathways—secondary results from a randomized controlled trial. *Support Care Cancer* 2016;24:3325–31.
- Thomas G, Tahir MR, Bongers BC, *et al*. Prehabilitation before major intra-abdominal cancer surgery: a systematic review of randomised controlled trials. *Eur J Anaesthesiol* 2019;36:933–45.
- Wiggins JM, Opoku-Acheampong AB, Baumfalk DR, *et al*. Exercise and the tumor microenvironment: potential therapeutic implications. *Exerc Sport Sci Rev* 2018;46:56–64.
- Zhang X, Ashcraft KA, Betof Warner A, *et al*. Can exercise-induced modulation of the tumor physiologic microenvironment improve antitumor immunity? *Cancer Res* 2019;79:2447–56.
- Shannon AM, Bouchier-Hayes DJ, Condron CM, *et al*. Tumour hypoxia, chemotherapeutic resistance and hypoxia-related therapies. *Cancer Treat Rev* 2003;29:297–307.
- Peixoto A, Fernandes E, Gaiteiro C, *et al*. Hypoxia enhances the malignant nature of bladder cancer cells and concomitantly antagonizes protein O-glycosylation extension. *Oncotarget* 2016;7:63138–57.
- Boström PJ, Thoms J, Sykes J, *et al*. Hypoxia marker GLUT-1 (glucose transporter 1) is an independent prognostic factor for survival in bladder cancer patients treated with radical cystectomy. *Bladder Cancer* 2016;2:101–9.
- Pocock SJ, Simon R. Sequential treatment assignment with balancing for prognostic factors in the controlled clinical trial. *Biometrics* 1975;31:103–15.
- Jin M, Polis A, Hartzel J. R package “minirand”. 0.1.3. ed. CRAN; 2020. Available: <https://cran.r-project.org/web/packages/Minirand/Minirand.pdf>
- Jin M, Polis A, Hartzel J. Algorithms for minimization randomization and the implementation with an R package. *Communications in Statistics - Simulation and Computation* 2021;50:3077–87.
- Campbell KL, Winters-Stone KM, Wiskemann J, *et al*. Exercise guidelines for cancer survivors: consensus statement from international multidisciplinary roundtable. *Med Sci Sports Exerc* 2019;51:2375–90.
- Rock CL, Thomson CA, Sullivan KR, *et al*. American cancer Society nutrition and physical activity guideline for cancer survivors. *CA Cancer J Clin* 2022;72:230–62.
- Cerantola Y, Valerio M, Persson B, *et al*. Guidelines for perioperative care after radical cystectomy for bladder cancer: enhanced recovery after surgery (ERAS®) society recommendations. *Clin Nutr* 2013;32:879–87.
- Jacobson E. You must relax. 1934.

- 36 Cermak NM, Res PT, de Groot LCPGM, *et al.* Protein supplementation augments the adaptive response of skeletal muscle to resistance-type exercise training: a meta-analysis. *Am J Clin Nutr* 2012;96:1454–64.
- 37 Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67:361–70.
- 38 Antonovsky H, Sagy S. The development of a sense of coherence and its impact on responses to stress situations. *J Soc Psychol* 1986;126:213–25.
- 39 Dindo D, Demartines N, Clavien P-A. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 2004;240:205–13.
- 40 Berkel AEM, Bongers BC, van Kamp M-JS, *et al.* The effects of rehabilitation versus usual care to reduce postoperative complications in high-risk patients with colorectal cancer or dysplasia scheduled for elective colorectal resection: study protocol of a randomized controlled trial. *BMC Gastroenterol* 2018;18:29.
- 41 Cooper R, Kaplan RS. Profit priorities from activity-based costing. *Harv Bus Rev* 1991;69:130–5.
- 42 Bouwmans *Cet al.* *Medical cost questionnaire (imcq)*. Rotterdam: iMTA, Erasmus Universiteit Rotterdam, 2013.
- 43 Bouwmans *Cet al.* *Productivity cost questionnaire*. Erasmus University Rotterdam: Institute for Medical Technology Assessment, 2013.
- 44 Uyl-de Groot CA, Rutten FF, Bonsel GJ. Measurement and valuation of quality of life in economic appraisal of cancer treatment. *Eur J Cancer* 1994;30A:111–7.
- 45 Dronkers J, Veldman A, Hoberg E, *et al.* Prevention of pulmonary complications after upper abdominal surgery by preoperative intensive inspiratory muscle training: a randomized controlled pilot study. *Clin Rehabil* 2008;22:134–42.
- 46 Dronkers JJ, Lamberts H, Reutelingsperger IMMD, *et al.* Preoperative therapeutic programme for elderly patients scheduled for elective abdominal oncological surgery: a randomized controlled pilot study. *Clin Rehabil* 2010;24:614–22.
- 47 Kulkarni SR, Fletcher E, McConnell AK, *et al.* Pre-operative inspiratory muscle training preserves postoperative inspiratory muscle strength following major abdominal surgery - a randomised pilot study. *Ann R Coll Surg Engl* 2010;92:700–7.
- 48 Barbalho-Moulim MC, Miguel GPS, Forti EMP, *et al.* Effects of preoperative inspiratory muscle training in obese women undergoing open bariatric surgery: respiratory muscle strength, lung volumes, and diaphragmatic excursion. *Clinics (Sao Paulo)* 2011;66:1721–7.
- 49 Soares SM de TP, Nucci LB, da Silva MM de C, *et al.* Pulmonary function and physical performance outcomes with preoperative physical therapy in upper abdominal surgery: a randomized controlled trial. *Clin Rehabil* 2013;27:616–27.
- 50 van Adrichem EJ, Meulenbroek RL, Plukker JTM, *et al.* Comparison of two preoperative inspiratory muscle training programs to prevent pulmonary complications in patients undergoing esophagectomy: a randomized controlled pilot study. *Ann Surg Oncol* 2014;21:2353–60.
- 51 Halliday LJ, Doganay E, Wynter-Blyth VA, *et al.* The impact of prehabilitation on post-operative outcomes in oesophageal cancer surgery: a propensity score matched comparison. *J Gastrointest Surg* 2021;25:2733–41.
- 52 Low DE, Kuppusamy MK, Alderson D, *et al.* Benchmarking complications associated with esophagectomy. *Ann Surg* 2019;269:291–8.
- 53 Wijburg CJ, Michels CTJ, Hannink G, *et al.* Robot-assisted radical cystectomy versus open radical cystectomy in bladder cancer patients: a multicentre comparative effectiveness study. *Eur Urol* 2021;79:609–18.
- 54 Barberan-Garcia A, Ubré M, Roca J, *et al.* Personalised prehabilitation in high-risk patients undergoing elective major abdominal surgery: a randomized blinded controlled trial. *Ann Surg* 2018;267:50–6.
- 55 Berkel AEM, Bongers BC, Kotte H, *et al.* Effects of community-based exercise prehabilitation for patients scheduled for colorectal surgery with high risk for postoperative complications: results of a randomized clinical trial. *Ann Surg* 2022;275:e299–306.
- 56 Zou G. A modified poisson regression approach to prospective studies with binary data. *Am J Epidemiol* 2004;159:702–6.
- 57 Cohen J. *Statistical power analysis for the behavioral sciences*. Routledge, 2013.
- 58 Briggs A, Sculpher M, Claxton K. *Decision modelling for health economic evaluation*. Oup Oxford, 2006.
- 59 Vallejo-Torres L, Steuten LMG, Buxton MJ, *et al.* Integrating health economics modeling in the product development cycle of medical devices: a Bayesian approach. *Int J Technol Assess Health Care* 2008;24:459–64.
- 60 Hakkaart-van Roijen L, Tan S, Bouwmans C. *Manual for cost research: methods and standard cost prices for economic evaluations in health care*. Diemen: Health Care Insurance Board, 2010.
- 61 Mauskopf JA, Sullivan SD, Annemans L, *et al.* Principles of good practice for budget impact analysis: report of the ISPOR Task force on good research practices -- budget impact analysis. *Value Health* 2007;10:336–47.
- 62 Gregg JR, Cookson MS, Phillips S, *et al.* Effect of preoperative nutritional deficiency on mortality after radical cystectomy for bladder cancer. *J Urol* 2011;185:90–6.
- 63 Barbosa ALA, Vermeulen SHHM, Aben KK, *et al.* Smoking intensity and bladder cancer aggressiveness at diagnosis. *PLoS One* 2018;13:e0194039.
- 64 Bellg AJ, Borrelli B, Resnick B, *et al.* Enhancing treatment fidelity in health behavior change studies: best practices and recommendations from the NIH behavior change Consortium. *Health Psychol* 2004;23:443–51.
- 65 van Waart H, van Harten WH, Buffart LM, *et al.* Why do patients choose (not) to participate in an exercise trial during adjuvant chemotherapy for breast cancer? *Psychooncology* 2016;25:964–70.
- 66 Vd Wiel HJ, Stuiver MM, May AM, *et al.* Characteristics of participants and nonparticipants in a blended internet-based physical activity trial for breast and prostate cancer survivors: cross-sectional study. *JMIR Cancer* 2021;7:e25464.
- 67 Agasi-Idenburg CS, Zuilen MK, Westerman MJ, *et al.* “I am busy surviving” - views about physical exercise in older adults scheduled for colorectal cancer surgery. *J Geriatr Oncol* 2020;11:444–50.
- 68 Ormel HL, van der Schoot GGF, Sluiter WJ, *et al.* Predictors of adherence to exercise interventions during and after cancer treatment: a systematic review. *Psychooncology* 2018;27:713–24.
- 69 Banerjee S, Semper K, Skarparis K, *et al.* Patient perspectives of vigorous intensity aerobic interval exercise prehabilitation prior to radical cystectomy: a qualitative focus group study. *Disabil Rehabil* 2021;43:1084–91.
- 70 van Wijk L, Bongers BC, Berkel AEM, *et al.* Improved preoperative aerobic fitness following a home-based bimodal prehabilitation programme in high-risk patients scheduled for liver or pancreatic resection. *Br J Surg* 2022;109:1036–9.
- 71 Krajewski W, Zdrojowy R, Tupikowski K, *et al.* How to lower postoperative complications after radical cystectomy—a review. *Cent European J Urol* 2016;69:370–6.
- 72 Mewes JC, Steuten LMG, Ijzerman MJ, *et al.* Effectiveness of multidimensional cancer Survivor rehabilitation and cost-effectiveness of cancer rehabilitation in general: a systematic review. *Oncologist* 2012;17:1581–93.
- 73 Lukez A, Baima J. The role and scope of prehabilitation in cancer care. *Semin Oncol Nurs* 2020;36:150976.
- 74 Moore SC, Lee I-M, Weiderpass E, *et al.* Association of leisure-time physical activity with risk of 26 types of cancer in 1.44 million adults. *JAMA Intern Med* 2016;176:816–25.
- 75 Friedenreich CM *et al.* Physical activity and cancer outcomes: A precision medicine approach. *Clin Cancer Res* 2016;22:4766–75.
- 76 Petrova V, Annicchiarico-Petruzzelli M, Melino G, *et al.* The hypoxic tumour microenvironment. *Oncogenesis* 2018;7:1–13.
- 77 Theodoropoulos VE, Lazaris AC, Sofras F, *et al.* Hypoxia-Inducible factor 1 alpha expression correlates with angiogenesis and unfavorable prognosis in bladder cancer. *Eur Urol* 2004;46:200–8.
- 78 Koelwyn GJ, Quail DF, Zhang X, *et al.* Exercise-dependent regulation of the tumour microenvironment. *Nat Rev Cancer* 2017;17:620–32.
- 79 Jones LW, Viglianti BL, Tashjian JA, *et al.* Effect of aerobic exercise on tumor physiology in an animal model of human breast cancer. *J Appl Physiol (1985)* 2010;108:343–8.
- 80 McCullough DJ, Nguyen LM-D, Siemann DW, *et al.* Effects of exercise training on tumor hypoxia and vascular function in the rodent preclinical orthotopic prostate cancer model. *J Appl Physiol (1985)* 2013;115:1846–54.
- 81 Betof AS, Lascola CD, Weitzel D, *et al.* Modulation of murine breast tumor vascularity, hypoxia and chemotherapeutic response by exercise. *J Natl Cancer Inst* 2015;107:djv040.
- 82 Schumacher O, Galvão DA, Taaffe DR, *et al.* Exercise modulation of tumour perfusion and hypoxia to improve radiotherapy response in prostate cancer. *Prostate Cancer Prostatic Dis* 2021;24:1–14.
- 83 Seet-Lee C, Yee J, Morahan H, *et al.* The effect of aerobic exercise on tumour blood delivery: a systematic review and meta-analysis. *Support Care Cancer* 2022;30:8637–53.
- 84 Guralnik JM, Simonsick EM, Ferrucci L, *et al.* A short physical performance battery assessing lower extremity function: association with self-reported disability and prediction of mortality and nursing home admission. *J Gerontol* 1994;49:M85–94.
- 85 Langius *et al.* *Meetprotocol handknijpkracht mbv hand dynamometer*. Nutritional Assessment Platform, 2016.

- 86 McAllister LS, Palombaro KM. Modified 30-second sit-to-stand test: reliability and validity in older adults unable to complete traditional sit-to-stand testing. *J Geriatr Phys Ther* 2020;43:153–8.
- 87 Bauer J, Capra S, Ferguson M. Use of the scored patient-generated subjective global assessment (PG-SGA) as a nutrition assessment tool in patients with cancer. *Eur J Clin Nutr* 2002;56:779–85.
- 88 Aaronson NK, Ahmedzai S, Bergman B, *et al.* The European organization for research and treatment of cancer QLQ-C30: a quality-of-life instrument for use in international clinical trials in oncology. *J Natl Cancer Inst* 1993;85:365–76.
- 89 Danna BJ, Metcalfe MJ, Wood EL, *et al.* Assessing symptom burden in bladder cancer: an overview of bladder cancer specific health-related quality of life instruments. *Bladder Cancer* 2016;2:329–40.
- 90 Smets EM, Garssen B, Bonke B, *et al.* The multidimensional fatigue inventory (MFI) psychometric qualities of an instrument to assess fatigue. *J Psychosom Res* 1995;39:315–25.
- 91 Wendel-Vos GCW, Schuit AJ, Saris WHM, *et al.* Reproducibility and relative validity of the short questionnaire to assess health-enhancing physical activity. *J Clin Epidemiol* 2003;56:1163–9.
- 92 Sangha O, Stucki G, Liang MH, *et al.* The self-administered comorbidity questionnaire: a new method to assess comorbidity for clinical and health services research. *Arthritis Rheum* 2003;49:156–63.
- 93 Antonovsky A. The structure and properties of the sense of coherence scale. *Soc Sci Med* 1993;36:725–33.
- 94 van Rooijen S, Carli F, Dalton S, *et al.* Multimodal prehabilitation in colorectal cancer patients to improve functional capacity and reduce postoperative complications: the first international randomized controlled trial for multimodal prehabilitation. *BMC Cancer* 2019;19:98.
- 95 Swain DP, Brawner CA, A.C.o.S. Medicine. *ACSM's resource manual for guidelines for exercise testing and prescription*. Wolters Kluwer Health/Lippincott Williams & Wilkins, 2014.