

## The exhumation of the westernmost Tien Shan (Tajikistan, Uzbekistan)

Sanaa Abdulhameed<sup>1</sup>, Lukasz Gagala<sup>1</sup>, Eva Enkelmann<sup>2</sup>, Raymond Jonckheere<sup>1</sup>, Lothar Ratschbacher<sup>1</sup>

<sup>1</sup>Geology, TU Bergakademie Freiberg, 09599 Freiberg, Germany

<sup>2</sup>Geology, University of Cincinnati, USA

Contact: Sanaa Abdulhameed

shameed84@hotmail.com

The westernmost segment of the Tien Shan and the Tajik-Afghan basin of Tajikistan and Uzbekistan (Central Asia) lie northwest of the Pamir and southeast of the Turan platform. The Tajik-Afghan Basin comprises up to 10 km thick, mostly clastic deposits, ranging in age from Lower Jurassic to Quaternary. A thin-skinned fold-and-thrust belt developed within the Cretaceous through Neogene sedimentary rocks; the basal detachment runs along the Lower Jurassic evaporates. In Uzbekistan, crystalline basement of the Ghissar unit of the westernmost Tien Shan is involved in the folding/thrusting of the Jurassic to Neogene sediments, proving thick-skinned deformation and suggesting basement involvement below the Jurassic evaporites underneath the Tajik-Afghan basin. Most of the thrusts in the region trend (NNE-SSW), proving an ~E-W shortening component during the India-Asia collision.

We collected more than forty samples from the basement rocks of the Tien Shan and the Cretaceous clastic rocks of the Tajik-Afghan Basin above the major thrusts to constrain the timing of deformation and exhumation of the region using apatite fission-track (AFT) and apatite (U-Th)/He (AHe) thermochronology. The samples from the Ghissar Range of Tajikistan and from the southern part of the Tajik-Afghan Basin yielded fully reset apatite ages, ranging between  $11.1 \pm 0.7$  and  $7.6 \pm 0.3$  Ma; we interpret these ages as recording cooling as result of shortening within the fold-and-thrust belt and coeval erosion. In the Kurgan section in the central Tajik-Afghan Basin, the samples are partially reset and the mean ages range between  $26.2 \pm 1.2$  and  $17.2 \pm 0.9$  Ma. Heavy ion irradiation and multiple edging provided a statistically significant amount of confined fission tracks to allow temperature-time history modeling. Cooling in this part of the Tajik-Afghan Basin likely started at  $\leq 20$  Ma.

The basement samples from the Tien Shan in Uzbekistan have AFT ages ranging between  $12.5 \pm 0.8$  and  $6.7 \pm 0.7$  Ma; temperature-time history models support rapid cooling within this period. The simplest interpretation of the combined data in the Tajik-Afghan Basin and in the crystalline basement of the Uzbek Tien Shan suggests a two-phase exhumation/cooling history: (1) Regional exhumation likely initiated at  $\leq 20$  Ma. (2) Exhumation/cooling by erosion initiated by deformation in the fold-and-thrust belt started at 13-7 Ma, reflecting slip along the Upper Jurassic evaporate detachment and faults in the sub-detachment basement.

-----  
Key words: Tajik Basin, Western Tien Shan, Thermochronology, Apatite fission track dating, (U-Th)/He dating.