

Evolution of Wifl: Expansion of Wifl Notation After Intervention

Virginie Blanchette^{1,2} , Malindu E. Fernando^{1,3,4}, Laura Shin¹, Vincent L. Rowe¹, Kenneth R. Ziegler¹, and David G. Armstrong¹

The International Journal of Lower Extremity Wounds
1–5

© The Author(s) 2022



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/15347346221122860

journals.sagepub.com/home/ijl



Abstract

Nearly a decade ago, the Society for Vascular Surgery (SVS)'s wound, ischemia, and foot Infection (Wifl) classification was first developed to help assess overall limb threat. However, managing conditions such as diabetic foot ulcer and chronic limb-threatening ischemia can be complex. For instance, certain investigative findings might initially be pending such as the level of ischemia or extent of infection before the final classification is established. In addition, wounds evolve rapidly, and the current classification does not allow for tracking their progression over time during treatment. Therefore, we propose a supplemental consistent notation for scoring Wifl re-assessment during treatment of a threatened limb inspired by the cancer staging before and after neoadjuvant treatment classification system. Thus, we describe the re-scoring system and how to use it. Our suggestion supports a coherent method to longitudinally communicate characteristics of a threatened limb. This has potential to support high quality interdisciplinary, patient-centered care and enhance the use of this classification in research. Further work is required to validate this modification of a common language of risk.

Keywords

diabetes, foot ulcer, wound healing, peripheral vascular disease, ischemia, classification, amputation

Diabetes-related foot complications such as diabetic foot ulcers (DFUs), peripheral arterial disease including critical limb-threatening ischemia (CLTI), Charcot neuroarthropathy and lower extremity amputations (LEAs) are a leading cause of global morbidity, mortality, reduced quality of life and direct and indirect healthcare costs.^{1–3} Indeed, 5-year mortality rates associated with these complications are greater compared to many cancers.⁴ Limb complications of diabetes increase in people with multimorbidity such as concomitant nephropathy and cardiovascular disease.^{4–6} People with an history of DFUs are at increased risk of ulcer recurrence.¹ In fact, 40% of those people will develop DFU recurrence within a year, 65% within 5 years, and greater than 90% within 10 years.^{1,7} Because the epidemiology of diabetes-related foot complications is comparable to that of cancer, and recurrence is common, after the initial healing of an index DFU, it is appropriate to refer to a person not as cured of DFU, but rather as being in “DFU remission”.⁸ In addition, as with cancer, the complexities associated with management require a comprehensive and organized team approach, including the patient, their family and caregivers, to achieve the best outcomes and high quality patient-centered care.^{9–12}

Nearly a decade ago, Mills, Conte, Armstrong and coworkers, in concert with the Society for Vascular Surgery (SVS) proposed an integrated lower extremity wound classification system related to LEA risk and revascularization benefit.¹³ Inspired by cancer research, it is intended to define the disease burden, analogous to the tumor, node, metastasis (TNM) system for cancer staging.^{13,14} Wifl consists of a graded scoring system for wound, ischemia, and foot infection. For any given

¹Southwestern Academic Limb Salvage Alliance (SALSA), Department of Surgery, Keck School of Medicine of University of Southern California, Los Angeles, CA, USA

²Department of Human Kinetics and Podiatric Medicine, Université du Québec à Trois-Rivières, Trois-Rivières, Québec, Canada

³Ulcer and wound Healing consortium (UHEAL), Queensland Research Centre for Peripheral Vascular Disease, College of Medicine and Dentistry, James Cook University, Townsville, Queensland, Australia

⁴Faculty of Health and Medicine, School of Health Sciences, University of Newcastle, Australia

Corresponding Author:

Virginie Blanchette, Southwestern Academic Limb Salvage Alliance (SALSA), Department of Surgery, Keck School of Medicine of University of Southern California, Los Angeles, CA, USA.

Email: Virginie.Blanchette@uqtr.ca

threatened limb, a severity grade of 0 to 3 (ie, none, mild, moderate, severe) is assigned to grade the severity and extent of wound, ischemia, and foot infection, respectively (Figure 1). On the basis of these three scores, patients are further assigned to four threatened limb clinical stages corresponding to estimated risk of LEA derived by an expert panel consensus. The underlying premise of Wifl is that the risk of LEA increases as the presenting disease burden progresses from clinical stage 1 (very low risk) to stage 4 (high risk).¹³ However, managing the threatened limb can be complex, and the impact of conservative, surgical, and medical interventions on the stabilization of DFUs is not reflected in the scoring system. Additionally, certain investigative findings might initially be pending such as the level of ischemia or extent of infection before the final classification is established. DFUs evolve rapidly, and the current classification does not allow for tracking limb progression over time during treatment and related to potential recurrence of the index DFU. Moreover, these limitations create a logistical challenge for utilizing Wifl for large scale reporting including in cohort studies and clinical trials. Therefore, consistency in notation is required for widespread application.

To continue the analogy of DFU with cancer staging, a recent publication suggested an updated classification to better reflect the situation after neoadjuvant therapies and to highlight the differences between clinical and pathological states.¹⁶ Given similar issues in categorizing DFUs post treatment such as revascularization and limb salvage, treatment of osteomyelitis or debridement or minor amputation, with a supplemental Wifl notation is appropriate and would assist the clinician as well as provide a consistent approach for use in clinical studies. A recent study demonstrated that




Figure 1. SWS Wifl classification system, adapted from Armstrong & Mills (2013).¹⁵

Wifl restaging is an important tool for predicting limb loss and assessing adequacy of intervention, more so than baseline Wifl alone.¹⁷ Therefore, the aim of this communication is to propose a revised, consistent notation for scoring Wifl re-assessment during treatment of a threatened limb for the clinical and research community with the intention to make it practical to use on a daily basis. Thus, we describe the re-scoring system.

Notation for re-Scoring Wifl

In order to address the issue of either upgrading or downgrading the Wifl score of a threatened limb, and to reflect the actual condition, and to allow the latitude of a pending score, identifying the timeline to the event and associated interventions is important. Thus, it will be possible to classify a DFU for example, according to the stage of management: before treatment, during/after treatment and at remission. This is separated in four phases: primary presentation (p), during therapy (y), healed (h) or recurrent (r).

When there is missing data to complete the final Wifl score, for example, when the clinician is awaiting the results of a pending vascular investigation to identify the ischemia status, or awaiting the results of histology/laboratory data to support the presence of bone involvement in infection, the number is temporarily replaced by the hashtag (#) until the score can be completed with certainty. In this case, it is still necessary to identify the index phase (Figure 2).

Thus, for an index ulcer, the assessor (ie, clinician or researcher) should always establish a score for the primary presentation that is associated with the risk of major LEA and/or the benefits of revascularization as a preliminary Wifl score or “pWifl” score. During and following the treatment, whatever the nature and number of treatments and the length of follow up, the scorer can define the classification as many times as needed by using a post treatment Wifl score or “tWifl” score. Considering that the goal is to stabilize (or to maintain in remission) the threatened limb condition, it is relevant to repeat the score when the condition is deemed healed (h). The repeated score is still relevant as there could still be an ischemic component to follow. Therefore, the user should note the status using “hWifl”.

A threatened limb, or healed limb is often evolving gradually, changing rapidly and is at high-risk of recurrence. Indeed, the healed Wifl score is not a static measurement and that it is in fact a cycle.

This cycle can then be restarted using the notation and this is based on the adapted clinical algorithm suggested by Conte and al. (2020) to implement limb staging with the Wifl as a part of the initial assessment of CLTI (Figure 3).¹⁸ On the other hand, the possibility that the cycle stops after the healing score is plausible. This indicates that the team has accomplished its mission to maximize




Figure 2. Schematization of the supplemental Wifl notation.

ulcer-free, hospital-free, and activity-rich days, the same way a cancer survivor is deemed to maximize cancer-free days.^{4,19} Alternatively, if limb salvage is not attained and limb loss ensues, then this is designated with a capital R to indicate limb loss due to an index ulcer as “RWifl” This is a comprehensive re-notation that can be preconize in research context.

Definitions

Primary Presentation (p)

This is the first Wifl score that is assigned when the diagnosis of the limb-threatening condition such DFU and CLTI. This score is made according to clinical criteria established by former guidelines. The pending score should be completed before proceeding to the next phase considering that it is the score that determines the LEA risk and revascularization risk according to the initial Wifl.¹³ This represents the initial state without treatment. This should be evaluated by a trained and competent health care provider that can diagnose the components of wound, ischaemia and infection within their scope of practice.

During Treatment/Therapy (t)

For our purpose, therapies, treatments, and cares are synonym to attempted remediation of the limb-threatening condition following the diagnosis. There are a multitude of treatments for the conditions that can supplement the Wifl score. Therefore, any action or way of treating a




Figure 3. Continuous evaluation and re-staging. Adapted clinical algorithm from Conte and al. 2020.²⁰ Abbreviation: Wifl: Wounds, Ischemia and foot Infection Classification; CLTI: Chronic Limb-Threatening Ischemia; DFU: Diabetic Foot Ulcer.

patient or a condition medically or surgically such as management and care to prevent, cure, ameliorate, or slow progression of a threatened limb condition is included in this phase. These treatments must be supported by the suitable

guidelines and respect the ethical principles of research with human subjects. This score can be used in the initial treatment strategy using the holistic patient, limb, anatomy (PLAN) framework or during the clinical reassessment and can be re-scored according to the clinical reassessment as needed.

Healed (or Controlled or Stable) (h)

Food and Drug Administration (FDA) defines healing as the 100% re-epithelialization of the wound surface with no discernible exudate and without drainage or dressing, confirmed at two visits two weeks apart.²¹ Moreover, a blinded adjudication for wound assessment is suggested in the case of a clinical trial.²² As discussed above, a DFU would be in remission instead of healed. The International working group on Diabetic Foot (IWGDF) defines it as an intact skin and absence of infection of the complete foot after healing of any DFUs.²³ Thus, the final score associated with this phase should be established according to the above conditions, confirmed at two week intervals by a final blinded wound assessment.²⁴

Recurrence (r)

The IWGDF defines DFU recurrence as a new foot ulcer in a person who has a history of DFU, irrespective of location and time, since previous DFU.²³ Nevertheless, with this notation, the recurrences are counted only if it is the same indexed DFU/wounds. If it is a new wound location, it is not a recurrence. This is supporting the complexity of the lower limb vascularization related to the healing potential and Wifl score.²⁵ For a new wound, it is then necessary to establish the score based on the primary presentation notation.

Limb Loss (R)

Limb loss is any major amputation defined as any resection proximal to the ankle.²³

Strengths and Limitation

This consistent notation for re-scoring tissue loss, level of ischemia, and severity of foot infection assessment during treatment aims to meet the needs of both the clinical and scientific community to enhance person-centered care and inform better practices. Moreover, it has the potential to promote better understanding, communication, and follow-up within the team but also in broader dissemination. In addition, it facilitates the use of the Wifl classification in a general way as we have established clear definitions for each phase according to the stage of clinical management. This addition does not distort the original SVS classification but can supplement the use of Wifl as needed. In the same

way, it allows longitudinal evaluation of the person with wounds in relation with treatments (ie, before and after) including the number of recurrences if necessary. This notation is flexible considering the use of a specific notation (#) while waiting for new data from investigations as an example. The “pending” result was previously ambiguous. As the initial Wifl score was intended to assess 1-year major amputation risk or benefit of revascularization, our suggestion evaluates progression rather than assessing risk or benefit. Thus, these predictive values can be considered only when assessing the Wifl score of the primary presentation (p) or at the initial assessment for a recurrence (r). Considering the variability of treatment options and the great heterogeneity in terms of dose, exposure, time, number, follow up, etc, attributable to this stage of management, a general predictive value seems unrealistic. However, it can open the way to evaluate predictive values of a specific care pathway or treatments, and this 2.0 Wifl notation can accompany efforts in this direction. Lastly, the revised scoring system we have suggested requires validation and evaluation for repeatability, which we plan to undertake. It is consistent with previous work implementing global guidelines for CLTI in clinical practice.²⁰

Conclusion

Our suggestion supports a coherent way to longitudinally communicate characteristics of a threatened limb. This has potential to support high quality interdisciplinary patient-centered care. This is one more improved way in the “toolbox” of clinicians and researchers to maximize ulcer-free, hospitalization-free, activity-rich days for people with limb-threatening conditions, in the same way the team and cancer survivors strive to maximize cancer-free days. We look forward to further efforts to validate this modification of a common language of risk.

Acknowledgements

None.

Declaration of Conflicting Interests

This study is partially supported by National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases Award Number 1R01124789-01A1.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

Ethics Approval and Consent to Participate

Not applicable.

Consent for Publication

Not applicable.

Availability of Data and Materials

Not applicable.

ORCID iD

Virginie Blanchette  <https://orcid.org/0000-0002-5901-8964>

References

1. Armstrong DG, Boulton AJ, Bus SA. Diabetic foot ulcers and their recurrence. *N Engl J Med*. 2017;376(24):2367–2375.
2. Jeffcoate WJ, Vileikyte L, Boyko EJ, Armstrong DG, Boulton AJ. Current challenges and opportunities in the prevention and management of diabetic foot ulcers. *Diabetes Care*. 2018;41(4):645–652.
3. Edmonds M, Manu C, Vas P. The current burden of diabetic foot disease. *J Clin Orthop Trauma*. 2021;17:88–93.
4. Armstrong DG, Swerdlow MA, Armstrong AA, Conte MS, Padula WV, Bus SA. Five year mortality and direct costs of care for people with diabetic foot complications are comparable to cancer. *J Foot Ankle Res*. 2020;13(1):1–4.
5. Thorud JC, Plemmons B, Buckley CJ, Shibuya N, Jupiter DC. Mortality after nontraumatic major amputation among patients with diabetes and peripheral vascular disease: a systematic review. *J Foot Ankle Surg*. 2016;55(3):591–599.
6. Fard B, Dijkstra PU, Voesten HG, Geertzen JH, Group NS. Mortality, reamputation, and preoperative comorbidities in patients undergoing dysvascular lower limb amputation. *Ann Vasc Surg*. 2020;64:228–238.
7. Huang ZH, Li SQ, Kou Y, Huang L, Yu T, Hu A. Risk factors for the recurrence of diabetic foot ulcers among diabetic patients: a meta-analysis. *Int Wound J*. 2019;16(6):1373–1382.
8. Armstrong DG, Mills JL. Toward a change in syntax in diabetic foot care: prevention equals remission. *J Am Podiatr Med Assoc*. 2013;103(2):161–162.
9. Blanchette V, Brousseau-Foley M, Cloutier L. Effect of contact with podiatry in a team approach context on diabetic foot ulcer and lower extremity amputation: systematic review and meta-analysis. *J Foot Ankle Res*. 2020;13(1):1–12.
10. Musuuza J, Sutherland BL, Kurter S, Balasubramanian P, Bartels CM, Brennan MB. A systematic review of multidisciplinary teams to reduce major amputations for patients with diabetic foot ulcers. *J Vasc Surg*. 2020Apr;71(4):1433–1446.e3.
11. Bernatchez J, Mayo A, Kayssi A. The epidemiology of lower extremity amputations, strategies for amputation prevention, and the importance of patient-centered care. *Elsevier*. 2021Mar; 34(1):54–58.
12. Khan T, Shin L, Woelfel S, Rowe V, Wilson BL, Armstrong DG. Building a scalable diabetic limb preservation program: four steps to success. *Diabet Foot Ankle*. 2018;9(1):1452513.
13. Mills Sr JL, Conte MS, Armstrong DG, et al. The society for vascular surgery lower extremity threatened limb classification system: risk stratification based on wound, ischemia, and foot infection (WIFI). *J Vasc Surg*. 2014;59(1): 220–234. e2.
14. Greene FL, Compton CC, Fritz AG, Shah JP, Winchester DP. *AJCC cancer staging atlas*. Springer; 2006.
15. Armstrong DG, Mills JL. Juggling risk to reduce amputations: the three-ring circus of infection, ischemia and tissue loss-dominant conditions. *Wound Medicine*. 2013;1:13–14.
16. Byrd DR, Brierley JD, Baker TP, Sullivan DC, Gress DM. Current and future cancer staging after neoadjuvant treatment for solid tumors. *CA Cancer J Clin*. 2021;71(2):140–148.
17. Leithead C, Spangler E, Passman M, et al. Importance of postprocedural wound, ischemia, and foot infection restaging in predicting limb salvage. *J Vasc Surg*. 2017;65(1):e7–e8.
18. Mills JL. An integrated approach for the effective management of limb-threatening ischaemia in the diabetic foot. In: *The Diabetic Foot Syndrome*. Karger Publishers; 2018: 147–160.
19. Khan T, Armstrong DG. *Ulcer-free, hospital-free and activity-rich days: Three key metrics for the diabetic foot in remission*. MA Healthcare London; 2018:S3–S4.
20. Conte MS, Mills JL, Bradbury AW, White JV. Implementing global chronic limb-threatening ischemia guidelines in clinical practice: utility of the society for vascular surgery threatened limb classification system (WIFI). *J Vasc Surg*. 2020;72(4): 1451–1452.
21. FDA Wound Healing Clinical Focus Group. Guidance for industry: chronic cutaneous ulcer and burn wounds – developing products for treatments. *Wound Repair Regen* 2001;9(4):258–268.
22. Gould L, Li WW. Defining complete wound closure: closing the gap in clinical trials and practice. *Wound Repair Regen*. 2019;27(3):201–224.
23. van Netten JJ, Bus SA, Apelqvist J, et al. Definitions and criteria for diabetic foot disease. *Diabetes Metab Res Rev*. 2020;36:e3268.
24. Association WM. World medical association declaration of Helsinki. Ethical principles for medical research involving human subjects. *Bull W H O*. 2001;79(4):373.
25. Weaver ML, Hicks CW, Canner JK, et al. The society for vascular surgery wound, ischemia, and foot infection (WIFI) classification system predicts wound healing better than direct angiosome perfusion in diabetic foot wounds. *J Vasc Surg*. 2018;68(5):1473–1481.