

Rehabilitation journal editors recognize the need for interventions targeted to improve the completeness of reporting, but there is heterogeneity in terms of strategies actually adopted: A cross-sectional web-based survey

Reporting guidelines (RGs) have been developed by researchers, methodologists, and journal editors to enhance the quality and transparency of scientific research reporting¹ and to facilitate readers' assessment of the internal and external validity of the studies. In 2014, a joint editorial² of 28 rehabilitation journals highlighted the need to use RGs to ensure the quality of reporting rehabilitation research, stating "that simultaneous implementation of this new reporting requirement will send a strong message to all disability and rehabilitation researchers about the need to adhere to the highest standards when performing and disseminating research."

While there is encouraging evidence that RGs improve reporting quality,³ there are also challenges to implementation and dissemination. Rehabilitation is a growing field with increasing attention to the quality of reporting. However, our previous studies show that in rehabilitation journals authors do not frequently declare the use of RGs,⁴ and the completeness of reporting of randomized controlled trials⁵ and systematic reviews⁶ is suboptimal.

Little is known about barriers and facilitators that affect the use of RGs and whether editors pay attention to RG use in the field of rehabilitation. Evaluating the adherence to RGs and their potential barriers (e.g., editors and peer-reviewers do not consider RGs-use by the) has been done in other biomedical fields, but not in rehabilitation.⁷ Therefore, we aim to investigate the perceived value, the potential impact of RG use, and the strategies that journals adopt to encourage RG use in rehabilitation. More specifically, we aimed at:

- Exploring editors' opinions and beliefs about the importance of RGs
- Mapping which methods journals use to check the use of RGs (e.g., use of automatized electronic systems, providing training to editors and peer-reviewers about the importance of the reporting).
- Exploring editors' thoughts and opinions about the importance of RGs and their use during the peer-review process.

The study protocol and the full version of the survey are available on the OSF Repository.⁸ We followed the guidance by Gaur et al.⁹

for the reporting of this manuscript. We conducted a web-based survey among all editorial board members (i.e., editors-in-chief, senior editors, and associate editors) of the 68 journals indexed under the "rehabilitation" category in InCites Journal Citation Report. This study complies with the Code of Ethics for Research Involving Human Participants Faculty of Science (BETHCIE), Vrije Universiteit Amsterdam; therefore, a formal ethics review is not required.¹⁰ All participants were explicitly informed that participation in the survey was anonymous and voluntary. A copy of informed consent is available in the OSF repository.⁸ Only the participants that explicitly gave their consent to be acknowledged were mentioned in the acknowledgment section. Further information on ethics and privacy are available in the published protocol.⁸

Several meetings were held among the study authors, including one editor-in-chief of a rehabilitation journal (RZP) and two members of editorial boards of high-impact rehabilitation journals (RO, AV) to discuss the survey. One investigator (TI) developed a preliminary version, which was then further reviewed and discussed to ensure that the questions adequately met the study aims (confirming face validity). Next, the clarity of the questions with appropriate response items was checked in a pilot test among the study authors (RZP, AV, AC, RO, SS, SG), and any duplications or coinciding themes were removed. The initial questionnaire was revised three times and the final questionnaire included closed, semiclosed and open-ended questions. First, journal editors were identified by visiting the editorial board web pages of the 68 journals and by searching the publicly available academic email addresses of these authors on the internet. Second, we directly asked the journals for those emails not found through this process. Third, the developed survey was sent to all available email addresses. The online survey was open from May 15 to August 1, 2022. Follow-up reminder emails were sent every 4 weeks. Only completed responses were included in the analysis. The study protocol⁸ describes further details.

We performed a descriptive analysis of the responses. Percentages were provided for binary data and count data. For Likert scale

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *Journal of Evidence-Based Medicine* published by Chinese Cochrane Center, West China Hospital of Sichuan University and John Wiley & Sons Australia, Ltd.

TABLE 1 Main results

| Knowledge, beliefs and attitudes among the editors | | |
|---|--------------------|------------------------------|
| Awareness of the existence of reporting guidelines before receiving the survey | % (n = 142) | number of respondents |
| No | 4.2% | 6 |
| Yes | 95.8% | 136 |
| Awareness of the existence of separate reporting guidelines for different study design | | |
| No | 2.8% | 4 |
| Yes | 97.2% | 138 |
| Awareness of the existence of EQUATOR Network | | |
| No | 33.8% | 48 |
| Yes | 66.2% | 94 |
| Awareness about specific reporting guidelines | | |
| PRISMA (systematic reviews and meta-analyses) | 76.8% | 109 |
| CONSORT (randomized controlled trials) | 67.6% | 96 |
| STROBE (observational studies—cohort, case-control, and cross-sectional) | 56.3% | 80 |
| COREQ (qualitative research) | 35.2% | 50 |
| STARD (diagnostic accuracy studies) | 22.5% | 32 |
| MOOSE (meta-analyses of observational studies in epidemiology) | 14.1% | 20 |
| All of the above | 21.8% | 31 |
| None | 0.00% | 0 |
| Where the respondents learned about reporting guidelines | | |
| Doing research in which a reporting guideline is needed | 70.4% | 100 |
| Attending general courses on research methodology | 35.9% | 51 |
| Attending specific courses targeted to the editors | 11.3% | 16 |
| Other | 23.9% | 34 |
| When the respondents think a reporting guideline should be used | | |
| During all stages of a research project | 70.4% | 100 |
| When designing a study | 47.9% | 68 |
| When writing up a study | 42.3% | 60 |
| When required by a funding body | 12.7% | 18 |
| Only when required by a journal | 1.4% | 2 |
| Other | 2.1% | 3 |
| How important is that authors follow reporting guidelines for improving the completeness of research reporting | | |
| 1—not very important/useful | 1.4% | 2 |
| 2 | 0.7% | 1 |
| 3 | 12.0% | 17 |
| 4 | 26.8% | 38 |
| 5—very important/useful | 58.5% | 83 |
| Don't know | 0.7% | 1 |
| How important is to have a mechanism for checking the appropriateness of a reporting guideline submitted | | |
| 1—not very important/useful | 0.7% | 1 |
| 2 | 1.4% | 2 |
| 3 | 11.3% | 16 |
| 4 | 27.5% | 39 |
| 5—very important/useful | 57.7% | 82 |
| Don't know | 1.4% | 2 |

(Continues)

TABLE 1 (Continued)

| Knowledge, beliefs and attitudes among the editors | | |
|--|-------|-----|
| How important do you think reporting guidelines are to peer-reviewers when evaluating a research report | | |
| 1—not very important/useful | 0.7% | 1 |
| 2 | 5.6% | 8 |
| 3 | 19.0% | 27 |
| 4 | 34.5% | 49 |
| 5—very important/useful | 39.4% | 56 |
| Don't know | 0.7% | 1 |
| Policy and routine activity of the editors and journals | | |
| Instruction for authors containing an explicit statement about reporting guidelines | | |
| Yes | 80.3% | 114 |
| No | 12.0% | 17 |
| Don't know | 7.7% | 11 |
| Explicit request about the mandatory upload of the reporting guideline during the submission process | | |
| Yes | 57.7% | 82 |
| No | 34.5% | 49 |
| Don't know | 7.7% | 11 |
| What the journals do when a manuscript satisfies the editorial criteria but the authors do not declare to follow the relevant reporting guideline | | |
| Ask the authors to resubmit the study alongside the appropriate reporting guideline | 49.3% | 70 |
| Send such studies out for peer-review without asking the authors to resubmit the appropriate reporting guideline | 32.4% | 46 |
| Other | 18.3% | 26 |
| How often the editors check for the correct use of the reporting guidelines | | |
| Sometimes | 35.2% | 50 |
| Always | 32.4% | 46 |
| I believe it is the responsibility of the peer-reviewers | 9.9% | 14 |
| No check | 9.1% | 13 |
| Often | 4.9% | 7 |
| Other | 8.4% | 12 |
| How the editors check for the correct use of the reporting guidelines | | |
| By consulting the specific reporting guideline for each study | 70.4% | 100 |
| Filling a generic sheet provided by your journal | 13.4% | 19 |
| Other | 16.2% | 23 |
| Journals that explicitly ask the peer-reviewers to check for the use of reporting guidelines in the manuscripts reviewed | | |
| No | 62.0% | 88 |
| Yes | 21.8% | 31 |
| Don't know | 16.2% | 23 |
| Barriers for editors and peer-reviewers to endorse and adhere to reporting guidelines | | |
| Lack of time for looking at the completeness of the reporting | 54.9% | 78 |
| Lack of knowledge about the importance of the reporting guideline | 44.4% | 63 |
| Lack of knowledge about the existence of the reporting guideline | 38.0% | 54 |
| Don't know | 15.5% | 22 |
| Other | 12.7% | 18 |
| The use of artificial Intelligence or other automated tools to evaluate the correct use of the reporting guidelines | | |
| No | 76.8% | 109 |
| Yes | 0.7% | 1 |
| Don't know | 22.5% | 32 |

data, medians with interquartile ranges were reported. For free-text responses, two authors (TI and SG) undertook independent qualitative analyses to identify categories and themes following a thematic analysis approach.¹¹ A consensus discussion between the two authors was used to resolve any discrepancies. Preplanned subgroup analyses were conducted depending on the role of an editor in a journal (e.g., editor-in-chief, senior editor, or associate editor) and on the journal impact factor quartile. Data were analyzed and graphs were drawn using Qualtrics, Provo, UT.

From the 507 email addresses collected, 28 produced an email failure message after two attempts. Of a total of 479, we received 154 responses. Twelve responses were identified from Qualtrics Software as partial responses, which were removed from the sample, meaning 142 complete responses (29.6%) were available for analyses. Table S1 reports detailed characteristics of the sample.

Almost all participants stated they were aware of what RGs are ($n = 136$; 95.8%), and of the existence of separate RGs for different study designs ($n = 138$; 97.2%). Approximately one-third of participants (33.8%) were not aware of the Enhancing the Quality and Transparency Of health Research (EQUATOR) Network.¹²

When we asked about the importance and usefulness of the use of RGs, most of the editors ($n = 83$; 58.5%) believed that it is “very important” that authors follow the RGs to ensure the completeness of reporting (median rating = 5; interquartile range—IQR = 5–4).

Regarding policy and routine activity of the editors and journals, most of the journals ($n = 114$; 80.3%) referred to the RGs in the instructions for the authors while 82 (57.7%) required mandatory submission of the RG by the authors during the submission. The lack of time ($n = 78$; 54.9%) was the most common barrier to the use of RGs. Summary results are reported in Table 1.

Comments and suggestions to improve the use of RGs given by participants in response to the open question were grouped into themes (Table S2). The most suggested theme is about actions targeted to the publishers, journals, or editorial process, and making the use of RGs mandatory during submission ($n = 40$; 28.2%) and the need for a standard approach among all rehabilitation journals ($n = 11$; 7.7%).

Table S3 reports the subgroup analyses.

In summary, we found that almost all editors of rehabilitation journals are aware of the importance of the RGs and recognize the need for interventions to improve the completeness of reporting. However, our study revealed a great heterogeneity in terms of what the journals and editors do to endorse the use of RGs. Only approximately half (49%) of the editors ask to resubmit the study alongside the appropriate RG when the authors do not declare to follow any RG in their manuscripts, and most of the participants (60%) do not ask the peer-reviewers to check the RG use.

The most common perceived barrier to endorsing and adhering to RGs is the lack of time. This corroborates with previous studies,^{7,13} that concluded that too many responsibilities fall on the shoulders of busy unpaid peer-reviewers who may not be fully equipped to carry out the role.

This study allowed us to identify several potential actions that could promote the widespread adoption of RGs (Figure S1), where publishers, journals, editors, peer-reviewers, and authors play an (active) important role. We encourage publishers and journals to adopt a standard approach to increase the attention paid to the reporting issues during the editorial and peer-review screening processes, providing explicit instruction to authors and peer-reviewers (e.g., by placing active links to the relevant RGs in the manuscript submission systems), and promoting the dialogue between editors and reviewers. Even if journals, editors, and peer-reviewers are gatekeepers to scientific publication, we firmly believe that primarily authors are responsible for complete reporting, in the same fashion as they should ensure the integrity of study design, data analysis and statistical methods, and avoiding research misconduct.

Our study has limitations. The response rate was low (29.6%). However, other surveys¹⁴ in the same field shows an even lower response rate, probably due to a gradual decrease in survey participation¹⁵ among health professionals. Moreover, participants in this survey come from journals pertaining to all quartiles, so it is reasonable to believe that it is a representative sample of the rehabilitation category.

In conclusion, rehabilitation journal editors are generally aware of the existence of RGs and recognize the need to promote the use of RGs to improve the quality and completeness of reporting. However, there is great heterogeneity in terms of what the journals and editors do to endorse the use of RGs. Moreover, we hope that journals will adopt strategies to increase the attention paid to the reporting issues during the editorial and peer-review screening processes, providing explicit instruction to peer-reviewers and promoting the dialogue between authors, editors, and reviewers.

ACKNOWLEDGMENTS

We would like to express a huge thanks to all the participants for the time they spent filling this survey. Since the survey was totally anonymous, we asked the participants to express their willingness to be explicitly acknowledged. The following gave their consent to be mentioned: Dr Catherine Backman, Dr Ronaldo Valdir Briani, Dr Isabelle Gagnon, Dr Cesar Hincapié, Dr Stefano Negrini, Dr Bruno Saragiotto, Dr Saurab Sharma, Dr Gavin Williams, Dr Rafael Zegarra-Parodi.

CONFLICT OF INTEREST STATEMENT

RO, AV, and RZP are on the editorial board of The Journal of Physiotherapy, The Journal of Orthopaedic & Sports Physical Therapy, Musculoskeletal Science & Practice or The Brazilian Journal of Physical Therapy, respectively.

Tiziano Innocenti^{1,2} 
 Raymond Ostelo^{1,3}
 Arianne Verhagen⁴
 Rafael Zambelli Pinto⁵
 Stefano Salvioli^{2,6}
 Silvia Giagio^{7,8}
 Alessandro Chiarotto^{1,9}

¹Faculty of Science, Department of Health Sciences, Vrije Universiteit Amsterdam, Movement Sciences, Amsterdam, Netherlands

²GIMBE Foundation, Bologna, Italy

³Department of Epidemiology and Data Science, Amsterdam UMC, Movement Sciences Research Institute, Amsterdam, Netherlands

⁴Discipline of Physiotherapy, Graduate School of Health, University of Technology Sydney, Sydney, Australia

⁵Department of Physical Therapy, Universidade Federal de Minas Gerais (UFMG), Belo Horizonte, Brazil

⁶Department of Neuroscience, Rehabilitation, Ophthalmology, Genetics, Maternal and Child Health, University of Genoa, Genoa, Italy

⁷Division of Occupational Medicine, IRCCS Azienda Ospedaliero-Universitaria di Bologna, Bologna, Italy

⁸Department of Biomedical and Neuromotor Sciences (DIBINEM), Alma Mater Studiorum, University of Bologna, Bologna, Italy

⁹Department of General Practice, Erasmus MC, University Medical Center, Rotterdam, Netherlands

Correspondence

Tiziano Innocenti, Faculty of Science, Department of Health Sciences, Vrije Universiteit Amsterdam, Amsterdam Movement Sciences, The Netherlands.

Email: t.innocenti@vu.nl

ORCID

Tiziano Innocenti  <https://orcid.org/0000-0003-4696-6989>

REFERENCES

1. Simera I, Moher D, Hoey J, Schulz KF, Altman DG. A catalogue of reporting guidelines for health research. *Eur J Clin Invest*. 2010; 40(1):35–53.
2. Chan L, Heinemann AW, Roberts J. Elevating the quality of disability and rehabilitation research: mandatory use of the reporting guidelines. *Am J Occup Ther*. 2014;68(2), 127–129.
3. O'leary JD, Crawford MW. Review article: reporting guidelines in the biomedical literature. *Canad J Anaesth*. 2013;60(8):813–821.
4. Innocenti T, Salvioli S, Giagio S, Feller D, Cartabellotta N, Chiarotto A. Declaration of use and appropriate use of reporting guidelines in high-impact rehabilitation journals is limited: a meta-research study. *J Clin Epidemiol*. 2021;131:43–50.
5. Innocenti T, Giagio S, Salvioli S, et al. The completeness of reporting is suboptimal in randomized controlled trials published in rehabilitation journals, with trials with low risk of bias displaying better reporting: a meta-research study. *Arch Phys Med Rehabil*. 2022;103(9):1839–1847.
6. Innocenti T, Feller D, Giagio S, et al. Adherence to the PRISMA statement and its association with risk of bias in systematic reviews published in rehabilitation journals: a meta-research study. *Braz J Phys Ther*. 2022;26(5):100450.
7. Prager R, Gagnon L, Bowdridge J, et al. Barriers to reporting guideline adherence in point-of-care ultrasound research: a cross-sectional survey of authors and journal editors. *BMJ Evid Based Med*. 2021. Online ahead of print.
8. Innocenti T, Ostelo R, Verhagen A, et al. What are barriers and facilitators in endorsing reporting guidelines in rehabilitation journals? Protocol for a Cross-Sectional Survey of Journal Editors. OSF. 2022.
9. Gaur PS, Zimba O, Agarwal V, Gupta L. Reporting survey based studies—a primer for authors. *J Korean Med Sci*. 2020;35(45):e398.
10. Research ethics review committee: Vrije Universiteit Amsterdam. Available from: <https://vu.nl/en/about-vu/more-about/research-ethics-review-committee-beta>
11. Bradley EH, Curry LA, Devers KJ. Qualitative data analysis for health services research: developing taxonomy, themes, and theory. *Health Serv Res*. 2007;42(4):1758–1772.
12. Altman DG, Simera I, Hoey J, Moher D, Schulz K. EQUATOR: reporting guidelines for health research. *Lancet*. 2008;371(9619):1149–1150.
13. Hirst A, Altman DG. Are peer reviewers encouraged to use reporting guidelines? A survey of 116 health research journals. *PLoS One*. 2012;7(4):e35621.
14. Price A, Schroter S, Clarke M, Mcaneney H. Role of supplementary material in biomedical journal articles: surveys of authors, reviewers and readers. *BMJ Open*. 2018;8(9):e021753.
15. Brtnikova M, Crane LA, Allison MA, Hurley LP, Beaty BL, Kempe A. A method for achieving high response rates in national surveys of U.S. primary care physicians. *PLoS One*. 2018;13(8):e0202755.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.