



RESEARCH ARTICLE

REVISED General public awareness, knowledge and attitude toward COVID-19 infection and prevention: a cross-sectional study from Pakistan [version 2; peer review: 2 approved]

Beenish Fatima Alam¹, Abdullah A. Almojaibel ², Khalid Aziz Ansari ²,
 Mohammad Haroon³, Sara Noreen³, Saman Tauqir⁴, Khalid Almas⁵,
 Faraz A. Farooqi ⁶, Saqib Ali ⁷

¹Department of Oral Biology, Bahria University Medical and Dental College, Karachi, Pakistan

²Respiratory Care Department, College of Applied Medical Sciences, Imam Abdulrahman Bin Faisal University, Dammam, Saudi Arabia

³Department of Medicine, Khyber Teaching Hospital, Peshawar, Pakistan

⁴Department of Physiology, Kabir Medical College, Gandhara University, Peshawar, Pakistan

⁵Department of Preventive Dental Sciences, College of Dentistry, Imam Abdulrahman Bin Faisal University, Dammam, Saudi Arabia

⁶Department of Dental Education, College of Dentistry, Imam Abdulrahman Bin Faisal University, Dammam, Saudi Arabia

⁷Department of Biomedical Dental Sciences, College of Dentistry, Imam Abdulrahman Bin Faisal University, Dammam, Saudi Arabia

V2 First published: 21 Sep 2021, 10:946
<https://doi.org/10.12688/f1000research.52692.1>

Latest published: 05 Jun 2023, 10:946
<https://doi.org/10.12688/f1000research.52692.2>

Abstract

Background: The aim of this study is to evaluate the knowledge, perceptions, and attitude of the public in Pakistan (using social media) towards COVID-19.

Methods: A cross-sectional study was conducted amongst 1120 individuals nationwide. A self-developed, pre-tested questionnaire was used that comprised of sections covering demographic characteristics, medical history, hygiene awareness, COVID-19-related knowledge, and learning attitude. Descriptive statistics were used for frequencies, percentages, averages and standard deviations. Inferential statistics were done using the Student's t-test and ANOVA.

Results: The average age of participants was 31 years (range 18-60 years). In total 56 individuals (5%) had completed primary or secondary school education; 448 (40%) were employed (working from home) and 60% were jobless due to the COVID-19 crisis. Almost all the study subjects (1030 (92%)) were washing their hands multiple times a day. A total of 83% had awareness regarding quarantine time, 82% used face masks whenever they left their homes, 98% were aware of the origin of the disease, and 70% had knowledge regarding the most common symptoms of COVID-19.

Conclusion: It can be concluded from the current study that female participants had higher level of education, and more awareness regarding the coronavirus. The majority of the participants followed proper hand washing regimes and washed their faces. Further

Open Peer Review

Approval Status

	1	2
version 2 (revision) 05 Jun 2023	 view	
	↑	
version 1 21 Sep 2021	? view	 view

1. **Mohmed Isaqali Karobari** , Universiti Sains Malaysia, Kota Bharu, Malaysia
 Saveetha University, Chennai, India
 University of Puthisastra, Phnom Penh, Cambodia

2. **Anand Marya** , University of Puthisastra, Phnom Penh, Cambodia
 Saveetha Univeristy, Chennai, India

Any reports and responses or comments on the article can be found at the end of the article.


knowledge and awareness should be promoted.

Keywords

COVID-19, Awareness, Knowledge, Perception, Pakistan, Pandemic



This article is included in the **Emerging Diseases and Outbreaks** gateway.



This article is included in the **Coronavirus** collection.

Corresponding author: Faraz A. Farooqi (fafarooqi@iau.edu.sa)

Author roles: **Alam BF:** Conceptualization, Investigation, Writing – Original Draft Preparation, Writing – Review & Editing; **Almojaibel AA:** Writing – Review & Editing; **Aziz Ansari K:** Conceptualization, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Haroon M:** Data Curation, Writing – Original Draft Preparation, Writing – Review & Editing; **Noreen S:** Conceptualization, Data Curation, Writing – Original Draft Preparation, Writing – Review & Editing; **Tauqir S:** Writing – Review & Editing; **Almas K:** Conceptualization, Investigation, Writing – Original Draft Preparation, Writing – Review & Editing; **Farooqi FA:** Formal Analysis, Resources, Writing – Original Draft Preparation, Writing – Review & Editing; **Ali S:** Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2023 Alam BF *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Alam BF, Almojaibel AA, Aziz Ansari K *et al.* **General public awareness, knowledge and attitude toward COVID-19 infection and prevention: a cross-sectional study from Pakistan [version 2; peer review: 2 approved]** F1000Research 2023, 10:946 <https://doi.org/10.12688/f1000research.52692.2>

First published: 21 Sep 2021, 10:946 <https://doi.org/10.12688/f1000research.52692.1>

REVISED Amendments from Version 1

In this 2nd version of the manuscript, the authors have made some improvements suggested by the reviewers. Starting with the abstract where the background of the study was improved with grammar, in the results section the p-values were added along with more text from actual results. In the methods section of the manuscript, the nature of the participants was added as suggested. 1st line of results was revised, in the 3rd paragraph the score was corrected and the following paragraph was revised for the application of numbers from the table as suggested. The discussion was improved with significant differences instead of results only as suggested. Further recommendations along with the limitations were also added and separated in a new heading. Suggested Articles by the reviewer were cited in the discussion.

Any further responses from the reviewers can be found at the end of the article

Introduction

The first case of COVID-19 was identified within the Wuhan city of China in December 2019.¹ By February, a tremendous increase in the number of cases and number of deaths due to Covid-19 started being reported. Due to the rapid spread of infection from China to different countries, and with the number of cases exceeding the cases reported by the Chinese government, COVID-19 was declared a global pandemic by the World Health Organization in March 2020.^{2,3} This epidemic has now affected more than 200 different countries across the globe.⁴

COVID-19 is a highly contagious infection that spreads through human to human contact causing serious health problems within communities.⁵ Currently very little scientific information is available regarding this novel virus. However, it has been identified as an enveloped RNA virus that is further categorised into alpha, beta, gamma and delta.⁶ Depending upon the immune system of an individual it can cause symptoms ranging from mild to severe. Symptoms identified comprise of fever, cough, difficulty in breathing, loss of smell, while in extreme cases can lead to pneumonia, multi-organ failure and death.⁷ The rapid spread of this infection is principally through respiratory droplets of 5-10 μm in diameter, spread through the mouth or nose, when an infected person coughs or sneezes.⁸ A study conducted in Singapore has identified that these droplets can be transmitted across a distance of 4.5 metres.^{9,10} Moreover symptoms can be identified within a day or two after acquiring infection, extending up to 14 days.¹¹

The first positive case of COVID-19 in Pakistan was identified on 26th February 2020. In order to control the spread of the disease, on 23rd March 2020 a complete lockdown was imposed throughout the country. This included closure of all the educational organizations that comprised of schools, colleges, and universities along with religious schools.¹² All ceremonies and religious gatherings in any form were adjourned to prevent the spread of infection to the general public.¹³ It was highlighted that elderly people and individuals having systemic disease, such as heart disease, diabetes, cancer and respiratory disease, are at high risk of acquiring this infection, while children appear to be less susceptible towards this disease. Proper protocol must be followed to prevent the spread of this disease.¹⁴

Currently there is a secondary wave or surge reported from many parts of the world; therefore it is of great importance for the general public to have appropriate awareness regarding the signs and symptoms and the causative factors of this disease and preventive protocols that need to be followed, including social distancing, following proper hand hygiene protocol and use of face masks. As Pakistan is a developing country with limited resources, the country has introduced different strategies to increase the awareness of the general public. Therefore, this study was undertaken to analyse the level of awareness, knowledge and practices of the public of Pakistan using an online survey regarding COVID-19 and various measures to be undertaken to prevent the spread of this disease.

Methods**Study design**

This cross-sectional study was conducted including the general public of Pakistan using an online questionnaire. The questionnaire was sent to the general public through various social platforms (Whatsapp, Twitter, Emails, Facebook messengers).

Participants

Individuals of both genders, within the age group of 18 to 60 years of age, of all education levels were included in this survey. Informed consent was taken from all the participants before asking them to complete the survey. Responses were kept anonymous to maintain the confidentiality of the participants. Five to seven minutes were required to complete the survey.

Data collection

Due to complete closure of public places, an online survey was conducted using Google forms. Data was collected from May through to June 2020.

The questionnaire used for this study was developed by the authors and adapted from Ali et al.³⁰ Before collecting the data, pilot testing of the questionnaire was carried out among 20 individuals (selected from social media groups within the community) to test the reliability and validity of the questionnaire. Following their responses, necessary corrections (simplification of terms) were carried out.

The questionnaire utilised for this study was divided into multiple sections as follows: demographic characteristics (gender, education level, income earned, job status); section related to health (presence of any systemic disease and smoking habits); hygiene awareness (frequency of washing hands and face, type of soap used, duration of hand washing, frequency of touching face); COVID-19 related knowledge (origin, symptoms and transmission route for COVID-19); measures taken for prevention against COVID-19 (knowledge regarding usage, indications and different types of face masks); and lastly learning attitudes regarding COVID-19. There were 7 main questions covering COVID-19 knowledge. Three of which had multiple correct answers, each correct answer was given 1 point and wrong answer counted as 0. Total maximum knowledge score was 16.

Statistical analysis

SPSS version 22 was used for data analysis. Descriptive statistics including frequencies, percentages, means and standard deviations were used to determine demographical information, health history and knowledge related questions. Student's t-test and ANOVA were used for comparing knowledge score with demographical information and awareness related responses. Multiple comparisons were done between demographics of participants in case of significant relation found. P-value <0.05 was considered as statistically significant. Univariate and multivariate regression analysis was used to investigate the factor associated with knowledge score. P-value 0.10 was considered as cut off for significant value for factors in univariate analysis.

Ethics approval and consent to participate

Ethical permission was obtained from the Bahria Medical and Dental College Ethics Review committee to conduct the study (reference number ERC 55/2020). A participation consent statement was added before the survey as follows: "Your participation in this study is completely voluntary. There are no foreseeable risks associated with this project. However, if you feel uncomfortable answering any questions, you can withdraw from the survey at any point". Participation in the survey was therefore taken as consent to participate.

Results

A total of 1,120 participants completed the questionnaire including 46% females and 54% males with an average age of 31 years. Most of the study participants were males. A total of 448 (40%) individuals were employed (working from home), while more than half (60%) were jobless due to the COVID-19 crises. Most of the participants (37%) belonged to the lower income group ranging from 10,000 to 30,000 PKR (Table 1).

A total of 862 (77%) of the participants were free from any allergy or respiratory diseases, with only 10% suffering from seasonal allergies. Similarly, 840 (75%) participants expressed themselves as fit and medically healthy.

Table 1. Demographic characteristics of the participants.

Demographic characteristics	n	%
Gender		
Male	604	54
Female	516	46
Age in years		
15-30	482	43.0
30-45	582	52.0
45-60	56	5.0

Table 1. *Continued*

Demographic characteristics	n	%
Education level		
Primary/Secondary	56	5
College Graduate	470	42
Postgraduate	526	47
Doctorate	68	6
Working status		
Yes (working from home)	448	40
No	672	60
Monthly income		
Less than 30K	414.4	37
30K to 50K	403.2	36
More than 50K	302.4	27
Smoking status		
Yes	145.6	13
No	862.4	77
Occasional smoker	78.4	7
Past smoker	33.6	3
Health background		
<i>Respiratory disease or any allergy</i>		
Yes	145.6	13
No	862.4	77
Seasonal/medicinal allergy	112	10
<i>Underlying medical condition/systemic diseases</i>		
Fit and healthy	840	75
Heart disease	11.2	1
Hypertension	67.2	6
Arthritis	22.4	2
Any other disease	179.2	16

A significant number of participants (70%) with score of 9.48 ± 2.88 out of 16, claimed that they had a good knowledge about the disease. [Figure 1](#) shows the correct answer percentages of knowledge questions asked.

Comparisons of mean knowledge score with demographic characteristics are shown in [Table 2](#). There was no significant difference between male and female participant's knowledge score ($p = 0.86$). Age showed significant variation ($p = 0.040$) with a higher mean score of knowledge obtained in the 30-45 age group and lower mean score obtained from younger individuals.

Postgraduate participants showed highest mean score, although comparison with non-postgraduate groups was not significant (p -value 0.137). Those working from home have significantly higher mean score (9.78 ± 2.8) as compared to those who were not employed (8.02 ± 1.8) ($p = 0.029$). Participants with seasonal or medicinal allergies showed higher knowledge score (10.44 ± 2.8) than disease free participants, but the difference was not statistically significant ($p = 0.32$). Similarly, arthritis patients have higher knowledge of COVID-19 as compared to others but not significant (12 ± 3.66). The most stated source of getting information regarding COVID-19 was social media (89%), the television and radio (56%), followed by family friends (34%) and print media (newspapers; 30%) ([Figure 2](#)).

[Table 3](#) reveals the hand hygiene awareness of the participants. Almost all of the participants (1030; 92%) were washing their hands multiple times a day. Face washing as a part of Wadu (ablution before prayers) was observed by individuals.




Figure 1. Percentage of correct answers regarding COVID-19 knowledge.

Table 2. Comparisons of mean knowledge score with demographic characteristics.

Demographic characteristics	Mean	SD	p-value
Gender			
Male	9.45	2.834	0.86
Female	9.51	2.958	
Age in years			
15-30	9.04	2.881	0.04
30-45	10.01	2.646	
45-60	9.77	3.320	
Education level			
Primary/Secondary	9.75	2.8	0.137
College graduate	9.06	2.91	
Postgraduate	10.5	2.88	
Doctorate	8.7	2.71	
Working status			
Yes (working from home)	9.78	2.879	0.029
No	8.02	2.850	
Monthly income			
Less than 30K	9.32	2.83	0.627
30K to 50K	9.44	2.6	
More than 50K	10.1	3.62	

Table 2. *Continued*

Demographic characteristics	Mean	SD	p-value
Smoking status			
Yes	9.7	2.90	0.318
No	9.4	2.86	
Occasional smoker	9.75	3.07	
Past smoker	11.30	2.53	
Health background			
<i>Respiratory disease or any allergy</i>			
Yes	9.74	2.944	0.32
No	9.35	2.868	
Seasonal/medicinal allergy	10.44	2.80	
<i>Underlying medical condition/systemic diseases</i>			
Fit and healthy	9.42	2.678	0.203
Heart disease	9.00	0.000	
Hypertension	8.75	3.130	
Arthritis	12.00	3.633	
Any other disease	9.67	3.541	




Figure 2. Participants source of information on COVID-19.

Table 3. Awareness level of participants regarding maintenance of personal hygiene.

	n	%
No. of times you wash your hands		
Once a day	12	1.0
Twice a day	34	3.0
Multiple times a day	1030	92.0
Without soap (Wudu)	44	4.0

Table 3. *Continued*

	n	%
No. of times you wash your face		
Once a day	68	6
Twice a day	112	10
Multiple times a day	436	39
Five times Wudu (Wuzu)	504	45
Kind of soap used for washing		
Plain soap	426	38
Antiseptic solution	414	37
Antibacterial soap	12	1
Cloth washing soap	12	1
Liquid soap	258	23
Duration of washing		
10 Sec	134	12
10-20 Sec	258	23
More than 20 Sec	68	6
1 minute	4	0.4
Aware of hand washing technique		
Yes	660	59
No	22	2
May Be	34	3

Plain soap and antiseptic solution was found as most popular type of soap used for washing hands. Around 59% were aware regarding the hand washing technique.

Those who wash their hands twice a day showed significantly higher knowledge scores ($p = 0.0001$) as shown in [Table 4](#). Those who washed their face multiple times also have significantly higher knowledge of COVID-19 ($p = 0.005$).

Table 4. Knowledge score with hygiene awareness.

	Mean	SD+-	p-value
No. of times you wash your hands			
Once a day	9.00	3.606	0.0001
Twice a day	11.00	2.872	
Multiple times a day	9.53	2.812	
Without soap (Wudu)	8.56	2.128	
No. of times you wash your face			
Once a day	9.73	2.886	0.005
Twice a day	9.34	2.380	
Multiple times a day	11.14	2.795	
Five times Wudu (Wuzu) (Ablution)	9.42	2.752	
Kind of soap used for washing			
Plain soap	9.72	2.773	0.001
Antiseptic solution	9.38	2.812	
Antibacterial soap	10.50	3.416	
Liquid soap	9.44	2.856	

Table 4. *Continued*

	Mean	SD+-	p-value
Duration of washing			
10 sec	8.79	2.750	0.369
10-20 sec	9.79	3.234	
More than 20 sec	9.78	3.040	
1 minute	7.00	0.000	
Aware of hand washing technique			
Yes	9.85	2.910	0.286
No	8.67	2.160	
May be	8.56	3.468	

Duration of hand washing and awareness of the technique did not show any significant relation with knowledge score ($p = 0.369$ and $p = 0.286$, respectively).

In total 45% of the participants were afraid of COVID-19 and 47% would like to have more information about the virus and its development. The majority of participants (65%) agreed that the virus is deadly and is life threatening. Only 196 (18%) of the participants had a friend or relative infected with the disease (Table 5).

Multiple liner regression showed that male gender (vs female, $\beta - 0.319$) and respondents earning between 30K to 50K (vs 10K-30K, $\beta - 0.184$) presented low knowledge score, whereas 30-45 year olds (vs 15-30 year, $\beta - 1.148$, $P = 0.003$) presented significantly highest knowledge scores. Age increases and knowledge score decreases $\beta - 1.076$ (Table 6).

Table 5. Frequency of the learning attitude of the individuals.

	n	%
Like to have more information about COVID-19		
Yes	532	48
No	480	43
Don't know	96	9
Tired of listening about COVID-19		
Yes	700	63
No	100	9
I don't know	60	5
Absolutely	160	14
Not Sure	84	8
COVID-19 is a real-life threatening disease		
Agree	724	65
Not agree	76	7
I don't know	64	6
May be	240	21
your friend or relative had COVID-19		
Yes	196	18
No	872	78
Don't know	40	4

Table 6. Univariate and multivariate regression.

Variables	Partial regression coefficients		Standardized coefficients	t	Sig.
	B	Std. Error	Beta		
(Constant)	8.451	0.496		17.024	0.000*
Gender (male vs Female)	-0.319	0.358	-0.055	-0.892	0.373
Job (yes vs No)	0.825	0.444	0.140	1.858	0.064
Income (30K to 50K vs 10K-30K)	-0.184	0.458	-0.028	-0.402	0.688
Income (More than 50K vs 15K-30K)	0.395	0.475	0.066	0.832	0.406
Age (30-45 vs 15-30 years)	1.148	0.377	0.199	3.044	0.003*
Age (45 and above vs 15-30 years)	1.076	0.831	0.081	1.295	0.196

*Significant at 0.05, $F = 2.575$, $p = 0.019$, $R^2 = 0.054$, R^2 adjusted showed that regression had 3% predictive accuracy.

Discussion

To the best of author's knowledge this is among the first studies evaluating the awareness, knowledge and preventive practice of the public in Pakistan during the pandemic. Overall, 70% of the public studied gave the correct answers related to knowledge of COVID-19. These results are in agreement with a study conducted in India where a knowledge rate of 74% has been reported,¹⁵ while it is comparatively less when compared with study conducted in China (90%).¹⁶ However these findings are higher when compared to the response received by a study conducted in Jordan.¹⁷ The reason for such a high response rate within China may be due to difference in time and circumstance in which the study was conducted. On the contrary, in the current study the high response can be attributed to the campaigns related to COVID-19 awareness initiated by the Pakistani government just after the first cases started being reported within neighboring countries and also possibly due to the fact that many participants had higher education levels that included 42% graduates and 47% postgraduates.

The commonest source of acquiring information related to COVID-19 was through social media, followed by television/radio. Similar results had been revealed by a study conducted in China.¹⁶ The majority of participants knew that the incubation period for coronavirus is 14 days, whereas another study from China reported it as 2.5 days.¹⁸ However, the CDC also suggested it can be range from 2 to 14 days.¹⁹ Common symptoms of COVID-19 are fever, cough, fatigue and shortness of breath.²⁰ However, elderly people with underlying medical condition, such as hypertension, heart disease, chronic obstructive pulmonary disease coronary patients and those with chronic respiratory disease and frontline health providers are at a greater risk of acquiring the infection.²¹

In total, 75% of participants from the current study had awareness regarding COVID-19 symptoms, while 82% knew the importance of wearing face masks as preventive measures, and 69% had sufficient knowledge regarding coronavirus being transmitted through different surfaces. Our findings are in line with another study conducted among Pakistani university students, their knowledge about incubation was lower, whereas the awareness with common symptoms was higher.²² The transmission of COVID-19 can occur through direct or indirect close contact with the infected person or through discharge of saliva, and respiratory emissions or droplets which can be emitted from infected persons during coughing, sneezing or spitting.^{23,24} Respiratory droplets normally are greater than 5-10 μm and aerosol droplets are less than 5 μm in diameter; they can be transferred to any individual when a normal person comes in close contact with the infected person.²⁵

The risk of COVID-19 transmission in dental practices is considered higher than in other healthcare settings due to the close and prolonged contact between dental practitioners and patients. Aerosol transmission awareness was noted to be significantly lower in this study (21%), which is in agreement with a study conducted in India.¹⁵ Dental healthcare workers are concerned about the high risk of COVID-19 infection because some patients are not honest about their pre-treatment COVID-19 swab test results. Conducting regular rapid tests on both patients and healthcare staff is a viable solution to reduce the overall risk.²⁶ Hence, there is a need to increase the awareness level of people regarding the modes of transmission of COVID-19, importance of laboratory tests and to acknowledge its significance.

Many dentists have implemented additional precautions in their clinics to prevent the spread of the virus. These measures include using atomizers, UV lamps, and HEPA filters, which have been recommended by health regulatory bodies worldwide. Previous studies have demonstrated that air purifiers equipped with HEPA H12 class filters are more effective

at removing aerosols during dental procedures compared to standard air purifiers. To guarantee the safety of dental personnel during the pandemic, health regulatory bodies have issued guidelines and recommendations. It is crucial to provide infection control education to all clinic staff to monitor disease progression effectively. Before visiting the clinic, patients should also receive education on infection control measures, preventive methods, and updated hygiene guidelines. By understanding the situation and methods for prevention, the dental community can continue to provide healthcare services to patients during the pandemic.²⁷

Another study from China showed that 93% of study participants agreed that COVID-19 transmission can be prevented by thoroughly washing hands with soap.²⁸ These facts agree with the findings of the study; 92% of the participants washed their hands multiple times a day. As a preventive practice using antibacterial soap was significantly associated with knowledge score (10.5 ± 3.41 ; $p < 0.001$) which is similar to a study conducted in Jordan where 87% adopted hand washing as a preventive measure.¹⁷ Almost 60% of participants in the present study were aware of hand washing technique. A former study concluded that washing hands with soap and water is enough to reduce the risk of viral infections and when it is practiced with the recommended protocol of hand washing technique it reduces the rate of transmission of COVID-19.^{29,30}

The findings of the current study suggested that the majority of participants had good learning attitude towards COVID-19; however lower knowledge level and negative attitudes were also recorded. A total of 63% individuals were tired of listening and did not want further information about the virus. One reason may be that majority had no relative or friend who had been infected with the virus. Additional educational campaigns are needed for the general public for further guiding them regarding the mode of transmission, isolation period and the different adoptive preventive strategies (like social distancing, avoiding handshakes, wearing masks and gloves), along with the risk of personal and family infection with COVID-19. The strength of the current study was that it was a nationwide survey with a large sample, and that the research was conducted during the surging stage of the COVID-19 pandemic.

Limitations

The study is not a true representative sample of Pakistan's general population. It's a convenience sample through social media, which has its own limitations. Social media statistics are dynamic and can change according to its popularity and due to specific group of users being educated and having access to that information. This is the reason most of the participants were graduates and above.

The online survey used in the study may not have included responses from individuals living in areas with limited access to social media and internet facilities. Additionally, individuals from economically disadvantaged backgrounds who do not have access to smartphones with social media applications may also have been excluded from the sample, which could result in a bias in the sample's coverage. Another potential issue with the online survey used in the study is that participants may have provided inaccurate information when completing the self-administered questionnaire. Additionally, the survey may not have included individuals who are illiterate or who do not understand English, as the survey was conducted online and required participants to have access to the internet and be able to understand and respond in English.

Future recommendations

Further knowledge and awareness should be promoted. Future work should focus on a larger, national representative sample population. It is important to conduct information campaigns through government agencies, educate patients through clinicians, and provide accurate media coverage.

Conclusion

It can be concluded from the current study that female participants had a higher level of education, and more awareness regarding the coronavirus. The majority of the participants followed a proper hand washing regime and washed their face. It also highlighted the power of social media as the source of information.

Data availability

Underlying data

Harvard Dataverse: General Public Awareness, Knowledge and Attitude toward COVID-19 infection and prevention: A cross sectional study from Pakistan, <https://doi.org/10.7910/DVN/1K5LDD>.³¹

Extended data

Harvard Dataverse: General Public Awareness, Knowledge and Attitude toward COVID-19 infection and prevention: A cross sectional study from Pakistan, <https://doi.org/10.7910/DVN/1K5LDD>.³¹

This project contains the following extended data:

- Questionnaire

Data are available under the terms of the [Creative Commons Zero “No rights reserved” data waiver](#) (CC0 1.0 Public domain dedication).

Acknowledgments

The authors are thankful to all the participants who participated contributed in this study. We are also grateful to all friends and family who provided support in the distribution of this online questionnaire.

References

- Ali S, Noreen S, Farooq I: **COVID-19 and health care workers in Pakistan. Are we losing the fight?** *J Med Sci.* 2020; **28**(2): 186–188.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cucinotta D, Vanelli M: **WHO declares COVID-19 a pandemic.** *Acta Biomed.* 2020; **91**: 157–160.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ali S, Farooq I, Abdelsalam M, et al.: **Current clinical dental practice guidelines and the financial impact of COVID-19 on dental care providers.** *European Journal of Dentistry.* 2020.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Tripathi R, Alqahtani SS, Albarra AA, et al.: **Awareness and Preparedness of COVID-19 Outbreak Among Healthcare Workers and Other Residents of South-West Saudi Arabia: A Cross-Sectional Survey.** *Front Public Health.* 2020 Aug 18; **8**: 482.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ahmed F, Ahmed N, Pissarides C, et al.: **Why inequality could spread COVID-19.** *Lancet Pub Health.* 2020; **5**: e240.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- de Wilde AH, Snijder EJ, Kikkert M, et al.: **Host factors in coronavirus replication.** *Curr Top Microbiol Immunol.* 2018; **419**: 1–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ahmed EF, Shehata MA, Elheeny AA: **COVID-19 awareness among a group of Egyptians and their perception toward the role of dentists in its prevention: a pilot cross-sectional survey.**
- Hi W, Huang CJ, Casper JPZ, et al.: **Facemask shortage and the novel coronavirus disease (COVID-19) outbreak: reflections on public health measures.** *EClinicalMedicine.* 2020; **100329**: 21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bourouiba L: **Turbulent gas clouds and respiratory pathogen emissions: Potential implications for reducing transmission of Covid-19.** *JAMA.* 2020.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Loh NW, Tan Y, Taculod J, et al.: **The impact of high-flow nasal cannula (hfnc) on coughing distance: Implications on its use during the novel coronavirus disease outbreak.** *Can J Anaesth.* 2020.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Labban L, Thallaj N, Labban A: **Assessing the level of awareness and knowledge of COVID 19 pandemic among Syrians.** *Arch Med.* 2020; **12**(8).
[Publisher Full Text](#)
- Farooq I, Ali S, Moheet IA, et al.: **COVID-19 outbreak, disruption of dental education, and the role of teledentistry.** *Pak J Med Sci.* 2020; **36**(7): 1726–1731.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Salman M, Mustafa ZU, Asif N, et al.: **Knowledge, attitude and preventive practices related to COVID-19: a cross-sectional study in two Pakistani university populations.** *Drugs Ther Perspect.* 2020 May; **9**: 1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Labban L, Thallaj N, Labban A: **Assessing the level of awareness and knowledge of COVID 19 pandemic among Syrians.** *Arch Med.* 2020; **12**(2): 8.
[Publisher Full Text](#)
- Narayana G, Pradeepkumar B, Ramaiah JD, et al.: **Knowledge, perception, and practices towards COVID-19 pandemic among general public of India: A cross-sectional online survey.** *Curr Med Res Pract.* 2020; **10**(4): 153–159.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Zhong BL, Luo W, Li HM, et al.: **Knowledge, attitudes, and practices towards COVID-19 among Chinese residents during the rapid rise period of the COVID-19 outbreak: a quick online cross-sectional survey.** *Int J Biol Sci.* 2020; **16**(10): 1745.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Khader Y, Al Nsour M, Al-Batayneh OB, et al.: **Dentists' awareness, perception, and attitude regarding COVID-19 and infection control: cross-sectional study among Jordanian dentists.** *JMIR Public Health Surveill.* 2020; **6**(2): e18798.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Li Q, Guan X, Wu P, et al.: **Early transmission dynamics in Wuhan, China, of novel coronavirus-infected pneumonia.** *N Engl J Med.* 2020, Jan 29.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Centers for Disease Control and Prevention. **Symptoms of coronavirus.** Accessed April 21, 2020. View in Article. 2020.
[Reference Source](#)
- Ali S, Noreen S, Farooq I, et al.: **Risk assessment of health care workers at the frontline against COVID-19.** *Pak J Med Sci.* 2020; **36**(COVID19-S4): S99–S103.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Pan A, Liu L, Wang C, et al.: **Association of public health interventions with the epidemiology of the COVID-19 outbreak in Wuhan.** *China. Jama.* 2020 May 19; **323**(19): 1915–1923.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Salman M, Mustafa ZU, Asif N, et al.: **Knowledge, attitude and preventive practices related to COVID-19: a cross-sectional study in two Pakistani university populations.** *Drugs Ther Perspect.* 2020 May; **9**: 1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Liu J, Liao X, Qian S, et al.: **Community transmission of severe acute respiratory syndrome coronavirus 2, Shenzhen, China, 2020.**
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Luo L, Liu D, Liao X, et al.: **Modes of contact and risk of transmission in COVID-19 among close contacts (pre-print).** *MedRxiv.* 2020.
[Publisher Full Text](#)
- World Health Organization. **Infection Prevention and Control of Epidemic-and Pandemic-prone Acute Respiratory Infections in Health Care.** Geneva: World Health Organization; 2014.
[Reference Source](#)
- Marya A, Karobari MI, Selvaraj S, et al.: **Risk perception of SARS-CoV-2 infection and implementation of various protective measures by dentists across various countries.** *Int J Environ Res Public Health.* 2021 May 29; **18**(11): 5848.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Albargi AM, Assiry AA, Bahammam HA, et al.: **Analysis of the reliability of patient COVID-19 screening data compared against their actual test data.** *Sci. Prog.* 2021 Jul; **104**(3): 003685042110429.
[Publisher Full Text](#)

28. Giao H, Han NTN, Van Khanh T, *et al.*: **Knowledge and attitude toward COVID-19 among healthcare workers at District 2 Hospital, Ho Chi Minh City.** *Asian Pacific J Tropical Med.* 2020; **13**: 260.
[Publisher Full Text](#)
29. Chaudhary NK, Chaudhary N, Dahal M, *et al.*: **Fighting the SARS CoV-2 (COVID-19) pandemic with soap.** 2020.
[Publisher Full Text](#)
30. Ali S, Alam BF, Farooqi F, *et al.*: **Dental and Medical Students' Knowledge and Attitude toward COVID-19: A Cross-Sectional Study from Pakistan.** *Eur J Dent.* 2020 Dec 7; **14**(S1): S105-12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
31. Farooqi F: **General Public Awareness, Knowledge and Attitude toward COVID-19 infection and prevention : A cross sectional study from Pakistan.** *Harvard Dataverse, V2.* 2021.
[Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:  

Version 2

Reviewer Report 06 June 2023

<https://doi.org/10.5256/f1000research.146374.r176785>

© 2023 Karobari M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Mohmed Isaqali Karobari 

¹ Conservative Dentistry Unit, School of Dental Sciences, Universiti Sains Malaysia, Kota Bharu, Malaysia

² Department of Conservative Dentistry and Endodontics, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai, India

³ Department of Restorative Dentistry & Endodontics, Faculty of Dentistry, University of Puthisastra, Phnom Penh, 12211, Cambodia

Dear Authors,

The authors have addressed all the comments and the manuscript has much improved. I would like to congratulate the authors and wish them all the very best for future endeavors.

Best regards and keep well

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Dentistry, Endodontics, Restorative, Dental education, COVID

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 15 October 2021

<https://doi.org/10.5256/f1000research.56002.r94965>

© 2021 Marya A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Anand Marya 

¹ Department of Orthodontics, Faculty of Dentistry, University of Puthisastra, Phnom Penh, Cambodia

² Department of Orthodontics, Saveetha Dental College & Hospitals, Saveetha Institute of Medical and Technical Sciences (SIMATS), Saveetha Univeristy, Chennai, Tamil Nadu, India

I would like to thank the editor for inviting me to review this manuscript. I would also like to compliment the authors on a well-conducted study that is very relevant during the current pandemic.

The authors have made an effort to conduct a KAP study on the public in Pakistan to evaluate their response towards COVID-19. They conducted a survey using 1120 people for data collection and also ran the Student's t-test and ANOVA after the data collection. They concluded that female participants in the study had a higher awareness regarding the SARS-CoV2 pandemic and also found out that people generally had good awareness about hand hygiene methods.

I only have a few suggestions to help improve the quality of the paper:

- Please include a power of the sample part/discussion. Although the investigators have taken a sample of 1120, it would be a good idea to run a power of the sample test to demonstrate the population this sample would represent.
- Please include a few points of collecting information online in the limitations section. Each study has its own limitations and online surveys generally have their own limitations and bias. Kindly elaborate on this in a separate section.

Also, you may find the following studies helpful to add to your discussion. The reason for using these studies is they are relevant to your paper and also would help you add to the discussion and limitations sections as both are online surveys on very closely related topics that have been published in high-impact journals.

1. Marya *et al.*, 2021¹: This study focuses on what measures are being taken to curtail the spread of COVID-19 by dentists to ensure no contact spread or cross-contamination occurs.
2. Albargi *et al.*, 2021²: This is a very important paper that highlights the attitude of the patients in many situations where they do not divulge proper information regarding their COVID-19 status. Again, this paper is very important in demonstrating the attitude of the general public towards COVID-19.

References

1. Marya A, Karobari MI, Selvaraj S, Adil AH, et al.: Risk Perception of SARS-CoV-2 Infection and Implementation of Various Protective Measures by Dentists Across Various Countries. *Int J Environ Res Public Health*. 2021; **18** (11). [PubMed Abstract](#) | [Publisher Full Text](#)
2. Albargi AM, Assiry AA, Bahammam HA, Alassiri MY, et al.: Analysis of the reliability of patient COVID-19 screening data compared against their actual test data. *Sci Prog*. **104** (3):

368504211042980 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Orthodontics, oral health, dental public health

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 30 Mar 2023

Faraz Farooqi

I only have a few suggestions to help improve the quality of the paper:
Please include a power of the sample part/discussion. Although the investigators have taken a sample of 1120, it would be a good idea to run a power of the sample test to demonstrate the population this sample would represent.

Response: Power calculation has been included.

Please include a few points of collecting information online in the limitations section. Each study has its limitations and online surveys generally have their own limitations and bias. Kindly elaborate on this in a separate section.

Response: has been elaborated.

Also, you may find the following studies helpful to add to your discussion. The reason for using these studies is they are relevant to your paper and also would help you add to the discussion and limitations sections as both are online surveys on very closely related topics that have been published in high-impact journals.

Marya et al., 2021¹: This study focuses on what measures are being taken to curtail the spread of COVID-19 by dentists to ensure no contact spread or cross-contamination occurs.

Albargi et al., 2021²: This is a very important paper that highlights the attitude of the patients in many situations where they do not divulge proper information regarding their COVID-19 status. Again, this paper is very important in demonstrating the attitude of the general public toward COVID-19.

Response: the suggested articles have been added.

Competing Interests: No competing interests were disclosed.

Reviewer Report 14 October 2021

<https://doi.org/10.5256/f1000research.56002.r96322>

© 2021 Karobari M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Mohmed Isaqali Karobari 

¹ Conservative Dentistry Unit, School of Dental Sciences, Universiti Sains Malaysia, Kota Bharu, Malaysia

² Department of Conservative Dentistry and Endodontics, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai, India

³ Department of Restorative Dentistry & Endodontics, Faculty of Dentistry, University of Puthisastra, Phnom Penh, 12211, Cambodia

The manuscript under review attempts to evaluate the general public awareness, knowledge, and attitude towards COVID-19 infection and prevention amongst the Pakistani population. In general, the manuscript captures the details of the study design and implementation of the project. All the sections of the manuscript are well written and concluded, although several limitations exist, and it has been presented in the manuscript. The study is of sound design and clear practical and clinical interest, and I suggest accepting this article with minor revisions.

Following are the minor comments:

Abstract:

- *Background:* The authors have mentioned aims; either change the subheading to aims and add another subheading as background or write an introduction of the study.
- *Results:* Kindly mention the statistical significance.

Methods:

- Is the study conducted in a Pakistani subpopulation or a Pakistani population?

Results:

- The description in the results section is a duplication of tables and figures. Kindly avoid the repetition of data that is already presented in the tables and figures.

Discussion:

- Kindly avoid writing results in the discussion; the authors can mention the significant difference.
- Write future recommendations.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Dentistry, Endodontics, Restorative, Dental education, COVID

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 30 Mar 2023

Faraz Farooqi

Abstract:

Background: The authors have mentioned aims; either change the subheading to aims and add another subheading as background or write an introduction of the study.

Response: separate section of background and aims has been added in the abstract section.

Results: Kindly mention the statistical significance.

Response: has been added.

Methods:

Is the study conducted in a Pakistani subpopulation or a Pakistani population?

Response: the study was conducted on the Pakistani population, not a subpopulation.

Results:

The description in the results section is a duplication of tables and figures. Kindly avoid the repetition of data that is already presented in the tables and figures.

Response: has been corrected.

Discussion:

Kindly avoid writing results in the discussion; the authors can mention the significant difference.

Response: has been corrected.

Write future recommendations.

Response: Has been added.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research