

Ketone bodies : A double-edged sword for mammalian life span.

著者	TOMITA Issei, TSURUTA Hiroaki, YASUDA-YAMAHARA Mako, YAMAHARA Kosuke, KUWAGATA Shogo, TANAKA-SASAKI Yuki, KANASAKI-CHIN Masami, FUJITA Yukihiro, NISHI Eiichiro, KATAGIRI Hideki, MAEGAWA Hiroshi, KUME Shinji
journal or publication title	Aging cell
year	2023-04-14
URL	http://hdl.handle.net/10422/00013569

doi: 10.1111/accel.13833(<https://doi.org/10.1111/accel.13833>)



Ketone bodies: A double-edged sword for mammalian life span

Issei Tomita¹ | Hiroaki Tsuruta¹ | Mako Yasuda-Yamahara¹ | Kosuke Yamahara¹ | Shogo Kuwagata¹ | Yuki Tanaka-Sasaki¹ | Masami Chin-Kanasaki¹ | Yukihiro Fujita¹ | Eiichiro Nishi² | Hideki Katagiri³ | Hiroshi Maegawa¹ | Shinji Kume¹

¹Department of Medicine, Shiga University of Medical Science, Tsukinowa-cho, Seta, Otsu, Shiga 520-2192, Japan

²Department of Pharmacology, Shiga University of Medical Science, Tsukinowa-cho, Seta, Otsu, Shiga 520-2192, Japan

³Department of Metabolism and Diabetes, Tohoku University Graduate School of Medicine, 2-1 Seiryomachi, Aoba-ku, Sendai, Miyagi 980-8574, Japan

Correspondence

Shinji Kume, Department of Medicine, Shiga University of Medical Science, Tsukinowa-cho, Seta, Otsu, Shiga 520-2192, Japan.
Email: skume@belle.shiga-med.ac.jp

Funding information

Japan Science and Technology Agency, Grant/Award Number: Moonshot R&D; Grant Number JPMJMS2023; Japan Society for the Promotion of Science, Grant/Award Number: 21H03353 and 22H03128; Takeda Science Foundation, Grant/Award Number: grant to E.N.

Abstract

Accumulating evidence suggests health benefits of ketone bodies, and especially for longevity. However, the precise role of endogenous ketogenesis in mammalian life span, and the safety and efficacy of the long-term exogenous supplementation of ketone bodies remain unclear. In the present study, we show that a deficiency in endogenous ketogenesis, induced by whole-body *Hmgcs2* deletion, shortens life span in mice, and that this is prevented by daily ketone body supplementation using a diet containing 1,3-butanediol, a precursor of β -hydroxybutyrate. Furthermore, feeding the 1,3-butanediol-containing diet from early in life increases midlife mortality in normal mice, but in aged mice it extends life span and prevents the high mortality associated with atherosclerosis in *ApoE*-deficient mice. By contrast, an ad libitum low-carbohydrate ketogenic diet markedly increases mortality. In conclusion, endogenous ketogenesis affects mammalian survival, and ketone body supplementation may represent a double-edged sword with respect to survival, depending on the method of administration and health status.

KEYWORDS

Hmgcs2, ketone body, longevity, low-carbohydrate ketogenic diet

1 | INTRODUCTION

Ketone bodies (KBs) represent an important glucose-sparing energy source during fasting in mammals (Cahill et al., 1970; Reichard et al., 1974), and they have been suggested to be involved in the prolongation of life span induced by calorie restriction and low-carbohydrate ketogenic diet (LCKD)-feeding (Newman et al., 2017; Roberts et al., 2017; Stekovic et al., 2019). However, their precise role in mammalian longevity has long been debated because such dietary interventions cause many metabolic changes in addition to ketogenesis.

In the present study, the survival of *Hmgcs2*^{-/-} mice that we generated previously (Tomita et al., 2020), which are incapable of endogenous ketogenesis, was assessed to evaluate the role of KBs in mammalian life span (Figure 1a). *Hmgcs2* deficiency increased the mortality rate in old age, and this was accompanied by lower blood β -hydroxybutyrate (β -OHB) concentration, but no differences in body weight or blood glucose concentration (Figure 1b–e, Table S1). The reduction in life span of the *Hmgcs2*^{-/-} mice was prevented by the dietary administration of 1,3-butanediol (1,3-BD), a chemical precursor of β -OHB (Figure 1a,e). Higher mortality of *Hmgcs2*^{-/-} mice during the weaning period has previously been reported⁷.

Issei Tomita and Hiroaki Tsuruta authors contributed equally.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Aging Cell* published by Anatomical Society and John Wiley & Sons Ltd.




FIGURE 1 Effects of endogenous ketogenesis and dietary KB supplementation on mouse survival rate. (a) Study protocol for determining the life span of *Hmgcs2*^{+/+} mice fed a normal diet (ND; *n*=20) and *Hmgcs2*^{-/-} mice fed either an ND (*n*=15) or a 1,3-butanediol (1,3-BD)-containing diet (*n*=15). (b) β -OHB concentration during the study period. (c) Change in body weight. (d) Blood glucose concentration. (e) Cumulative survival rate. (f) Study protocol for determining the life span of *Hmgcs2*^{+/+} mice fed an ND (*n*=20), a 1,3-BD diet (*n*=20), or a low-carbohydrate ketogenic diet (LCKD), which contained 4.5% carbohydrate, 80.8% fat, and 14.7% protein (calorie %) (*n*=15), from 24 weeks of age. (g) β -OHB concentration. (h) Change in body weight. (i) Blood glucose concentration. (j) Cumulative survival rate. **p*<0.01, †*p*<0.05; ns, not significant.

Therefore, the present findings emphasize the importance of KBs for mammalian survival and imply that they have significant effects on life span during two time windows: just after birth and in old age.

The health benefits of KBs have recently been gaining attention (Cheng et al., 2019; Cotter et al., 2014; Cox et al., 2016; Fan et al., 2021; Mujica-Parodi et al., 2020; Nielsen et al., 2019; Tomita et al., 2020; Torres et al., 2019). Therefore, there is an urgent need to determine whether long-term KB supplementation is safe

and extends life span. To answer this question, the effects on the life span of control *Hmgcs2*^{+/+} mice of an LCKD, which contained 4.5% carbohydrate, 80.8% fat, and 14.7% protein (calorie %), and a 1,3-BD-containing diet from 24 weeks of age were examined (Figure 1f). LCKD-fed mice exhibited higher mortality in old age, which was accompanied by rapid decreases in body weight and blood glucose concentration after midlife (Figure 1g–j, Table S2). The survival rate of the 1,3-BD diet-fed mice was also lower, but there

were differences in mortality from the LCKD-fed mice at various time points (Figure 1j). The 1,3-BD diet was associated with higher midlife mortality, but this difference disappeared when the normal diet (ND)-fed mice began to die (between 90 and 110 weeks of age), such that their mortality in old age was similar to that of ND-fed mice (Figure 1j).

This finding led us to hypothesize that KB supplementation commencing just prior to old age might extend life span. To evaluate this

possibility, the 1,3-BD diet or LCKD was fed to C57BL/6J mice from 72 weeks of age (Figure 2a). Both interventions increased the blood β -OHB concentration similarly, and had no marked effects on body weight or blood glucose levels (Figure 2b–d, Table S3). However, the 1,3-BD diet extended the life span, whereas the LCKD shortened it (Figure 2e).

These findings led us to further hypothesize that the 1,3-BD diet might extend life span, even in young mice, if they have organ




FIGURE 2 Effects of KB supplementation on the survival rate of aged mice and young *ApoE*^{-/-} mice. (a) Study protocol for determining the life span of aged wild-type (WT) C57BL/6J mice fed a normal diet (ND; *n* = 15), a 1,3-butanediol (1,3-BD)-containing diet (*n* = 15), or a low-carbohydrate ketogenic diet (LCKD), which contained 4.5% carbohydrate, 80.8% fat, and 14.7% protein (calorie %) (*n* = 15). (b) β -OHB concentration during the study period. (c) Change in body weight. (d) Blood glucose concentration. (e) Cumulative survival rate. (f) Study protocol for determining the life span of *ApoE*^{-/-} mice fed either an ND (*n* = 20), a 1,3-BD diet, (*n* = 20) or an LCKD (*n* = 15). (g) β -OHB concentration. (h) Change in body weight. (i) Blood glucose concentration. (j) Cumulative survival rate. **p* < 0.01, †*p* < 0.05; ns, not significant; na, not available.



damage. Therefore, we next determined the effects of KB administration on the life span of high-fat diet (HFD)-fed *ApoE*^{-/-} mice with atherosclerosis-related organ damage (Figure 2f). The 1,3-BD diet and LCKD induced similar metabolic changes to those identified in the aged mice (Figure 2g–i, Table S4). The high mortality of the HFD-fed *ApoE*^{-/-} mice was reduced by the 1,3-BD diet but worsened by the LCKD (Figure 2j).

The present results suggest that the timing and method of KB supplementation and health status have significant influences on the effects of KBs on mammalian life span. Given that *Hmgcs2*^{-/-} mice had a lower survival rate in old age and that 1,3-BD extended the life span of aged and *ApoE*^{-/-} mice, KBs may have an important role in tissue repair, as suggested by recent studies (Cheng et al., 2019; Tomita et al., 2020). By contrast, 1,3-BD diet from early life was associated with higher mortality, although the reason for this remains unclear. Thus, the potential for KBs to be used to promote human health (Chen et al., 2021) and their safety, particularly in younger individuals, requires further investigation. Also, further analysis regarding the cause of death is needed to elucidate the mechanisms by which KBs affect the life span.

The health benefits of LCKD remain the subject of debate. Time-restricted or energy-controlled LCKD feeding was reported to extend life span in mice (Newman et al., 2017; Roberts et al., 2017), whereas ad libitum LCKD feeding leading to much calorie intake shortened their life span in our study. Thus, the unrestricted LCKD seems to cancel out the health benefits of KBs and to be harmful. Because LCKDs are often used for weight management (Foster et al., 2003), the effects of long-term LCKD consumption require careful monitoring.

In conclusion, it is possible that endogenous ketogenesis plays an important role in mammalian survival, although there may be sex differences in the effects. Furthermore, KB supplementation represents a double-edged sword, with their effects depending on the method of administration and health status. The present findings provide a further ray of hope, but also a new challenge, for the use of KBs to prolong healthy life span.

2 | METHODS

2.1 | Ethics

The experimental protocols were approved by the Gene Recombination Experiment Safety Committee and Research Center for Animal Life Science of Shiga University of Medical Science.

2.2 | Animal studies

Details on the creation of each transgenic mouse, the dietary intervention, and the measurement of each parameter are provided in the [Supplemental Information](#).

2.3 | Statistical analysis

Two-way ANOVA followed by Tukey's *post hoc* test was used to determine the effects of genotype and treatment among multiple groups, and the unpaired Student's *t* test was used to compare two groups. Survival rates were determined using the Kaplan–Meier method and compared among the groups using the log-rank test, with the Bonferroni correction. R statistical software version 1.55 (Vienna, Austria) was used for the analysis. *p* < 0.05 was considered to represent statistical significance. Each experiment was conducted twice, and the combined data were statistically analyzed.

AUTHOR CONTRIBUTIONS

I.T., S.K., and H.M. designed the study. I.T. and H.T. performed the experiments. I.T., S.K., M.Y.-Y., K.Y., S.K., Y.T.-S., M.C.-K., Y.F., E.N., and H.K. discussed and analyzed the data. I.T. and S.K. drafted the manuscript. All the authors revised the manuscript and approved the final version.

ACKNOWLEDGMENTS

We thank Naoko Y., Keiko K., and the Central Research Laboratory of SUMS for their technical assistance. This study was supported by the JSPS [21H03353 and 22H03128], JST [Moonshot R&D; Grant Number JPMJMS2023], and the Takeda Science Foundation [grant to E.N.].

CONFLICT OF INTEREST STATEMENT

The authors declare no competing interests.

DATA AVAILABILITY STATEMENT

The datasets are available from the corresponding author on reasonable request.

ORCID

Shinji Kume  <https://orcid.org/0000-0001-6937-9715>

REFERENCES

- Cahill, G. F., Jr. (1970). Starvation in man. *The New England Journal of Medicine*, 282(12), 668–675. <https://doi.org/10.1056/NEJM197003192821209>
- Chen, O., Blonquist, T. M., Mah, E., Sanoshy, K., Beckman, D., Nieman, K. M., Winters, B. L., Anthony, J. C., Verdin, E., Newman, J. C., & Stubbs, B. J. (2021). Tolerability and safety of a novel ketogenic Ester, bis-hexanoyl (R)-1,3-butanediol: A randomized controlled trial in healthy adults. *Nutrients*, 13(6), 2066. <https://doi.org/10.3390/nu13062066>
- Cheng, C. W., Biton, M., Haber, A. L., Gunduz, N., Eng, G., Gaynor, L. T., Tripathi, S., Calibasi-Kocal, G., Rickelt, S., Butty, V. L., Moreno-Serrano, M., Iqbal, A. M., Bauer-Rowe, K. E., Imada, S., Ulutas, M. S., Mylonas, C., Whary, M. T., Levine, S. S., Basbinar, Y., ... Yilmaz, O. H. (2019). Ketone body signaling mediates intestinal stem cell homeostasis and adaptation to diet. *Cell*, 178(5), 1115–1131 e15. <https://doi.org/10.1016/j.cell.2019.07.048>
- Cotter, D. G., Ercal, B., Huang, X., Leid, J. M., d'Avignon, D. A., Graham, M. J., Dietzen, D. J., Brunt, E. M., Patti, G. J., & Crawford, P. A. (2014). Ketogenesis prevents diet-induced fatty liver injury and



- hyperglycemia. *The Journal of Clinical Investigation*, 124(12), 5175–5190. <https://doi.org/10.1172/JCI76388>
- Cox, P. J., Kirk, T., Ashmore, T., Willerton, K., Evans, R., Smith, A., Murray, A. J., Stubbs, B., West, J., McLure, S. W., King, M. T., Dodd, M. S., Holloway, C., Neubauer, S., Drawer, S., Veech, R. L., Griffin, J. L., & Clarke, K. (2016). Nutritional ketosis alters fuel preference and thereby endurance performance in athletes. *Cell Metabolism*, 24(2), 256–268. <https://doi.org/10.1016/j.cmet.2016.07.010>
- Fan, S. Z., Lin, C. S., Wei, Y. W., Yeh, S. R., Tsai, Y. H., Lee, A. C., Lin, W.-S., & Wang, P. Y. (2021). Dietary citrate supplementation enhances longevity, metabolic health, and memory performance through promoting ketogenesis. *Aging Cell*, 20(12), e13510. <https://doi.org/10.1111/accel.13510>
- Foster, G. D., Wyatt, H. R., Hill, J. O., McGuckin, B. G., Brill, C., Mohammed, B. S., Szapary, P. O., Rader, D. J., Edman, J. S., & Klein, S. (2003). A randomized trial of a low-carbohydrate diet for obesity. *The New England Journal of Medicine*, 348(21), 2082–2090. <https://doi.org/10.1056/NEJMoa022207>
- Mujica-Parodi, L. R., Amgalan, A., Sultan, S. F., Antal, B., Sun, X., Skiena, S., Lithen, A., Adra, N., Ratai, E.-M., Weistuch, C., Govindarajan, S. T., Strey, H. H., Dill, K. A., Stufflebeam, S. M., Veech, R. L., & Clarke, K. (2020). Diet modulates brain network stability, a biomarker for brain aging, in young adults. *Proceedings of the National Academy of Sciences of the United States of America*, 117(11), 6170–6177. <https://doi.org/10.1073/pnas.1913042117>
- Newman, J. C., Covarrubias, A. J., Zhao, M., Yu, X., Gut, P., Ng, C. P., Huang, Y., Haldar, S., & Verdin, E. (2017). Ketogenic diet reduces midlife mortality and improves memory in aging mice. *Cell Metabolism*, 26(3), 547–557 e548. <https://doi.org/10.1016/j.cmet.2017.08.004>
- Nielsen, R., Moller, N., Gormsen, L. C., Tolbod, L. P., Hansson, N. H., Sorensen, J., Harms, H. J., Frøkiær, J., Eiskjaer, H., Jespersen, N. R., Mellemkjaer, S., Lassen, T. R., Pryds, K., Bøtker, H. E., & Wiggers, H. (2019). Cardiovascular effects of treatment with the ketone body 3-hydroxybutyrate in chronic heart failure patients. *Circulation*, 139(18), 2129–2141. <https://doi.org/10.1161/CIRCULATIONAHA.118.036459>
- Reichard, G. A., Jr., Owen, O. E., Haff, A. C., Paul, P., & Bortz, W. M. (1974). Ketone-body production and oxidation in fasting obese humans. *The Journal of Clinical Investigation*, 53(2), 508–515. <https://doi.org/10.1172/JCI107584>
- Roberts, M. N., Wallace, M. A., Tomilov, A. A., Zhou, Z., Marcotte, G. R., Tran, D., Perez, G., Gutierrez-Casado, E., Koike, S., Knotts, T. A., Imai, D. M., Griffey, S. M., Kim, K., Hagopian, K., McMackin, M. Z., Haj, F. G., Baar, K., Cortopassi, G. A., Ramsey, J. J., & Lopez-Dominguez, J. A. (2017). A ketogenic diet extends longevity and Healthspan in adult mice. *Cell Metabolism*, 26(3), 539–546 e535. <https://doi.org/10.1016/j.cmet.2017.08.005>
- Stekovic, S., Hofer, S. J., Tripolt, N., Aon, M. A., Royer, P., Pein, L., Stadler, J. T., Pendl, T., Prietl, B., Url, J., Schroeder, S., Tadic, J., Eisenberg, T., Magnes, C., Stumpe, M., Zuegner, E., Bordag, N., Riedl, R., Schmidt, A., & Madeo, F. (2019). Alternate day fasting improves physiological and molecular markers of aging in healthy, non-obese humans. *Cell Metabolism*, 30(3), 462–476 e466. <https://doi.org/10.1016/j.cmet.2019.07.016>
- Tomita, I., Kume, S., Sugahara, S., Osawa, N., Yamahara, K., Yasuda-Yamahara, M., Takeda, N., Chin-Kanasaki, M., Kaneko, T., Mayoux, E., Mark, M., Yanagita, M., Ogita, H., Araki, S.-I., & Maegawa, H. (2020). SGLT2 inhibition mediates protection from diabetic kidney disease by promoting ketone body-induced mTORC1 inhibition. *Cell Metabolism*, 32(3), 404–419 e406. <https://doi.org/10.1016/j.cmet.2020.06.020>
- Torres, J. A., Kruger, S. L., Broderick, C., Amarlkhagva, T., Agrawal, S., Dodam, J. R., Mrug, M., Lyons, L. A., & Weimbs, T. (2019). Ketosis ameliorates renal cyst growth in polycystic kidney disease. *Cell Metabolism*, 30(6), 1007–1023 e1005. <https://doi.org/10.1016/j.cmet.2019.09.012>

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Tomita, I., Tsuruta, H., Yasuda-Yamahara, M., Yamahara, K., Kuwagata, S., Tanaka-Sasaki, Y., Chin-Kanasaki, M., Fujita, Y., Nishi, E., Katagiri, H., Maegawa, H., & Kume, S. (2023). Ketone bodies: A double-edged sword for mammalian life span. *Aging Cell*, 00, e13833. <https://doi.org/10.1111/accel.13833>