

Accepted Manuscript

# *British Journal of General Practice*

## Adapting domestic abuse training to remote delivery during the COVID-19 pandemic: perspectives from general practice and support services

Emsley, Elizabeth; Dowrick, Anna; Szilassy, Eszter; Dixon, Sharon; De Simoni, Anna; Downes, Lucy; Johnson, Medina; Feder, Gene; Griffiths, Chris; Panovska-Griffiths, Jasmina; Barbosa, EC; Wileman, Vari

DOI: <https://doi.org/10.3399/BJGP.2022.0570>

To access the most recent version of this article, please click the DOI URL in the line above.

Received 16 November 2022

Revised 27 March 2023

Accepted 21 April 2023

© 2023 The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0/>). Published by British Journal of General Practice. For editorial process and policies, see: <https://bjgp.org/authors/bjgp-editorial-process-and-policies>

When citing this article please include the DOI provided above.

### **Author Accepted Manuscript**

This is an 'author accepted manuscript': a manuscript that has been accepted for publication in British Journal of General Practice, but which has not yet undergone subediting, typesetting, or correction. Errors discovered and corrected during this process may materially alter the content of this manuscript, and the latest published version (the Version of Record) should be used in preference to any preceding versions

**Adapting domestic abuse training to remote delivery during the COVID-19  
pandemic: perspectives from general practice and support services**

Elizabeth Emsley BMBS, MRCGP (her): **Corresponding author**  
Clinical Teaching Fellow  
Bristol Medical School  
University of Bristol  
ORCID ID: 0009-0005-2060-1545  
[elizabeth.emsley@bristol.ac.uk](mailto:elizabeth.emsley@bristol.ac.uk)

Anna Dowrick PhD, MSc, BA (her)  
Senior Researcher,  
Nuffield Department of Primary Care Health Sciences,  
University of Oxford  
ORCID ID: 0000-0002-2937-6728

Eszter Szilassy PhD, MA (her)  
Senior Research Fellow  
Bristol Medical School,  
University of Bristol  
ORCID ID: 0000-0002-6267-0531

Sharon Dixon BA (Hons) Cantab, MBBS, MRCGP, MSc (her)  
NIHR Doctoral Research Fellow and General Practitioner,  
Nuffield Department of Primary Care Health Sciences,  
University of Oxford  
ORCID ID: 0000-0002-7469-6093

Anna De Simoni MBBS, PhD, FHEA, MRCGP (her)  
Clinical Reader and General Practitioner,  
Wolfson Institute of Population Health,  
Queen Mary University of London  
ORCID ID: 0000-0001-6955-0885

Lucy Downes MSc, BA Hons (her)  
IRIS Network Director,  
IRISi

Medina Johnson MA, PGCE, BA Hons (her)  
Chief Executive,  
IRISi

Gene Feder MBBS, MD, FRCGP (him)  
Professor of Primary Care and General Practitioner,  
Bristol Medical School,  
University of Bristol  
ORCID ID: 0000-0002-7890-3926

Chris Griffiths MA, DPhil, FRCGP (him)  
Professor of Primary Care and General Practitioner,  
Wolfson Institute of Population Health,  
Queen Mary University of London  
ORCID ID: 0000-0001-7935-8694

Jasmina Panovska-Griffiths DPhil, MMath, BA (her)  
Senior Research Fellow and Lecturer in Probability and Statistics,  
The Pandemic Sciences Institute and the Big Data Institute,  
University of Oxford  
ORCID ID: 0000-0002-7720-1121

Estela Capelas Barbosa PhD (her)  
Senior Research Fellow  
City, University of London  
ORCID ID: 0000-0001-8282-131X

Vari Wileman PhD, CPsychol (her)  
Senior Research Associate and Teaching Fellow,  
School of Mental Health and Psychological Sciences  
Institute of Psychiatry and Neuroscience,  
King's College London  
ORCID ID: 0000-0002-9673-8047

## **Abstract**

Background: Identifying and responding to patients affected by domestic violence and abuse (DVA) is vital in primary care. There may have been a rise in the reporting of DVA cases during the COVID-19 pandemic and associated lockdown measures. Concurrently general practice adopted remote working which extended to training and education. IRIS (Identification and Referral to Improve Safety) is an example of an evidence-based UK healthcare training support and referral programme, focusing on DVA. IRIS transitioned to remote delivery during the pandemic.

Aim: We aimed to understand the adaptations and impact of remote DVA training in IRIS-trained general practices, by exploring perspectives of those delivering and receiving training.

Design and setting: Qualitative interviews and observation of remote training of general practice teams in England.

Method: Semi-structured interviews with 21 participants (three practice managers, three reception and administrative staff, eight general practice clinicians and seven specialist DVA staff), alongside observation of 11 remote training sessions. Analysis using a Framework approach.

Results: Remote DVA training in UK general practice widened access to learners. However, it may have reduced learner engagement compared with face-to-face

training. DVA training is integral to the partnership between general practice and specialist DVA services, and reduced engagement risks weakening this partnership.

Conclusion: We recommend a hybrid DVA training model for general practice, including remote information delivery alongside a structured face-to-face element.

This has broader relevance for other specialist services providing training and education in primary care.

### **Keywords**

Domestic violence, COVID-19, SARS-CoV-2, general practice, qualitative research, training activities

### **How this fits in**

There was a shift to remote DVA training in UK general practice during the COVID-19 pandemic. This qualitative study explored lessons learned from general practice and DVA specialist services regarding the transition to remote DVA IRIS training, with consideration of the impact on the overall DVA referral pathway between general practice and specialist DVA services. While remote delivery can improve access to training for learners, this study has found concerns regarding a trade-off between the gain in access and a loss of learner engagement. Connectivity between general practice and specialist DVA services is important in establishing an effective referral pathway between both services.

## **Main text**

### **Introduction**

Domestic violence and abuse (DVA) is a violation of human rights challenging global public health policy and clinical practice. DVA damages mental and physical health for victims and their children.<sup>1</sup> While DVA can affect everyone, the experience of DVA is gendered, with research often focusing on women.<sup>2</sup> Women experiencing DVA have increased health service use and are more likely to contact health services than those unaffected by abuse.<sup>3</sup> In general, patients attending general practice also have a higher lifetime prevalence of DVA compared to the wider population.<sup>4-6</sup> Therefore, the general practice response to DVA is crucial in supporting patients and provides a vital link to specialist services. During the COVID-19 pandemic, there was a rise in reporting of DVA cases occurring alongside a transition to remote general practice consulting.<sup>7,8</sup> Remote working also extended to training and education, including DVA training.<sup>9</sup>

IRIS (Identification and Referral to Improve Safety) is a DVA training, support and referral programme, developed by IRISi,<sup>10</sup> and delivered by local DVA services. IRIS transitioned to remote delivery during the pandemic. DVA training for the whole general practice team is part of a broader intervention aiming to establish a referral pathway between general practice and DVA support services. Advocate Educators (AEs) are specialist DVA workers in the IRIS programme.<sup>11</sup> The AE role includes training general practices and receiving patient referrals for expert advocacy.<sup>11</sup> The IRIS programme was trialled in 2007-2008,<sup>12</sup> and was recommended in high-level

UK government documents, with over 1,275 general practices having received training.<sup>13-16</sup> However, challenges have included difficulties in scheduling training and maintaining general practice engagement.<sup>17</sup> General practice engagement is essential when building a strong pathway of support for patients between general practice and specialist DVA services.<sup>17</sup> Although general practice engagement is an essential part of the IRIS programme, practices are facing competing challenges including increasing scarcity of resources and recruitment challenges.<sup>18</sup> This state of flux has worsened during the pandemic.<sup>19</sup>

Remote DVA training to general practice teams may continue beyond the pandemic. Remote education allows scalable, cost-effective and standardised learning, overcoming geographic and time constraints, and widening access.<sup>20 21</sup> However this may result in learner isolation, technical problems and reduced visual cues which risk misunderstandings and represent a cognitive demand on trainers.<sup>21 22</sup> Spontaneous conversation between trainers and learners, as well as between learners, may also be limited.<sup>22</sup> Remote DVA training could have additional challenges due to the sensitive and emotionally challenging nature of its content, and participants may have lived experience of abuse. There is limited evidence on the impact of remotely delivering this content, and perspectives from those receiving and delivering training are important in deciding the future direction of remote DVA training.

We aimed to understand the adaptations and impact of the shift to remote domestic abuse training during the COVID-19 pandemic in IRIS trained general practices, by exploring perspectives of those delivering and receiving training. We also considered

the broader impacts of remote DVA training on the relationship between general practice and specialist DVA services. This study used the IRIS programme as a case study and is part of a larger study exploring the primary care response to DVA in the COVID-19 pandemic (PRECODE).<sup>23 24</sup>

## Methods

There are multiple actors who support patients affected by DVA in general practice and in DVA specialist services. Any member of the whole general practice team can raise a DVA concern including GPs, nurses and reception or administration staff, however it is typically clinicians who refer patients to specialist DVA services where they are then supported by an AE. AEs work in partnership with clinical leads to deliver training together and connect with general practices.<sup>11</sup> These connections help to establish a working relationship between general practice and specialist DVA services, which underpins the support for patients affected by DVA. For this reason, we considered perspectives from multiple standpoints, by conducting interviews with general practice professionals and IRIS AEs, as well as direct observation of training sessions.

Six IRIS-commissioned sites were selected using a multi-stage sampling framework. The first stage of sampling involved identification of ten geographically diverse sample areas by IRISi regional managers (who support local implementation of IRIS), with six of these areas agreeing to participate. With the involvement of AEs in these six areas, and guided by IRISi regional managers, we used 'snowballing' to recruit individuals from general practices and AEs involved in the IRIS programme.



Participants were approached using a range of methods including email and face-to-face. Data collection occurred between 20 April 2021 and 12 April 2022.

*Interviews with advocate educators and general practice staff*

The qualitative study team conducted interviews with AEs from local domestic abuse agencies delivering the IRIS programme to explore their experience of delivering remote DVA training. We also conducted interviews with general practice staff to understand their perspective of receiving training, with some GPs interviewed also having experience of co-delivering training with AEs. Prior to the interview, participants received an information sheet explaining the researcher's reasons for conducting the research (Supplementary File 1). Interviews were conducted by VW (Psychologist, PhD), ES (Health Services Researcher, PhD), AdS (Academic GP, PhD), AD (Sociologist, PhD), SD (Academic GP) and EE (GP Academic Clinical Fellow). Interviews were semi-structured, used a flexible topic guide (included as Supplementary File 2) and took place remotely, due to COVID-19 social distancing restrictions. The option of a telephone or online platform interview (Microsoft Teams or Zoom), was offered. All participants provided informed consent which was audio-recorded before the interview, and contextual data was collected from participants in the interview. Only participants and researchers were present in interviews. No repeat interviews were carried out. EE had a pre-existing professional relationship with a participant interviewed but this was not felt to affect overall findings.

We used rapid qualitative analysis methods which were modified from McNall and Foster-Fishman,<sup>25</sup> and Vindrola-Padros et al.<sup>26</sup> Data-driven inductive methods involved an interview summary using a rapid assessment procedure (RAP) sheet, which we reviewed regularly in the research team to progress towards iterative data analysis.

All interviews were audio-recorded and encrypted, then transferred to an approved university transcription provider. Verbatim transcripts were anonymised. A Framework approach was applied,<sup>27</sup> which involved a first round of independent manual coding by EE, with the subsequent development of a 'working analytical framework' (included as Supplementary File 3) which was applied to all transcripts. The data was charted into a framework matrix in Microsoft Excel, which allowed connections to be made across the data and key themes to be generated.

#### *Observation of training sessions*

The use of observation methods, such as researchers being immersed in a setting where respondents receive training, provided additional perspectives to those self-reported by participants in interviews. We observed remotely delivered IRIS training sessions to evaluate enablers and barriers to training, content delivery and receipt, and to consider the potential for relationship building between the general practices and IRIS AEs. Sessions were independently observed by VW, AD, SD, AdS, ES and EE. Observers joined the session as a participant on Microsoft Teams with microphone silenced. During the session, we documented the context and dynamics of training sessions, variations in training delivery and demonstration of participants'

engagement. We applied an observation framework informed by Spradley (1980) to record observations' (included as Supplementary File 4). The framework refers to the space and layout of the virtual setting, objects used, how time is managed, the actors involved, how they engage and communicate and what activities they engage in.

We recognised that observations for remote training may have limitations compared with face-to-face settings, for example the possibility of training participants having their cameras off, thereby impacting group interactions. However we aimed for discreet observation recording, analytical detachment and minimising the influence of the researcher on the training environment, recognising the risk of potential observer bias, and the impact of the presence of the observer may have on primary data by influencing group elements.<sup>28</sup>

#### *Patient and Public Involvement*

We have been guided by a group of women advisers with lived experience and perspectives of accessing healthcare services for domestic abuse. Regular meetings helped to inform our research approach and overall application of research findings to support general practice in identifying and responding to domestic abuse.

#### *Ethics*

Ethical approval for the study was received from the Health Research Authority and Health and Care Research Wales (reference 20/HRA/5873) and University of Bristol Faculty of Health Science Research Ethics Approval (reference 113044).

## Results

### Participants' characteristics

#### Interviews

We interviewed 21 participants purposively sampled from NHS general practices and domestic abuse agencies delivering the IRIS programme from four regions in England and Wales, with participant characteristics described in table 1. We interviewed seven AEs involved in training delivery and 14 general practice staff (three practice managers, eight GPs, three reception team members). The average interview length was approximately 38 minutes. At least two GPs interviewed had experience in delivering DVA training. Training was one aspect of the interview topic guide and the most detailed perspectives on training were offered by ten participants (seven AEs, two practice managers, one GP). AEs delivering training most commonly shared their views regarding the shift to remote training and this is reflected in the data below.

**Table 1: Interview participants characteristics**

Characteristic	N (%)
<b>Type of healthcare professional</b>	
Advocate educator	7 (33)
GP	8 (38)
Practice manager	3 (14)
Reception team	3 (15)

<b>Gender</b>	
Female	20 (95)
Male	1 (5)
<b>Years spent working in sector</b>	
0-3	2 (10)
4-9	9 (43)
10-20	2 (10)
20+	3 (14)
Not specified	5 (24)

### Training sessions

We observed eight IRIS training sessions delivered remotely by 11 AEs to general practice teams in England and Wales. 52 participants attended training across five groups of clinical staff (GPs, practice nurses and general practice based psychological well-being practitioners) and three groups of support staff (receptionists, administrators, and practice managers). Participant group sizes ranged from four to nine and were diverse in terms of age, gender, and ethnicity with varied time in role (<1year to >20years). Training sessions were up to two hours long for clinical staff and one hour for support staff. AEs sometimes delivered the sessions independently (5 sessions) and sometimes in pairs (3 sessions), if AEs preferred to share delivery and communication.

### **Themes**

Results were grouped into four themes identified from both the observation of the training sessions and the interviews, with key themes below weaving across both datasets:

- 1) Constants and adaptations,
- 2) The access-engagement trade-off,
- 3) Sensitive content in remote sessions,
- 4) Is remote training here to stay?

In Supplementary Table S1 we highlight the main codes relevant to each theme. To demonstrate how data from interviews and observations link to individual themes, we have created Supplementary Table S2.

#### 1. Constants and adaptations

The shift towards remote training highlighted issues which previously existed in face-to-face training. These included challenges in scheduling DVA training with busy general practices facing competing demands. Major adaptations in the transition to remote training during the pandemic included changes in DVA training content, as well as adjusting to novel technology for those delivering and receiving training.

When scheduling training sessions with general practices, a variability in commitment amongst practices, predating the pandemic, was reflected: *“There is always going to be some practices that are not as- they have never been as easy to get onboard as others. Some of them waiting for their update training they are, not resistant, but they are not good referrers, they don’t seem to be quite as onboard as some of the others.”* AE1

One AE explained why arranging training with practices is essential for the relationship between general practice and specialist DVA services: *"...If they haven't had the training then the communication is really poor. They just don't have a clue who I am, what's going on, who IRIS is, why the patient would be referred...I'm very confident in which surgeries I have that good rapport with..."* AE6

There have also been major adaptations in the shift to remote training delivery: *"We talked about the increase in referrals through the COVID 19 pandemic... that there had been more homicides due to domestic violence during the pandemic... But then a huge part of the training is exactly that of asking how to ask the questions of domestic abuse over phone and video consultation, how to do that safely. Because obviously women are at home and, their perpetrator could be right beside them or behind the computer, listening on the call."* AE3

A focus on clinician empowerment was also highlighted by this GP who delivers training: *"I am giving doctors, clinicians and nurses and things confidence that they need to actually get in there when they are on the phone and actually be curious and ask questions. And also work around issues of safety which are much more concerning if there is somebody in the background..."* GP1

Adapting to novel technology was another major adjustment for those delivering and receiving training. When observing training sessions, our study team noticed technical problems across all sessions; screen camera and microphone faults occurred and some attendees were unable to connect.

Some AEs struggled with the new technology: *“Oh and I have had some disasters. I think once my electricity tripped and I got thrown out of the session and... You get back in, people understand and yes I was the one sweating, they were all quite fine waiting for me.”* AE1

Other AEs were surprised at how well they adapted: *“Yes, I was surprised by it. I don’t know maybe I was surprised because it was- I doubted my own abilities to deliver training online but yes, it is just I can’t sit in front of a screen and talk but I actually can.”* AE1

General practices receiving training also needed to adapt to novel, unfamiliar technology: *“This practice, they didn’t have the right equipment, they didn’t have enough webcams and laptops for everyone to be sat... This practice were really keen on doing it all together in one room with just one camera for all of them.”* AE2

Having multiple attendees in one room, sharing a screen, could be confusing for the AE trainer: *“That was a nightmare because you just couldn’t hear anyone. You couldn’t see people. Rather than everyone being a little icon, it was one shot with everyone in. You had no idea who was saying anything. Usually, it flashes up with someone’s name when they speak. With this, you couldn’t tell if it was Doctor X or Doctor Y who was saying something.”* AE2

Amongst general practice staff receiving training, there was diverse technical ability: *“Speaking as a manager, we learn differently. So, I don’t mind learning on Teams.*



*I've done quite a lot of training in the last 12 months on Teams, courses, that kind of thing. Or being on meetings like this. So, I'm used to it. A lot of the satellite receptionists, they've always done face-to-face training. They've never had to use Microsoft Teams. They've never had to use Zoom so a lot of them might find it a bit unusual and it might not sink in the same way."* Practice manager 2

## 2. The access-engagement trade off

A marked improvement in access to training was noted by AEs delivering training, with training capturing practice staff who otherwise would not have been able to attend, for example participants joining from their home or on the school run.

One AE was surprised at how effective remote delivery was in widening access to training: *"When people are at home with the children. You see them shooing them out of the room but they were still able to do the training so yes... it was quite phenomenal; it blew me away... Because I thought, "Wow we wouldn't get these people at the training."* AE1

Remote training also allowed multiple practices to attend training at once: *"And we now can reach people from all 49 practices, so that training, where we were having different people from different practices, was a lot easier for us, a lot less time-consuming."* AE4

However, with a gain in widened access, came a perceived loss in overall engagement, impacting on opportunities for learning. Our study team noticed just

under half the participants' screen cameras were turned off during a training session and were only switched on for the initial introductions or when responding to, or asking, a question. This could affect the communication of non-verbal cues between trainer and participants, non-verbal cues which might indicate understanding, engagement, and emotional response, important when discussing a subject such as DVA.

Participants were noticeably 'multitasking' during training: answering calls, eating lunch, and using the chat function to explain when they needed to attend to an urgent task and pause training. In contrast to previous face-to-face delivery which involved regular participant interaction, in remote sessions AE trainers were the main speakers for the majority of the session time and participants were largely passive recipients.

We observed AEs using a variety of techniques to engage trainees, for example videos to illustrate key behaviours seen in DVA or 'ice-breaker' tasks. Despite these interventions there were still limits to engagement.

The trade-off in widening access to training at the expense of engagement was a concern for some AEs. One GP who is involved in training delivery explained the loss of visual feedback from participants can be challenging for the facilitator: *"I would say that it feels very removed. You can't make eye contact in the same way, you can't read the responses that GP's are having, or clinicians are having to what they are being told, which is hard stuff"*. GP1

At times the reduced engagement, compounded by technological challenges of training delivery, created stress and isolation for the facilitator: *"...with IRIS training, you play some videos and there is some kind of group work there too. And that was really hard to try and adapt to video because, you have to figure out how to kind of play the video and how can they hear it. Those little things really impact the stress levels when you're trying to do training, and especially if no one is kind of- If no one has their screens on, it feels very odd to kind of just talk to yourself."* AE3

### 3. Sensitive content in remote sessions

The challenge of discussing a sensitive subject such as DVA in a remote setting was explored. In observation of training sessions, participants were reminded that the content could potentially be distressing and were reassured that they can simply switch off their screens/sound to 'leave the room' if they needed to. This is especially important given the possibility of training participants having lived experience themselves. However, this limited the trainer's ability to "read the room" and gauge a response from the audience, meaning that there was a risk trainees could be overwhelmed by the content, with no-one close by to support them if needed.

AEs acknowledged that this was challenging content to discuss in a virtual environment: *"That's what the practice manager gave me as feedback is that perhaps, that wasn't thought about enough. You know, that it is quite a lot of overwhelming information and really, there was no way to gauge that for me...because you can't read the room literally."* AE7

One GP who delivers training shared concerns on establishing a safe remote learning environment: *“So, I think it is much easier to gauge how engaged your trainees, or the doctors you are teaching are, when you are in a room with them. And it is also much easier to set up a feeling of trust, and safety, people can ask whatever they want and there is no judgement than it is possible to do remotely. I think that probably is the main issue...”* GP1

4. Is remote training here to stay?

There was diversity of opinion amongst interview participants regarding the longevity of remote training. Remote training was viewed as an efficient option given the busy work schedules of practice staff, and for some AEs there was worry around reverting back to face-to-face training. On the other hand, some preferred a return to face-to-face training to enhance learning and improve absorption of information.

To their own surprise, one AE felt remote training had significant benefits and actually could achieve a similar experience to face-to-face: *“I am a real convert and I can't believe I am even saying that. I can't believe these words are coming out of my mouth... So, feedback-wise, I think it was good. It was as it always has been. They were asking the same sort of questions that they have always asked. So yes, I was actually really impressed with it, if I am honest. Seems to have gone really all right, yes.”* AE1

For another AE, the thought of returning to face-to-face training caused worry: *“I have built up an anxiety around going back to doing that face-to-face because it has*

*been easy to do it via Teams because I'm at home...For example, I can have my notes and it's not so formal."* AE6

Although acknowledging reduced engagement in remote training, one practice manager felt that remote training offered a significant advantage. It was felt to be a time-efficient option, complimentary to work schedules of general practice staff: *"I think it works better in some ways...One, you haven't got interaction with people there and then, but for time for people, they are not having to go places so they've got more time, then they can do the meetings, do the training, and then get on with their work."* Practice manager 1

However, another practice manager expressed that they felt remote training was inferior to face-to-face training based on attendees absorbing less information: *"So, I think it's different and not everybody will take on as much as they would if they were face-to-face, I don't think. You can't beat face-to-face."* Practice manager 2

## **Discussion**

### *Summary*

In the transition to remote DVA training in general practices during the COVID-19 pandemic, we found examples of successful adaptation to existing training content and delivery, which was well received by both general practice staff participants and trainers.

However, AEs found it challenging to schedule DVA training with busy practices and there were major adaptations in content of DVA training, with a new and proactive focus on safely addressing potential or disclosed DVA in remote consultations alongside a need to adjust to novel technology. Variable technical ability amongst general practice staff and AEs was reported, and at times some general practices did not have the required technical infrastructure. This resulted in a halfway house of remote training delivery, with practice staff sharing rooms and screens, making it more difficult for AEs to connect with staff as individuals and complicating group communication. These reports from interviews were consistent with study team observations in training sessions.

The remote model widened access to training for those based away from the practice however there was a concurrent overall loss of participant engagement. Not meeting in person reduced important non-verbal cues for the trainer to pick up on and address. This is especially important given the sensitive content of training. Interaction between participants was variable and this potentially limited opportunities for learners to discuss emotionally challenging content as a group, including the sharing of experiences and reflections. Reduced learner engagement may have limited personal interactions between AEs and general practice staff, potentially weakening the working partnership which is critical in supporting DVA referrals from general practice to specialist DVA services.

This trade-off between access and engagement, created mixed feelings amongst AEs delivering training: from surprise at the reach remote training could have, to stressful and isolating experiences for AEs facilitating the training. In our

observations it appeared that remote training was helpful in allowing general practice staff to subtly step away from sensitive content; but in interviews there were concerns that the loss of an ability to read the room in remote training, made it more difficult to gauge training, and there was a risk of overwhelming the audience. This is of particular concern, given that staff may have personal DVA experience or know others who have lived-experience.

Divergent opinions existed regarding the future of remote DVA training in general practice. Although some felt there was little to distinguish it from face-to-face training, others felt that the practice were likely to come away with less than they would have done in face-to-face settings.

#### *Comparison with existing literature*

Our findings provide insight into the experiences of delivering and receiving remote DVA training in general practice during the pandemic. They are consistent with previously reported advantages of remote training offering valuable inclusivity and widened access,<sup>21</sup> alongside concerns regarding learner isolation, technical problems and reduced engagement.<sup>21 22</sup> Through our analysis of trainer perspectives, we contribute the additional insight of the experience of trainer isolation. There is limited research in this area, although this has been reported amongst teachers in a non-medical distance learning setting during the pandemic.<sup>29</sup>

Reduced engagement creates distance between AEs and general practice staff attending remote training. This is concerning because previous research has shown

that personal interactions between GPs and AEs mean that AEs are viewed more positively.<sup>17</sup> Therefore, reduced opportunity for personal interactions between AEs and general practice staff, could negatively impact on the relationship between the general practice team and DVA specialist services. In contrast to some other types of training sessions, which may be a stand-alone or one-off session, the impacts of these DVA training sessions extend beyond the session itself and can affect the ongoing partnership between general practice and DVA specialist services.

Our findings also raise questions regarding whether remote training is an appropriate forum to discuss sensitive and emotional content, such as DVA. More research is needed in this area, although concerns have already been raised about managing safeguarding in remote general practice consultations.<sup>30</sup>

### *Strength and limitations*

This is a novel area of research and observation of training alongside interviews with those delivering and receiving training, allowed us to triangulate key findings. Generalisability of findings is limited by the study only being performed in the context of the IRIS training, support and referral programme. The general practices had specific guidance and training, as well as a referral pathway to a named specialist in a DVA service. Our findings may be less generalisable to remote training which does not link to referral pathways or are one off events. The study benefitted from valuable perspectives of AEs who deliver training and general practice staff who predominantly receive training. The study had an imbalance in the number of AEs and general practice staff interviewed, and in future would benefit from more



perspectives of the general practice teams receiving training. An additional limitation of the study was that a predominance of quotations came from two participants and the study would be strengthened by the inclusion of a greater range of perspectives.

### *Implications for practice*

Building strong relationships between general practice and DVA services is crucial in supporting patients affected by DVA. The IRIS training, support and referral programme can help to achieve this, with DVA training to general practice staff being a key component. Although remote training widens access to sessions, the loss of engagement risks weakening the partnership between general practice and DVA services. We suggest a mixed DVA training model, which includes remote information delivery alongside a supplemented face-to-face element. This would take advantage of the improved access from remote training whilst building on the general practice and DVA support service partnership in face-to-face settings.

### **Conclusion**

Further research is needed regarding remote training in primary care, including how to support those delivering training, achieving learner engagement and the safeguarding of remote learners who are DVA survivors. Our findings have relevance for other organisations and services forming partnerships with primary care, for example social prescribing and social services.

## **Funding**

This study is part of PRECODE as a 12-month cross university collaborative project funded by the UKRI (UK Research and Innovation) Rapid Response Call and the MRC (MR/V041533/1).

## **Competing interests**

MJ is chief executive of IRISi. LD is IRIS Network Director for IRISi. GF is a non-executive IRISi board member. ECB worked part-time as a data scientist for IRISi between June 2019 and March 2022.

## **Acknowledgements**

We are most grateful to the practitioners delivering IRIS in each local area, including the general practice staff, IRIS clinical leads, advocate educators, their managers, the third sector DVA host agencies and commissioners based within Primary Care, Public Health and Local Authorities within the participating boroughs. We would like to thank IRISi for supporting, guiding, and facilitating our data collection.

JPG's work is in part supported by funding from the UK Health Security Agency (UKHSA) and the UK Department of Health and Social Care. *The funders had no role in the study design, data analysis, data interpretation, or writing of this paper.*

GF and ECB receive salary contributions from the VISION research, supported by the **UK Prevention Research Partnership** (Violence, Health and Society; MR-

VO49879/1), funded by the British Heart Foundation, Chief Scientist Office of the Scottish Government Health and Social Care Directorates, Engineering and Physical Sciences Research Council, Economic and Social Research Council, Health and Social Care Research and Development Division (Welsh Government), Medical Research Council, National Institute for Health and Care Research, Natural Environment Research Council, Public Health Agency (Northern Ireland), The Health Foundation, and Wellcome. *The views expressed are those of the researchers and not necessarily those of the UK Prevention Research Partnership or any other funder.*

SD is supported by NIHR DRF (Sharon NIHR301787). The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care.

## References

1. World Health Organisation. Violence against women: a global health problem of epidemic proportions (2013), <https://www.who.int/news-room/fact-sheets/detail/violence-against-women> (accessed 27 Mar 2023).
2. Dheensa S, McLindon E, Spencer C, et al. Healthcare Professionals' Own Experiences of Domestic Violence and Abuse: A Meta-Analysis of Prevalence and Systematic Review of Risk Markers and Consequences. *Trauma Violence Abuse* 2022;15248380211061771. doi: 10.1177/15248380211061771 [published Online First: 20220103].
3. García-Moreno C, Hegarty K, d'Oliveira AFL, et al. The health-systems response to violence against women. *Lancet* 2015;385(9977):1567-79. doi: 10.1016/S0140-6736(14)61837-7.
4. Feder G, Ramsay J, Dunne D, et al. How far does screening women for domestic (partner) violence in different health-care settings meet criteria for a screening programme? Systematic reviews of nine UK National Screening Committee criteria. *Health Technol Assess*. 2009;13:16. doi: 10.3310/hta13160.
5. Richardson J, Coid J, Petrukevitch A, et al. Identifying domestic violence: cross sectional study in primary care. *BMJ* 2002;324(7332):274. doi: 10.1136/bmj.324.7332.274.
6. Britton A. Homicides, Firearm Offences and Intimate Violence 2010/11 Supplementary volume 2 to crime in England and Wales 2010/11. Home Office statistical bulletin: London: Home Office (2012), [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/116483/hosb0212.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/116483/hosb0212.pdf) (accessed 27 Mar 2023)

7. Panovska-Griffiths J SE, Johnson M, Dixon S, De Simoni A, Wileman V, Dowrick A, Emsley E, Griffiths C, Capelas Barbosa E, Feder G. Impact of the first national COVID-19 lockdown on referral of women experiencing domestic violence and abuse in England and Wales. *BMC Public Health* 2022;22(1):504. doi: 10.1186/s12889-022-12825-6.
8. Havard T. Domestic abuse and Covid-19: A year into the pandemic. House of Commons Library: UK Parliament (2021), <https://commonslibrary.parliament.uk/domestic-abuse-and-covid-19-a-year-into-the-pandemic/> (accessed 27 Mar 2023).
9. Szilassy E, Barbosa EC, Dixon S, et al. Primary care response to domestic violence and abuse in the COVID-19 pandemic (PRECODE): protocol of a rapid mixed-methods study in the UK. *BMC Fam Pract* 2021;22(1):91. doi: 10.1186/s12875-021-01447-3 [published Online First: 20210512].
10. IRIS Interventions. What we do, <https://irisi.org/what-we-do/> (accessed 27 Mar 2023).
11. Bolchover L. IRIS: Identification and Referral to Improve Safety. Improving the general practice response to domestic violence and abuse: how the IRIS programme can support your practice and your patients. IRISi: University of Bristol, (2018), [https://irisi.org/wp-content/uploads/2019/10/IRIS\\_Booklet.pdf](https://irisi.org/wp-content/uploads/2019/10/IRIS_Booklet.pdf) (accessed 27 Mar 2023).
12. Feder G, Davies RA, Baird K, et al. Identification and Referral to Improve Safety (IRIS) of women experiencing domestic violence with a primary care training and support programme: a cluster randomised controlled trial. *Lancet* 2011;378(9805):1788-95. doi: 10.1016/S0140-6736(11)61179-3.
13. Department of Health and Social Care. The Government's Mandate to NHS England for 2016-2017 (2017). [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/600604/NHSE\\_Mandate\\_2016-17.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/600604/NHSE_Mandate_2016-17.pdf).
14. Home Office. Ending Violence against Women and Girls (2016). [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/522166/VAWG\\_Strategy\\_FINAL\\_PUBLICATION\\_MASTER\\_vRB.PDF](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/522166/VAWG_Strategy_FINAL_PUBLICATION_MASTER_vRB.PDF).
15. IRIS Interventions. IRIS Programme: find your local site. <https://irisi.org/find-your-local-iris-site/> (accessed 27 Mar 2023).
16. Barbosa E.C MM, D'avo G, Downes L, Johnson M, Howell A. Identification & Referral to Improve Safety: Improving the General Practice Response to Domestic Violence and Abuse. A review of IRIS Programmes in England, Wales, the Channel Islands and Northern Ireland to March 2021. <https://irisi.org/wp-content/uploads/2022/03/IRIS-National-Report-2020-2021.pdf>: IRISi, 2022.
17. Dowrick A, Kelly M, Feder G. Boundary spanners: Negotiating connections across primary care and domestic violence and abuse services. *Soc Sci Med*. 2020;245:112687. doi: <https://doi.org/10.1016/j.socscimed.2019.112687>.
18. Sonsale A, Thorne T, Kalia R, et al. The future of general practice: brave new dawn or dark skies ahead? *Br J Gen Pract*. 2018;68(666):4. doi: 10.3399/bjgp17X693953.
19. Cooper M. Lockdown and the Future of General Practice. *BJGP Life*. 2021 (accessed 13 Apr 2022).
20. Beetham H SR. Rethinking pedagogy for a digital age: Abingdon, Oxon: Routledge 2007.
21. Reeves S, Fletcher S, McLoughlin C, et al. Interprofessional online learning for primary healthcare: findings from a scoping review. *BMJ Open* 2017;7(8):e016872. doi: 10.1136/bmjopen-2017-016872 [published Online First: 20170804].
22. Houghton N HW, Yates S, Badley B, Kneebone R. Cross-disciplinary perspectives on the transition to remote education. *BMJ Simul Technol Enhanc Learn*. 2021;7(6):586-89.
23. Dixon S, Szilassy E, et al. General Practice wide adaptations to support patients affected by DVA during the COVID-19 pandemic: a rapid qualitative study. *PREPRINT (Version 1) available at Research Square [https://doi.org/10.21203/rs.3.rs-2069950/v1]* 2022.
24. Panovska-Griffiths J, Szilassy E, Johnson M, et al. Impact of the first national COVID-19 lockdown on referral of women experiencing domestic violence and abuse in England and Wales. *BMC*

- Public Health* 2022;22(1):504. doi: 10.1186/s12889-022-12825-6 [published Online First: 20220315].
25. McNall M, Foster-Fishman PG. Methods of rapid evaluation, assessment, and appraisal. *Am J Eval*. 2007;28:151-68.
  26. Vindrola-Padros C, Chisnall G, Cooper S, et al. Carrying Out Rapid Qualitative Research During a Pandemic: Emerging Lessons From COVID-19. *Qual Health Res* 2020;30(14):2192-204. doi: 10.1177/1049732320951526 [published Online First: 20200831].
  27. Gale NK, Heath G, Cameron E, et al. Using the framework method for the analysis of qualitative data in multi-disciplinary health research. *BMC Med Res Methodol* 2013;13:117. doi: 10.1186/1471-2288-13-117 [published Online First: 20130918].
  28. Marks DF YL. *Research Methods for Clinical and Health Psychology*: SAGE Publications, Limited 2003.
  29. Parte L, Herrador-Alcaide T. Teaching Disruption by COVID-19: Burnout, Isolation, and Sense of Belonging in Accounting Tutors in E-Learning and B-Learning. *Int J Environ Res Public Health* 2021;18(19) doi: 10.3390/ijerph181910339 [published Online First: 20210930].
  30. Dixon S, Frost L, Feder G, et al. Challenges of safeguarding via remote consulting during the COVID-19 pandemic: a qualitative interview study. *Br J Gen Pract*. 2022;72(716):e199-e208. doi: 10.3399/bjgp.2021.0396