## POLITECNICO DI TORINO Repository ISTITUZIONALE

Seasonal compartmentalisation of water in a grassland at 2600 m a.s.l

Original Seasonal compartmentalisation of water in a grassland at 2600 m a.s.l / Gentile, Alessio; Bevilacqua, Ivan; Canone, Davide; Ceperley, Natalie; Gisolo, Davide; N'Sassila, Mesmer; Previati, Maurizio; Zuecco, Giulia; Schaefli, Bettina; Ferraris, Stefano (2022). (Intervento presentato al convegno EGU General Assembly 2022 tenutosi a Vienna nel 2 Maggio 2022) [10.5194/egusphere-egu22-9190].	
Availability: This version is available at: 11583/2977680 since: 2023-03-31T13:32:23Z	
Publisher: EGU	
Published DOI:10.5194/egusphere-egu22-9190	
Terms of use: openAccess	
This article is made available under terms and conditions as specified in the corresponding bibliographic description the repository	in
Publisher copyright	
(Article begins on next page)	

12 June 2023



EGU22-9190, updated on 31 Mar 2023 https://doi.org/10.5194/egusphere-egu22-9190 EGU General Assembly 2022 © Author(s) 2023. This work is distributed under the Creative Commons Attribution 4.0 License.



## Seasonal compartmentalisation of water in a grassland at 2600 m a.s.l.

**Alessio Gentile**<sup>1</sup>, Ivan Bevilacqua<sup>1</sup>, Davide Canone<sup>1</sup>, Natalie Ceperley<sup>2,3,4</sup>, Davide Gisolo<sup>1</sup>, Mesmer N'Sassila<sup>1</sup>, Maurizio Previati<sup>1</sup>, Giulia Zuecco<sup>5</sup>, Bettina Schaefli<sup>2,3,4</sup>, and Stefano Ferraris<sup>1</sup>

Interuniversity Department of Regional and Urban Studies and Planning, Politecnico and Università degli Studi di Torino,

'interuniversity Department of Regional and Orban Studies and Planning, Politecnico and Università degli Studi di Torino Torino, Italy (alessio.gentile@polito.it)

High Alpine catchments are precious water-resources since they act as natural storage reservoirs, storing water in the snow cover and in the subsurface and thereby providing water during the dry seasons. Thus, a deeper knowledge of the hydrological functioning of these systems is necessary, in particular to make climate change projections. The role of seasonality is crucial in these catchments that generally exhibit a snow-dominated hydro-climatic regime.

Here we use high-frequency observations of stable isotopes of water to identify the seasonal origin of streamwater in a high-elevation Alpine catchment located in the Valle d'Aosta Region, Italy. We quantify the relative contribution of winter and summer precipitation reaching the stream through the Seasonal Origin Index ( $SOI_Q$ ), calculated using the  $\delta^{18}O$  values and the volumes of precipitation and streamflow. Highly negative  $SOI_Q$  values are obtained suggesting that streamwater is mainly composed of winter precipitation. Conversely, the Seasonal Origin Index for evapotranspiration ( $SOI_{ET}$ ), which can be directly inferred from  $SOI_Q$ , returns a positive value reflecting that plants preferentially take up water deriving from summer precipitation.

These findings allow us to develop a conceptual model of this Alpine system. This conceptual model suggests:

- a deep infiltration component, mainly composed by snowmelt water, reaching the stream through a preferential flow.
- a shallow infiltration component, predominantly represented by summer rainfall, that dominates the shallow soils and that is used by plants.

Therefore, we presume a seasonal compartmentalisation of water in this high-elevation catchment.

<sup>&</sup>lt;sup>2</sup>Institute of Earth Surface Dynamics, Faculty of Geosciences and Environment, University of Lausanne, Lausanne, Switzerland

<sup>&</sup>lt;sup>3</sup>Institute of Geography, Faculty of Science, University of Bern, Bern, Switzerland

<sup>&</sup>lt;sup>4</sup>Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland

<sup>&</sup>lt;sup>5</sup>Department of Land, Environment, Agriculture and Forestry, University of Padova, Legnaro, Italy

Nevertheless, a previous study in Switzerland revealed  $SOI_Q \approx 0$  for the Allenbach and Dischmabach snow-dominated catchments, indicating that similar fractions of summer and winter precipitation become streamflow. This different result achieved in systems with an apparently similar functioning highlights the need for a deep insight into the flow paths governing high-elevation catchments and it opens the way for new challenges to understand the hydrological processes hidden behind this difference.