

2020

Lung nodule management practice patterns - A survey of Indian physicians

V. Mehta

P. Makkar

K. Bhatia

S. Mehta

S. Raouf

Zucker School of Medicine at Hofstra/Northwell

Follow this and additional works at: <https://academicworks.medicine.hofstra.edu/publications>




Part of the [Pulmonology Commons](#)

Recommended Citation

Mehta V, Makkar P, Bhatia K, Mehta S, Raouf S. Lung nodule management practice patterns - A survey of Indian physicians. . 2020 Jan 01; 37(4):Article 7311 [p.]. Available from: <https://academicworks.medicine.hofstra.edu/publications/7311>. Free full text article.

This Article is brought to you for free and open access by Donald and Barbara Zucker School of Medicine Academic Works. It has been accepted for inclusion in Journal Articles by an authorized administrator of Donald and Barbara Zucker School of Medicine Academic Works. For more information, please contact academicworks@hofstra.edu.



Lung India. 2020 Jul-Aug; 37(4): 350–351.

PMCID: PMC7507916

Published online 2020 Jul 1.

PMID: [32643650](#)

doi: 10.4103/lungindia.lungindia_41_20: 10.4103/lungindia.lungindia_41_20

Lung nodule management practice patterns – A survey of Indian physicians

[Vishisht Mehta](#),¹ [Priyanka Makkar](#),¹ [Karishma Bhatia](#),² [Sanjeev Mehta](#),³ and [Suhail Raof](#)⁴

¹*Department of Medicine, Division of Pulmonary Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA. E-mail: pri.makkar@gmail.com*

²*Department of Pulmonary Critical Care, Mount Sinai St. Luke's/Beth Israel, New York, NY, USA*

³*The Chest Center, Mumbai, Maharashtra, India*

⁴*Department of Pulmonary and Critical Care Medicine, Lenox Hill Hospital, New York, NY, USA*

Received 2020 Jan 21; Revised 2020 Jan 28; Accepted 2020 Jan 30.

[Copyright](#) : © 2020 Indian Chest Society

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Sir,

A lung nodule is a rounded opacity, well or poorly defined, measuring up to 3 cm in diameter.[1] The majority of such lesions are benign, but malignancy needs to be excluded in most. Nodules may be detected incidentally or by screening. Indeed, lung cancer screening has demonstrated mortality benefits in the national lung screening trial in the United States.[2] Applying these recommendations to the Indian population is controversial since there are no evidence-based guidelines on lung nodule detection and management that take into consideration indigenous factors such as prevalence of lung cancer and available local resources. It may be speculated that implementing consensus guidelines for lung nodule management in Asia[3] may more closely mimic the Indian population. We conducted a survey in three metropolitan cities across India to identify perspectives of physicians managing lung nodules.

We surveyed 210 participants, of whom 165 (78.5%) answered every question. Of those answering, 144 (72.3%), 18 (9%), and 12 (6%) respondents were practicing pulmonologists, internists, and general practitioners, respectively. Furthermore, 121 (73.3%), 108 (58%), and 140 (67.3%) respondents were aware of the size ranges of micronodules, nodules, and masses, respectively. Further, 165 (78.5%) respondents answered that they saw up to five patients a month with incidental nodules. In addition, 140 (70%) physicians felt on clinical grounds that up to 10% of their patients had multiple nodules attributable to the prior granulomatous disease. [Table 1](#) indicates the distribution of responses by physicians of percentage of patients with an eventual histologic diagnosis of the nodules. Majority of respondents reported using either chest or Fleischner Society guidelines for nodule management. Majority of nodules

were visualized using physical films ($n = 131$; 74%) for radiographic imaging. A total of 70 (35.9%) respondents utilized a multidisciplinary approach (1 pulmonologist and any two specialists from radiology, medical oncology and/or surgery).

Through this survey, we aim to highlight the heterogeneity in practice patterns of physicians managing lung nodules in India. Intuitively, formulating guidelines for lung cancer screening and incidental nodules should reduce this variability. However, the situation is complex with a great deal of variability in patient education, perspectives, preferences, and available resources. Furthermore, the prevalence of nodules may vary greatly due to endemic granulomatous diseases, which may make it more difficult to make recommendations for tissue diagnosis.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Hansell DM, Bankier AA, MacMahon H, McLoud TC, Müller NL, Remy J. Fleischner society: Glossary of terms for thoracic imaging. *Radiology*. 2008;246:697–722. [PubMed: 18195376]
2. National Lung Screening Trial Research Team. Aberle DR, Adams AM, Berg CD, Black WC, Clapp JD, et al. Reduced lung-cancer mortality with low-dose computed tomographic screening. *N Engl J Med*. 2011;365:395–409. [PMCID: PMC4356534] [PubMed: 21714641]
3. Bai C, Choi CM, Chu CM, Anantham D, Chung-Man Ho J, Khan AZ, et al. Evaluation of pulmonary nodules: Clinical practice consensus guidelines for asia. *Chest*. 2016;150:877–93. [PubMed: 26923625]

Figures and Tables

Table 1

Eventual histologic diagnosis of lung nodules - Distribution of responses

Percentage patients with histologic diagnosis	Number of respondents, <i>n</i> (%)
0-5	74 (39.3)
6-10	49 (26)
11-15	19 (10.1)
>15	46 (24.4)

Articles from Lung India : Official Organ of Indian Chest Society are provided here courtesy of **Wolters Kluwer -- Medknow Publications**