

2019

Myocardial fibrosis and arrhythmogenesis in elite athletes

S. A. Ahmad

N. Khalid

E. Shlofmitz

Zucker School of Medicine at Hofstra/Northwell

L. Chhabra

Follow this and additional works at: <https://academicworks.medicine.hofstra.edu/publications>



Part of the [Cardiology Commons](#)

Recommended Citation

Ahmad SA, Khalid N, Shlofmitz E, Chhabra L. Myocardial fibrosis and arrhythmogenesis in elite athletes. . 2019 Jan 01; 42(9):Article 4882 [p.]. Available from: <https://academicworks.medicine.hofstra.edu/publications/4882>. Free full text article.

This Article is brought to you for free and open access by Donald and Barbara Zucker School of Medicine Academic Works. It has been accepted for inclusion in Journal Articles by an authorized administrator of Donald and Barbara Zucker School of Medicine Academic Works. For more information, please contact academicworks@hofstra.edu.

LETTER TO THE EDITOR

Myocardial fibrosis and arrhythmogenesis in elite athletes

To the Editor,

Cuspidi et al present an excellent meta-analysis of nine studies involving 403 elite athletes and 297 controls investigating volumetric and strain analysis of left atrial (LA) phasic function.¹ Salient conclusions of their analysis include that the LA volume is higher and LA reservoir and contractile functions are impaired in elite athletes during active training when compared to the controls. The authors also conclude that the LA remodeling and contractile dysfunction may provide a substrate for the development of atrial and ventricular tachyarrhythmias. We wish to highlight a few additional points.

In addition to the LA remodeling that contributes to arrhythmogenesis, there is substantial evidence that endurance athletes are prone to develop myocardial fibrosis (patchy, interstitial, or dense scar) which can also trigger arrhythmias.^{2,3} The myocardial fibrosis can be focal or diffuse and can develop in the right or left ventricle, septum, papillary muscles, or bundle branches.² Cardiac magnetic resonance imaging using late gadolinium enhancement has detected myocardial fibrosis in endurance athletes.⁴ In addition, these findings have been supported by murine models and small post-mortem studies on human subjects.² Cardiac fibroblasts are normal cell types occurring in healthy hearts and are responsible for extracellular matrix homeostasis, but upon exposure to injury, mechanical stress, or inflammation, these cells may transform into myofibroblasts and induce cardiac fibrosis.³ Sequelae of this pathologic remodeling may include myocardial stiffness (increasing ventricular end-diastolic pressures), chamber dilation, hypertrophy, and ultimately heart failure.^{2,3}


The cardiovascular benefits of exercise are well accepted but the tipping point where the physiologic remodeling becomes pathologic is yet somewhat unclear. The authors are to be commended for adding to our fund of knowledge for this burgeoning field. Characterization of at-risk elite athletes is important because of the risk of arrhythmias and consequently, sudden cardiac death as these may be preventable. A risk prediction model should be established incorporating LA volumetric and strain data from the present study, myocardial fibrosis from the imaging studies, and other established criteria. Future studies should also aim to provide a risk prediction clinical tool to decrease the arrhythmia related morbidity and mortality in elite athletes.

CONFLICT OF INTEREST

The authors declare no potential conflict of interests.

ORCID

Nauman Khalid  <https://orcid.org/0000-0003-4159-555X>

Sarah A. Ahmad¹
Nauman Khalid² 
Evan Shlofmitz²
Lovely Chhabra³

¹Department of Cardiovascular Surgery, Saint Francis Medical Center, Monroe, Louisiana

²Section of Interventional Cardiology, MedStar Washington Hospital Center, Washington, District of Columbia

³Department of Cardiology, Heartland Regional Medical Center, Marion, Illinois

Correspondence

Nauman Khalid, MD, MedStar Washington Hospital Center, 110 Irving St., NW, Suite 4B-1X, Washington DC 20010.
Email: nauman.khalid@medstar.net

REFERENCES

1. Cuspidi C, Tadic M, Sala C, Gherbesi E, Grassi G, Mancia G. Left atrial function in elite athletes: a meta-analysis of two-dimensional speckle tracking echocardiographic studies. *Clin Cardiol*. 2019;42:579-587.
2. van de Schoor FR, Aengevaeren VL, Hopman MT, et al. Myocardial fibrosis in athletes. *Mayo Clin Proc*. 2016;91:1617-1631.
3. Travers JG, Kamal FA, Robbins J, Yutzey KE, Blaxall BC. Cardiac fibrosis: the fibroblast awakens. *Circ Res*. 2016;118:1021-1040.
4. Breuckmann F, Möhlenkamp S, Nassenstein K, et al. Myocardial late gadolinium enhancement: prevalence, pattern, and prognostic relevance in marathon runners. *Radiology*. 2009;251:50-57.