



Outcomes of coronary revascularization versus optimal medical therapy alone for ischemic left ventricular dysfunction: A meta-analysis of randomized controlled trials

Authors: Kamil Bujak, Bartosz Hudzik, Łukasz Pyka, Michał Skrzypek, Salvatore Brugaletta,
Manel Sabaté, Mateusz Tajstra, Jacek Legutko, Wojciech Wojakowski, Mariusz Gąsior

Article type: Short communication

Received: February 18, 2023

Accepted: April 14, 2023

Early publication date: June 2, 2023

This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

Outcomes of coronary revascularization versus optimal medical therapy alone for ischemic left ventricular dysfunction: A meta-analysis of randomized controlled trials

Short title: Revascularization in patients with ischemic left ventricular dysfunction

Kamil Bujak^{1, 2}, Bartosz Hudzik¹, Łukasz Pyka¹, Michał Skrzypek³, Salvatore Brugaletta², Manel Sabaté², Mateusz Tajstra¹, Jacek Legutko^{4, 5}, Wojciech Wojakowski⁶, Mariusz Gąsior¹

¹3rd Department of Cardiology, Faculty of Medical Sciences in Zabrze, Medical University of Silesia, Katowice, Poland

²Hospital Clinic, Cardiovascular Clinic Institute, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), University of Barcelona, Barcelona, Spain

³Department of Biostatistics, Faculty of Health Sciences in Bytom, Medical University of Silesia, Katowice, Poland

⁴Clinical Department of Interventional Cardiology, John Paul II Hospital, Kraków, Poland

⁵Department of Interventional Cardiology, Institute of Cardiology, Jagiellonian University Medical College, Kraków, Poland

⁶Department of Cardiology and Structural Heart Diseases, Medical University of Silesia, Katowice, Poland

Correspondence to:

Kamil Bujak, MD,

3rd Department of Cardiology,

Faculty of Medical Sciences in Zabrze,

Medical University of Silesia,

M Skłodowskiej-Curie 9, 41–800 Zabrze, Poland,

phone: +48 32 37 33 860,

e-mail: k.bujak@sccs.pl

INTRODUCTION

Coronary artery disease (CAD) is the most frequent cause of heart failure with reduced ejection fraction worldwide. The observations that a large proportion of patients with ischemic heart failure have areas of dysfunctional-yet-viable myocardium have led to the hypothesis that

coronary revascularization might improve left ventricular function and outcomes in this population [1].

Randomized controlled trials (RCTs) published in recent years did not demonstrate a significant advantage of routine coronary revascularization in patients with stable CAD over optimal medical therapy (OMT) [2]. However, patients with left ventricular systolic dysfunction (LVSD), who might potentially benefit the most from revascularization, were mainly excluded from these trials. Only a few RCTs compared coronary revascularization with OMT alone in patients with severe LVSD.

To the best of our knowledge, there is no meta-analysis that synthesizes the results of these trials. Therefore, we aimed to perform a meta-analysis comparing outcomes following coronary revascularization (both percutaneous and surgical) in comparison to OMT alone in patients with LVSD based on the latest available evidence from RCTs.

METHODS

This systematic review was prospectively registered in the PROSPERO (The International Prospective Register of Systematic Reviews) database (CRD42022340212) and conforms to the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines [3].

PubMed and Scopus were systematically searched for original articles published in English until December 8, 2022. The search strategy is presented in Supplementary material, *Table S1*. Articles were eligible for inclusion in this meta-analysis if they presented results of RCTs comparing coronary revascularization (coronary bypass surgery [CABG] or percutaneous coronary intervention [PCI]) vs. OMT alone in patients with severe LVSD (left ventricular ejection fraction of 35% or less). If multiple reports from the same RCTs were available, papers presenting the longest follow-up were included in the meta-analysis.

The following data have been extracted from eligible reports: clinical trial name, publication year, sample size, inclusion and exclusion criteria, mode of revascularization, data on the baseline and angiographic characteristics, events rates, hazard ratios (HRs) with corresponding 95% confidence intervals (CI). Subsequently, the included studies were assessed for bias using the Cochrane risk-of-bias tool for randomized trials Version 2 (RoB 2). Any discrepancies between two co-authors who independently searched for eligible papers, extracted data, and assessed data for bias were resolved by consensus.

The primary outcome of interest was cardiovascular death. Secondary outcomes included death from any cause, and death from any cause or hospitalization for heart failure. All analyzed endpoints were defined according to the studies' protocol.

Statistical analysis

Random effects inverse variance meta-analysis was conducted based on estimates (i.e., log HR) and standard errors. Log HR and standard errors were calculated from HRs and corresponding 95% CI extracted from analyzed reports. If HRs and 95% CIs were unavailable, estimates and standard errors were calculated using reconstructed individual patient data from Kaplan-Meier survival curves using the freely available online tool: *IPDfromKM Shiny app* (<https://www.trialdesign.org/one-page-shell.html#IPDfromKM>). Heterogeneity was tested using Cochrane Q statistics. Publication bias was not assessed due to the small number of included studies. All statistical analyses were performed in R version 4.2.0 (R Core Team. R: A Language and Environment for Statistical Computing, <https://www.r-project.org>) with package *meta*. Relative treatment effects were presented as HR with 95% CI. A two-tailed *P*-value of <0.05 was considered significant.

RESULTS AND DISCUSSION

An electronic search revealed 4 762 records, and after removing duplicates, the titles and abstracts of 3 499 records were screened for eligibility. Nineteen records were selected for full-text assessment, and 3 RCTs that enrolled 2 050 patients followed up for a weighted mean of 7.3 years fulfilled the inclusion criteria of this meta-analysis [4–6]. PRISMA flow chart is presented in Supplementary material, *Figure S1*, and details on the included studies are presented in Supplementary material, *Table S2*. The risk of bias was low in all included studies. The baseline characteristics of patients included in these trials are summarized in Supplementary material, *Table S3*.

Two of the three included reports provided data on the primary endpoint of cardiovascular death. Coronary revascularization was associated with a reduced risk of primary endpoint compared to OMT alone (HR, 0.81; 95% CI, 0.70–0.94; *P* <0.01); (**Figure 1A**). There was also a trend toward a lower risk of death from any cause in patients who underwent revascularization (HR, 0.88; 95% CI, 0.78–1.01; *P* = 0.06); (**Figure 1B**). However, there was no difference between treatment strategies regarding the composite endpoint of death from any cause or hospitalization for heart failure (**Figure 1C**). Event rates according to study groups are

presented in Supplementary material, *Table S4*. No significant statistical heterogeneity was identified regarding any of the analyzed outcomes.

The main finding of our meta-analysis is that coronary revascularization might be associated with improved survival, mainly driven by reduced cardiovascular mortality in patients with severe LVSD. This finding is in line with the data from observational studies, which were summarized in the recent meta-analysis [1]. However, some important limitations should be acknowledged. First, the results of only three RCTs comparing revascularization with OMT have been published to date. The STICHES trial, an extended follow-up study of the STICH trial, which had the most significant impact on the pooled estimates for all analyzed endpoints in this meta-analysis, evaluated only surgical revascularization. This trial demonstrated a reduced mortality rate in revascularized patients at ten years of follow-up. The REVIVED-BCIS2 trial, which compared OMT to PCI, demonstrated similar efficacy in terms of the primary endpoint of death from any cause or hospitalization for heart failure. Only the HEART trial studied both modes of revascularization in the invasive strategy arm but enrolled only 138 of the planned 800 patients because of the withdrawal of funding.

An open question remains whether the benefit of both modes of revascularization in patients with LVSD is similar. Contemporary RCTs have shown the superiority of CABG over PCI in patients with higher disease burden and lesion complexity, which is often the case in patients with ischemic heart failure [7]. However, patients with severe LVSD were underrepresented or excluded from these trials. Because severe LVSD and high comorbidity burden accompanying heart failure strongly increase perioperative risks, the results of these trials should not be translated to patients with severely impaired ventricular function. Unfortunately, no RCTs compared PCI against CABG in this population to date. The only available evidence comes from observational studies, which showed similar all-cause mortality in patients treated with PCI using drug-eluting stents in comparison to CABG [1].

Second, most of the analyzed patients in this meta-analysis were enrolled in the RCTs over a decade ago. Meanwhile, substantial progress in OMT was made. This might diminish the potential benefits of a revascularization strategy. On the other hand, the outcomes of patients treated invasively, mainly with PCI, improved as well, owing to broader utilization of newer generation stents and physiology- and imaging-guided revascularization [8].

Finally, considering the small number of included RCTs, statistical tools used in meta-analysis might be underpowered to assess between-study heterogeneity. For the same reason, we were unable to perform any meta-regression or subgroup analyses to identify the groups of patients who benefit the most from revascularization.

In conclusion, coronary revascularization on top of OMT seems to be associated with reduced cardiovascular mortality in patients with severely impaired left ventricular function. However, whether this effect is independent of the mode of revascularization remains unclear.

Supplementary material

Supplementary material is available at https://journals.viamedica.pl/kardiologia_polska.

Article information

Acknowledgments: The first author (K.B.) was supported by the 2021 EAPCI Education and Training Grant. The authors have not received any specific funding for conducting this review.

Conflict of interest: None declared.

Funding: None.


Open access: This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, which allows downloading and sharing articles with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially. For commercial use, please contact the journal office at kardiologiapolska@ptkardio.pl.

REFERENCES


1. Yokoyama Y, Fukuhara S, Mori M, et al. Network meta-analysis of treatment strategies in patients with coronary artery disease and low left ventricular ejection fraction. *J Card Surg.* 2021; 36(10): 3834–3842, doi: [10.1111/jocs.15850](https://doi.org/10.1111/jocs.15850), indexed in Pubmed: [34310729](https://pubmed.ncbi.nlm.nih.gov/34310729/).
2. Nguyen DD, Spertus JA, Alexander KP, et al. Management of Coronary Disease in Patients with Advanced Kidney Disease. *N Engl J Med.* 2020; 382(17): 1608–1618, doi: [10.1056/NEJMoa1915925](https://doi.org/10.1056/NEJMoa1915925), indexed in Pubmed: [32227756](https://pubmed.ncbi.nlm.nih.gov/32227756/).
3. Page MJ, McKenzie JE, Bossuyt PM, et al. Updating guidance for reporting systematic reviews: development of the PRISMA 2020 statement. *J Clin Epidemiol.* 2021; 134(3): 103–112, doi: [10.1016/j.jclinepi.2021.02.003](https://doi.org/10.1016/j.jclinepi.2021.02.003), indexed in Pubmed: [33577987](https://pubmed.ncbi.nlm.nih.gov/33577987/).
4. Perera D, Clayton T, O'Kane PD, et al. Percutaneous revascularization for ischemic left ventricular dysfunction. *N Engl J Med.* 2022; 387(15): 1351–1360, doi: [10.1056/NEJMoa2206606](https://doi.org/10.1056/NEJMoa2206606), indexed in Pubmed: [36027563](https://pubmed.ncbi.nlm.nih.gov/36027563/).

5. Velazquez EJ, Lee KL, Jones RH, et al. Coronary-Artery bypass surgery in patients with ischemic cardiomyopathy. *N Engl J Med.* 2016; 374(16): 1511–1520, doi: [10.1056/NEJMoa1602001](https://doi.org/10.1056/NEJMoa1602001), indexed in Pubmed: [27040723](https://pubmed.ncbi.nlm.nih.gov/27040723/).
6. Cleland JGF, Calvert M, Freemantle N, et al. The Heart Failure Revascularisation Trial (HEART). *Eur J Heart Fail.* 2011; 13(2): 227–233, doi: [10.1093/eurjhf/hfq230](https://doi.org/10.1093/eurjhf/hfq230), indexed in Pubmed: [21156659](https://pubmed.ncbi.nlm.nih.gov/21156659/).
7. Thuijs DJ, Davierwala P, Milojevic M, et al. Percutaneous coronary intervention versus coronary artery bypass grafting in patients with three-vessel or left main coronary artery disease: 10-year follow-up of the multicentre randomised controlled SYNTAX trial. *Lancet.* 2019; 394(10206): 1325–1334, doi: [10.1016/S0140-6736\(19\)31997-X](https://doi.org/10.1016/S0140-6736(19)31997-X), indexed in Pubmed: [31488373](https://pubmed.ncbi.nlm.nih.gov/31488373/).
8. Banning AP, Serruys P, De Maria GL, et al. Five-year outcomes after state-of-the-art percutaneous coronary revascularization in patients with de novo three-vessel disease: final results of the SYNTAX II study. *Eur Heart J.* 2022; 43(13): 1307–1316, doi: [10.1093/eurheartj/ehab703](https://doi.org/10.1093/eurheartj/ehab703), indexed in Pubmed: [34617993](https://pubmed.ncbi.nlm.nih.gov/34617993/).

Death from cardiovascular causes



Death from any cause



Death from any cause or hospitalization for heart failure




Figure 1. Forest plots presenting the meta-analysis results for primary (A) and secondary outcomes (B, C)

Abbreviations: HEART, The Heart Failure Revascularisation Trial; REVIVED-BCIS2, Revascularization for Ischemic Ventricular Dysfunction Trial; seTE, standard error of treatment estimate; STICHES, The Surgical Treatment for Ischemic Heart Failure Extension Study; TE, estimate of treatment effect