



Positive left atrial remodeling in patients with paroxysmal atrial fibrillation after a successful radiofrequency pulmonary vein isolation

Authors: Joanna Wieczorek, Katarzyna Mizia-Stec, Małgorzata Cichoń, Piotr Wieczorek,
Iwona Woźniak-Skowarska, Hoffmann Andrzej, Anna M Wnuk-Wojnar, Krzysztof Szydło

Article type: Original article

Received: October 7, 2022

Accepted: March 13, 2023

Early publication date: April 25, 2023

This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

Positive left atrial remodeling in patients with paroxysmal atrial fibrillation after a successful radiofrequency pulmonary vein isolation

Short title: Left atrial remodeling after a successful pulmonary vein isolation

Joanna Wieczorek, Katarzyna Mizia-Stec, Małgorzata Cichoń, Piotr Wieczorek, Iwona Woźniak-Skowerska, Andrzej Hoffmann, Anna M Wnuk-Wojnar, Krzysztof Szydło

1st Department of Cardiology, School of Medicine in Katowice, Medical University of Silesia, Katowice, Poland

Correspondence to:

Joanna Wieczorek, MD, PhD,
1st Department of Cardiology,
School of Medicine in Katowice,
Medical University of Silesia,
Ziolowa 47, 40-635 Katowice, Poland,
phone: +48 32 359 88 90,
e-mail: wie.joanna@gmail.com

WHAT'S NEW?

The aim was to determine the influence of initial and after follow-up standard and novel echocardiographic parameters of left atrial (LA) morphology and function on the effectiveness of the pulmonary vein isolation with RF energy (PVI) in patients with paroxysmal atrial fibrillation (AF). The presented study is unique in terms of the comprehensiveness of the detailed echocardiographic assessment of the left atrium (both morphology and deformations - segmental and global) in the entire study group. Although some of the echocardiographic parameters describing the morphology of the LA improve after a successful PVI procedure, there is still no single parameter to clearly define the possibility of recurrence of arrhythmia after PVI. Echocardiographic assessment should be complex, using various available imaging techniques and potentially useful parameters to assess the morphology and function of the heart chambers. Our observations might be reflected in a daily clinical practice and turn out to be a point of interest for a large group of the physicians and investigators focused on AF, especially

in connection with long-term effectiveness of ablation treatment and detailed echocardiographic examination.

ABSTRACT

Background: The potential relationship between the initial left atrial (LA) echocardiographic parameters and LA remodeling after pulmonary vein isolation with RF energy (PVI) with effectiveness of this treatment is discussed.

Aim: To determine the influence of initial and after follow-up transthoracic echocardiography derived predictors of successful PVI in patients with paroxysmal atrial fibrillation (AF).

Methods: 80 patients with paroxysmal AF (58 [interquartile range, IQR], 50–63] years, males: 58 [IQR, 50–63]), hospitalized for the first time PVI procedure were included. Before and after a minimum of 6 months of follow-up period a clinical and echocardiographic evaluation were performed. LA morphological parameters (diameter, volumes and other detailed LA parameters), as well as LA peak segmental and global longitudinal strains (PLS) and LA wall strain synchrony were assessed.

Results: In the whole group after a follow-up period, patients presented higher mean LA Vol_{conduit}. Patients with no AF recurrences had lower post-PVI LA volumes, higher LA ejection fraction and LA expansion index, when compared to the patients after ineffective PVI. Patients who maintained sinus rhythm after PVI procedure were characterized by a higher initial segmental strains: LA PLS_{basal-inferior} and PLS_{apical-septal}, as well as higher LA wall strain dispersion in time.

Conclusions: Some echocardiographic parameters related to LA morphology improve after successful PVI treatment. LA strains and wall strain dispersion in time are not related to LA remodeling after successful PVI procedure. However the baseline LA standard and novel echocardiographic parameters cannot be used as a remote evaluation of the effectiveness of the PVI procedure.

Key words: atrial fibrillation, left atrium, pulmonary vein isolation, strain

INTRODUCTION

Atrial fibrillation (AF) is a common supraventricular arrhythmia resulting mainly from progressive unfavourable remodeling of the left atrium (LA). Additionally, in the course of AF, the negative electrical and structural remodeling of LA is consolidated and intensified at the same time [1, 2]. LA remodeling is therefore a complex process, simply defined as a change in the size and function of LA. The observed structural changes in the LA do not always correlate

with the changes observed during myocardial electrical remodeling in patients with recurrent AF [3]. It was found that LA dysfunction assessed using the 2-dimensional speckle-tracking echocardiography (STE) method may precede visible structural changes in the atria, also in potentially healthy individuals [4–6].

The complete isolation of pulmonary veins by linear lesions around their antrum is currently recognized as the most effective method of AF treatment, which is reflected in the growing number of procedures performed, also in Poland [7, 8]. Actually, in the light of the latest 2020 European Society of Cardiology (ESC) guidelines on the treatment of AF, several randomized controlled trials and observational studies, pulmonary vein isolation with radiofrequency energy (PVI) or cryoballoon ablation are comparable methods of AF treatment in terms of effectiveness and possible complications, mostly as the first procedure [8, 9].

The pre-procedural echocardiographic assessment, taking into account both the LA morphology (analysis of dimensions and volume: passive, active and phase) and LA function (analysis of global and segmental strains), may be helpful in assessing the relationship between the effectiveness of the PVI procedure and early LA dysfunction in the group of patients with paroxysmal AF [10–13]. In turn, post-procedural assessment will contribute to a more complete understanding of LA remodeling processes depending on the maintaining sinus rhythm.

The aim of the study was to determine the influence of initial and after follow-up transthoracic echocardiography derived predictors of successful PVI in patients with paroxysmal AF undergoing ablation procedure for the first time.

METHODS

Eighty patients with diagnosed paroxysmal non-valvular AF, that were hospitalized in the Department of Cardiology between 2013 and 2017 in order to perform for the first-time PVI procedure were enrolled into the study.

The standard inclusion criteria were: documented paroxysmal symptomatic non-valvular AF (European Heart Rhythm Association [EHRA] scale, IIb–III) despite the optimal treatment and qualified for the PVI, adequate anticoagulant therapy before admission, maintaining sinus rhythm during hospitalization, preserved left ventricular systolic function (LVEF \geq 50%), written informed consent and $>$ 18 years of age.

We excluded patients with: a history of any artery pathology (stenosis (defined as arterial stenosis \geq 50% in the NASCET [14], vasculitis or dissection), connective tissue disease, a history of stroke or transient ischemic attack (TIA) in the past, structural heart disease (cardiomyopathies, significant valvular heart disease), states connected with hypercoagulation

or a predisposition to systemic embolism, a history of PVI in the past, pregnancy, refusal to participate, acute kidney disease, chronic kidney disease with glomerular filtration (GFR) <30 ml/min/1.73 m².

Informed written consent was obtained from each patient. The study protocol was approved by the Bioethical Committee of the Medical University of Silesia and performed according to the ethical guidelines of the 1975 Declaration of Helsinki.

The study group was evaluated during hospitalization before the procedure and after a follow-up period. In the current study, up to six months after the ablation procedure, a telephone conversation was conducted with the patient to assess the arrhythmia sensation and the pharmacotherapy used. Due to the assessment of multiple endpoints, clinical evaluation was performed after a minimum of 6 months to maximum 12 months after PVI. This is the time necessary, among other things, for full healing of damaged tissue after PVI with RF energy. On the other hand, It was necessary due to the technical conduct of Holter ECG monitoring for 7 days in each patient and other examinations resulting from multi-endpoint study design.

On admission, a detailed medical history that included the current course of the disease, the main symptoms (classified in EHRA class), concomitant diseases (including coronary artery disease, type 2 diabetes mellitus, arterial hypertension, hyperlipidemia, peripheral artery disease), a familial history of arrhythmia, current pharmacotherapy (especially compliance using oral anticoagulants) and tobacco smoking was collected from each subject. We also collected physical examination parameters: weight, height and body mass index (BMI), body surface area (BSA). Before ablation procedure each patient underwent in-hospital Holter electrocardiogram (ECG) monitoring (24 hours), as well as transthoracic and transesophageal echocardiography.

After a minimum of 6 months of follow-up period transthoracic echocardiography and Holter ECG home monitoring (7 days) were performed in all subjects. Furthermore, medical history, including possible recurrence of arrhythmia and pharmacotherapy was taken.

Holter ECG monitoring

Holter ECG recordings were made using Lifecard CF recorders and the recordings were assessed using the Del Mar Reynolds Sentinel system. The registration was made during the day immediately preceding the PVI procedure. Registration after the observation period was carried out at least 7 months after the procedure using the 7-day option with ECG recording at home. An episode of AF was considered to be this arrhythmia lasting >30 seconds, two episodes

of AF at the same time separated by sinus rhythm lasting <30 seconds were considered as one episode.

Transthoracic and transesophageal echocardiography

On admission, ECG-gated transthoracic and transesophageal echocardiography were performed in all patients. An experienced physician took all of the measurements during sinus rhythm, using the same investigation protocol and techniques in order to reduce inter- and intra-observer variability. The echocardiography investigation was performed using a VIVID 7 echocardiographic devices — General Healthcare equipment with a 2.5 MHz sector ultrasound transducer for transthoracic, while a 2–7 MHz for transesophageal echocardiography. The examination was stored for further analysis.

Evaluation of LA echocardiographic parameters of morphology and function

The LA_{diameter} was measured in the LAX view, while all LA volumes were measured in the 4CH apical view.

Assessed LA passive volumes included:

- Pre-atrial contraction LA volume — LA Vol_{preA};
- Minimal LA volume — LA Vol_{min};
- Maximal LA volume — LA Vol_{max};

Assessed LA active volumes included:

- LA reservoir volume — LA Vol_{reservoir};
- LA conduit volume — LA Vol_{conduit};
- LA passive emptying volume — LA Vol_{passive emptying};
- LA contractile volume — LA Vol_{contractile};

Other parameters were calculated from the volumes of passive and active LA:

- LA ejection fraction — LA EF;
- LA expansion index LA — LA_{expansion index};
- LA active emptying fraction — LA_{active empt frac};
- LA passive emptying fraction — LA_{passive empt frac}.

Previously recorded echocardiographic images in DICOM format were analyzed using an external workstation equipped with EchoPAC PC Dimension software (version 7.1.2 by General Electrics Healthcare), enabling semi-automatic deformation analysis.

Grayscale imaging for two chamber (2CH) and four chamber (4CH) projections were obtained with a frame rate of 60–80 Hz per second. The line along the endocardium was manually drawn starting at the endocardial border of the mitral ring and along the endocardial-lumen boundary of LA excluding the pulmonary veins to the opposite side of the mitral ring. An additional epicardium line was automatically generated by the software, creating a region of interest (ROI). After manually adjusting the ROI shape, in both the for four and two chamber projections, the software split the LA into six segments and generated longitudinal strain curves. In the curve analysis, the zero point was considered to be the beginning of the P-wave timing [15, 16]. During the systolic phase of LA, segmental and global longitudinal peak LA strains were assessed.

Maximum LA peak longitudinal strain (LA PLS) corresponding to LA segments and maximum LA global PLS (LA PGLS), were measured in the LA contraction phase in the two chamber — 2CH (Figure 1) and four chamber — 4CH (Figure 2) projections:

- Basal for the inferior wall (basal-inferior) — $LA\ PLS_{bas-inf\ 2CH}$
- Medial for the inferior wall (medial-inferior) — $LA\ PLS_{med-inf\ 2CH}$
- Apical for the inferior wall (apical-inferior) — $LA\ PLS_{api-inf\ 2CH}$
- Basal for the anterior wall (basal-anterior) — $LA\ PLS_{bas-ant\ 2CH}$
- Medial for the anterior wall (medial-anterior) — $LA\ PLS_{med-ant\ 2CH}$
- Apical for the anterior wall (apical-anterior) — $LA\ PLS_{api-ant\ 2CH}$
- global for all segments — $LA\ PGLS_{2CH}$
- Basal for the lateral wall (basal-lateral) — $LA\ PLS_{bas-lat\ 4CH}$
- Medial for the lateral wall (medial-lateral) — $LA\ PLS_{med-lat\ 4CH}$
- Apical for the lateral wall (apical-lateral) — $LA\ PLS_{api-lat\ 4CH}$
- Basal for the septal wall (basal-septal) — $LA\ PLS_{bas-sept\ 4CH}$
- Medial for the septal wall (medial-septal) — $LA\ PLS_{med-sept\ 4CH}$
- Apical for the septal wall (apical-septal) — $LA\ PLS_{api-sept\ 4CH}$
- global for all segments — $LA\ PGLS_{4CH}$

In addition, the LA segmental wall strain dispersion in time [ms] (LA synchrony) in the 2CH and 4CH views were assessed and defined as the difference between the earliest and the latest maximum longitudinal LA strains for individual segments.

PVI procedure

At the beginning of the PVI procedure, rotational LA angiography was performed. Then, after transseptal puncture, 3-dimensional electro-anatomical mapping was performed using the CARTO®3 system (Biosense Webster, Diamond Bar, CA, US). PVI was obtained by RF radiofrequency ablation with a ThermoCool® SmartTouch® SF catheter (Biosense Webster, Diamond Bar, CA, US). The procedure was performed using a Lasso electrode (Biosense Webster, Diamond Bar, CA, US) or Achieve (Medtronic, MN, US).

Immediately after transseptal puncture, all patients received a continuous infusion of unfractionated heparin (2000 IU/h) (preceded by an intravenous bolus of unfractionated heparin (100 IU/kg)) to obtain activated clotting time (ACT) above 300 seconds.

Statistical analysis

Statistical analysis was performed using STATISTICA software (version 13.1 PL). All data was collected in a Microsoft Office Excel spreadsheet (version 2016 PL). A *P*-value of less than 0.05 was considered to indicate statistical significance. Results for continuous variables are presented as the mean with standard deviation for normal distributions or the median with interquartile range for non-normal distributions. The normality of the distribution of continuous variables was verified with the Shapiro-Wilk test. Ordinal variables in tables are shown as absolute numbers and percentages. Depending on the distribution of variables, for pre- and post-PVI comparisons, parametric (T test for dependent variables) and non-parametric (Wilcoxon matched pairs signed rank test) tests for dependent variables were used. For separate groups (effective/ineffective PVI) parametric (T test) and non-parametric (Mann–Whitney U test) tests for independent variables were used.

RESULTS

The study group characteristics

The median (IQR) follow-up was 9.9 (7.6–11.8) months. After the observation period, the effectiveness of the PVI procedure (sinus rhythm maintenance confirmed by 7-day Holter ECG examination) was confirmed in 53.8% of patients. **Table 1** presents the basic parameters characterizing the study group. Patent foramen ovale was found in 23 patients (28.2%), while patients with LA appendage thrombus were excluded from the study.

Echocardiographic evaluation before PVI procedure and after the observation period

The analyzed LA echocardiographic parameters are summarized in **Table 2**.

After the follow-up period, patients had statistically significant higher mean LA Vol_{conduit}. Moreover, a trend towards the difference in the LA Vol_{passive emptying} parameter was shown — after PVI procedure, lower values were obtained as compared to the initial results ($P = 0.05$). Additionally, in the study group a difference in left ventricular end-systolic diameter (LV ESD) was demonstrated — lower LV ESD was observed before PVI procedure compared to the results after the observation period (30 [28–33] vs. 32 [29–34]; $P = 0.02$). There were no statistically significant differences between LV ejection fraction (LVEF), other LV dimensions and volumes, as well as LV stroke volume initially and after follow-up period.

LA echocardiographic parameters and the effectiveness of PVI procedure

The analysis of the influence of the pre- and post-PVI procedure echocardiographic parameters on the effectiveness of procedure (patients with and without AF recurrence after observation period) was performed (Table 3 and 4).

Patients after successful PVI procedure were characterized by a statistically significant higher initial segmental deformations: LAPLS_{bas-inf 2CH} and LAPLS_{api-sept 4CH}, as well as a higher initial LA wall strain dispersion in time in the 2CH view (Table 3). In the analysis performed after the follow-up period, revealed lower LA Vol_{min}, LA Vol_{max} and LAVI_{max} and higher LA EF and LA_{expansion index} in patients who underwent successful PVI treatment (Table 4).

DISCUSSION

This publication represents part of a single-center, non-randomized, prospective study of a population consisting of relatively young patients with a history of paroxysmal, symptomatic AF, without significant structural heart disease and with a low score obtained in the CHA₂DS₂-Vasc scale that were classified to the first-time PVI procedure with RF energy.

In the whole group, after a follow-up period, patients presented statistically significant higher mean LA Vol_{conduit}. In the analysis of LA echocardiographic parameters after follow-up period, patients with no AF recurrences had statistically significant lower LA volumes (minimal, maximal and maximal indexed), higher LA ejection fraction and LA expansion index, when compared to the patients after ineffective PVI treatment. Patients who maintained sinus rhythm after PVI procedure were characterized by statistically significant higher initial segmental strains: LA PLS_{bas-inf} and PLS_{api-sept}, as well as higher LA wall strain dispersion in time in the 2-chamber projection.

In recent years, there has been a growing interest in the use of a novel techniques for assessing function of the heart chambers, in that case LA strains in clinical practice. More and more

information about the 2D LA PGLS can be found in the available literature [11, 17–19]. However, the assessment of deformation of individual LA segments may provide more valuable information about risk of arrhythmia, a detailed assessment of potential wall fibrosis / weakening with taking into account LA symmetric and asymmetric remodeling, an additional assessment of cardioembolic risk or an evaluation of the effectiveness in time of a sinus rhythm recovery procedures. In this study, in the analysis of the pre-procedural LA PLS, both global and segmental, two initial segmental strains that had a statistically significant impact on the effectiveness after the observational period were identified. The lower (better) pre-procedural LA PLS_{bas-inf} assessed in 2CH view and LA PLS_{api-sept} in 4CH view values were associated with the maintenance of a sinus rhythm after follow-up. Moreover, statistical significance was obtained for the LA wall strain dispersion in time — initially, greater LA dyssynchrony was observed in patients in whom PVI treatment proved to be effective in time. These results seem to be rather random.

PVI procedure is currently considered to be the most effective therapy to restore sinus rhythm in AF patients. In this study, patients who maintain sinus rhythm after a successful first-time PVI procedure are characterized by a LA positive remodeling. In the study group, after the observational period, a significant reduction in the LA volumes — minimum, maximum and maximum indexed to BSA were observed. These are similar results, consistent with published meta-analysis of Augustine Njoku et al. [12], analyzing twenty one studies (3822 subjects), where patients with AF recurrence after PVI treatment have a higher mean LA volume/LA volume indexed to BSA, when compared to patients without AF recurrences.

On the other hand, in this study no statistically significant difference in the LAVI_{max} was found, assessed in the whole group before and after PVI treatment, without taking into account the effectiveness of the procedure. It is the parameter commonly considered to be prognostic for AF recurrences [12]. However, in the available literature, there is a cut-off point of LAVI_{max} <34.4 ml/m², that is associated with the best AF ablation outcome [20, 21], while Shin et al. [22] found that LAVI_{max} of 34 ml/m² showed a sensitivity of 70% and a specificity of 91% to predict AF recurrence. In the current study patients in whom PVI proved to be effective in the observational period had a lower mean value of the LAVI_{max} parameter when compared to the patients with AF recurrences – assessed before the PVI procedure (the mean [SD] 36.9 [9.8] ml vs. 41.9 [13.6] ml; trend towards statistical significance — **P 0.07**) and after the PVI procedure (34.5 [7] ml vs. 41.4 [13] ml; *P* <0.001).

At the same time, among patients without post-PVI AF recurrence, an improvement of LA function measured by an increase of LA_{expansion index} and thus higher LA EF were observed. A

lower median of LA Vol_{passive emptying} after PVI (trend towards significance) was also noted, compared to the results obtained before procedure. This demonstrates positive LA remodeling in these patients, which in turn may favor the continued maintenance of sinus rhythm in the future. It is puzzling that there is no coincident improvement of LA function as measured by the evaluation of LA global and segmental strains and LA wall strain synchrony. It should be noted, however, that the study population consists of a selected, relatively young patients with initially non-sustained form of atrial arrhythmia, not burdened with significant cardiovascular disease. Perhaps they did not have significant LA dysfunction at baseline, and the lack of statistical variability in the post-procedural evaluation should be treated as a success and the absence of complications related to the procedure itself, e.g. narrowing of the pulmonary veins or complications following transseptal puncture.

There are several studies indicating that LA strain has higher predictive value than LA size obtained from conventional echocardiography [13, 23]. Data published so far, strongly suggests that the longitudinal 2D strains of LA may be useful in predicting AF recurrence after PVI procedures [11, 17, 23–26]. 2D STE analysis enables the detection of decreased LA reservoir function and as a pump in patients with paroxysmal AF, even before changes in LA volume are revealed [14], which may be potentially associated with greater sensitivity of the method in detecting and predicting AF recurrence. Mirza et al. [23] showed that regardless of LA enlargement, pre-procedural strain of the LA lateral wall can be considered an independent determinant of AF recurrence after PVI. However, the effectiveness of the treatment was assessed 18 months after the procedure, using the TomTec software, which differs significantly from the methodology presented in the current study. On the other hand, in the work of Hammerstingl et al. [13], independent predictors of AF recurrence after PVI procedure were global LA strains obtained in the 2CH and 4CH projections and regional LA septal wall strain. Researchers also used TomTec software to analyze STE in 76 patients with paroxysmal and 27 patients with persistent AF, as well as in a 30-person control group. Similarly, as in the author's own work, the effectiveness of the PVI procedure was assessed after a minimum 6 months follow-up period.

Recently, there has been an increase in data on new cardiac visualization techniques assessing the volumes and strains of the LA using 3D techniques. What's more, it turns out that these techniques are potentially more accurate in patients with AF and surpass the 2D visualizations widely used so far [27–29]. For example, in a group of 348 patients with symptomatic paroxysmal or persistent AF, Montserrat et al. [30] showed none of the echocardiographic parameters considered, including LAVI, was associated with AF recurrence after PVI

procedure. Only volumetric assessment of LA with 3D rotational angiography showed in a multivariate analysis that LAVI is the only independent predictor for AF recurrence. LA volume measured with this method may be superior to TTE assessment and to AF history in predicting arrhythmia recurrence after PVI. There are also reports showing that LA strain determined by 3D STE is a novel and better predictor of AF recurrence after PVI than that determined by 2D STE or other known predictor [27]. These new 3D techniques, which were not specifically evaluated in the present study and although less common in routine echocardiographic assessment, may help to more accurately assess LA morphology and function.

It should be certainly borne in mind that there is no single parameter to clearly define the possibility of recurrence of arrhythmia after PVI. Echocardiographic assessment should be complex, using various available imaging techniques and potentially useful parameters to assess the morphology and function of the heart chambers, which may help improve the prediction of rhythm-control strategy success in AF [31].

Limitations

The analyzed group was relatively small and the statistical power of this study is limited. The conclusions should be used in relation to the population of relatively young patients with a history of paroxysmal, symptomatic AF, without significant structural heart disease and with a low score obtained in the CHA₂DS₂-Vasc scale who underwent the first-time PVI procedure with RF energy. Increasing of the population group could influence the demonstration of other relationships, especially in the case of parameters where only trend towards significance was achieved. On the other hand, the size of the group in the presented study did not differ significantly from other studies on related topics. Additional study limitation is associated with the absence of a control group. At the time of the study, no software was available to evaluate strains of LA and there were no guidelines for evaluating LA strains in patients during AF. Additionally, a significant limitation of the presented work is the lack of LV deformation analysis, the more so that the LV EF change significantly influenced the effectiveness of the PVI procedure.

We are aware of data gaps. Around 20% of the firstly described population is not included in some of the analyses, which actually reduces the size of the analyzed population. We precisely showed population numbers which differ a bit from the general population numbers because in each major test missing data were eliminated casewise.

CONCLUSIONS

1. Some echocardiographic parameters related to LA morphology improve after successful PVI treatment, which may be associated with positive LA remodeling.
2. LA strains and wall strain dispersion in time are not related to LA remodeling after successful PVI procedure.
3. The baseline LA standard and novel echocardiographic parameters cannot be used as a remote evaluation of the effectiveness of the PVI procedure. Further research is needed in this area, especially taking into account the limitations of the study.

Article information

Conflict of interest: None declared.

Funding: None.

Open access: This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, which allows downloading and sharing articles with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially. For commercial use, please contact the journal office at kardiologiapolska@ptkardio.pl.

REFERENCES

1. Thomas L, Abhayaratna WP. Left Atrial Reverse Remodeling: Mechanisms, Evaluation, and Clinical Significance. *JACC Cardiovasc Imaging*. 2017; 10(1): 65–77, doi: [10.1016/j.jcmg.2016.11.003](https://doi.org/10.1016/j.jcmg.2016.11.003), indexed in Pubmed: [28057220](https://pubmed.ncbi.nlm.nih.gov/28057220/).
2. Liżewska-Springer A, Dąbrowska-Kugacka A, Lewicka E, et al. Echocardiographic assessment in patients with atrial fibrillation (AF) and normal systolic left ventricular function before and after catheter ablation: If AF begets AF, does pulmonary vein isolation terminate the vicious circle? *Cardiol J*. 2020; 27(2): 126–135, doi: [10.5603/CJ.a2019.0004](https://doi.org/10.5603/CJ.a2019.0004), indexed in Pubmed: [30701515](https://pubmed.ncbi.nlm.nih.gov/30701515/).
3. Lo LW, Chen SA, Tsao HM, et al. Characterization of the dynamic function of the pulmonary veins before and after atrial fibrillation ablation using multi-detector computed tomographic images. *Int J Cardiovasc Imaging*. 2011; 27(7): 1049–1058, doi: [10.1007/s10554-010-9752-7](https://doi.org/10.1007/s10554-010-9752-7), indexed in Pubmed: [21110102](https://pubmed.ncbi.nlm.nih.gov/21110102/).
4. Silveira MM, Cabral JV, Xavier AT, et al. Detection of atrial fibrosis using echocardiographic strain: a new pathway. *Rev Assoc Med Bras (1992)*. 2022; 68(6): 866–870, doi: [10.1590/1806-9282.20211244](https://doi.org/10.1590/1806-9282.20211244), indexed in Pubmed: [35766703](https://pubmed.ncbi.nlm.nih.gov/35766703/).

5. Mandoli GE, D'Ascenzi F, Vinco G, et al. Novel Approaches in Cardiac Imaging for Non-invasive Assessment of Left Heart Myocardial Fibrosis. *Front Cardiovasc Med.* 2021; 8: 614235, doi: [10.3389/fcvm.2021.614235](https://doi.org/10.3389/fcvm.2021.614235), indexed in Pubmed: [33937354](https://pubmed.ncbi.nlm.nih.gov/33937354/).
6. Her AY, Choi EY, Shim CY, et al. Prediction of left atrial fibrosis with speckle tracking echocardiography in mitral valve disease: a comparative study with histopathology. *Korean Circ J.* 2012; 42(5): 311–318, doi: [10.4070/kcj.2012.42.5.311](https://doi.org/10.4070/kcj.2012.42.5.311), indexed in Pubmed: [22701133](https://pubmed.ncbi.nlm.nih.gov/22701133/).
7. Wojdyła-Hordyńska A, Baran J, Mazurek M, et al. Results of a survey concerning atrial fibrillation ablation strategies in Poland. *Kardiol Pol.* 2020; 78(10): 974–981, doi: [10.33963/KP.15407](https://doi.org/10.33963/KP.15407), indexed in Pubmed: [32486629](https://pubmed.ncbi.nlm.nih.gov/32486629/).
8. Hindricks G, Potpara T, Dagres N, et al. Corrigendum to: 2020 ESC Guidelines for the diagnosis and management of atrial fibrillation developed in collaboration with the European Association for Cardio-Thoracic Surgery (EACTS): The Task Force for the diagnosis and management of atrial fibrillation of the European Society of Cardiology (ESC) Developed with the special contribution of the European Heart Rhythm Association (EHRA) of the ESC. *Eur Heart J.* 2021; 42(40): 4194, doi: [10.1093/eurheartj/ehab648](https://doi.org/10.1093/eurheartj/ehab648), indexed in Pubmed: [34520521](https://pubmed.ncbi.nlm.nih.gov/34520521/).
9. Hachem AH, Marine JE, Tahboub HA, et al. Radiofrequency Ablation versus Cryoablation in the Treatment of Paroxysmal Atrial Fibrillation: A Meta-Analysis. *Cardiol Res Pract.* 2018; 2018: 6276241, doi: [10.1155/2018/6276241](https://doi.org/10.1155/2018/6276241), indexed in Pubmed: [29805800](https://pubmed.ncbi.nlm.nih.gov/29805800/).
10. Nielsen AB, Skaarup KG, Lassen MC, et al. Usefulness of left atrial speckle tracking echocardiography in predicting recurrence of atrial fibrillation after radiofrequency ablation: a systematic review and meta-analysis. *Int J Cardiovasc Imaging.* 2020; 36(7): 1293–1309, doi: [10.1007/s10554-020-01828-2](https://doi.org/10.1007/s10554-020-01828-2), indexed in Pubmed: [32248332](https://pubmed.ncbi.nlm.nih.gov/32248332/).
11. Yasuda R, Murata M, Roberts R, et al. Left atrial strain is a powerful predictor of atrial fibrillation recurrence after catheter ablation: study of a heterogeneous population with sinus rhythm or atrial fibrillation. *Eur Heart J Cardiovasc Imaging.* 2015; 16(9): 1008–1014, doi: [10.1093/ehjci/jev028](https://doi.org/10.1093/ehjci/jev028), indexed in Pubmed: [25750193](https://pubmed.ncbi.nlm.nih.gov/25750193/).
12. Njoku A, Kannabhiran M, Arora R, et al. Left atrial volume predicts atrial fibrillation recurrence after radiofrequency ablation: a meta-analysis. *Europace.* 2018; 20(1): 33–42, doi: [10.1093/europace/eux013](https://doi.org/10.1093/europace/eux013), indexed in Pubmed: [28444307](https://pubmed.ncbi.nlm.nih.gov/28444307/).
13. Hammerstingl C, Schwekendiek M, Momcilovic D, et al. Left atrial deformation imaging with ultrasound based two-dimensional speckle-tracking predicts the rate of

recurrence of paroxysmal and persistent atrial fibrillation after successful ablation procedures. *J Cardiovasc Electrophysiol.* 2012; 23(3): 247–255, doi: [10.1111/j.1540-8167.2011.02177.x](https://doi.org/10.1111/j.1540-8167.2011.02177.x), indexed in Pubmed: [21955059](https://pubmed.ncbi.nlm.nih.gov/21955059/).

14. Moneta GL, Edwards JM, Chitwood RW, et al. Correlation of North American Symptomatic Carotid Endarterectomy Trial (NASCET) angiographic definition of 70% to 99% internal carotid artery stenosis with duplex scanning. *J Vasc Surg.* 1993; 17(1): 152–7; discussion 157, doi: [10.1067/mva.1993.42888](https://doi.org/10.1067/mva.1993.42888), indexed in Pubmed: [8421332](https://pubmed.ncbi.nlm.nih.gov/8421332/).
15. Todaro MC, Choudhuri I, Belohlavek M, et al. New echocardiographic techniques for evaluation of left atrial mechanics. *Eur Heart J Cardiovasc Imaging.* 2012; 13(12): 973–984, doi: [10.1093/ehjci/jes174](https://doi.org/10.1093/ehjci/jes174), indexed in Pubmed: [22909795](https://pubmed.ncbi.nlm.nih.gov/22909795/).
16. Rosca M, Mondillo S, O'Connor K. Left atrium. In: Lancellotti P, Zamorano JL, Habib G, Badano L. ed. *The EACVI Textbook of Echocardiography* (2 edn). Oxford University Press 2017: 162–170.
17. Ma XX, Boldt LH, Zhang YL, et al. Clinical Relevance of Left Atrial Strain to Predict Recurrence of Atrial Fibrillation after Catheter Ablation: A Meta-Analysis. *Echocardiography.* 2016; 33(5): 724–733, doi: [10.1111/echo.13184](https://doi.org/10.1111/echo.13184), indexed in Pubmed: [26857344](https://pubmed.ncbi.nlm.nih.gov/26857344/).
18. Parwani AS, Morris DA, Blaschke F, et al. Left atrial strain predicts recurrence of atrial arrhythmias after catheter ablation of persistent atrial fibrillation. *Open Heart.* 2017; 4(1): e000572, doi: [10.1136/openhrt-2016-000572](https://doi.org/10.1136/openhrt-2016-000572), indexed in Pubmed: [28674624](https://pubmed.ncbi.nlm.nih.gov/28674624/).
19. Liżewska-Springer A, Dąbrowska-Kugacka A, Lewicka E, et al. Echocardiographic predictors of atrial fibrillation recurrence after catheter ablation: A literature review. *Cardiol J.* 2020; 27(6): 848–856, doi: [10.5603/CJ.a2018.0067](https://doi.org/10.5603/CJ.a2018.0067), indexed in Pubmed: [29924375](https://pubmed.ncbi.nlm.nih.gov/29924375/).
20. Zhuang J, Wang Yi, Tang K, et al. Association between left atrial size and atrial fibrillation recurrence after single circumferential pulmonary vein isolation: a systematic review and meta-analysis of observational studies. *Europace.* 2012; 14(5): 638–645, doi: [10.1093/europace/eur364](https://doi.org/10.1093/europace/eur364), indexed in Pubmed: [22117033](https://pubmed.ncbi.nlm.nih.gov/22117033/).
21. J Albano A, Bush J, L Parker J, et al. Left Atrial Volume Index Predicts Arrhythmia-Free Survival in Patients with Persistent Atrial Fibrillation Undergoing Cryoballoon Ablation. *J Atr Fibrillation.* 2019; 12(2): 2192, doi: [10.4022/jafib.2192](https://doi.org/10.4022/jafib.2192), indexed in Pubmed: [32002112](https://pubmed.ncbi.nlm.nih.gov/32002112/).

22. Shin SH, Park MY, Oh WJ, et al. Left atrial volume is a predictor of atrial fibrillation recurrence after catheter ablation. *J Am Soc Echocardiogr.* 2008; 21(6): 697–702, doi: [10.1016/j.echo.2007.10.022](https://doi.org/10.1016/j.echo.2007.10.022), indexed in Pubmed: [18187293](https://pubmed.ncbi.nlm.nih.gov/18187293/).
23. Mirza M, Caracciolo G, Khan U, et al. Left atrial reservoir function predicts atrial fibrillation recurrence after catheter ablation: a two-dimensional speckle strain study. *J Interv Card Electrophysiol.* 2011; 31(3): 197–206, doi: [10.1007/s10840-011-9560-6](https://doi.org/10.1007/s10840-011-9560-6), indexed in Pubmed: [21424845](https://pubmed.ncbi.nlm.nih.gov/21424845/).
24. Montserrat S, Gabrielli L, Bijns B, et al. Left atrial deformation predicts success of first and second percutaneous atrial fibrillation ablation. *Heart Rhythm.* 2015; 12(1): 11–18, doi: [10.1016/j.hrthm.2014.08.032](https://doi.org/10.1016/j.hrthm.2014.08.032), indexed in Pubmed: [25172009](https://pubmed.ncbi.nlm.nih.gov/25172009/).
25. Yu ZX, Yang W, Yin WS, et al. Clinical utility of left atrial strain in predicting atrial fibrillation recurrence after catheter ablation: An up-to-date review. *World J Clin Cases.* 2022; 10(23): 8063–8075, doi: [10.12998/wjcc.v10.i23.8063](https://doi.org/10.12998/wjcc.v10.i23.8063), indexed in Pubmed: [36159552](https://pubmed.ncbi.nlm.nih.gov/36159552/).
26. Marchandise S, Garnir Q, Scavée C, et al. Prediction of Left Atrial Fibrosis and Success of Catheter Ablation by Speckle Tracking Echocardiography in Patients Imaged in Persistent Atrial Fibrillation. *Front Cardiovasc Med.* 2022; 9: 856796, doi: [10.3389/fcvm.2022.856796](https://doi.org/10.3389/fcvm.2022.856796), indexed in Pubmed: [35694674](https://pubmed.ncbi.nlm.nih.gov/35694674/).
27. Mochizuki A, Yuda S, Fujito T, et al. Left atrial strain assessed by three-dimensional speckle tracking echocardiography predicts atrial fibrillation recurrence after catheter ablation in patients with paroxysmal atrial fibrillation. *J Echocardiogr.* 2017; 15(2): 79–87, doi: [10.1007/s12574-017-0329-5](https://doi.org/10.1007/s12574-017-0329-5), indexed in Pubmed: [28155065](https://pubmed.ncbi.nlm.nih.gov/28155065/).
28. Kobayashi Y, Okura H, Kobayashi Y, et al. Assessment of atrial synchrony in paroxysmal atrial fibrillation and impact of pulmonary vein isolation for atrial dyssynchrony and global strain by three-dimensional strain echocardiography. *J Am Soc Echocardiogr.* 2014; 27(11): 1193–1199, doi: [10.1016/j.echo.2014.08.004](https://doi.org/10.1016/j.echo.2014.08.004), indexed in Pubmed: [25240493](https://pubmed.ncbi.nlm.nih.gov/25240493/).
29. Bossard M, Knecht S, Aeschbacher S, et al. Conventional versus 3-D Echocardiography to Predict Arrhythmia Recurrence After Atrial Fibrillation Ablation. *J Cardiovasc Electrophysiol.* 2017; 28(6): 651–658, doi: [10.1111/jce.13202](https://doi.org/10.1111/jce.13202), indexed in Pubmed: [28301685](https://pubmed.ncbi.nlm.nih.gov/28301685/).
30. Montserrat S, Gabrielli L, Borrás R, et al. Left atrial size and function by three-dimensional echocardiography to predict arrhythmia recurrence after first and repeated

ablation of atrial fibrillation. *Eur Heart J Cardiovasc Imaging*. 2014; 15(5): 515–522, doi: [10.1093/ehjci/jet194](https://doi.org/10.1093/ehjci/jet194), indexed in Pubmed: [24168909](https://pubmed.ncbi.nlm.nih.gov/24168909/).

31. Kupczyńska K, Mandoli GE, Cameli M, et al. Left atrial strain - a current clinical perspective. *Kardiol Pol*. 2021; 79(9): 955–964, doi: [10.33963/KP.a2021.0105](https://doi.org/10.33963/KP.a2021.0105), indexed in Pubmed: [34599503](https://pubmed.ncbi.nlm.nih.gov/34599503/).

Table 1. Characteristics of the study group

		Study group (n = 80)
Demographics	Age, year	58 (50–63)
	Male sex	50 (62.5%)
	0–5 years	37 (46%)
AF duration	5–10 years	27 (34%)
	>10 years	16 (20%)
Baseline biometric evaluation	Height, cm	174.2 (9.2)
	Weight, kg	88.3 (15.2)
	BMI, kg/m ²	29.1 (4.25)
	BSA, m ²	2.06 (0.22)
Functional class and risk scale	EHRA, score	3 (2–3)
	CHA ₂ DS ₂ -Vasc, score	2 (1–2.5)
	Coronary artery disease	15 (19%)
Co-morbidities	Arterial hypertension	56 (70%)
	Diabetes mellitus	15 (19%)
	Hyperlipidemia	55 (69%)
	Obesity	27 (34%)
	Tobacco smoking	

Never	46 (57.5%)
In the past	22 (27.5%)
Active	12 (15%)

Results are given as the mean with standard deviation (SD) for normal distributions or the median with interquartile range (IQR) for non-normal distributions or absolute numbers and percentages (%)

Abbreviations: AF, atrial fibrillation; BMI, body mass index; BSA, body mas area

Table 2. LA echocardiographic parameters before PVI procedure and after the observation period

Parameters		Before (n = 66)	After (n = 66)	PVI P-value
	LA _{diameter} , mm	39.6	—	—
	LA Vol _{preA} , ml	53 (35–64)	51 (37–66)	0.76
	LA Vol _{min} , ml	36.5 (25–49)	33 (26–51)	0.57
	LA Vol _{max} , ml	74.5 (60–103)	72.5 (62–88)	0.15
	LAVI _{max} , ml/m ²	36.3 (29.4–49.5)	35.7 (30–45.4)	0.16
	LA Vol _{reservoir} , ml	38 (32–49)	37 (32–44)	0.21
LA morphological parameters	LA Vol _{conduit} , ml	25.4 (15.1)	30.1 (13.3)	0.01
	LA Vol _{passive emptying} , ml	21 (16–35)	20 (15–26)	0.05
	LA Vol _{contractile} , ml	14.5 (8–20)	16 (11–22)	0.56
	LAEF, %	52 (46–58)	54 (44–59)	0.33
	LA _{expansion index}	1.17 (0.48)	1.15 (0.46)	0.75
	LA _{active empt frac} , %	31 (17–38)	33 (26–39)	0.63
	LA _{passive empt frac} , %	32 (13)	30 (12)	0.3
LA strains and LA wall strain dispersion	LA PLS _{bas-inf 2CH} , %	–17.5 (–21.8 to 13.4)	–19.4 (–22.3 to –13)	0.32
	LA PLS _{med-inf 2CH} , %	–15.2 (–18 to 11.8)	–14.6 (–18.3 to –11.5)	0.64

LA PLS _{api-inf} 2CH, %	-10.9 (-14.3 to -6.4)	-10.1 (-12.6 to -6.8)	0.25
LA PLS _{bas-ant} 2CH, %	-14.6 (-19.5 to 9.8)	-14.7 (-19.4 to -10.3)	0.97
LA PLS _{med-ant} 2CH, %	-11.6 (-17.1 to 7)	-13.1 (-16.7 to -8.6)	0.83
LA PLS _{api-ant} 2CH, %	-10.7 (-16.3 to 6.3)	-10.2 (-14.1 to -6.1)	0.12
LA PGLS _{2CH} , %	-12.8 (4.7)	-12.4 (3.8)	0.53
LA wall strain dispersion _{2CH} , ms	98.5 (45–182)	97 (63–164)	0.17
LA PLS _{bas-lat} 4CH, %	-14.2 (-19 to 9.1)	-16.9 (-19.7 to -12.4)	0.26
LA PLS _{med-lat} 4CH, %	-11.6 (-16.6 to 8.6)	-12.6 (-17.3 to -9.8)	0.8
LA PLS _{api-lat} 4CH, %	-10.1 (-14.2 to 5.6)	-10.7 (-14.2 to -7)	0.92
LA PLS _{bas-sept} 4CH, %	-14.1 (-18 to 10.6)	-16.8 (-19.9 to -14.5)	0.5
LA PLS _{med-sept} 4CH, %	-14.7 (-18.1 to 9.9)	-15.6 (-19 to -11.8)	0.87
LA PLS _{api-sept} 4CH, %	-11.6 (-16.3 to 7)	-12.4 (-17.9 to -7)	0.57
LA PGLS _{4CH} , %	-12 (4.8)	-12.7 (4.3)	0.37
LA wall strain dispersion _{4CH} , ms	101.5 (44–177)	97 (49–165.5)	0.35

Results are given as the mean with standard deviation (SD) for normal distributions or the median with interquartile range (IQR) for non-normal distributions

Abbreviations: api-ant, apical-anterior; api-inf, apical-inferior; api-lat, apical-lateral; api-sept, apical-septal; bas-ant, basal-anterior; bas-inf, basal-inferior; bas-lat, basal-lateral; bas-sept, basal-septal; empt frac, emptying fraction; LA, left atrium; PGLS, maximum left atrial peak global longitudinal strain; PLS, maximum left atrial peak longitudinal strain; max, maximal;

min, minimal; med-ant, medial-anterior; med-inf, medial-inferior; med-lat, medial-basal; med-sept, medial-septal; RF, radiofrequency pulmonary vein isolation; preA, preatrial; PVI, pulmonary vein isolation with RF energy; TTE, transthoracic echocardiography; Vol, volume; 2CH, two chamber view; 4CH, four chamber view

Table 3. Initial LA echocardiographic parameters depending on the effectiveness of the PVI procedure

Parameters	Effective	Ineffective	P-value
	PVI (n = 43)	PVI (n = 35)	
LA _{diameter} , mm	39.4 (4.4)	39.7 (4.1)	0.78
LA Vol _{preA} , ml	51.6 (17.3)	58.2 (24.7)	0.26
LA Vol _{min} , ml	33 (24–48)	40 (30–55)	0.18
LA Vol _{max} , ml	70 (57–96)	82 (67–104)	0.36
LAVI _{max} , ml/m ²	36.9 (9.8)	41.9 (13.6)	0.07
LA Vol _{reservoir} , ml	40.6 (12.9)	39.9 (11.4)	0.83
LA Vol _{conduit} , ml	28 (12–39)	22 (16–32)	0.37
LA Vol _{passive emptying} , ml	24 (16–37)	21 (17–34)	1
LA Vol _{contractile} , ml	14.9 (9.2)	14.7 (8.1)	0.95
LA EF, %	54 (47–60)	49 (40–60)	0.15
LA _{expansion index}	1.16 (0.9–1.5)	0.98 (0.8–1.3)	0.15
LA _{active empt frac} , %	33 (17–40)	24 (20–40)	0.45
LA _{passive empt frac} , %	33 (10)	31 (10)	0.6
LA PLS _{bas-inf 2CH} , %	–19.4 (4.7)	–15.8 (6.2)	0.02
LA PLS _{med-inf 2CH} , %	–15.6 (4.7)	–13 (5.6)	0.06
LA PLS _{api-inf 2CH} , %	–10.5 (–14.6 to 6.1)	–11.4 (–13.9 to –7)	0.8
LA PLS _{bas-ant 2CH} , %	–14.6 (–19.4 to –10.6)	–15.3 (–19.5 to –8)	0.8

LA PLS _{med-ant} 2CH, %	-11.4 (-17.7 to -8.6)	-11.8 (-15.4 to -6.6)	0.75
LA PLS _{api-ant} 2CH, %	-11.7 (7.9)	-11.2 (6)	0.76
LA PGLS _{2CH} , %	-12.4 (5)	-11.8 (4.6)	0.59
LA wall strain dispersion _{2CH} , ms	115 (50–200)	63 (38–147)	0.02
LA PLS _{bas-lat} 4CH, %	-13.8 (-18.4 to -8.9)	-14.9 (-21.7 to -9.2)	0.62
LA PLS _{med-lat} 4CH, %	-11.6 (-16 to 8.4)	-13.1 (-17 to 9.6)	0.62
LA PLS _{api-lat} 4CH, %	-10.3 (-14 to 5.7)	-10.1 (-17.3 to -5.8)	0.78
LA PLS _{bas-sept} 4CH, %	-15.5 (5.6)	-14 (6.6)	0.37
LA PLS _{med-sept} 4CH, %	-15.9 (-17.9 to -12)	-11.1 (-18.6 to -7.7)	0.05
LA PLS _{api-sept} 4CH, %	-13.4 (-16.3 to -9.6)	-8.6 (-13.3 to 6)	0.02
LA PGLS _{4CH} , %	-12 (4.3)	-11 (5.4)	0.45
LA wall strain dispersion _{4CH} , ms	86.5 (32.5–171)	133 (59–179)	0.34

Results are given as the mean with standard deviation (SD) for normal distributions or the median with interquartile range (IQR) for non-normal distributions.

Abbreviations: see [Table 2](#)

Table 4. LA echocardiographic parameters after follow-up period depending on the effectiveness of the PVI procedure

Parameters	Effective	Ineffective	P-value
	PVI (n = 43)	PVI (n = 35)	
LA Vol _{preA} , ml	47.5 (37–57)	55 (37–78)	0.27
LA Vol _{min} , ml	30 (25–43)	45 (27–61)	0.02

	LA Vol _{max} , ml	72.6 (18.8)	84.1 (20.8)	0.02
	LAVI _{max} , ml/m ²	34.5 (7)	41.4 (13)	<0.001
	LA Vol _{reservoir} , ml	38 (33–46)	35.5 (28.5–41)	0.22
	LA Vol _{conduit} , ml	29.9 (14.6)	31.3 (13.5)	0.7
	LA Vol _{passive emptying} , ml	22.2 (9.8)	19.1 (6.4)	0.14
	LA Vol _{contractile} , ml	16 (11–19)	14 (11–24)	0.97
	LA EF, %	57 (49–60)	46 (38–56)	0.002
	LA _{expansion index}	1.3 (0.4)	0.9 (0.45)	0.004
	LA _{active empt frac} , %	30 (10)	30 (10)	0.4
	LA _{passive empt frac} , %	30 (13)	30 (10)	0.14
	LA _{passive empt frac} , %	33 (10)	31 (10)	0.6
LA echocardiographic parameters after follow-up period	LA PLS _{bas-inf 2CH} , %	-19.7 (-23.5 to 12.5)	-19.4 (-21.8 to -15.5)	0.84
	LA PLS _{med-inf 2CH} , %	-15.6 (-18.4 to 11.5)	-14.1 (-17.5 to -11.2)	0.62
	LA PLS _{api-inf 2CH} , %	-10.1 (-12.5 to -4.8)	-9.7 (-12.7 to -7.9)	0.47
	LA PLS _{bas-ant 2CH} , %	-15.6 (-20.1 to 11.1)	-12.9 (-18.4 to -9.6)	0.41
	LA PLS _{med-ant 2CH} , %	-13.6 (5.3)	-12.1 (5.6)	0.3
	LA PLS _{api-ant 2CH} , %	-11 (-13.3 to -7.2)	-9.1 (-15.5 to -6)	0.94
	LA PGLS _{2CH} , %	-12.2 (3.5)	-12.1 (4)	0.3
	LA wall strain dispersion _{2CH} , ms	100 (56–154)	89 (67–198)	0.76
	LA PLS _{bas-lat 4CH} , %	-17.2 (-19.7 to 11.5)	-16.4 (-19.7 to -13.7)	0.36
	LA PLS _{med-lat 4CH} , %	-12.4 (-19.7 to -9.8)	-13.2 (-16.9 to -8.6)	0.87

LA PLS _{api-lat} 4CH, %	-10.3 (-16.6 to -6.8)	-11.5 (-12.9 to -7.1)	0.85
LA PLS _{bas-sept} 4CH, %	-16.3 (-19.1 to -13.6)	-18 (-21.4 to -15.6)	0.07
LA PLS _{med-sept} 4CH, %	-15.3 (-18 to -10.8)	-16.1 (-20 to -12)	0.29
PLS _{api-sept} 4CH, %	-12.8 (-17.7 to -7.4)	-12.3 (-18.2 to -6.2)	0.75
LA PGLS _{4CH} , %	-12.6 (4.6)	-13.6 (4.8)	0.4
LA wall strain dispersion _{4CH} , ms	105 (70-167)	87 (32-164)	0.38

Results are given as the mean with standard deviation (SD) for normal distributions or the median with interquartile range (IQR) for non-normal distributions

Abbreviations: see [Table 2](#)




Figure 1. Division of the LA into segments corresponding to strains in the 2CH view
 Abbreviations: LA, left atrium; 2CH, two chamber




Figure 2. Division of the LA into segments corresponding to strains in 4CH view
Abbreviations: LA, left atrium; 4CH, four chamber