

Living Without Beth and Craig: Definitions and Interpolants in Description and Modal Logics with Nominals and Role Inclusions

Alessandro Artale¹, Jean Christoph Jung², Andrea Mazzullo¹, Ana Ozaki³, and Frank Wolter⁴

¹ Free University of Bozen-Bolzano ² TU Dortmund University
³ University of Bergen ⁴ University of Liverpool

Abstract. The Craig interpolation property (CIP) states that an interpolant for an implication exists iff it is valid. The projective Beth definability property (PBDP) states that an explicit definition exists iff a formula stating implicit definability is valid. Thus, the CIP and PBDP reduce potentially hard existence problems to entailment in the underlying logic. Description (and modal) logics with nominals and/or role inclusions do not enjoy the CIP nor the PBDP, but interpolants and explicit definitions have many applications, in particular in concept learning, ontology engineering, and ontology-based data management. In this article we show that, even without Beth and Craig, the existence of interpolants and explicit definitions is decidable in description logics with nominals and/or role inclusions such as \mathcal{ALCCO} , \mathcal{ALCH} and \mathcal{ALCHOI} and corresponding hybrid modal logics. However, living without Beth and Craig makes these problems harder than entailment: the existence problems become 2EXPTIME -complete in the presence of an ontology or the universal modality, and CONEXPTIME -complete otherwise. We also analyze explicit definition existence if all symbols (except the one that is defined) are admitted in the definition. In this case the complexity depends on whether one considers individual or concept names. Finally, we consider the problem of computing interpolants and explicit definitions if they exist and turn the complexity upper bound proof into an algorithm computing them, at least for description logics with role inclusions.

Keywords: Description logic, Modal logic, Craig interpolants, Beth definability, Explicit definitions, Computational complexity

1 Introduction

The *Craig Interpolation Property* (CIP) for a logic \mathcal{L} states that an implication $\varphi \Rightarrow \psi$ is valid in \mathcal{L} iff there exists a formula χ in \mathcal{L} using only the common symbols of φ and ψ such that $\varphi \Rightarrow \chi$ and $\chi \Rightarrow \psi$ are both

valid in \mathcal{L} . The intermediate formula χ is then called an \mathcal{L} -interpolant for $\varphi \Rightarrow \psi$ [27]. The CIP is generally regarded as one of the most important and useful properties in formal logic [95], with numerous applications ranging from formal verification [74] and software specification [29] to theory combinations [25, 39, 21, 22] and query reformulation and rewriting in databases [91, 13]. A particularly important consequence of the CIP is the *projective Beth definability property* (PBDP), which states that a relation is implicitly definable using a signature Σ of symbols iff it is explicitly definable using Σ . If Σ is the set of all symbols distinct from that relation, then we speak of the (non-projective) *Beth definability property* (BDP) [16].

In this paper, we investigate interpolants and explicit definitions in description logics (DLs), and we also highlight consequences in modal logic. In DLs, one distinguishes essentially two forms of interpolation, both of which are relevant and have their applications. Given an entailment $\mathcal{O} \models C \sqsubseteq D$, that is, C is subsumed by D w.r.t. some background knowledge in the form of a DL ontology \mathcal{O} , one might either be interested in an interpolant between the concepts C and D or in an interpolant between \mathcal{O} and the concept inclusion (CI) $C \sqsubseteq D$. In the first case, the interpolant is a concept, whereas in the second case, the interpolant is an ontology. We refer with *CI-interpolation* to the latter form and call the interpolant a *CI-interpolant*. The CIP for CI-interpolation has been shown to be the most important logical property that ensures the robust behaviour of ontology modules and decompositions [55, 54].

In this article, we mostly focus on interpolation (in the former sense of an interpolating concept) and only derive some corollaries for CI-interpolation. Hence, unless stated otherwise, here and in what follows we speak about interpolating concepts and the corresponding CIP. For explicit definability, one asks for definitions of concepts, possibly with respect to an ontology; these explicit definitions are strongly related to interpolants and as stated above BDP and PBDP follow from the CIP. In DLs, the BDP and PBDP have been used in ontology engineering to extract explicit definitions of concepts and obtain equivalent acyclic terminologies from ontologies [89, 90], they have been investigated in ontology-based data management to equivalently rewrite ontology-mediated queries [85, 92, 35, 34, 93], and they have been proposed to support the construction of alignments between ontologies [47]. Interpolants have been used to study P/NP dichotomies in ontology-based query answering [69].

The CIP, PBDP, and BDP are so powerful because potentially very hard existence questions are reduced to straightforward entailment ques-

tions: an interpolant *exists* iff an implication is valid and an explicit definition *exists* iff a straightforward formula stating implicit definability is valid. The existence problems are thus not harder than validity. Many basic DLs such as \mathcal{ALC} , \mathcal{ALCI} , and \mathcal{ALCIQ} enjoy the CIP and PBDP [90], and consequently the existence of an interpolant or an explicit definition can be decided in EXPTIME simply because entailment checking in these DLs is in EXPTIME (and without ontology even in PSPACE). Unfortunately, the CIP and the PBDP fail to hold for some important DLs. The most basic examples are the extension \mathcal{ALCO} of \mathcal{ALC} with nominals (concepts of the form $\{a\}$ with a an individual name), the extension \mathcal{ALCH} of \mathcal{ALC} with role inclusions (inclusions $r \sqsubseteq s$ between binary relations/role names r and s), and all standard DLs containing either \mathcal{ALCO} or \mathcal{ALCH} [55, 90]. It follows that for these DLs the existence of interpolants and explicit definitions cannot be reduced (directly) to entailment checking.

The aim of this article is to explore the consequences of the failure of the CIP and PBDP for interpolant and explicit definition existence. To this end, we investigate the complexity of deciding the existence of interpolants and explicit definitions for the set DL_{nr} of DLs containing \mathcal{ALCO} , \mathcal{ALCH} , and their extensions by inverse roles and/or the universal role. We discuss next two more applications of interpolants and explicit definitions for \mathcal{ALCO} and its extensions.

Data Separability and Concept Learning. We show that interpolants are essentially the same as concepts separating positive and negative data examples in DL knowledge bases (KBs). Recall that a DL KB is a pair $(\mathcal{O}, \mathcal{D})$ with \mathcal{O} a DL ontology and \mathcal{D} a set of data items of the form $A(a)$ and $r(a, b)$ with a, b individuals, A a concept name, and r a role name. Let \mathcal{O} be an ontology and P and N sets of positively and negatively labelled pairs (\mathcal{D}, a) with \mathcal{D} a set of data items and a an individual in \mathcal{D} . Then the aim of supervised concept learning is to determine a concept C in a signature Σ of relevant symbols such that C separates P and N in the sense that $(\mathcal{O}, \mathcal{D}) \models C(a)$ for all positive examples $(\mathcal{D}, a) \in P$ and $(\mathcal{O}, \mathcal{D}) \models \neg C(a)$ for all negative examples $(\mathcal{D}, a) \in N$.¹ Concept learning has received significant interest over the past 15 years, where the focus has been on developing and analyzing refinement based algorithms for finding separating concepts [62, 63, 65, 84, 30, 64, 82]. Prominent concept

¹ This condition is called strong separation in [36, 49, 50]. A weaker version, called weak separation, only demands that $(\mathcal{O}, \mathcal{D}) \not\models C(a)$ for all negative examples $(\mathcal{D}, a) \in N$. Concept learning systems have been developed for both the weak and the strong notion.

learning systems include the DL LEARNER [20, 19], DL-FOIL [31] and its extension DL-FOCL [83], SPaCEL [94], YINYANG [45]. The existence problem for separating concepts has been investigated in [36, 48–50]. For DLs extending \mathcal{ALCO} , we establish a one-to-one correspondence between interpolants and separating concepts, modulo a rather straightforward polynomial time translation. Hence the existence of separating concepts reduces to the existence of interpolants and finding small such concepts or concepts of a certain syntactic shape, as is often useful in supervised learning, also reduces to the same task for interpolants. We emphasize that the presence of nominals in the DL is critical as they are required to encode the individuals used in \mathcal{D} into concepts.

Referring Expressions. The computation of explicit definitions of concept names has been explored in detail since at least [89], see also [7]. Only recently, the focus on defining concept names has been extended to defining individual names, also called referring expression generation in computational linguistics and data management [60, 3, 17]. In fact, it has been convincingly argued that very often in applications the individual names used in ontologies or data sets are insufficient “to allow humans to figure out what real-world objects they refer to” [18]. A natural way to address this problem is to check for such an individual name a whether there exists a concept C over a set of relevant symbols Σ that provides an explicit definition of $\{a\}$ and present such a concept C to the human user. Observe that one has to work with DLs extending \mathcal{ALCO} to formulate this problem as an explicit definition existence problem.

To conclude, data separation, concept learning, and referring expression generation are challenging research problems which directly benefit from a better understanding of interpolant and explicit definition existence in extensions of \mathcal{ALCO} . We now discuss the main results of this article, formulated in an informal way. Precise formulations are given later. Recall that DL_{nr} is the set of DLs \mathcal{ALCO} , \mathcal{ALCH} , and their extensions with inverse roles and the universal role, and that we assume the presence of a background DL ontology. Our first main result is as follows.

Theorem 1. *Let $\mathcal{L} \in DL_{nr}$. Then \mathcal{L} -interpolant existence and \mathcal{L} -definition existence are 2EXPTIME-complete.*

Theorem 1 confirms the suspicion that interpolant and definition existence are much harder problems than entailment if one has to live without Beth and Craig. On the positive side, these problems are still decidable. Interestingly, for DLs in DL_{nr} with nominals, the 2EXPTIME lower bound for

definition existence already holds if one asks for an explicit definition of an individual over the signature containing all symbols distinct from that individual. In contrast, the same problem for concept names is shown to be EXPTIME-complete and thus not harder than entailment. Hence, in contrast to concept name definitions, referring expression existence does not become less complex in the non-projective case when all symbols are allowed in definitions.

We next consider the same problems if the background ontology is empty, or, in the case of DLs in DL_{nr} without nominals, if the ontology contains only role inclusions. Observe that if the DL admits the universal role or both nominals and inverse roles, then the ontology can be encoded as a concept using spy points [1], so nothing changes compared to the case with ontologies covered in Theorem 1. For the remaining cases we show the following.

Theorem 2. (1) *If $\mathcal{L} \in \{\mathcal{ALCO}, \mathcal{ALCHO}\}$, then for the empty ontology and ontologies containing role inclusions only, \mathcal{L} -interpolant existence and \mathcal{L} -definition existence are both coNEXPTIME-complete;*

(2) *If $\mathcal{L} \in \{\mathcal{ALCH}, \mathcal{ALCHI}\}$, then for ontologies containing role inclusions only \mathcal{L} -interpolant existence and \mathcal{L} -definition existence are both coNEXPTIME-complete.*

It follows that without ontology and ontologies containing role inclusions only interpolant existence and explicit definition existence are still harder than entailment which is PSPACE-complete.

The proofs of Theorems 1 and 2 can be adapted to also obtain results about CI-interpolation and interpolation in modal logic. Regarding the former we show that for the DLs \mathcal{L} which extend \mathcal{ALCO} with the universal role or with the universal role and inverse roles the problem of deciding the existence of a CI-interpolant for $\mathcal{O} \models C \sqsubseteq D$ is 2EXPTIME-complete. It follows that again failure of the CIP leads to an exponentially harder interpolant existence problem than entailment. We conjecture that the same can be proved for all DLs in DL_{nr} , but leave a proof for future work.

In modal logic, the CIP and PBDP have been investigated for many years. In fact, the CIP and PBDP of DLs such as \mathcal{ALC} and \mathcal{ALCI} follows rather directly from earlier results on the CIP and PBDP in modal logic [71, 72, 81]. Also the fact that nominals lead to failure of the CIP and PBDP, and how this could be repaired by adding logical connectives, was first analyzed in depth in the literature on hybrid modal logic, in particular [2, 86]. In our investigation of interpolant existence in modal logic, we first consider basic modal logic with nominals and show that as a direct

consequence of Theorem 2 the problem of deciding interpolant existence is CONEXPTIME-complete for the standard local consequence relation. We also show using Theorem 1 that if one adds the universal modality, then interpolant existence becomes 2EXPTIME-complete. In modal logic, nominals are often considered in tandem with the @-operator, where $@_a\varphi$ states that formula φ holds at the world denoted by nominal a . The resulting language is more expressive than modal logic with nominals and less expressive than modal logic with nominals and the universal modality. We show that for the modal logic with both nominals and the @-operator interpolant existence is still CONEXPTIME-complete. Our complexity results also hold for the modal language with a single modal operator (and the universal modality, if present).

While the focus in this article is on the decision problem, we also make initial observations regarding the problem of actually computing interpolants or explicit definitions if they exist. More specifically, for DLs in DL_{nr} that do not admit nominals, we present a modification of the decision procedure from the proof of Theorem 1 that returns in double exponential time the DAG representation of an interpolant (if it exists). This corresponds to interpolants of worst case triple exponential size which we conjecture to be optimal.

Overview of the Paper. In the following Section 2, we discuss further related work. In Sections 3 and 4, we introduce the preliminaries on description logics and Craig interpolation and Beth definability, respectively. In Section 5, we provide model-theoretic characterizations of the definition and interpolation existence problems and formulate our main results in detail. The subsequent four sections are devoted to the proofs of these main results. In more detail, Section 6 provides the upper bound proof for the case with ontologies and Section 7 provides the matching lower bounds. Sections 8 and 9 cover the ontology-free case and the case of ontologies containing only role inclusions. In Section 10, we investigate the problem of actually computing interpolants and explicit definitions in case they exist, and in Section 11 we draw the connections of our results on DLs to modal logic. Finally, we conclude and point out directions for future work in Section 12.

An appendix available as supplementary material provides a few proofs that were left out of the paper. Here we prove, in particular, our main result about the computational complexity of non-projective definition existence of concept names.

2 Related Work

This paper is an extended version of [5, 6, 51]. We include detailed proofs and additionally discuss the link to concept learning, interpolants between ontologies and concept inclusions, and applications to modal logic.

Related work on Craig interpolation and the Beth definability property has been discussed already in the introduction. We therefore focus on work on deciding interpolant and explicit definition existence. These decision problems have only very recently been investigated. A notable exception is linear temporal logic, LTL, for which the CIP fails and for which decidability of interpolant existence has been shown both over finite linear orderings [41, 42] and over the natural numbers [79]. Note that these results are formulated as separability results for formal languages of finite and, respectively, infinite words: given two regular languages R_1 and R_2 , does there exist a first-order definable language L separating R_1 and R_2 in the sense that $R_1 \subseteq L$ and $L \cap R_2 = \emptyset$. Neither LTL nor Craig interpolation are mentioned in [79, 41, 42]. Using the fact that regular languages are projectively LTL definable and that LTL and first-order logic are equivalent over the natural numbers, it is, however, easy to see that interpolant existence is the same problem as separability of regular languages in first-order logic, modulo the representation of the inputs. We note that this result is just one instance of an ongoing exploration of separation between languages in automata theory. The problem of deciding separation is interesting in this context because obtaining an algorithm for separation yields a far deeper understanding of the class under consideration than just membership [78, 80]. We conjecture that deciding interpolant existence could well play a similar role for understanding fragments of first-order logic.

Indeed, interpolant existence has recently also been studied for the guarded fragment (GF), the two-variable fragment (FO^2) of FO [52], for Horn description logics extending \mathcal{EL} [33], and for first-order modal logics [61]. While GF is a good generalization of modal and description logic in many respects, it neither enjoys the CIP [44] nor the PBDP [10]. Failure of the CIP for FO^2 was shown using algebraic [26, 76] and model-theoretic techniques [73]. Using techniques that are similar to those introduced in this article it is shown in [52] that, in GF, explicit definability and interpolant existence are both 3EXPTIME -complete in general, and 2EXPTIME -complete if the arity of relation symbols is bounded by a constant $c \geq 3$. In FO^2 , explicit definability and interpolant existence are in coN2EXPTIME and 2EXPTIME -hard [52]. Failure of the CIP and PBDP

for first-order modal logics with constant domain is shown in [32, 73]. Both properties also fail for their otherwise well-behaved one-variable and monodic fragments [38]. In [61], the complexity of interpolant existence is investigated for first-order S5 with one variable (and some monodic fragments) and for first-order K with one variable. For S5, explicit definability and interpolant existence turn out to be in coN2EXPTIME and 2EXPTIME -hard while for K only a non-elementary upper bound is shown. These results confirm that for many logics not enjoying the CIP and PBDP, interpolant and explicit definition existence are harder than entailment.

It turns out that this is not always the case. It is shown in [33] that extensions of the description logic \mathcal{EL} with any combination of the universal role, nominals, or inverse roles do not enjoy the CIP nor PBDP, but that interpolant existence and explicit definition existence still have the same complexity as entailment (in PTIME for those that do not admit inverse roles and EXPTIME -complete for those that admit inverse roles). The proofs are rather different from those given in this paper, as they make use of the universal/canonical model that only exists for Horn logics.

We note that for logics that do not enjoy the CIP nor PBDP it is also of interest to look for “small” extensions that enjoy the CIP and PBDP and are decidable. For example, the guarded negation fragment of FO is a decidable extension of GF that enjoys the CIP and the PBDP [11, 15, 14, 12]. Also the two-variable fragment of GF is a decidable extension of \mathcal{ALCH} enjoying both properties [44, 43]. In both cases the complexity of entailment does not increase for the extension (2EXPTIME -complete for the guarded negation fragment and EXPTIME -complete for the two-variable fragment of FO). On the other hand, under mild conditions there is no decidable extension of \mathcal{ALCO} with the universal role nor of modal logic with nominals and the @-operator enjoying the CIP [86].

While the problem of deciding interpolant and explicit definition existence for logics that neither enjoy the CIP nor the PBDP has only been considered rather recently, the problem of computing and deciding the existence of uniform interpolants for logics that do not enjoy the uniform interpolation property (UIP) has been investigated before. Recall that uniform interpolants generalize Craig interpolants in the sense that a uniform interpolant is an interpolant for a fixed φ and all ψ which are entailed by φ and share with φ a fixed set of symbols. First-order logic enjoys the CIP but not the UIP. Propositional intuitionistic logic, local modal logic, and the modal mu-calculus are examples of expressive logics that enjoy the UIP [77, 96, 28], see [59, 46] for more recent results. In de-

scription logic, uniform interpolants of ontologies (extending what we call CI-interpolants in this article) are of particular importance but do not always exist for any standard description logic, including \mathcal{ALC} . The complexity of deciding their existence has been investigated in [68, 70], their size has been considered in [75, 58], and various approaches to computing them have been developed and implemented [56–58, 97].

3 Preliminaries

We introduce the syntax and semantics of the relevant description logics, see also [9]. Let \mathbf{N}_C , \mathbf{N}_R , and \mathbf{N}_I be mutually disjoint and countably infinite sets of *concept*, *role*, and *individual names*. A *role* is a role name s , or an *inverse role* s^- , with s a role name and $(s^-)^- = s$. We use u to denote the *universal role*. A *nominal* takes the form $\{a\}$, with a an individual name. An \mathcal{ALCOT}^u -*concept* is defined according to the syntax rule

$$C, D ::= \top \mid A \mid \{a\} \mid \neg C \mid C \sqcap D \mid \exists r.C$$

where A ranges over concept names, a ranges over individual names, and r over roles and the universal role. We use $C \sqcup D$ as abbreviation for $\neg(\neg C \sqcap \neg D)$, $C \rightarrow D$ for $\neg C \sqcup D$, $C \leftrightarrow D$ for $(C \rightarrow D) \sqcap (D \rightarrow C)$, and $\forall r.C$ for $\neg \exists r.\neg C$. We use several fragments of \mathcal{ALCOT}^u , including \mathcal{ALCOT} , obtained by dropping the universal role, \mathcal{ALCO}^u , obtained by dropping inverse roles, \mathcal{ALCO} , obtained from \mathcal{ALCO}^u by dropping the universal role, and \mathcal{ALC} , obtained from \mathcal{ALCO} by dropping nominals. If \mathcal{L} is any of the DLs above, then an \mathcal{L} -*concept inclusion* (\mathcal{L} -CI) takes the form $C \sqsubseteq D$ with C and D \mathcal{L} -concepts. An \mathcal{L} -*ontology* is a finite set of \mathcal{L} -CIs. We also consider DLs with *role inclusions* (RIs), expressions of the form $r \sqsubseteq s$, where r and s are roles. As usual, the addition of RIs is indicated by adding the letter \mathcal{H} to the name of the DL, where inverse roles occur in RIs only if the DL admits inverse roles. Thus, for example, \mathcal{ALCH} -*ontologies* are finite sets of \mathcal{ALC} -CIs and RIs not using inverse roles and \mathcal{ALCHOT}^u -*ontologies* are finite sets of \mathcal{ALCOT}^u -CIs and RIs. In what follows we use DL_{nr} to denote the set of DLs \mathcal{ALCO} , \mathcal{ALCOT} , \mathcal{ALCH} , \mathcal{ALCHI} , \mathcal{ALCHO} , \mathcal{ALCHOI} , and their extensions with the universal role. To simplify notation we do not drop the letter \mathcal{H} when speaking about the concepts and CIs of a DL with RIs. Thus, for example, we sometimes use the expressions \mathcal{ALCHO} -concept and \mathcal{ALCHO} -CI to denote \mathcal{ALCO} -concepts and CIs, respectively. An *RI-ontology* is an ontology containing RIs only.

The semantics is defined in terms of *interpretations* $\mathcal{I} = (\Delta^{\mathcal{I}}, \cdot^{\mathcal{I}})$, where $\Delta^{\mathcal{I}}$ is a non-empty set, called *domain* of \mathcal{I} , and $\cdot^{\mathcal{I}}$ is a function mapping every $A \in \mathbf{N}_{\mathcal{C}}$ to a subset of $\Delta^{\mathcal{I}}$, every $s \in \mathbf{N}_{\mathbf{R}}$ to a subset of $\Delta^{\mathcal{I}} \times \Delta^{\mathcal{I}}$, the universal role u to $\Delta^{\mathcal{I}} \times \Delta^{\mathcal{I}}$, and every $a \in \mathbf{N}_{\mathbf{I}}$ to an element in $\Delta^{\mathcal{I}}$. Given a role name $s \in \mathbf{N}_{\mathbf{R}}$, we set $(s^-)^{\mathcal{I}} = \{(d, e) \in \Delta^{\mathcal{I}} \times \Delta^{\mathcal{I}} \mid (e, d) \in s^{\mathcal{I}}\}$. Moreover, the *extension* $C^{\mathcal{I}}$ of an \mathcal{L} -concept C in \mathcal{I} is defined as follows, where r ranges over roles and the universal role:

$$\begin{aligned} \top^{\mathcal{I}} &= \Delta^{\mathcal{I}}, \\ \{a\}^{\mathcal{I}} &= \{a^{\mathcal{I}}\}, \\ \neg C^{\mathcal{I}} &= \Delta^{\mathcal{I}} \setminus C^{\mathcal{I}}, \\ (C \sqcap D)^{\mathcal{I}} &= C^{\mathcal{I}} \cap D^{\mathcal{I}}, \\ (\exists r.C)^{\mathcal{I}} &= \{d \in \Delta^{\mathcal{I}} \mid \text{there exists } e \in C^{\mathcal{I}} : (d, e) \in r^{\mathcal{I}}\}. \end{aligned}$$

An interpretation \mathcal{I} *satisfies* an \mathcal{L} -CI $C \sqsubseteq D$ if $C^{\mathcal{I}} \subseteq D^{\mathcal{I}}$ and an RI $r \sqsubseteq s$ if $r^{\mathcal{I}} \subseteq s^{\mathcal{I}}$. We say that \mathcal{I} is a *model* of an ontology \mathcal{O} if it satisfies all inclusions in it. We say that an inclusion α follows from an ontology \mathcal{O} , in symbols $\mathcal{O} \models \alpha$, if every model of \mathcal{O} satisfies α . We write $\mathcal{O} \models C \equiv D$ if $\mathcal{O} \models C \sqsubseteq D$ and $\mathcal{O} \models D \sqsubseteq C$. We drop \mathcal{O} if it is empty and write $\models C \sqsubseteq D$ for $\emptyset \models C \sqsubseteq D$. A concept C is *satisfiable* w.r.t. an ontology \mathcal{O} if there is a model \mathcal{I} of \mathcal{O} with $C^{\mathcal{I}} \neq \emptyset$. We use a few well known complexity bounds for reasoning in DLs from DL_{nr} . The \mathcal{L} -*subsumption problem* is the problem to decide for any \mathcal{L} -ontology \mathcal{O} and \mathcal{L} -CI $C \sqsubseteq D$ whether $\mathcal{O} \models C \sqsubseteq D$. The *ontology-free \mathcal{L} -subsumption problem* and the *RI-ontology \mathcal{L} -subsumption problem* are the sub-problems of the \mathcal{L} -subsumption problem in which the ontology is empty or an RI-ontology, respectively. For any $\mathcal{L} \in \text{DL}_{\text{nr}}$, the \mathcal{L} -subsumption problem is EXPTIME-complete [8, 1]. If \mathcal{L} admits the universal role or both inverse roles and nominals, then ontologies can be encoded in concepts and so ontology-free \mathcal{L} -subsumption and RI-ontology \mathcal{L} -subsumption are also EXPTIME-complete. In the remaining cases, that is for \mathcal{ALCO} , \mathcal{ALCH} , \mathcal{ALCHO} , and \mathcal{ALCHI} , \mathcal{L} -subsumption becomes PSPACE-complete [8, 1].

A *signature* Σ is a set of concept, role, and individual names, uniformly referred to as *symbols*. Following standard practice we do not regard the universal role as a symbol but as a logical connective. Thus, the universal role is not contained in any signature. We use $\text{sig}(X)$ to denote the set of symbols used in any syntactic object X such as a concept or an ontology. An $\mathcal{L}(\Sigma)$ -*concept* is an \mathcal{L} -concept C with $\text{sig}(C) \subseteq \Sigma$. A Σ -role r is a role with $\text{sig}(r) \subseteq \Sigma$. The *size* of a (finite) syntactic object X , denoted $\|X\|$, is the number of symbols needed to represent it as a word.

[AtomC]	for all $(d, e) \in S$: $d \in A^{\mathcal{I}}$ iff $e \in A^{\mathcal{J}}$
[AtomI]	for all $(d, e) \in S$: $d = a^{\mathcal{I}}$ iff $e = a^{\mathcal{J}}$
[Forth]	if $(d, e) \in S$ and $(d, d') \in r^{\mathcal{I}}$, then there is a e' with $(e, e') \in r^{\mathcal{J}}$ and $(d', e') \in S$.
[Back]	if $(d, e) \in S$ and $(e, e') \in r^{\mathcal{J}}$, then there is a d' with $(d, d') \in r^{\mathcal{I}}$ and $(d', e') \in S$.

Fig. 1. Conditions on $S \subseteq \Delta^{\mathcal{I}} \times \Delta^{\mathcal{J}}$.

We next recall model-theoretic characterizations when elements in interpretations are indistinguishable by concepts formulated in one of the DLs \mathcal{L} introduced above. A *pointed interpretation* is a pair \mathcal{I}, d with \mathcal{I} an interpretation and $d \in \Delta^{\mathcal{I}}$. For pointed interpretations \mathcal{I}, d and \mathcal{J}, e and a signature Σ , we write $\mathcal{I}, d \equiv_{\mathcal{L}, \Sigma} \mathcal{J}, e$ and say that \mathcal{I}, d and \mathcal{J}, e are $\mathcal{L}(\Sigma)$ -*equivalent* if $d \in C^{\mathcal{I}}$ iff $e \in C^{\mathcal{J}}$, for all $\mathcal{L}(\Sigma)$ -concepts C .

As for the model-theoretic characterizations, we start with \mathcal{ALC} . Let Σ be a signature. A relation $S \subseteq \Delta^{\mathcal{I}} \times \Delta^{\mathcal{J}}$ is an $\mathcal{ALC}(\Sigma)$ -*bisimulation* if conditions [AtomC], [Forth], and [Back] from Figure 1 hold, where A and r range over all concept and role names in Σ , respectively. We write $\mathcal{I}, d \sim_{\mathcal{ALC}, \Sigma} \mathcal{J}, e$ and call \mathcal{I}, d and \mathcal{J}, e $\mathcal{ALC}(\Sigma)$ -*bisimilar* if there exists an $\mathcal{ALC}(\Sigma)$ -bisimulation S such that $(d, e) \in S$. For \mathcal{ALCO} , we define $\sim_{\mathcal{ALCO}, \Sigma}$ analogously, but now demand that, in Figure 1, also condition [AtomI] holds for all individual names $a \in \Sigma$. For languages \mathcal{L} with inverse roles, we demand that, in Figure 1, r additionally ranges over inverse roles. For languages \mathcal{L} with the universal role we extend the respective conditions by demanding that the domain $\text{dom}(S)$ and range $\text{ran}(S)$ of S contain $\Delta^{\mathcal{I}}$ and $\Delta^{\mathcal{J}}$, respectively. If a DL \mathcal{L} has RIs, then we use $\mathcal{I}, d \sim_{\mathcal{L}, \Sigma} \mathcal{J}, e$ to state that $\mathcal{I}, d \sim_{\mathcal{L}', \Sigma} \mathcal{J}, e$ for the fragment \mathcal{L}' of \mathcal{L} without RIs.

The next lemma summarizes the model-theoretic characterizations for all relevant DLs [67, 40]. For the definition of ω -saturated structures, we refer the reader to [24].

Lemma 3. *Let \mathcal{I}, d and \mathcal{J}, e be pointed interpretations and ω -saturated. Let $\mathcal{L} \in \text{DL}_{nr}$ and Σ a signature. Then*

$$\mathcal{I}, d \equiv_{\mathcal{L}, \Sigma} \mathcal{J}, e \quad \text{iff} \quad \mathcal{I}, d \sim_{\mathcal{L}, \Sigma} \mathcal{J}, e.$$

For the “if”-direction, the ω -saturatedness condition can be dropped.

4 Craig Interpolation and Beth Definability

We introduce interpolants and the Craig interpolation property (CIP) as well as implicit and explicit definitions and the (projective) Beth definabil-

ity property ((P)BDP). Recall from the introduction that there are two forms of interpolants, one pertaining to concepts and the other pertaining to concept inclusions. We start the discussion here with the former one, and discuss CI-interpolants later. For concept interpolants, we establish a close link between interpolants and separators of positive and negative data examples, show that logics in DL_{nr} do not enjoy the CIP nor PBDP, and determine which DLs in DL_{nr} enjoy the BDP.

Let \mathcal{O} be an \mathcal{L} -ontology, C_1, C_2 be \mathcal{L} -concepts, and let Σ be a signature. Then, an \mathcal{L} -concept D is an $\mathcal{L}(\Sigma)$ -interpolant for $C_1 \sqsubseteq C_2$ under \mathcal{O} , if $\text{sig}(D) \subseteq \Sigma$, $\mathcal{O} \models C_1 \sqsubseteq D$, and $\mathcal{O} \models D \sqsubseteq C_2$. If \mathcal{O} is empty, then we drop it and call an $\mathcal{L}(\Sigma)$ -interpolant for $C_1 \sqsubseteq C_2$ under \mathcal{O} simply an $\mathcal{L}(\Sigma)$ -interpolant for $C_1 \sqsubseteq C_2$. Observe that Σ is arbitrary in this definition, so it does not follow from $\mathcal{O} \models C_1 \sqsubseteq C_2$ that an $\mathcal{L}(\Sigma)$ -interpolant for $C_1 \sqsubseteq C_2$ under \mathcal{O} exists. If \mathcal{O} is empty, then we obtain the standard definition of the Craig interpolation property by demanding that for $\Sigma = \text{sig}(C_1) \cap \text{sig}(C_2)$ from $\models C_1 \sqsubseteq C_2$ it follows that there exists an $\mathcal{L}(\Sigma)$ -interpolant for $C_1 \sqsubseteq C_2$. The obvious generalization of this definition to non-empty ontologies, however, does not work. Consider, for instance, $\mathcal{O} = \{A_1 \sqsubseteq A_2, A_2 \sqsubseteq A_3\}$ and $A_1 \sqsubseteq A_3$. Then for $\Sigma = \text{sig}(A_1) \cap \text{sig}(A_3) = \emptyset$, we have $\mathcal{O} \models A_1 \sqsubseteq A_3$, but there does not exist any $\mathcal{L}(\Sigma)$ -interpolant for $A_1 \sqsubseteq A_3$ under \mathcal{O} . In fact, to generalize the Craig interpolation property to non-empty ontologies, one has to split the ontology \mathcal{O} into two. Hence, we adopt here the following definition of the Craig interpolation property in DLs from [90]. We set $\text{sig}(\mathcal{O}, C) = \text{sig}(\mathcal{O}) \cup \text{sig}(C)$, for any ontology \mathcal{O} and concept C .

Definition 4. *A DL \mathcal{L} enjoys the Craig interpolation property (CIP) if for any \mathcal{L} -ontologies $\mathcal{O}_1, \mathcal{O}_2$ and \mathcal{L} -concepts C_1, C_2 such that $\mathcal{O}_1 \cup \mathcal{O}_2 \models C_1 \sqsubseteq C_2$ there exists an $\mathcal{L}(\Sigma)$ -interpolant for $C_1 \sqsubseteq C_2$ under $\mathcal{O}_1 \cup \mathcal{O}_2$, where $\Sigma = \text{sig}(\mathcal{O}_1, C_1) \cap \text{sig}(\mathcal{O}_2, C_2)$. If $\mathcal{O}_1, \mathcal{O}_2$ range over \mathcal{L} -ontologies containing RIs only or $\mathcal{O}_1 = \mathcal{O}_2 = \emptyset$, then we say that \mathcal{L} enjoys the CIP for RI-ontologies and the CIP for the empty ontology, respectively.*

Note that the CIP for the empty ontology coincides with the standard definition of the CIP mentioned before. It is shown in [90] that the DLs \mathcal{ALC} and \mathcal{ALCI} and their extensions with qualified number restrictions and the universal role all enjoy the CIP. In contrast, no DL in DL_{nr} enjoys the CIP. This is implicitly proved in [90] and is shown in Theorem 12 below. The following illustrating example is folklore and shows that this holds even for the empty ontology for logics admitting nominals.

Example 5. Consider $C_1 = \{a\} \sqcap \exists r.\{a\}$ and $C_2 = \{b\} \rightarrow \exists r.\{b\}$. Then $\models C_1 \sqsubseteq C_2$ but there does not exist any $\mathcal{ALCHOI}^u(\{r\})$ -interpolant for $C_1 \sqsubseteq C_2$. Intuitively, no such interpolant D exists as it would have to be true in exactly the elements x with $r(x, x)$ and no such $\mathcal{ALCHOI}^u(\{r\})$ -concept exists. A formal proof is given in Example 22 below. \dashv

As discussed in the introduction, the close link between separation of data examples and interpolants is one of our main motivations for studying interpolants for DLs with nominals. We next formalize this link. A *database* \mathcal{D} is a finite set of assertions of the form $A(a)$ and $r(a, b)$ with a, b individuals, A a concept name, and r a role name. By $\text{ind}(\mathcal{D})$ we denote the set of individual names in \mathcal{D} . A *knowledge base* (KB) is a pair $\mathcal{K} = (\mathcal{O}, \mathcal{D})$ consisting of an ontology \mathcal{O} and database \mathcal{D} . An interpretation \mathcal{I} is a *model of* \mathcal{K} if it is a model of \mathcal{O} , $a^{\mathcal{I}} \in A^{\mathcal{I}}$ for all $A(a) \in \mathcal{D}$, and $(a^{\mathcal{I}}, b^{\mathcal{I}}) \in r^{\mathcal{I}}$ for all $r(a, b) \in \mathcal{D}$. An assertion $C(a)$ with C a concept and a an individual *follows from* \mathcal{K} , in symbols $\mathcal{K} \models C(a)$, if every model \mathcal{I} of \mathcal{K} satisfies $a^{\mathcal{I}} \in C^{\mathcal{I}}$. A *labelled data set* consists of two sets, P and N , of positive and negative examples each containing pairs (\mathcal{D}, a) with $a \in \text{ind}(\mathcal{D})$. Let \mathcal{O} be an ontology. An $\mathcal{L}(\Sigma)$ -*separator* for \mathcal{O}, P, N is an $\mathcal{L}(\Sigma)$ -concept C such that $(\mathcal{O}, \mathcal{D}) \models C(a)$ for all $(\mathcal{D}, a) \in P$ and $(\mathcal{O}, \mathcal{D}) \models \neg C(a)$ for all $(\mathcal{D}, a) \in N$. The following result establishes a one-to-one correspondence between interpolants and separators, modulo straightforward polynomial time reductions. Note that we do not require the frequent assumption that the database is uniform across the examples in the sense that $\mathcal{D} = \mathcal{D}'$ for all $(\mathcal{D}, a), (\mathcal{D}', a') \in P \cup N$ [50].

Theorem 6. *Let $\mathcal{L} \in DL_{nr}$ admit nominals. Then one can construct for any ontology \mathcal{O} , labelled data sets P, N , and signature Σ with $\Sigma \cap \{a \mid (\mathcal{D}, a) \in P \cup N\} = \emptyset$ in polynomial time \mathcal{L} -ontologies $\mathcal{O}_1, \mathcal{O}_2$ and \mathcal{L} -concepts C_1, C_2 such that $\Sigma = \text{sig}(\mathcal{O}_1, C_1) \cap \text{sig}(\mathcal{O}_2, C_2)$ and the following conditions are equivalent for all \mathcal{L} -concepts C :*

1. C is an $\mathcal{L}(\Sigma)$ -separator for \mathcal{O}, P, N ;
2. C is an $\mathcal{L}(\Sigma)$ -interpolant for $C_1 \sqsubseteq C_2$ under $\mathcal{O}_1 \cup \mathcal{O}_2$.

Conversely, assume that \mathcal{L} -ontologies $\mathcal{O}_1, \mathcal{O}_2$, and \mathcal{L} -concepts C_1, C_2 are given. Then one can construct in polynomial time an ontology \mathcal{O} and labelled data sets P, N such that Conditions (1) and (2) are equivalent for $\Sigma = \text{sig}(\mathcal{O}_1, C_1) \cap \text{sig}(\mathcal{O}_2, C_2)$.

Proof. Assume \mathcal{O}, P, N , and Σ are given. Let $P = \{(\mathcal{D}_1, a_1), \dots, (\mathcal{D}_n, a_n)\}$ and $N = \{(\mathcal{D}_{n+1}, a_{n+1}), \dots, (\mathcal{D}_{n+m}, a_{n+m})\}$. If $\mathcal{L} \in DL_{nr}$ admits nominals and the universal role, then a pair (\mathcal{D}, a) can be represented using the

\mathcal{L} -concept $C_{\mathcal{D},a} = \{a\} \sqcap \exists u. C_{\mathcal{D}}$, where $C_{\mathcal{D}}$ is the conjunction of all $\{b\} \sqcap A$ with $A(b) \in \mathcal{D}$ and $\{b\} \sqcap \exists r. \{c\}$ with $r(b,c) \in \mathcal{D}$. Pick for any symbol X not in Σ a fresh copy X' . Let $\mathcal{O}_1 = \mathcal{O}$ and obtain \mathcal{O}_2 from \mathcal{O} by replacing all symbols not in Σ by their copies. Let $C_1 = C_{\mathcal{D}_1,a_1} \sqcup \dots \sqcup C_{\mathcal{D}_n,a_n}$ and obtain C_2 from $\neg(C_{\mathcal{D}_{n+1},a_{n+1}} \sqcup \dots \sqcup C_{\mathcal{D}_{n+m},a_{n+m}})$ by replacing all symbols not in Σ by their copies. If \mathcal{L} admits the universal role, then $\mathcal{O}_1, \mathcal{O}_2$ and C_1, C_2 are as required. Otherwise replace the universal role in any $C_{\mathcal{D}_i,a_i}$ by fresh role names not in Σ . The resulting C_1, C_2 are still as required.

Conversely, assume that $\mathcal{O}_1, \mathcal{O}_2$ and C_1, C_2 are given. Introduce fresh individual names a, b and fresh concept names A, B and let $P = \{(\{A(a)\}, a)\}$, $N = \{(\{B(b)\}, b)\}$ and $\mathcal{O} = \mathcal{O}_1 \cup \mathcal{O}_2 \cup \{A \sqsubseteq C_1, B \sqsubseteq \neg C_2\}$. Then \mathcal{O}, P, N is as required. \square

We next introduce the relevant definability notions. Let \mathcal{O} be an ontology and C, C_0 be concepts. Let Σ be a signature. An $\mathcal{L}(\Sigma)$ -concept D is an *explicit $\mathcal{L}(\Sigma)$ -definition of C_0 under \mathcal{O} and C* if $\mathcal{O} \models C \sqsubseteq (C_0 \leftrightarrow D)$. We call C_0 *explicitly $\mathcal{L}(\Sigma)$ -definable under \mathcal{O} and C* if there is an explicit $\mathcal{L}(\Sigma)$ -definition of C_0 under \mathcal{O} and C . If $C = \top$ or \mathcal{O} is empty, then we drop \mathcal{O} and C , respectively. For instance, an $\mathcal{L}(\Sigma)$ -concept D is an explicit $\mathcal{L}(\Sigma)$ -definition of C_0 under \mathcal{O} if $\mathcal{O} \models C_0 \equiv D$. The following example illustrates the link between explicit definitions of nominals and referring expressions discussed in the introduction and also indicates that often one can single out an individual from a set of individuals using an explicit definition without being able to provide an ‘absolute’ explicit definition of that individual.

Example 7. Let \mathbb{L} be an abbreviation for the \mathcal{ALCO} -concept

$$\{\mathcal{ALC}\} \sqcup \{\mathcal{ALCO}\} \sqcup \{\mathcal{ALCO}^u\} \sqcup \{\text{ML}\} \sqcup \{\text{ML}_n\} \sqcup \{\text{ML}_n^u\},$$

where ML , ML_n , and ML_n^u are modal logics introduced below in Section 11. Let \mathcal{O} be the ontology consisting of the following CIs:

$$\begin{aligned}
\{\mathcal{ALC}\} \sqcup \{\mathcal{ALCO}\} \sqcup \{\mathcal{ALCO}^u\} &\sqsubseteq \exists \text{hasOperator.DLOperator} \sqcap \\
&\quad \neg \exists \text{hasOperator.MLOperator}, \\
\{\text{ML}\} \sqcup \{\text{ML}_n\} \sqcup \{\text{ML}_n^u\} &\sqsubseteq \exists \text{hasOperator.MLOperator} \sqcap \\
&\quad \neg \exists \text{hasOperator.DLOperator}, \\
\{\mathcal{ALC}\} \sqcup \{\text{ML}\} &\sqsubseteq \text{EnjoysCIP}, \\
\text{EnjoysCIP} &\sqsubseteq \text{EnjoysPBDP}, \\
\text{EnjoysPBDP} &\sqsubseteq \text{EnjoysBDP}, \\
\{\mathcal{ALCO}^u\} \sqcup \{\text{ML}_n^u\} &\sqsubseteq \text{EnjoysBDP} \sqcap \neg \text{EnjoysPBDP}, \\
\{\mathcal{ALCO}\} \sqcup \{\text{ML}_n\} &\sqsubseteq \neg \text{EnjoysBDP}.
\end{aligned}$$

Then $\mathcal{O} \models \text{L} \sqsubseteq (\{\mathcal{ALC}\} \leftrightarrow \text{EnjoysCIP} \sqcap \exists \text{hasOperator.DLOperator})$. Hence, $\{\mathcal{ALC}\}$ is explicitly $\mathcal{ALCO}(\Sigma_0)$ -definable under \mathcal{O} and L , with $\Sigma_0 = \{\text{EnjoysCIP}, \text{DLOperator}, \text{hasOperator}\}$. However, $\{\mathcal{ALC}\}$ is not explicitly $\mathcal{ALCO}(\Sigma)$ -definable under \mathcal{O} , for any signature Σ with $\mathcal{ALC} \notin \Sigma$, since there does not exist an $\mathcal{ALCO}(\Sigma)$ -concept C such that $\mathcal{O} \models \{\mathcal{ALC}\} \equiv C$.
 \dashv

We next define when a concept is implicitly definable. For a signature Σ , the Σ -reduct $\mathcal{I}_{|\Sigma}$ of an interpretation \mathcal{I} coincides with \mathcal{I} except that no non- Σ symbol is interpreted in $\mathcal{I}_{|\Sigma}$. A concept C_0 is called *implicitly Σ -definable under \mathcal{O} and C* if the Σ -reduct of any pointed model \mathcal{I}, d with \mathcal{I} a model of \mathcal{O} and $d \in C^{\mathcal{I}}$ determines whether $d \in C_0^{\mathcal{I}}$. More formally, C_0 is implicitly Σ -definable under \mathcal{O} and C if the following holds for all models \mathcal{I} and \mathcal{J} of \mathcal{O} and $d \in \Delta^{\mathcal{I}} = \Delta^{\mathcal{J}}$: if $\mathcal{I}_{|\Sigma} = \mathcal{J}_{|\Sigma}$ and $d \in C^{\mathcal{I}}$, then $d \in C_0^{\mathcal{I}}$ iff $d \in C_0^{\mathcal{J}}$. If $C = \top$, then we drop C and say that C_0 is implicitly Σ -definable under \mathcal{O} . To illustrate, observe that in Example 7, $\{\mathcal{ALC}\}$ is not implicitly Σ -definable, for any Σ such that $\mathcal{ALC} \notin \Sigma$ under \mathcal{O} . Implicit definability can be reformulated as a standard reasoning problem as follows: a concept C_0 is implicitly Σ -definable under \mathcal{O} and C iff

$$\mathcal{O} \cup \mathcal{O}_{\Sigma} \models C \sqcap C_0 \sqsubseteq C_{\Sigma} \rightarrow C_{0\Sigma} \quad (1)$$

where \mathcal{O}_{Σ} , C_{Σ} , and $C_{0\Sigma}$ are obtained from \mathcal{O} , C and, respectively, C_0 , by replacing every non- Σ symbol uniformly by a fresh symbol. If a concept is explicitly $\mathcal{L}(\Sigma)$ -definable under \mathcal{O} and C , then it is implicitly Σ -definable under \mathcal{O} and C , for any language \mathcal{L} . A logic enjoys the projective Beth definability property if the converse implication holds as well.

Definition 8. A DL \mathcal{L} enjoys the projective Beth definability property (PBDP) if for any \mathcal{L} -ontology \mathcal{O} , \mathcal{L} -concepts C and C_0 , and signature $\Sigma \subseteq \text{sig}(\mathcal{O}, C)$ the following holds: if C_0 is implicitly Σ -definable under \mathcal{O} and C , then C_0 is explicitly $\mathcal{L}(\Sigma)$ -definable under \mathcal{O} and C . If \mathcal{O} ranges over \mathcal{L} -ontologies containing RIs only or $\mathcal{O} = \emptyset$, then we say that \mathcal{L} enjoys the PBDP for RI-ontologies and the PBDP for the empty ontology, respectively.

The DLs \mathcal{ALC} , \mathcal{ALCI} , and their extensions with qualified number restrictions and the universal role all enjoy the PBDP [90]. The following example shows that, in contrast, \mathcal{ALCH} does not.

Example 9. Consider $\mathcal{O} = \{r \sqsubseteq r_1, r \sqsubseteq r_2\}$ and let

$$C = ((\neg \exists r. \top \sqcap \exists r_1. A) \rightarrow \forall r_2. \neg A) \sqcap ((\neg \exists r. \top \sqcap \exists r_1. \neg A) \rightarrow \forall r_2. A).$$

Let $\Sigma = \{r_1, r_2\}$ and $C_0 = \exists r. \top$. Then the concept $D = \exists r_1 \sqcap r_2. \top$ is an explicit definition of C_0 under \mathcal{O} and C in the extension of \mathcal{ALCH} with role intersection (the semantics of $r_1 \sqcap r_2$ is defined in the obvious way). Hence C_0 is implicitly Σ -definable under \mathcal{O} and C . There does not exist an explicit $\mathcal{ALCH}(\Sigma)$ -definition of C_0 under \mathcal{O} and C , however. Intuitively, the reason is that role intersection cannot be expressed in \mathcal{ALCH} (see Example 24 below for a proof).

Note that an example without "background concept" C can be obtained by taking the ontology

$$\mathcal{O}' = \{ r \sqsubseteq r_1, \quad r \sqsubseteq r_2, \\ \neg \exists r. \top \sqcap \exists r_1. A \sqsubseteq \forall r_2. \neg A, \quad \neg \exists r. \top \sqcap \exists r_1. \neg A \sqsubseteq \forall r_2. A \}$$

and asking for an explicit $\mathcal{ALCH}(\{r_1, r_2\})$ -definition of $\exists r. \top$ under \mathcal{O}' . \dashv

It is known that the CIP and PBDP are tightly linked [90]. We state the inclusion for logics in DL_{nr} only, but the proof shows that it holds under rather mild conditions.

Lemma 10. *If $\mathcal{L} \in \text{DL}_{nr}$ enjoys the CIP, then \mathcal{L} enjoys the PBDP.*

Proof. Assume that an \mathcal{L} -concept C_0 is implicitly Σ -definable under an \mathcal{L} -ontology \mathcal{O} and \mathcal{L} -concept C , for some signature Σ . Then (1) holds. Take an $\mathcal{L}(\Sigma)$ -interpolant D for $C \sqcap C_0 \sqsubseteq C_\Sigma \rightarrow C_{0\Sigma}$ under $\mathcal{O} \cup \mathcal{O}_\Sigma$. Then D is an explicit $\mathcal{L}(\Sigma)$ -definition of C_0 under \mathcal{O} and C . \square

An important special case of explicit definability is the explicit definability of a concept name A from $\text{sig}(\mathcal{O}, C) \setminus \{A\}$ under an ontology \mathcal{O} and concept C . For this case, we also consider the following *non-projective version* of the Beth definability property.

Definition 11. *A DL \mathcal{L} enjoys the Beth definability property (BDP) if for any \mathcal{L} -ontology \mathcal{O} , concept C , and any concept name A the following holds: if A is implicitly $(\text{sig}(\mathcal{O}, C) \setminus \{A\})$ -definable*

under \mathcal{O} and C , then A is explicitly $\mathcal{L}(\text{sig}(\mathcal{O}, C) \setminus \{A\})$ -definable under \mathcal{O} and C . If \mathcal{O} ranges over \mathcal{L} -ontologies containing RIs only or $\mathcal{O} = \emptyset$, then we say that \mathcal{L} enjoys the BDP for RI-ontologies and the BDP for the empty ontology, respectively.

Clearly the PBDP entails the BDP, but we will see below that the converse direction does not always hold. In fact, the following theorem states that no DL in DL_{nr} enjoys the CIP or PBDP, but that quite a few DLs in DL_{nr} enjoy the BDP. Moreover, all DLs in DL_{nr} enjoy the BDP for RI-ontologies and for the empty ontology.

As mentioned before, the theorem is mostly folklore and therefore proved in the appendix.

Theorem 12. *The following statements hold.*

- (1) *No $\mathcal{L} \in \text{DL}_{\text{nr}}$ enjoys the CIP nor the PBDP. The CIP and PBDP also do not hold for RI-ontologies and, if \mathcal{L} admits nominals, the empty ontology.*
- (2) *All $\mathcal{L} \in \text{DL}_{\text{nr}} \setminus \{\mathcal{ALCCO}, \mathcal{ALCHO}\}$ enjoy the BDP. \mathcal{ALCCO} and \mathcal{ALCHO} do not enjoy the BDP.*
- (3) *All $\mathcal{L} \in \text{DL}_{\text{nr}}$ enjoy the BDP for RI-ontologies and the BDP for the empty ontology.*

We have seen that all $\mathcal{L} \in \text{DL}_{\text{nr}} \setminus \{\mathcal{ALCCO}, \mathcal{ALCHO}\}$ enjoy the BDP. One might be tempted to conjecture that this holds as well if concept names are replaced by nominals; that is to say, a nominal $\{a\}$ that is implicitly definable using symbols distinct from a is explicitly definable using symbols distinct from a . Rather surprisingly, the following example shows that this is not the case for any DL in DL_{nr} with nominals (for DLs without nominals this notion is clearly meaningless).

Example 13. Let $\mathcal{L} \in \text{DL}_{\text{nr}}$ admit nominals and assume that

$$\begin{aligned} \mathcal{O} &= \{\{a\} \sqsubseteq \exists r.\{a\}, \\ &\quad A \sqcap \neg\{a\} \sqsubseteq \forall r.(\neg\{a\} \rightarrow \neg A), \\ &\quad \neg A \sqcap \neg\{a\} \sqsubseteq \forall r.(\neg\{a\} \rightarrow A)\}. \end{aligned}$$

Thus, \mathcal{O} implies that a is r -reflexive and that no element distinct from a is r -reflexive. Let $\Sigma = \{r, A\}$. Then $\{a\}$ is implicitly Σ -definable under \mathcal{O} since we have the following explicit definition in first-order logic:

$$\mathcal{O} \models \forall x((x = a) \leftrightarrow r(x, x)),$$

but one can show that $\{a\}$ is not explicitly $\mathcal{L}(\Sigma)$ -definable under \mathcal{O} for any $\mathcal{L} \in \text{DL}_{\text{nr}}$ with nominals. Indeed, the interpretation \mathcal{I} in Figure 2 (where $a^{\mathcal{I}} = a$) is a model of \mathcal{O} and the relation $S = \Delta^{\mathcal{I}} \times \Delta^{\mathcal{I}}$ is an $\mathcal{L}(\Sigma)$ -bisimulation on \mathcal{I} . Thus, Lemma 3 implies $\mathcal{I}, a^{\mathcal{I}} \equiv_{\mathcal{L}, \Sigma} \mathcal{I}, d$ and there is no explicit $\mathcal{L}(\Sigma)$ -definition for $\{a\}$ under \mathcal{O} , as any such definition would apply to d as well. \dashv




Fig. 2. Failure of the PBDP for $\mathcal{L} \in \text{DL}_{\text{nr}}$ with nominals.

The CIP defined above is concerned with interpolating concepts. In the context of modular ontologies and forgetting there is also an interest in interpolating concept inclusions [55]. For simplicity, we only consider DLs without RIs. Let \mathcal{L} not admit RIs and let \mathcal{O} and \mathcal{O}' be \mathcal{L} -ontologies. We write $\mathcal{O} \models \mathcal{O}'$ if $\mathcal{O} \models \alpha$ for all $\alpha \in \mathcal{O}'$. Then an \mathcal{L} -ontology \mathcal{O}'' is called an \mathcal{L} -CI interpolant for \mathcal{O} and \mathcal{O}' if $\text{sig}(\mathcal{O}'') \subseteq \text{sig}(\mathcal{O}) \cap \text{sig}(\mathcal{O}')$, $\mathcal{O} \models \mathcal{O}''$, and $\mathcal{O}'' \models \mathcal{O}'$. If the particular language \mathcal{L} is clear from the context or not important we drop it and call \mathcal{L} -CI interpolants simply CI-interpolants.

Definition 14. *Let \mathcal{L} be a DL that does not admit RIs. Then \mathcal{L} has the CI-interpolation property if for all \mathcal{L} -ontologies \mathcal{O} and \mathcal{O}' such that $\mathcal{O} \models \mathcal{O}'$ there exists an \mathcal{L} -CI-interpolant for \mathcal{O} and \mathcal{O}' .*

Observe that for any $\mathcal{L} \in \text{DL}_{\text{nr}}$ that does not admit RIs and \mathcal{L} -ontology \mathcal{O} one can construct in linear time an \mathcal{L} -concept D such that \mathcal{O} and $\{\top \sqsubseteq D\}$ are equivalent in the sense that $\mathcal{O} \models \top \sqsubseteq D$ and $\{\top \sqsubseteq D\} \models \mathcal{O}$. Hence \mathcal{L} has the CI-interpolation property if for all \mathcal{L} -ontologies \mathcal{O} and \mathcal{L} -CIs $C \sqsubseteq D$ such that $\mathcal{O} \models C \sqsubseteq D$ there exists an \mathcal{L} -CI-interpolant for \mathcal{O} and $\{C \sqsubseteq D\}$. It is known that \mathcal{ALC} and its extensions with inverse roles, qualified number restrictions, and the universal role enjoy the CI-interpolation property [55]. The following example shows that no DL in DL_{nr} that does not admit RIs enjoys the CI-interpolation property.

Example 15. We modify the ontology given in Example 13. Let

$$\mathcal{O}' = \{A \sqsubseteq \forall r. \neg A, \neg A \sqsubseteq \forall r. A\}.$$

We have that $\mathcal{O}' \models \{a\} \sqsubseteq \neg \exists r. \{a\}$, but there does not exist an \mathcal{ALCCOT}^u -CI-interpolant for \mathcal{O}' and $\{a\} \sqsubseteq \neg \exists r. \{a\}$, since one cannot express using an $\mathcal{ALCCOT}^u(\{r\})$ -CI that $\forall x \neg r(x, x)$. Indeed, assume for a proof by contradiction that there exists an \mathcal{ALCCOT}^u -CI-interpolant \mathcal{O}'' for \mathcal{O}' and $\{\{a\} \sqsubseteq \neg \exists r. \{a\}\}$. Consider the interpretations $\mathcal{I}_1, \mathcal{I}_2$ in Figure 3, where $\mathcal{I}_1 \models \mathcal{O}'$, $a^{\mathcal{I}_2} \in (\{a\} \sqcap \exists r. \{a\})^{\mathcal{I}_2}$, and $\mathcal{I}_1, d \sim_{\mathcal{ALCCOT}^u, \{r\}} \mathcal{I}_2, a^{\mathcal{I}_2}$. Since $\mathcal{I}_1 \models \mathcal{O}'$, we have that $\mathcal{I}_1 \models \mathcal{O}''$, where \mathcal{O}'' can be assumed to be of the form $\{\top \sqsubseteq D\}$, with $\text{sig}(D) \subseteq \{r\}$. Thus, $d \in (\forall u. D)^{\mathcal{I}_1}$ and, from $\mathcal{I}_1, d \sim_{\mathcal{ALCCOT}^u, \{r\}} \mathcal{I}_2, a^{\mathcal{I}_2}$, we obtain by Lemma 3 that $a^{\mathcal{I}_2} \in (\forall u. D)^{\mathcal{I}_2}$. This implies $\mathcal{I}_2 \models \mathcal{O}''$, and hence $\mathcal{I}_2 \models \{a\} \sqsubseteq \neg \exists r. \{a\}$, contrary to the assumption that $a^{\mathcal{I}_2} \in (\{a\} \sqcap \exists r. \{a\})^{\mathcal{I}_2}$. \dashv




Fig. 3. Failure of the CI-interpolation property for $\mathcal{L} \in \text{DL}_{nr}$ without RIs.

In this article we focus on interpolating concepts and not CIs. The main reasons are that the corresponding notion of an explicit definition of an ontology appears to be less useful than definitions of concepts and nominals and that while the CI-interpolation property is crucial for robust decompositions of ontologies and for robust forgetting [55], checking the existence of an interpolant or computing it for concrete ontologies and CIs appears to not have found any applications yet. Regarding the first point, observe that CI-interpolants correspond to the following notion of an explicit CI-definition of an ontology. Let Σ be a signature and \mathcal{O} and \mathcal{O}' ontologies. Then an $\mathcal{L}(\Sigma)$ -ontology \mathcal{O}'' is called an *explicit $\mathcal{L}(\Sigma)$ -definition of \mathcal{O}' under \mathcal{O}* if $\mathcal{O} \cup \mathcal{O}' \models \mathcal{O}''$ and $\mathcal{O} \cup \mathcal{O}'' \models \mathcal{O}'$. In particular, if \mathcal{O} is empty then one asks for an ontology using symbols in Σ only that is equivalent to \mathcal{O} . While the existence of such ontologies is an interesting theoretical question that could well have applications in the future, investigating this problem is beyond the focus of this article. In what follows we only consider aspects of CI-interpolants that are closely related to concept interpolants, leaving their detailed investigation for future work.

5 Main Results

The failure of CIP and (P)BDP reported in Theorem 12 imply that interpolant existence and projective and non-projective definition existence cannot be directly polynomially reduced to subsumption checking. This motivates studying the respective decision problems of interpolant existence and projective and non-projective definition existence. In this section we introduce the decision problems, formulate model-theoretic characterizations of the problems that play a fundamental role in our proofs, and we formulate the main results.

We start with interpolant existence for which we take the definition used in the formulation of the CIP.

Definition 16. *Let \mathcal{L} be a DL. Then \mathcal{L} -interpolant existence is the problem to decide for any \mathcal{L} -ontologies $\mathcal{O}_1, \mathcal{O}_2$ and \mathcal{L} -concepts C_1, C_2 , whether there exists an $\mathcal{L}(\Sigma)$ -interpolant for $C_1 \sqsubseteq C_2$ under $\mathcal{O}_1 \cup \mathcal{O}_2$, where $\Sigma = \text{sig}(\mathcal{O}_1, C_1) \cap \text{sig}(\mathcal{O}_2, C_2)$.*

In our proofs, we actually focus on a more general version of interpolant existence which has been discussed in the previous section and in which we do not split \mathcal{O} into two ontologies and in which Σ is arbitrary.

Definition 17. *Let \mathcal{L} be a DL. Then generalized \mathcal{L} -interpolant existence is the problem to decide for any \mathcal{L} -ontology \mathcal{O} , \mathcal{L} -concepts C_1, C_2 , and signature Σ whether there exists an $\mathcal{L}(\Sigma)$ -interpolant for $C_1 \sqsubseteq C_2$ under \mathcal{O} .*

We also consider (generalized) \mathcal{L} -interpolant existence with empty ontologies, called *ontology-free (generalized) \mathcal{L} -interpolant existence*, and with RI-ontologies, called *RI-ontology (generalized) \mathcal{L} -interpolant existence*, both defined in the obvious way. Observe that in the ontology-free case there is no difference between generalized interpolant existence and interpolant existence. In fact, also with ontologies generalized interpolant existence and interpolant existence are interreducible.

Lemma 18. *Let $\mathcal{L} \in DL_{nr}$. There are mutual polynomial time reductions between generalized \mathcal{L} -interpolant existence and \mathcal{L} -interpolant existence.*

Proof. The reduction from \mathcal{L} -interpolant existence to generalized \mathcal{L} -interpolant existence is trivial: for input $\mathcal{O}_1, \mathcal{O}_2, C_1, C_2$ to \mathcal{L} -interpolant existence, set $\mathcal{O} = \mathcal{O}_1 \cup \mathcal{O}_2$ and $\Sigma = \text{sig}(\mathcal{O}_1, C_1) \cap \text{sig}(\mathcal{O}_2, C_2)$.

For the converse reduction from generalized interpolant existence to interpolant existence, assume that an \mathcal{L} -ontology \mathcal{O} , \mathcal{L} -concepts C_1, C_2 , and Σ are given. Then there exists an $\mathcal{L}(\Sigma)$ -interpolant for $C_1 \sqsubseteq C_2$ under

\mathcal{O} iff there exists an $\mathcal{L}(\Sigma)$ -interpolant for $C_1 \sqsubseteq C_{2\Sigma}$ under $\mathcal{O} \cup \mathcal{O}_\Sigma$, where \mathcal{O}_Σ and $C_{2\Sigma}$ are obtained from \mathcal{O} and C_2 by replacing every non- Σ symbol uniformly by a fresh symbol. The latter is an instance of \mathcal{L} -interpolant existence. \square

Note that the reduction above works for all standard DLs including \mathcal{ALC} . Recall that interpolant existence reduces to checking $\mathcal{O}_1 \cup \mathcal{O}_2 \models C_1 \sqsubseteq C_2$ for logics with the CIP. Hence, for DLs which enjoy the CIP such as \mathcal{ALC} , interpolant existence and generalized interpolant existence are EXPTIME-complete and ontology-free interpolant existence and generalized ontology-free interpolant existence are PSPACE-complete.

We next introduce the relevant definition existence problems.

Definition 19. *Let \mathcal{L} be a DL. Projective \mathcal{L} -definition existence is the problem to decide for any \mathcal{L} -ontology \mathcal{O} , \mathcal{L} -concepts C and C_0 , and signature Σ , whether there exists an explicit $\mathcal{L}(\Sigma)$ -definition of C_0 under \mathcal{O} and C .*

(Non-projective) \mathcal{L} -definition existence of concept names (nominals) is the sub-problem where C_0 ranges only over concept names A (nominals $\{a\}$) and $\Sigma = \text{sig}(\mathcal{O}, C) \setminus \{A\}$ (and $\Sigma = \text{sig}(\mathcal{O}, C) \setminus \{a\}$, respectively).

We also consider the (projective) \mathcal{L} -definition existence problems with empty ontologies, called *ontology-free (projective) \mathcal{L} -definition existence*, and with RI-ontologies, called *RI-ontology (projective) \mathcal{L} -definition existence*, both defined in the obvious way. Similar to the case of interpolant existence, definition existence reduces to checking implicit definability for logics with the PBDP. We provide model-theoretic characterizations for the non-existence of generalized interpolants and explicit definitions in terms of bisimulations.

Definition 20 (Joint consistency). *Let $\mathcal{L} \in DL_{nr}$. Let \mathcal{O} be an \mathcal{L} -ontology, C_1, C_2 be \mathcal{L} -concepts, and Σ a signature. Then C_1 and C_2 are called jointly consistent under \mathcal{O} modulo $\mathcal{L}(\Sigma)$ -bisimulations if there exist pointed interpretations \mathcal{I}_1, d_1 and \mathcal{I}_2, d_2 such that \mathcal{I}_i is a model of \mathcal{O} , $d_i \in C_i^{\mathcal{I}_i}$, for $i = 1, 2$, and $\mathcal{I}_1, d_1 \sim_{\mathcal{L}, \Sigma} \mathcal{I}_2, d_2$.*

The associated decision problem, *joint consistency modulo \mathcal{L} -bisimulations*, is defined in the expected way. The following result characterizes the existence of interpolants using joint consistency modulo $\mathcal{L}(\Sigma)$ -bisimulations. The proof uses Lemma 3.

Theorem 21. *Let $\mathcal{L} \in DL_{nr}$. Let \mathcal{O} be an \mathcal{L} -ontology, C_1, C_2 be \mathcal{L} -concepts, and Σ a signature. Then the following conditions are equivalent:*

1. there is no $\mathcal{L}(\Sigma)$ -interpolant for $C_1 \sqsubseteq C_2$ under \mathcal{O} ;
2. C_1 and $\neg C_2$ are jointly consistent under \mathcal{O} modulo $\mathcal{L}(\Sigma)$ -bisimulations.

Proof. The proof is standard and we refer the reader to [40] for similar proofs. We only provide a sketch.

“1 \Rightarrow 2”. Assume there is no $\mathcal{L}(\Sigma)$ -interpolant for $C_1 \sqsubseteq C_2$ under \mathcal{O} . Let

$$\Gamma = \{D \mid \mathcal{O} \models C_1 \sqsubseteq D, D \in \mathcal{L}(\Sigma)\}.$$

Then $\mathcal{O} \not\models D \sqsubseteq C_2$, for any $D \in \Gamma$. As Γ is closed under conjunction and by compactness (recall that \mathcal{ALCHOT}^u is a fragment of first-order logic), there exists a model \mathcal{J} of \mathcal{O} and an element $d \in \Delta^{\mathcal{J}}$ such that $d \in D^{\mathcal{J}}$ for all $D \in \Gamma$ but $d \notin C_2^{\mathcal{J}}$. Consider the set $t_{\mathcal{J}}(d) = \{D \in \mathcal{L}(\Sigma) \mid d \in D^{\mathcal{J}}\}$. Then, using again compactness, there exists a model \mathcal{I} of \mathcal{O} and an element $e \in \Delta^{\mathcal{I}}$ such that $e \in C_1^{\mathcal{I}}$ and $e \in D^{\mathcal{I}}$ for all $D \in t_{\mathcal{J}}(d)$. Thus $\mathcal{I}, e \equiv_{\mathcal{L}, \Sigma} \mathcal{J}, d$. For every interpretation \mathcal{I} there exists an ω -saturated elementary extension \mathcal{I}' of \mathcal{I} [24]. Thus, it follows from the fact that \mathcal{ALCHOT}^u is a fragment of first-order logic that we may assume that both \mathcal{I} and \mathcal{J} are ω -saturated. By Lemma 3, $\mathcal{I}, e \sim_{\mathcal{L}, \Sigma} \mathcal{J}, d$.

“2 \Rightarrow 1”. Assume an $\mathcal{L}(\Sigma)$ -interpolant D for $C_1 \sqsubseteq C_2$ under \mathcal{O} exists. Assume that Condition 2 holds, that is, there are models \mathcal{I}_1 and \mathcal{I}_2 of \mathcal{O} and $d_i \in \Delta^{\mathcal{I}_i}$ for $i = 1, 2$ such that $d_1 \in C_1^{\mathcal{I}_1}$ and $d_2 \notin C_2^{\mathcal{I}_2}$ and $\mathcal{I}_1, d_1 \sim_{\mathcal{L}, \Sigma} \mathcal{I}_2, d_2$. Then, by Lemma 3, $\mathcal{I}_1, d_1 \equiv_{\mathcal{L}, \Sigma} \mathcal{I}_2, d_2$. But then from $d_1 \in C_1^{\mathcal{I}_1}$ we obtain $d_1 \in D^{\mathcal{I}_1}$ and so $d_2 \in D^{\mathcal{I}_2}$ which implies $d_2 \in C_2^{\mathcal{I}_2}$, a contradiction. \square

Example 22. Consider again $C_1 = \{a\} \sqcap \exists r.\{a\}$ and $C_2 = \{b\} \rightarrow \exists r.\{b\}$ from Example 5 and set $\Sigma = \{r\}$. The interpretations $\mathcal{I}_1, \mathcal{I}_2$ depicted in Figure 4 (where we set $a^{\mathcal{I}_i} = a$ and $b^{\mathcal{I}_i} = b$, for $i = 1, 2$) show that C_1 and $\neg C_2$ are jointly consistent modulo $\mathcal{ALCO}(\Sigma)$ -bisimulations. By extending the bisimulation in Figure 4 to a relation S such that $(b^{\mathcal{I}_1}, a^{\mathcal{I}_2}) \in S$ (so that the domain and range of S contain $\Delta^{\mathcal{I}_1}$ and $\Delta^{\mathcal{I}_2}$, respectively), one can show that C_1 and $\neg C_2$ are jointly consistent modulo $\mathcal{ALCO}^u(\Sigma)$ -bisimulations. Moreover, by introducing an element e in \mathcal{I}_2 so that $(e, b^{\mathcal{I}_2}) \in r^{\mathcal{I}_2}$ and $(e, e) \in r^{\mathcal{I}_2}$, and further extending S by adding $(a^{\mathcal{I}_1}, e) \in S$, it can be seen that C_1 and $\neg C_2$ are jointly consistent modulo $\mathcal{ALCHOT}^u(\Sigma)$ -bisimulations (and hence $\mathcal{ALCHOT}^u(\Sigma)$ -bisimulations). \dashv

The following characterization of the existence of explicit definitions can be proved similarly to Theorem 21.




Fig. 4. Interpretations \mathcal{I}_1 and \mathcal{I}_2 illustrating Example 22.

Theorem 23. Let $\mathcal{L} \in DL_{nr}$. Let \mathcal{O} be an \mathcal{L} -ontology, C and C_0 \mathcal{L} -concepts, and $\Sigma \subseteq \text{sig}(\mathcal{O}, C)$ a signature. Then the following conditions are equivalent:

1. there is no explicit $\mathcal{L}(\Sigma)$ -definition of C_0 under \mathcal{O} and C ;
2. $C \sqcap C_0$ and $C \sqcap \neg C_0$ are jointly consistent under \mathcal{O} modulo $\mathcal{L}(\Sigma)$ -bisimulations.

Example 24. Consider \mathcal{O} , C , and Σ from Example 9. The interpretations $\mathcal{I}_1, \mathcal{I}_2$ depicted in Figure 5 show that $C \sqcap \exists r. \top$ and $C \sqcap \neg \exists r. \top$ are jointly consistent under \mathcal{O} modulo $\mathcal{ALCH}(\Sigma)$ -bisimulations. Note that the $\mathcal{ALCH}(\Sigma)$ -bisimulation in Figure 5 is also an $\mathcal{ALCH}^u(\Sigma)$ -bisimulation, but it is not an $\mathcal{ALCHI}(\Sigma)$ -bisimulation, since e_1 has both an r_1 - and an r_2 -predecessor, whereas e_2 and e'_2 lack an r_2 - and an r_1 -predecessor, respectively. To repair this, we replace \mathcal{I}_2 with an interpretation \mathcal{J} that is obtained by taking the union of \mathcal{I}_2 with a copy $\bar{\mathcal{I}}_1$ of \mathcal{I}_1 , and further adding $(\bar{d}_1, e_2) \in r_2^{\mathcal{J}}$ and $(\bar{d}_1, e'_2) \in r_1^{\mathcal{J}}$ (where \bar{d}_1 is the copy of d_1 in \mathcal{J}). Then, we extend the $\mathcal{ALCH}(\Sigma)$ -bisimulation in Figure 5 to a relation S that also connects the elements of \mathcal{I}_1 with the respective copies in \mathcal{J} . It can be seen that \mathcal{J} is a model of \mathcal{O} , $d_2 \in (C \sqcap \neg \exists r. \top)^{\mathcal{J}}$, and $(d_1, d_2) \in S$, where S is an $\mathcal{ALCHI}^u(\Sigma)$ -bisimulation. \dashv




Fig. 5. Interpretations \mathcal{I}_1 and \mathcal{I}_2 illustrating Example 24.

Interpolant existence and explicit definition existence are closely linked. We use Theorems 21 and 23 to show the following reductions.

Lemma 25. *Let $\mathcal{L} \in DL_{nr}$, \mathcal{O} be an \mathcal{L} -ontology, C, C_0, C_1 , and C_2 be \mathcal{L} -concepts, and Σ a signature. Then the following conditions are equivalent:*

1. *there is an explicit $\mathcal{L}(\Sigma)$ -definition of C_0 under \mathcal{O} and C ;*
2. *there is an $\mathcal{L}(\Sigma)$ -interpolant for $C \sqcap C_0 \sqsubseteq C \rightarrow C_0$ under \mathcal{O} .*

Conversely, the following conditions are also equivalent:

1. *there is an $\mathcal{L}(\Sigma)$ -interpolant for $C_1 \sqsubseteq C_2$ under \mathcal{O} ;*
2. *$\mathcal{O} \models C_1 \sqsubseteq C_2$ and there is an explicit $\mathcal{L}(\Sigma)$ -definition of C_2 under \mathcal{O} and $C_2 \rightarrow C_1$.*

Proof. We show the second equivalence. Assume that (1) does not hold. To show that (2) does not hold, assume $\mathcal{O} \models C_1 \sqsubseteq C_2$ (otherwise we are done). By Theorem 21 there exist pointed interpretations \mathcal{I}_1, d_1 and \mathcal{I}_2, d_2 such that \mathcal{I}_i is a model of \mathcal{O} , $d_1 \in C_1^{\mathcal{I}_1}$, $d_2 \notin C_2^{\mathcal{I}_2}$, and $\mathcal{I}_1, d_1 \sim_{\mathcal{L}, \Sigma} \mathcal{I}_2, d_2$. But then $d_1 \in ((C_2 \rightarrow C_1) \sqcap C_2)^{\mathcal{I}_1}$ and $d_2 \in ((C_2 \rightarrow C_1) \sqcap \neg C_2)^{\mathcal{I}_2}$ which shows that (2) does not hold by Theorem 23. The other direction is shown similarly. \square

Hence we obtain the following corollary.

Theorem 26. *Let $\mathcal{L} \in DL_{nr}$. Then there is a polynomial time reduction of projective \mathcal{L} -definition existence to \mathcal{L} -interpolant existence (and thus to generalized \mathcal{L} -interpolant existence). Conversely, there is a polynomial time reduction of generalized \mathcal{L} -interpolant existence (and thus \mathcal{L} -interpolant existence) to projective \mathcal{L} -definition existence if an oracle for \mathcal{L} -subsumption is admitted.*

Both reductions also exist for the ontology-free case and for RI-ontologies.

We now formulate the main complexity results proved in this article.

Theorem 27. *Let $\mathcal{L} \in DL_{nr}$. Then \mathcal{L} -interpolant existence, generalized \mathcal{L} -interpolant existence, and projective \mathcal{L} -definition existence are all 2EXPTIME-complete.*

It follows that interpolant existence and projective definition existence are one exponential harder than subsumption for logics in DL_{nr} . Our lower bound proofs rely on the presence of ontologies. To understand the ontology-free case (and the case with RI-ontologies) we first recall from our introduction of DLs in DL_{nr} above that for DLs with the universal role

or with both inverse roles and nominals, the ontology can be encoded in a concept and so interpolant existence and projective definition existence are still 2EXPTIME -complete with empty ontologies and RI-ontologies, respectively. For the remaining DLs in DL_{nr} , interpolant existence and projective definition existence become CONEXPTIME -complete, however. Thus, less complex than with ontologies, but still harder than subsumption (which is PSPACE -complete), under standard complexity theoretic assumptions.

Theorem 28. *Let $\mathcal{L} \in \text{DL}_{\text{nr}}$.*

1. *If \mathcal{L} admits nominals and the universal role, or nominals and inverse roles, then ontology-free \mathcal{L} -interpolant existence, generalized \mathcal{L} -interpolant existence, and projective \mathcal{L} -definition existence are all 2EXPTIME -complete.*
2. *If \mathcal{L} admits the universal role and RIs, then RI-ontology \mathcal{L} -interpolant existence, generalized \mathcal{L} -interpolant existence, and projective \mathcal{L} -definition existence are all 2EXPTIME -complete.*
3. *If $\mathcal{L} \in \{\mathcal{ALCO}, \mathcal{ALCHO}\}$, then ontology-free and RI-ontology \mathcal{L} -interpolant existence, generalized \mathcal{L} -interpolant existence, and projective \mathcal{L} -definition existence are all CONEXPTIME -complete.*
4. *If $\mathcal{L} \in \{\mathcal{ALCH}, \mathcal{ALCHI}\}$, then RI-ontology \mathcal{L} -interpolant existence, generalized \mathcal{L} -interpolant existence, and projective \mathcal{L} -definition existence are all CONEXPTIME -complete.*

We have seen that with the exception of \mathcal{ALCO} and \mathcal{ALCHO} , all DLs in DL_{nr} enjoy the non-projective Beth definability property. Hence checking the existence of a non-projective definition of a concept name is polynomial time reducible to subsumption checking and so EXPTIME -complete in the presence of an ontology. The following result states that even for \mathcal{ALCO} and \mathcal{ALCHO} checking the existence of non-projective definitions of concept names is not harder than subsumption.

Theorem 29. *Let $\mathcal{L} \in \{\mathcal{ALCO}, \mathcal{ALCHO}\}$. Then non-projective \mathcal{L} -definition existence of concept names is EXPTIME -complete.*

Interestingly, Theorem 29 is the only result where the lack of either the CIP or (P)BDP does not lead to an increase in complexity of the interpolant/explicit definition existence problem. We show Theorem 29 in the appendix provided as supplementary material as it uses techniques that are slightly different from our other main results.

We next consider the non-projective explicit definability of nominals. We have seen in Example 13 above that for nominals even the non-projective Beth definability property does not hold for any DL in DL_{nr} .

In fact, the following result states that the non-projective definability of nominals is as hard as their projective definability.

Theorem 30. *Let $\mathcal{L} \in DL_{nr}$ admit nominals. Then non-projective \mathcal{L} -definition existence of nominals is 2EXPTIME-complete.*

Observe that the characterizations given in Theorems 21 and 23 provide mutual polynomial time reductions of generalized interpolant and definition existence to the complement of joint consistency modulo \mathcal{L} -bisimulations. Hence, to prove Theorems 27 to 30, it suffices to prove the corresponding complexity bounds for joint consistency.

We finally discuss an interesting consequence for CI-interpolants. Let \mathcal{L} be a DL in DL_{nr} that does not admit RIs. The *CI-interpolant existence problem in \mathcal{L}* is the problem to decide for \mathcal{L} -ontologies \mathcal{O} and \mathcal{O}' whether there exists an \mathcal{L} -CI-interpolant for \mathcal{O} and \mathcal{O}' .

Theorem 31. *Let $\mathcal{L} \in \{ALCO^u, ALCOI^u\}$. Then CI-interpolant existence in \mathcal{L} is 2EXPTIME-complete.*

Observe that the 2EXPTIME upper bound is an immediate consequence of Point 1 of Theorem 28 as we can give a polynomial time reduction of CI-interpolant existence to ontology-free interpolant existence. Assume \mathcal{L} -ontologies \mathcal{O} and \mathcal{O}' are given. Let $\Sigma = \text{sig}(\mathcal{O}) \cap \text{sig}(\mathcal{O}')$. We find \mathcal{L} -concepts D and D' such that \mathcal{O} is equivalent to $\{\top \sqsubseteq D\}$ and \mathcal{O}' is equivalent to $\{\top \sqsubseteq D'\}$, respectively. Then there exists a \mathcal{L} -CI-interpolant for \mathcal{O} and \mathcal{O}' iff there exists an \mathcal{L} -interpolant for $\forall u.D \sqsubseteq \forall u.D'$.

The 2EXPTIME lower bound is proved in Section 7 (Lemma 44) by adapting the 2EXPTIME lower bound proof for interpolant existence in \mathcal{L} .

6 Upper Bound Proofs With Ontology

We show the double exponential upper bound of Theorem 27 (and thus of Theorem 30) using a new mosaic elimination procedure that decides joint consistency modulo \mathcal{L} -bisimulations, for all $\mathcal{L} \in DL_{nr}$.

Theorem 32. *Let $\mathcal{L} \in DL_{nr}$. Then joint consistency modulo \mathcal{L} -bisimulations is in 2EXPTIME.*

To motivate our approach, reconsider Example 22. Notice that in interpretations $\mathcal{I}_1, \mathcal{I}_2$ witnessing joint consistency of C_1 and $\neg C_2$, $a^{\mathcal{I}_1}$ is bisimilar to both $b^{\mathcal{I}_2}$ and d . Moreover, it can be easily verified that there are no witnessing interpretations where $a^{\mathcal{I}_1}$ is bisimilar to a single element in \mathcal{I}_2 .

Using an ontology, one can extend this example so that $a^{\mathcal{I}_1}$ is enforced to be bisimilar to exponentially many elements in \mathcal{I}_2 in any interpretations $\mathcal{I}_1, \mathcal{I}_2$ witnessing joint consistency of two concepts (in fact, this will be the basis for showing the lower bound in the subsequent section). Thus, we cannot consider (pairs of) elements in isolation, but instead need to consider sets of elements. As usual in DLs, we abstract elements in interpretations by *types*, which syntactically describe the behavior of these elements by listing the relevant concepts that are satisfied there. Correspondingly, sets of elements are abstracted to sets of types. Since we need to coordinate two interpretations $\mathcal{I}_1, \mathcal{I}_2$, we thus consider *mosaics*, which are pairs (T_1, T_2) of sets of types. The intuitive meaning of such a pair is that it describes collections of elements in two interpretations \mathcal{I}_1 and \mathcal{I}_2 which realize precisely the types in T_1 and T_2 , respectively, and are all mutually bisimilar. Naturally, not all possible mosaics (T_1, T_2) can be realized in this way and the goal is to determine the realizable ones. For this task, we use an elimination procedure. We start with the set of all possible mosaics and drop the ‘bad’ ones until a fixed point is reached. We will see that the elimination conditions extend the conditions known from standard type elimination procedures in a relatively natural way to mosaics. Then, concepts C_1, C_2 will be jointly consistent under an ontology \mathcal{O} modulo bisimulations if there is a surviving mosaic (T_1, T_2) such that C_1 is contained in some type in T_1 and C_2 is contained in some type in T_2 .

We will now formalize our approach and start by introducing the relevant notions. Assume $\mathcal{L} \in \text{DL}_{\text{nr}}$ and consider an \mathcal{L} -ontology \mathcal{O} , \mathcal{L} -concepts C_1, C_2 , and a signature Σ . Let $\Xi = \text{sub}(\mathcal{O}, C_1, C_2)$ denote the closure under single negation of the set of subconcepts of concepts in \mathcal{O}, C_1, C_2 . A Ξ -type t is a subset of Ξ such that there exists a model \mathcal{I} of \mathcal{O} and $d \in \Delta^{\mathcal{I}}$ with $t = \text{tp}_{\Xi}(\mathcal{I}, d)$, where

$$\text{tp}_{\Xi}(\mathcal{I}, d) = \{C \in \Xi \mid d \in C^{\mathcal{I}}\}$$

is the Ξ -type realized at d in \mathcal{I} . Let $\text{Tp}(\Xi)$ denote the set of all Ξ -types. We remark that the number of Ξ -types is at most exponential in $\|\mathcal{O}\| + \|C_1\| + \|C_2\|$ and, moreover, the set of all Ξ -types can be computed in time exponential in $\|\mathcal{O}\| + \|C_1\| + \|C_2\|$ for all considered logics [8, 1]. A *mosaic* is a pair (T_1, T_2) of sets of types $T_1, T_2 \subseteq \text{Tp}(\Xi)$. For interpretations $\mathcal{I}_1, \mathcal{I}_2$ and $i \in \{1, 2\}$, the *mosaic defined by* $d \in \Delta^{\mathcal{I}_i}$ in $\mathcal{I}_1, \mathcal{I}_2$ is the pair $(T_1(d), T_2(d))$ where

$$T_j(d) = \{\text{tp}_{\Xi}(\mathcal{I}_j, e) \mid e \in \Delta^{\mathcal{I}_j}, \mathcal{I}_i, d \sim_{\mathcal{L}, \Sigma} \mathcal{I}_j, e\},$$

for $j = 1, 2$. We say that a pair (T_1, T_2) of sets T_1, T_2 of types is a *mosaic defined by $\mathcal{I}_1, \mathcal{I}_2$* if there exists $d \in \Delta^{\mathcal{I}_1} \cup \Delta^{\mathcal{I}_2}$ such that $(T_1, T_2) = (T_1(d), T_2(d))$. Clearly, there are at most doubly exponentially many mosaics.

Example 33. Recall C_1, C_2, Σ , and $\mathcal{I}_1, \mathcal{I}_2$ from Example 22, and let $\mathcal{O} = \emptyset$. The set Ξ consists of the concepts $\{a\}, \exists r.\{a\}, \{b\}, \exists r.\{b\}, C_1, C_2$, and negations thereof. We have that:

$$\begin{aligned} \text{tp}_\Xi(\mathcal{I}_1, a^{\mathcal{I}_1}) &= \{\{a\}, \exists r.\{a\}, \neg\{b\}, \neg\exists r.\{b\}, C_1, C_2\} \\ \text{tp}_\Xi(\mathcal{I}_2, b^{\mathcal{I}_2}) &= \{\neg\{a\}, \neg\exists r.\{a\}, \{b\}, \neg\exists r.\{b\}, \neg C_1, \neg C_2\} \\ \text{tp}_\Xi(\mathcal{I}_2, d) &= \{\neg\{a\}, \neg\exists r.\{a\}, \neg\{b\}, \neg\exists r.\{b\}, \neg C_1, C_2\} \end{aligned}$$

The mosaic defined by $a^{\mathcal{I}_1}$ in $\mathcal{I}_1, \mathcal{I}_2$ is $(T_1(a^{\mathcal{I}_1}), T_2(a^{\mathcal{I}_1}))$, where

$$T_1(a^{\mathcal{I}_1}) = \{\text{tp}_\Xi(\mathcal{I}_1, a^{\mathcal{I}_1})\} \quad \text{and} \quad T_2(a^{\mathcal{I}_1}) = \{\text{tp}_\Xi(\mathcal{I}_2, b^{\mathcal{I}_2}), \text{tp}_\Xi(\mathcal{I}_2, d)\}.$$

⊣

As announced above, the aim of the mosaic elimination procedure is to determine all mosaics (T_1, T_2) such that all $t \in T_1 \cup T_2$ can be realized in mutually $\mathcal{L}(\Sigma)$ -bisimilar elements of models $\mathcal{I}_1, \mathcal{I}_2$ of \mathcal{O} . In order to formulate the elimination conditions, we define several compatibility conditions between types and between mosaics, similar to the compatibility conditions that are used in standard type elimination procedures. Throughout the rest of the section, we treat the universal role u as a role name contained in Σ , in case \mathcal{L} admits the universal role. Note that u^- is equivalent to u , and that $\mathcal{O} \models r \sqsubseteq u$, for every role r .

Let t_1, t_2 be Ξ -types. We call t_1, t_2 *u-equivalent* if $\exists u.C \in t_1$ iff $\exists u.C \in t_2$, for every $\exists u.C \in \Xi$. Notice that the condition is trivially satisfied if \mathcal{L} does not admit the universal role. For a role r , we call t_1, t_2 *r-coherent for \mathcal{O}* , in symbols $t_1 \rightsquigarrow_r t_2$, if t_1, t_2 are u -equivalent and the following conditions hold for all roles s with $\mathcal{O} \models r \sqsubseteq s$: (1) if $\neg\exists s.C \in t_1$, then $C \notin t_2$ and (2) if $\neg\exists s^-.C \in t_2$, then $C \notin t_1$. Note that $t \rightsquigarrow_r t'$ iff $t' \rightsquigarrow_{r^-} t$. We lift the definition of r -coherence from types to mosaics $(T_1, T_2), (T'_1, T'_2)$. Specifically, we call $(T_1, T_2), (T'_1, T'_2)$ *r-coherent*, in symbols $(T_1, T_2) \rightsquigarrow_r (T'_1, T'_2)$, if for $i = 1, 2$:

- for every $t \in T_i$ there exists a $t' \in T'_i$ such that $t \rightsquigarrow_r t'$, and
- if \mathcal{L} admits inverse roles, then for every $t' \in T'_i$, there is a $t \in T_i$ such that $t \rightsquigarrow_r t'$.

Note that $(T_1, T_2) \rightsquigarrow_r (T'_1, T'_2)$ iff $(T'_1, T'_2) \rightsquigarrow_{r^-} (T_1, T_2)$ in case \mathcal{L} admits inverse roles. Also notice that $(T_1, T_2) \rightsquigarrow_r (T'_1, T'_2)$ implies $(T_1, T_2) \rightsquigarrow_u (T'_1, T'_2)$, for every role r .

Example 34. Consider again interpretations $\mathcal{I}_1, \mathcal{I}_2$ from Example 22 and the types $t_1 = \text{tp}_\Xi(\mathcal{I}_1, a^{\mathcal{I}_1})$, $t_2 = \text{tp}_\Xi(\mathcal{I}_2, b^{\mathcal{I}_2})$, and $t_3 = \text{tp}_\Xi(\mathcal{I}_2, d)$. Then, $t_1 \rightsquigarrow_r t_1$, $t_2 \rightsquigarrow_r t_3$, and $t_3 \rightsquigarrow_r t_3$. Moreover, the mosaic (T_1, T_2) defined by $a_1^{\mathcal{I}}$ in $\mathcal{I}_1, \mathcal{I}_2$ satisfies $(T_1, T_2) \rightsquigarrow_r (T_1, T_2)$. \dashv

We are now in the position to formulate the mosaic elimination conditions. Let $\mathcal{S} \subseteq 2^{\text{Tp}(\Xi)} \times 2^{\text{Tp}(\Xi)}$ be a set of mosaics. We call $(T_1, T_2) \in \mathcal{S}$ *bad* if it violates one of the following conditions.

- Σ -concept name coherence** $A \in t$ iff $A \in t'$, for every concept name $A \in \Sigma$ and every $t, t' \in T_1 \cup T_2$;
- Existential saturation** for $i = 1, 2$ and $\exists r.C \in t \in T_i$, there exists $(T'_1, T'_2) \in \mathcal{S}$ such that (1) there exists $t' \in T'_i$ with $C \in t'$ and $t \rightsquigarrow_r t'$ and (2) if $\mathcal{O} \models r \sqsubseteq s$ for a Σ -role s , then $(T_1, T_2) \rightsquigarrow_s (T'_1, T'_2)$.

For didactic purposes and because we need it later in Section 10, we first give the mosaic elimination procedure for logics \mathcal{L} that do not admit nominals. The procedure starts with the set \mathcal{S}_0 of all mosaics. Then obtain, for $i \geq 0$, \mathcal{S}_{i+1} from \mathcal{S}_i by eliminating all mosaics (T_1, T_2) that are bad in \mathcal{S}_i . Let \mathcal{S}^* be where the sequence stabilizes. The elimination procedure decides joint consistency in the following sense.

Lemma 35. *If \mathcal{L} does not admit nominals, the following conditions are equivalent:*

1. C_1, C_2 are jointly consistent under \mathcal{O} modulo $\mathcal{L}(\Sigma)$ -bisimulations;
2. there exist $(T_1, T_2) \in \mathcal{S}^*$ and Ξ -types $t_1 \in T_1, t_2 \in T_2$ with $C_1 \in t_1$ and $C_2 \in t_2$.

We refrain from giving the proof of Lemma 35 since it will follow from Lemma 36 below. We note, however, that for \mathcal{L} as in the lemma, Theorem 32 is an immediate consequence of the procedure: there are only double exponentially many mosaics, so the elimination terminates after at most double exponentially steps. It remains to observe that every elimination step can be executed in double exponential time.

This relatively straightforward elimination procedure does not quite work in the presence of nominals. Intuitively, the reason is that in any two interpretations $\mathcal{I}_1, \mathcal{I}_2$, every nominal a is realized (modulo bisimulation) in exactly one mosaic. Now, if the set \mathcal{S} contains several mosaics mentioning a , they possibly witness existential saturation of each other

which, however, cannot be reflected in an interpretation. Thus, for the mosaic elimination procedure to work (in the sense of Lemma 35) one has to “guess” for every nominal a exactly one mosaic that describes a .

To formalize this idea, let us call a set \mathcal{S} of mosaics *good for nominals* if for every individual name $a \in \text{sig}(\Xi)$ and $i = 1, 2$ there exists exactly one t_a^i with $\{a\} \in t_a^i \in \bigcup_{(T_1, T_2) \in \mathcal{S}} T_i$ and exactly one pair $(T_1, T_2) \in \mathcal{S}$ with $t_a^i \in T_i$. Moreover, if $a \in \Sigma$, then that pair takes the form

- $(\{t_a^1\}, \{t_a^2\})$ in case \mathcal{L} admits the universal role, and
- $(\{t_a^1\}, \{t_a^2\}), (\{t_a^1\}, \emptyset)$, or $(\emptyset, \{t_a^2\})$, otherwise.

We can now formulate the more general lemma.

Lemma 36. *The following conditions are equivalent:*

1. C_1, C_2 are jointly consistent under \mathcal{O} modulo $\mathcal{L}(\Sigma)$ -bisimulations;
2. there exists a set \mathcal{S}^* of mosaics that is good for nominals and does not contain a bad mosaic, such that there exist $(T_1, T_2) \in \mathcal{S}^*$ and Ξ -types $t_1 \in T_1, t_2 \in T_2$ with $C_1 \in t_1$ and $C_2 \in t_2$.

Proof. “1 \Rightarrow 2”. Let $\mathcal{I}_1, d_1 \sim_{\mathcal{L}, \Sigma} \mathcal{I}_2, d_2$ for models \mathcal{I}_1 and \mathcal{I}_2 of \mathcal{O} such that d_1, d_2 realize Ξ -types t_1, t_2 and $C_1 \in t_1, C_2 \in t_2$. Let \mathcal{S}^* be the set of all mosaics defined by $\mathcal{I}_1, \mathcal{I}_2$. It is routine to show that no (T_1, T_2) in \mathcal{S}^* is bad and that \mathcal{S}^* is good for nominals. Now, the mosaic (T_1, T_2) defined by $d_1^{\mathcal{I}_1}$ in $\mathcal{I}_1, \mathcal{I}_2$ witnesses Condition (2).

“2 \Rightarrow 1”. Suppose there exist a good set \mathcal{S}^* of mosaics and $(S_1, S_2) \in \mathcal{S}^*$ and Ξ -types $s_1 \in S_1, s_2 \in S_2$ with $C_1 \in s_1$ and $C_2 \in s_2$. Let \mathcal{I}_i , for $i = 1, 2$ be interpretations defined by setting:

$$\begin{aligned} \Delta^{\mathcal{I}_i} &:= \{(t, (T_1, T_2)) \mid (T_1, T_2) \in \mathcal{S}^*, t \in T_i, \text{ and} \\ &\quad \text{if } \mathcal{L} \text{ admits the universal role, then } (S_1, S_2) \rightsquigarrow_u (T_1, T_2) \\ &\quad \text{and } t, s_i \text{ are } u\text{-equivalent}\} \\ r^{\mathcal{I}_i} &:= \{((t, p), (t', p')) \in \Delta^{\mathcal{I}_i} \times \Delta^{\mathcal{I}_i} \mid t \rightsquigarrow_r t' \text{ and for all } \Sigma\text{-roles } s : \\ &\quad ((\mathcal{O} \models r \sqsubseteq s) \Rightarrow p \rightsquigarrow_s p')\} \\ A^{\mathcal{I}_i} &:= \{(t, p) \in \Delta^{\mathcal{I}_i} \mid A \in t\} \\ a^{\mathcal{I}_i} &:= (t, (T_1, T_2)) \in \Delta^{\mathcal{I}_i}, \{a\} \in t \in T_i \end{aligned}$$

Note that the interpretation of nominals is well-defined since \mathcal{S}^* is good for nominals.

We verify that interpretations \mathcal{I}_1 and \mathcal{I}_2 witness Condition (1).

Claim 1. For $i = 1, 2$, all $C \in \Xi$, and all $(t, p) \in \Delta^{\mathcal{I}_i}$, we have $(t, p) \in C^{\mathcal{I}_i}$ iff $C \in t$.

Proof of Claim 1. Let $i \in \{1, 2\}$. The proof is by induction on the structure of concepts in Ξ .

- The claim holds for concept names $C = A$ and all nominals $C = \{a\}$, by definition of \mathcal{I}_i .
- The Boolean cases, $\neg C$ and $C \sqcap C'$, are immediate consequences of the hypothesis.
- Let $C = \exists r.D$. (Recall that r is possibly the universal role u .)
“if”: Suppose $\exists r.D \in t$. By existential saturation, there is a $p' = (T'_1, T'_2) \in \mathcal{S}^*$ such that (1) there exists $t' \in T'_i$ with $D \in t'$ and $t \rightsquigarrow_r t'$, and (2) if $\mathcal{O} \models r \sqsubseteq s$ for some Σ -role s , then $p \rightsquigarrow_s p'$. Note that t, t' are thus also u -equivalent, so $(t', p') \in \Delta^{\mathcal{I}_i}$. We distinguish cases:
 - If r is a role name, then by definition of $r^{\mathcal{I}_i}$, we have that $((t, p), (t', p')) \in r^{\mathcal{I}_i}$. Since $D \in t'$, induction yields $(t', p') \in D^{\mathcal{I}_i}$. Overall, we get $(t, p) \in (\exists r.D)^{\mathcal{I}_i}$.
 - If $r = r_0^-$ is an inverse role, then (1) and (2) above imply (1') $t' \rightsquigarrow_{r_0} t$ and (2') if $\mathcal{O} \models r_0 \sqsubseteq s$ for some Σ -role s , then $p \rightsquigarrow_s p'$. As before, we can then conclude that $((t', p'), (t, p)) \in r_0^{\mathcal{I}_i}$. Since $D \in t'$, induction yields $(t', p') \in D^{\mathcal{I}_i}$. Overall, we get $(t, p) \in (\exists r_0^- . D)^{\mathcal{I}_i}$.
“only if”: Suppose $(t, p) \in (\exists r.D)^{\mathcal{I}_i}$. Then, there is $(t', p') \in \Delta^{\mathcal{I}_i}$ with $((t, p), (t', p')) \in r^{\mathcal{I}_i}$ and $(t', p') \in D^{\mathcal{I}_i}$. By induction, the latter implies $D \in t'$. We distinguish cases:
 - If r is a role name, then by definition of $r^{\mathcal{I}_i}$, $t \rightsquigarrow_r t'$ and thus $\exists r.D \in t$.
 - If $r = r_0^-$ is an inverse role, then by definition of $r_0^{\mathcal{I}_i}$, $t' \rightsquigarrow_{r_0} t$. Thus, also $\exists r_0^- . D \in t$.

This finishes the proof of Claim 1. Claim 1 implies that $(s_1, (S_1, S_2)) \in C_1^{\mathcal{I}_1}$ and $(s_2, (S_1, S_2)) \in C_2^{\mathcal{I}_2}$. Claim 1 also implies that the type realized by (t, p) in \mathcal{I}_i is t , for all $(t, p) \in \Delta^{\mathcal{I}_i}$. Since types are, by definition, realized in models of \mathcal{O} , it follows that both \mathcal{I}_1 and \mathcal{I}_2 are models of \mathcal{O} .

Claim 2. The relation R defined by

$$R = \{((t, p), (t', p)) \mid (t, p) \in \Delta^{\mathcal{I}_1}, (t', p) \in \Delta^{\mathcal{I}_2}\}$$

is an $\mathcal{L}(\Sigma)$ -bisimulation.

Proof of Claim 2. Clearly, R satisfies Condition [AtomC] due to Σ -concept name coherence. Condition [AtomI] follows from the fact that \mathcal{S}^* is good for nominals in case \mathcal{L} admits nominals.

For Condition [Forth], let $((t, p), (t', p)) \in R$ and $((t, p), (t_1, p_1)) \in r^{\mathcal{I}_1}$, for some Σ -role r , and let $p = (T_1, T_2)$ and $p_1 = (T'_1, T'_2)$. We distinguish cases:

- If r is a role name, then by definition of $r^{\mathcal{I}_1}$, we have (1) $t \rightsquigarrow_r t_1$ and (2) for all Σ -roles s with $\mathcal{O} \models r \sqsubseteq s$, we have $p \rightsquigarrow_s p_1$. Since $t' \in T_2$ and $p \rightsquigarrow_r p_1$ there is some $t'' \in T'_2$ with $t' \rightsquigarrow_r t''$. Thus, in particular, t'' is u -equivalent to t' (and thus to s_2), which implies $(t'', p_1) \in \Delta^{\mathcal{I}_2}$. The definition of $r^{\mathcal{I}_2}$ then implies that $((t', p), (t'', p_1)) \in r^{\mathcal{I}_2}$. It remains to note that the definition of R yields $((t_1, p_1), (t'', p_1)) \in R$.
- If $r = r_0^-$ is an inverse role, then by definition of $r_0^{\mathcal{I}_1}$, we have (1) $t_1 \rightsquigarrow_{r_0} t$ and (2) for all Σ -roles s with $\mathcal{O} \models r_0 \sqsubseteq s$, we have $p_1 \rightsquigarrow_s p$. Since $t' \in T_2$ and $p_1 \rightsquigarrow_{r_0} p$ there is some $t'' \in T'_2$ with $t'' \rightsquigarrow_{r_0} t'$. Thus, in particular, t'' is u -equivalent to t' (and thus to s_2), which implies $(t'', p_1) \in \Delta^{\mathcal{I}_2}$. The definition of $r_0^{\mathcal{I}_2}$ then implies that $((t'', p_1), (t', p)) \in r_0^{\mathcal{I}_2}$. It remains to note that the definition of R yields $((t_1, p_1), (t'', p_1)) \in R$.

Condition [Back] is dual.

Finally, we verify that R and R^- are surjective if \mathcal{L} admits the universal role. Let $(t, (T_1, T_2)) \in \Delta^{\mathcal{I}_1}$. Then, $(S_1, S_2) \rightsquigarrow_u (T_1, T_2)$, by definition of $\Delta^{\mathcal{I}_1}$. This implies that there is a type $t' \in T_2$ which is u -equivalent to s_2 and thus $(t', (T_1, T_2)) \in \Delta^{\mathcal{I}_2}$. The definition of R implies $((t, (T_1, T_2)), (t', (T_1, T_2))) \in R$. The other direction is dual.

This finishes the proof of Claim 2. By definition of R , we have $((s_1, (S_1, S_2)), (s_2, (S_1, S_2))) \in R$, and thus $\mathcal{I}_1, (s_1, (S_1, S_2)) \sim_{\mathcal{L}, \Sigma} \mathcal{I}_2, (s_2, (S_1, S_2))$. \square

It remains to argue that we can find in double exponential time a set \mathcal{S}^* as in Condition (2) of Lemma 36. We use a suitable variant of the elimination procedure described after Lemma 35.

Lemma 37. *Let $\mathcal{L} \in DL_{nr}$. Then it is decidable in time double exponential in $\|\mathcal{O}\| + \|C_1\| + \|C_2\|$ whether for an \mathcal{L} -ontology \mathcal{O} , \mathcal{L} -concepts C_1, C_2 , and a signature $\Sigma \subseteq \text{sig}(\Xi)$ there exists some \mathcal{S}^* satisfying Condition (2) of Lemma 36.*

Proof. Let $\mathcal{L} \in DL_{nr}$, and assume \mathcal{O}, C_1, C_2 , and Σ are given. We can enumerate in double exponential time the maximal good sets $\mathcal{U} \subseteq 2^{T(\Xi)} \times 2^{T(\Xi)}$ by picking, for each nominal $a \in \text{sig}(\Xi)$ and $i = 1, 2$, a type t_a^i , and a mosaic (T_1, T_2) with $t_a^i \in T_i$. In doing so, we make sure that $(\{t_a^1\}, \{t_a^2\})$ is selected in case $a \in \Sigma$. Crucially, there are only double exponentially many possibilities to make this choice. Remove all mosaics that mention a nominal and have not been selected. The resulting set is good for nominals.

Then we eliminate from any set \mathcal{U} obtained in that process recursively all bad mosaics. Let $\mathcal{S}_{\mathcal{U}} \subseteq \mathcal{U}$ be the largest fixpoint of that procedure. Then one can easily show that there exists a set \mathcal{S}^* satisfying Condition (2) of Lemma 36 iff there exists a set \mathcal{U} that can be obtained by the process described above such that the largest fixpoint $\mathcal{S}_{\mathcal{U}}$ satisfies Condition (2) of Lemma 36. Since elimination terminates after double exponential time, and there are only double exponentially many possible choices for \mathcal{U} , the lemma follows. \square

Theorem 32 is a direct consequence of Lemmas 36 and 37.

7 Lower Bound Proofs With Ontology

The goal of this section is to provide the proofs of the lower bounds in Theorems 27, 30, and 31. We start with the former two. By Lemma 25 and Theorem 23, it suffices to consider joint consistency. We will provide two reductions: in Section 7.1, we provide the reduction for DLs in DL_{nr} that admits nominals and, in Section 7.3, the one for DLs that admits role inclusions. In Section 7.2, we will investigate the shape of the interpolants / explicit definitions that arise in the preceding lower bound proof. In Section 7.4, we then show how to adapt the lower bound proof from Section 7.1 to the case of CI-interpolant existence. In all cases we reduce the word problem for languages recognized by exponentially space bounded alternating Turing machines, which we introduce next.

An *alternating Turing machine (ATM)* is a tuple

$$M = (Q, \Theta, \Gamma, q_0, \Delta)$$

where $Q = Q_{\exists} \uplus Q_{\forall}$ is a finite set of states partitioned into *existential states* Q_{\exists} and *universal states* Q_{\forall} . Further, Θ is the input alphabet and Γ is the tape alphabet that contains a *blank symbol* $\square \notin \Theta$, $q_0 \in Q_{\forall}$ is the *initial state*, and $\Delta \subseteq Q \times \Gamma \times Q \times \Gamma \times \{L, R\}$ is the *transition relation*. We assume without loss of generality that the set $\Delta(q, a) := \{(q', a', M) \mid (q, a, q', a', M) \in \Delta\}$ contains exactly two or zero elements for every $q \in Q$ and $a \in \Gamma$. Moreover, the state q' must be from Q_{\forall} if $q \in Q_{\exists}$ and from Q_{\exists} otherwise, that is, existential and universal states alternate. Acceptance of ATMs is defined in a slightly unusual way, without using accepting states. Intuitively, an ATM accepts if it runs forever on all branches and rejects otherwise. More formally, a *configuration* of an ATM is a word wqw' with $w, w' \in \Gamma^*$ and $q \in Q$. We say that wqw' is *existential* if q is, and likewise for *universal*. *Successor configurations* are defined in the usual way. Note

that every configuration has exactly zero or two successor configurations. A *computation tree* of an ATM M on input w is a (possibly infinite) tree whose nodes are labeled with configurations of M such that

- the root is labeled with the initial configuration q_0w ;
- if a node is labeled with an existential configuration wqw' , then it has a single successor which is labeled with a successor configuration of wqw' ;
- if a node is labeled with a universal configuration wqw' , then it has two successors which are labeled with the two successor configurations of wqw' .

An ATM M *accepts* an input w if there is a computation tree of M on w . Note that we can convert any ATM M in which acceptance is based on accepting states to our model by assuming that M terminates on any input and then modifying it to enter an infinite loop from the accepting states. It is well-known that there are 2^n -space bounded ATMs which recognize a 2EXPTIME-hard language [23], where n is the length of the input w .

7.1 DLs with Nominals

We start with DLs supporting nominals. By Theorem 23, it suffices to prove the following result.

Lemma 38. *Let $\mathcal{L} \in DL_{nr}$ admit nominals. It is 2EXPTIME-hard to decide for an \mathcal{L} -ontology \mathcal{O} , individual name b , and signature $\Sigma \subseteq \text{sig}(\mathcal{O}) \setminus \{b\}$ whether $\{b\}$ and $\neg\{b\}$ are jointly consistent under \mathcal{O} modulo $\mathcal{L}(\Sigma)$ -bisimulations. This is true even if b is the only individual in \mathcal{O} and $\Sigma = \text{sig}(\mathcal{O}) \setminus \{b\}$.*

As announced, we reduce the word problem for 2^n -space bounded ATMs. Let us fix such an ATM $M = (Q, \Theta, \Gamma, q_0, \Delta)$ and an input $w = a_0 \dots a_{n-1}$ of length n . We first provide the reduction for $\mathcal{L} = \mathcal{ALCO}$ using an ontology \mathcal{O} and a signature Σ such that \mathcal{O} contains concept names that are not in Σ and uses two role names r, s , and show later how to adapt this proof to $\Sigma = \text{sig}(\mathcal{O}) \setminus \{b\}$ and DLs supporting inverses and/or the universal role.

The idea of the reduction is as follows. We aim to construct an ontology \mathcal{O} such that M accepts w iff $\{b\}$ and $\neg\{b\}$ are jointly consistent under \mathcal{O} modulo $\mathcal{L}(\Sigma)$ -bisimulations, where

$$\Sigma = \{r, s, Z, B_\forall, B_{\exists}^1, B_{\exists}^2\} \cup \{A_\sigma \mid \sigma \in \Gamma \cup (Q \times \Gamma)\}.$$

The ontology \mathcal{O} enforces that $r(b, b)$ holds in any model \mathcal{O} using the concept inclusion $\{b\} \sqsubseteq \exists r.\{b\}$. Moreover, it enforces that any element distinct from $b^{\mathcal{I}}$ with an r -successor lies on an infinite r -path ρ enforced by the concept inclusions:

$$\neg\{b\} \sqcap \exists r.\top \sqsubseteq I_s \qquad I_s \sqsubseteq \exists r.\top \sqcap \forall r.I_s$$

with I_s a concept name. Thus, if there exist models \mathcal{I}, \mathcal{J} of \mathcal{O} with $\mathcal{I}, b^{\mathcal{I}} \sim_{\mathcal{ALCO}, \pm} \mathcal{J}, d$ for some $d \neq b^{\mathcal{J}}$ and $d \in (\exists r.\top)^{\mathcal{J}}$, it follows that all elements on the path ρ are $\mathcal{ALCO}(\pm)$ -bisimilar to $b^{\mathcal{I}}$ and thus mutually $\mathcal{ALCO}(\pm)$ -bisimilar. The situation is depicted in Figure 6, where the trees T_* and $T_i, i \geq 0$ starting in $b^{\mathcal{I}}$ and on the path elements, respectively, are also mutually $\mathcal{ALCO}(\pm)$ -bisimilar. These trees shall represent the computation tree of M on input w (using symbols from Σ) as follows, cf. Figure 7 which shows the skeleton of a single tree T_i . Configurations of M are represented as paths of length 2^n over a role s in which every element is labeled with a symbol $A_\sigma, \sigma \in \Gamma \cup (Q \times \Gamma)$ that represents the content of a single tape cell (omitted in the figure for the sake of readability). In Figure 7, the start of a configuration is indicated by \circ and the s -path between consecutive \circ has length 2^n . Every configuration is marked as existential or universal using concept names $B_\forall, B_{\exists}^1, B_{\exists}^2$; the superscript \cdot^1/\cdot^2 indicates which successor is chosen for an existential configuration. Existential configurations have a single successor configuration and universal configurations have two successor configurations.




Fig. 6. Enforced bisimulation in lower bound

The structure of this computation tree can easily be enforced in \mathcal{ALCO} using standard techniques (as we detail below). The difficulty is to achieve

synchronization between successor configurations in the tree. That is, if a configuration c in the computation tree is followed by another configuration c' , then c' is actually a successor configuration of c according to M . To achieve this, we first ensure that in T_i , for $i \leq 2^n$, the $(2^n - i)$ -th cell of each configuration in the computation tree is synchronized with the $(2^n - i)$ -th cell of the next configuration(s), as indicated by the dotted lines in Figure 7. This can be realized in \mathcal{ALC} using a set of concept names not in Σ . Then we exploit the fact that the trees T_i are mutually $\mathcal{ALC}(\pm)$ -bisimilar which implies that in *all* T_i *all* cells of *all* configurations are synchronized. In more detail, we do this by using several counters modulo 2^n as follows.




Fig. 7. Computation tree of M

The first counter counts modulo 2^n along the path ρ using concept names not in Σ . As announced, each point of ρ starts an infinite tree along role s that is supposed to mimick the computation tree of M on input w . Along this tree, two more counters are maintained:

- one counter starting at 0 and counting modulo 2^n , and
- another counter starting at *the value of the counter on ρ* and also counting modulo 2^n .

The first counter is used to divide the tree into configurations of length 2^n and the second counter is used to link the $(2^n - i)$ -th cell of successive configurations in T_i as described above.

We will next provide the concept inclusions in \mathcal{O} in more detail. The counter along ρ is realized using concept names A_i , $0 \leq i < n$ and by including the following (standard) concept inclusions, for every i with

$0 \leq i < n$:

$$\begin{array}{ll}
I_s \sqcap A_i \sqcap \prod_{j < i} A_j \sqsubseteq \forall r. \neg A_i & I_s \sqcap \neg A_i \sqcap \prod_{j < i} A_j \sqsubseteq \forall r. A_i \\
I_s \sqcap A_i \sqcap \bigsqcup_{j < i} \neg A_j \sqsubseteq \forall r. A_i & I_s \sqcap \neg A_i \sqcap \bigsqcup_{j < i} \neg A_j \sqsubseteq \forall r. \neg A_i
\end{array}$$

Using again the concept name I_s , we start the s -trees with two counters, realized using concept names U_i and V_i , $0 \leq i < n$, and initialized to 0 and the value of the A -counter, respectively, by including the following concept inclusions for every j with $0 \leq j < n$:

$$\begin{array}{l}
I_s \sqsubseteq (U = 0) \\
I_s \sqsubseteq A_j \leftrightarrow V_j \\
\top \sqsubseteq \exists s. \top
\end{array}$$

Here, $(U = 0)$ is an abbreviation for the concept $\prod_{i=0}^{n-1} \neg U_i$; we use similar abbreviations below without further notice. The counters U_i and V_i are incremented along s in the same way as A_i is incremented along r , so we omit details. Configurations of M are represented between two consecutive points having U -counter value 0. We next enforce the structure of the computation tree (recall that $q_0 \in Q_\forall$):

$$\begin{array}{ll}
I_s \sqsubseteq B_\forall & (\dagger) \\
(U < 2^n - 1) \sqcap B_\forall \sqsubseteq \forall s. B_\forall & \\
(U < 2^n - 1) \sqcap B_\exists^i \sqsubseteq \forall s. B_\exists^i & i \in \{1, 2\} \\
(U = 2^n - 1) \sqcap B_\forall \sqsubseteq \forall s. (B_\exists^1 \sqcup B_\exists^2) & \\
(U = 2^n - 1) \sqcap B_\exists^i \sqsubseteq \forall s. B_\forall & i \in \{1, 2\} \\
(U = 2^n - 1) \sqcap B_\forall \sqsubseteq \exists s. Z \sqcap \exists s. \neg Z &
\end{array}$$

These concept inclusions enforce that all points which represent a configuration satisfy one of $B_\forall, B_\exists^1, B_\exists^2$ indicating the kind of configuration and, if existential, also a choice of the transition function. The symbol $Z \in \Sigma$ enforces the branching.

We next set the initial configuration, for input $w = a_0, \dots, a_{n-1}$.

$$\begin{array}{ll}
I_s \sqsubseteq A_{q_0, a_0} & \\
I_s \sqsubseteq \forall s^k. A_{a_k} & 0 < i < n \\
I_s \sqsubseteq \forall s^n. \text{Blank} & \\
\text{Blank} \sqsubseteq A_\square & \\
\text{Blank} \sqcap (U < 2^n - 1) \sqsubseteq \forall s. \text{Blank} &
\end{array}$$

To coordinate successor configurations, we associate with M functions f_i , $i \in \{1, 2\}$ that map the content of three consecutive cells of a configuration to the content of the middle cell in the i -the successor configuration (assuming an arbitrary order on the set $\Delta(q, a)$, for all q, a). In what follows, we ignore the corner cases that occur at the border of configurations; they can be treated in a similar way. Clearly, for each possible triple $(\sigma_1, \sigma_2, \sigma_3) \in (\Gamma \cup (Q \times \Gamma))^3$, the \mathcal{ALC} -concept $C_{\sigma_1, \sigma_2, \sigma_3} = A_{\sigma_1} \sqcap \exists s.(A_{\sigma_2} \sqcap \exists s.A_{\sigma_3})$ is true at an element a of the computation tree iff a is labeled with A_{σ_1} , an s -successors b of a is labeled with A_{σ_2} , and an s -successors c of b is labeled with A_{σ_3} . In each configuration, we synchronize elements with V -counter 0 by including for every $(\sigma_1, \sigma_2, \sigma_3)$ and $i \in \{1, 2\}$ the following concept inclusions:

$$\begin{aligned} (V = 2^n - 1) \sqcap (U < 2^n - 2) \sqcap C_{\sigma_1, \sigma_2, \sigma_3} \sqcap B_{\forall} &\sqsubseteq \forall s.A_{f_1(\sigma_1, \sigma_2, \sigma_3)}^1 \sqcap \\ &\quad \forall s.A_{f_2(\sigma_1, \sigma_2, \sigma_3)}^2 \\ (V = 2^n - 1) \sqcap (U < 2^n - 2) \sqcap C_{\sigma_1, \sigma_2, \sigma_3} \sqcap B_{\exists}^i &\sqsubseteq \forall s.A_{f_i(\sigma_1, \sigma_2, \sigma_3)}^i \end{aligned}$$

At this point, the importance of the superscript in B_{\exists}^* becomes apparent: since different cells of a configuration are synchronized in different trees T_k the superscript makes sure that all trees rely on the same choice for existential configurations. The concept names A_{σ}^i are used as markers (not in Σ) and are propagated along s for 2^n steps, exploiting the V -counter. The superscript $i \in \{1, 2\}$ determines the successor configuration that the symbol is referring to. After crossing the end of a configuration, the symbol σ is propagated using concept names A'_{σ} (the superscript is not needed anymore because the branching happens at the end of the configuration, based on Z).

$$\begin{aligned} (U < 2^n - 1) \sqcap A_{\sigma}^i &\sqsubseteq \forall s.A_{\sigma}^i \\ (U = 2^n - 1) \sqcap B_{\forall} \sqcap A_{\sigma}^1 &\sqsubseteq \forall s.(Z \rightarrow A'_{\sigma}) \\ (U = 2^n - 1) \sqcap B_{\forall} \sqcap A_{\sigma}^2 &\sqsubseteq \forall s.(\neg Z \rightarrow A'_{\sigma}) \\ (U = 2^n - 1) \sqcap B_{\exists}^i \sqcap A_{\sigma}^i &\sqsubseteq \forall s.A'_{\sigma} & i \in \{1, 2\} \\ (V < 2^n - 1) \sqcap A'_{\sigma} &\sqsubseteq \forall s.A'_{\sigma} \\ (V = 2^n - 1) \sqcap A'_{\sigma} &\sqsubseteq \forall s.A_{\sigma} \end{aligned}$$

For those (q, a) with $\Delta(q, a) = \emptyset$, we add the concept inclusion

$$A_{q, a} \sqsubseteq \perp.$$

The following lemma establishes correctness of the reduction.

Lemma 39. *The following conditions are equivalent:*

1. M accepts w ;
2. there exist models \mathcal{I} and \mathcal{J} of \mathcal{O} such that $\mathcal{I}, b^{\mathcal{I}} \sim_{\mathcal{ALCO}, \pm} \mathcal{J}, d$, for some $d \neq b^{\mathcal{J}}$.

Proof. “1 \Rightarrow 2”. If M accepts w , there is a computation tree of M on w . We construct a single interpretation \mathcal{I} with $\mathcal{I}, b^{\mathcal{I}} \sim_{\mathcal{ALCO}, \pm} \mathcal{I}, d$ for some $d \neq b^{\mathcal{I}}$ as follows. Let $\widehat{\mathcal{J}}$ be the infinite tree-shaped interpretation that represents the computation tree of M on w as described above, that is, configurations are represented by sequences of 2^n elements linked by role s and labeled by $B_V, B_{\exists}^1, B_{\exists}^2$ depending on whether the configuration is universal or existential, and in the latter case the superscript indicates which choice has been made for the existential state. Finally, the first element of the first successor configuration of a universal configuration is labeled with Z . Observe that $\widehat{\mathcal{J}}$ interprets only the symbols in Σ as non-empty. Now, we obtain interpretations $\mathcal{I}_k, k < 2^n$ from $\widehat{\mathcal{J}}$ by interpreting non- Σ -symbols as follows:

- the root of \mathcal{I}_k satisfies I_s ;
- the U -counter starts at 0 at the root and counts modulo 2^n along each s -path;
- the V -counter starts at k at the root and counts modulo 2^n along each s -path;
- the auxiliary concept names of the shape A_{σ}^i and A'_{σ} are interpreted in a minimal way so as to satisfy the concept inclusions starting from concept inclusion (\dagger). Note that, by definition of these concept inclusions, there is a unique result.

Now obtain \mathcal{I} from $\widehat{\mathcal{J}}$ and the \mathcal{I}_k by creating an infinite outgoing r -path ρ from some element $d \neq b^{\mathcal{I}}$ (with the corresponding A -counter) and adding $\mathcal{I}_k, k < 2^n$ to every element with A -counter value k on the r -path, identifying the roots of the \mathcal{I}_k with the element on the path. Additionally, include $(b^{\mathcal{I}}, b^{\mathcal{I}}) \in r^{\mathcal{I}}$ and add $\widehat{\mathcal{J}}$ to \mathcal{I} by identifying $b^{\mathcal{I}}$ with the root of $\widehat{\mathcal{J}}$. It should be clear that \mathcal{I} is as required. In particular, the reflexive, transitive, and symmetric closure of

- all pairs $(b^{\mathcal{I}}, e)$, with e on ρ , and
- all pairs (e, e') , with e in $\widehat{\mathcal{J}}$ and e' a copy of e in some tree \mathcal{I}_k

is an $\mathcal{ALCO}(\pm)$ -bisimulation S on \mathcal{I} with $(b^{\mathcal{I}}, d) \in S$.

“2 \Rightarrow 1”. Assume that $\mathcal{I}, b^{\mathcal{I}} \sim_{\mathcal{ALCO}, \pm} \mathcal{J}, d$ for some $d \neq b^{\mathcal{J}}$. As argued above, due to the r -self loop at $b^{\mathcal{I}}$, from d there has to be an outgoing

infinite r -path on which all s -trees are $\mathcal{ALCCO}(\pm)$ -bisimilar. Since \mathcal{I} is a model of \mathcal{O} , all these s -trees are additionally labeled with some auxiliary concept names not in Σ , depending on the distance from their roots on ρ . Using the concept inclusions in \mathcal{O} and the arguments given in their description, it can be shown that all s -trees contain a computation tree of M on input w (which is solely represented with concept names in Σ). \square

The same ontology \mathcal{O} can be used for the remaining DLs with nominals. For \mathcal{ALCCO}^\square , exactly the same proof works; in particular, note that both the bisimulation S constructed in “1 \Rightarrow 2” and its inverse are surjective. For the DLs with inverse roles the (one-way) infinite r -path ρ has to be replaced by a two-way infinite path in “1 \Rightarrow 2”.

Using the ontology \mathcal{O} defined above we define a new ontology \mathcal{O}' to obtain the 2EXPTIME lower bound for signatures $\Sigma' = \text{sig}(\mathcal{O}') \setminus \{b\}$. Fix a role name r_E for any concept name $E \in \text{sig}(\mathcal{O}) \setminus \Sigma$. Now replace in \mathcal{O} any occurrence of $E \in \text{sig}(\mathcal{O}) \setminus \Sigma$ by $\exists r_E.\{b\}$ and denote the resulting ontology by \mathcal{O}' .

Lemma 40. *The following conditions are equivalent:*

1. M accepts w ;
2. there exist models \mathcal{I} and \mathcal{J} of \mathcal{O}' such that $\mathcal{I}, b^{\mathcal{I}} \sim_{\mathcal{ALCCO}, \pm'} \mathcal{J}, d$, for some $d \neq b^{\mathcal{J}}$.

Proof. “1 \Rightarrow 2”. We modify the interpretation \mathcal{I} defined in the proof of Lemma 39 in such a way that we obtain a model of \mathcal{O}' and such that the $\mathcal{ALCCO}(\pm)$ -bisimulation S on \mathcal{I} defined in that proof is, in fact, an $\mathcal{ALCCO}(\pm')$ -bisimulation on the new interpretation. Formally, obtain \mathcal{I}' from \mathcal{I} by interpreting every r_E , $E \in \text{sig}(\mathcal{O}) \setminus \Sigma$ as follows:

- (i) there is an r_E -edge from e to $b^{\mathcal{I}}$, for all $e \in E^{\mathcal{I}}$;
- (ii) there is an r_E -edge from e to all elements on the path ρ , for all $(e, e') \in S$ and $e' \in E^{\mathcal{I}}$;
- (iii) there are no more r_E -edges.

Note that, by (i), \mathcal{I}' is a model of \mathcal{O}' . By (ii), the relation S defined in the proof of Lemma 39 is an $\mathcal{ALCCO}(\pm')$ -bisimulation. In particular, by (i), elements $e' \in E^{\mathcal{I}}$ have now an r_E -edge to $b^{\mathcal{I}}$, so any element e bisimilar to e' , that is, $(e, e') \in S$, needs an r_E -successor to some element bisimilar to $b^{\mathcal{I}}$. Since all elements on the path ρ are bisimilar to $b^{\mathcal{I}}$, these r_E -successors exist due to (ii).

“2 \Rightarrow 1”. This direction remains the same as in the proof of Lemma 39. \square

The extension to DLs with inverse roles and the universal role and the restriction to a single role name are again straightforward.

We conclude the section with an observation that will be relevant for the application of our results to modal logic in Section 11. More specifically, we strengthen the lower bound for the case of $\mathcal{L} = \mathcal{ALCCO}^\square$ as follows:

Lemma 41. *Let $\mathcal{L} \in \{\mathcal{ALCCO}, \mathcal{ALCCO}^\square\}$. Then, it is 2EXPTIME-hard to decide for an \mathcal{L} -ontology \mathcal{O} , individual name b , and signature $\Sigma \subseteq \text{sig}(\mathcal{O}) \setminus \{b\}$ whether $\{b\}$ and $\neg\{b\}$ are jointly consistent under \mathcal{O} modulo $\mathcal{L}(\Sigma)$ -bisimulations, even if \mathcal{O} is allowed to use only a single role name.*

Proof. We modify the ontology \mathcal{O} and signature Σ used in the proof of Lemma 38. Let \mathcal{O}' be the ontology obtained from \mathcal{O} by:

- replacing every subconcept of the shape $\exists r.C$ with $\exists r.(X_r \sqcap C)$ and
- replacing every subconcept of the shape $\exists s.C$ with $\exists s.(X_s \sqcap C)$,

for fresh concept names X_r, X_s , and set $\Sigma' = \Sigma \cup \{X_r, X_s\}$. It is routine to verify that Lemma 39 holds for \mathcal{O}', Σ' instead of \mathcal{O}, Σ . In particular, we can obtain an interpretation \mathcal{I}' from \mathcal{I} as constructed in “1 \Rightarrow 2” as follows.

- replace all s -connections by r -connections;
- every element that has an s -predecessor in \mathcal{I} satisfies X_s in \mathcal{I}' , that is, $X_s^{\mathcal{I}'} = (\exists s^-. \top)^{\mathcal{I}}$;
- $b^{\mathcal{I}}$ and every element on the infinite r -path ρ in \mathcal{I} satisfy X_r in \mathcal{I}' , that is, $X_r^{\mathcal{I}'} = (\exists r. \top)^{\mathcal{I}}$ (the root of the infinite path has to satisfy X_r since it is bisimilar to $b^{\mathcal{I}}$ which satisfies X_r).

□

7.2 Shape of Explicit Definitions in the Lower Bound

The goal of this subsection is to provide some intuition on the shape of the explicit definitions that arise in the proof of Lemma 38. We note first that $r(x, x)$ is an explicit FO(Σ)-definition of $\{b\}$ under \mathcal{O} , regardless of whether the ATM accepts its input or not. This means that interpolant and explicit definition existence is 2EXPTIME-hard even under the promise that a fixed FO-definition / FO-interpolant exists.

We now analyze the $\mathcal{ALCCO}(\pm)$ -definitions that arise in the proof of Lemma 38. Recall that such a definition exists iff the ATM M does not accept its input w . So, for the rest of the discussion we assume the latter.

Instead of directly providing an explicit $\mathcal{ALCO}(\pm)$ -definition of $\{b\}$, we give a definition $C_{\neg b}$ of $\neg\{b\}$, since the definition of $C_{\neg b}$ is close to the intuitions provided in the proof of Lemma 38. Obviously, $\neg C_{\neg b}$ will be the desired definition of $\{b\}$. Let n be the length of the input word w and let $k = |\Gamma \cup (Q \times \Gamma)|$ be the number of possible labelings of a cell in some configuration of the ATM. Moreover, set $K = k^{2^n} + 2^n$.

The concept $C_{\neg b}$ takes the shape

$$C_{\neg b} = \exists r. \top \rightarrow (C_{\text{tree}} \sqcap C_{\text{start}} \sqcap \neg C_{\text{stop}} \sqcap \bigsqcup_{i=0}^{2^n-1} C_i).$$

To understand the structure $\exists r. \top \rightarrow C'$ of $C_{\neg b}$, recall that the proof of Lemma 38 relies on the assumption that an element $d \neq b^{\mathcal{I}}$ has an r -successor. The concepts $C_{\text{tree}}, C_{\text{stop}}, C_{\text{start}}, C_i$ provide an ‘‘approximation’’ of an accepting computation tree of the ATM M on its input w in the following sense. (Note that the definition of $\neg\{b\}$ cannot describe the full accepting computation since it is not entailed).

The concept C_{tree} enforces an s -tree of depth K that acts as the skeleton for encoding (an initial fragment of) a computation tree. It is labeled with concepts $Z, B_{\forall}, B_{\exists}^1, B_{\exists}^2$ in the expected way. Formally, C_{tree} is

$$\begin{aligned} & \text{Path}_{s, B_{\forall}}^{2^n} \sqcap \bigsqcap_{\substack{i=\ell \cdot 2^n - 1 \\ i < K}} \forall s^i. (B_{\forall} \rightarrow (\exists s. (Z \sqcap \text{Path}_{s, B_{\exists}^1}^{2^n}) \sqcap \exists s. (\neg Z \sqcap \text{Path}_{s, B_{\exists}^2}^{2^n}))) \\ & \sqcap (B_{\exists}^1 \sqcup B_{\exists}^2) \rightarrow \exists s. \text{Path}_{s, B_{\forall}}^{2^n} \end{aligned}$$

where $\text{Path}_{s, X}^m$ is a concept that enforces an s -path of length m with each element labeled with X . We refrain from giving the precise definitions of the remaining concepts, and rather provide the intuitions. C_{start} is a concept that enforces the initial configuration to be true in the computation tree, and C_{stop} is a concept that is true if some element within K s -steps is labeled with a concept name $A_{q,a}$ for which $\Delta(q, a) = \emptyset$. Moreover, each C_i is a concept with $\mathcal{O} \models I_s \sqcap (A = i) \sqsubseteq C_i$; recall that we denote with $(A = i)$ that the A -counter has value i . The disjunction over all possible C_i in $C_{\neg b}$ is needed since the A -counter can take any value between 0 and $2^n - 1$ at a given element in $d \neq b^{\mathcal{I}}$. More precisely, each C_i is a conjunction

$$C_i = \bigsqcap_{j=0}^{2^n-1} \forall r^j. C_{\text{sync}}^{i \oplus_{2^n} j},$$

where \oplus_m denotes addition modulo m , and for each m with $0 \leq m < 2^n$, C_{sync}^m is a concept that coordinates the content of the m -th cell in every

configuration in the computation tree with the same cell in the successor configuration(s). This can be easily realized using value restrictions $\forall s$.

Observe that $\mathcal{O} \models \neg\{b\} \sqsubseteq C_{\neg b}$ regardless of whether the ATM accepts w or not. In particular, in every model of \mathcal{O} , each element d satisfying $\neg\{b\} \sqcap \exists r.\top$ satisfies the concepts $C_{\text{tree}}, C_{\text{start}}$, and $\neg C_{\text{stop}}$. Moreover, d satisfies I_s and $(A = i)$ for some i , and thus d also satisfies C_i .

For the converse, $\mathcal{O} \models C_{\neg b} \sqsubseteq \neg\{b\}$, suppose that $C_{\neg b}$ is realizable in a model \mathcal{I} of \mathcal{O} in an element d with $(d, d) \in r^{\mathcal{I}}$. We thus also have $d \in (C_{\text{tree}} \sqcap C_{\text{start}} \sqcap \neg C_{\text{stop}})^{\mathcal{I}}$, and $d \in C_i^{\mathcal{I}}$, for some i . Due to the r -self loop, $d \in (C_{\text{sync}}^m)^{\mathcal{I}}$, for all m with $0 \leq m < 2^n$. But this means that at d starts the initial segment of a computation tree of M which is not labeled with a halting configuration, and all of whose cells are coordinated with the corresponding cell of the successor configuration(s). By the choice of K , on every path there is a configuration that occurs twice. We can thus extend the initial fragment of the computation tree to an infinite computation tree for the word w , in contradiction to the fact that M does not accept w .

We conclude with observing that the size of the definition $C_{\neg b}$ of $\neg\{b\}$ is double exponential in the length n of the input word, due to the depth K of the enforced tree. This is in stark contrast with the (constant!) size of the $\text{FO}(\Sigma)$ -definition. We conjecture that one can enforce explicit definitions of triple exponential size. For example, when using two roles s_1, s_2 instead of s for encoding the computation tree, already the concept C_{tree} will be of triple exponential size. We leave a detailed analysis for future work.

7.3 DLs with Role Inclusions

By Theorem 23, it suffices to prove the following.

Lemma 42. *Let $\mathcal{L} \in \text{DL}_{nr}$ admit role inclusions. It is 2EXPTIME-hard to decide for an \mathcal{L} -ontology \mathcal{O} , concept C , and signature $\Sigma \subseteq \text{sig}(\mathcal{O})$ whether C and $\neg C$ are jointly consistent under \mathcal{O} modulo $\mathcal{L}(\Sigma)$ -bisimulations.*

As in the proof of Lemma 38, we reduce the word problem for exponentially space bounded ATMs, so let M be a 2^n -space bounded ATM and $w = a_0 \dots a_{n-1}$ an input of length n . In fact, the only difference to the proof of Lemma 38 is the way in which we enforce that exponentially many elements are $\mathcal{L}(\Sigma)$ -bisimilar. We first provide the reduction for $\mathcal{L} = \mathcal{ALCH}$ and

$$\Sigma = \{r_1, r_2, s, Z, B_{\forall}, B_{\exists}^1, B_{\exists}^2\} \cup \{A_{\sigma} \mid \sigma \in \Gamma \cup (Q \times \Gamma)\}.$$

The symbols $s, Z, B_{\forall}, B_{\exists}^1, B_{\exists}^2$ and $A_{\sigma}, \sigma \in \Gamma \cup (Q \times \Gamma)$, play exactly the same role as above. The main difference is that we replace the nominal b by an r -chain of length n . The ontology \mathcal{O} contains the RIs $r \sqsubseteq r_1, r \sqsubseteq r_2$ and the CI $\neg \exists r^n. \top \sqcap \exists r_1^n. \top \sqsubseteq R$. As usual $\exists r^n$ abbreviates a sequence of n times $\exists r$.

To see how we use these inclusions, suppose there exist models \mathcal{I} and \mathcal{J} of \mathcal{O} and $d \in \Delta^{\mathcal{I}}, e \in \Delta^{\mathcal{J}}$ such that

- $d \in (\exists r^n. \top)^{\mathcal{I}}$;
- $e \in (\neg \exists r^n. \top)^{\mathcal{J}}$;
- $\mathcal{I}, d \sim_{\mathcal{ALCH}, \Sigma} \mathcal{J}, e$.

then it follows that $e \in R^{\mathcal{J}}$: due to $\mathcal{I}, d \sim_{\mathcal{ALCH}, \Sigma} \mathcal{J}, e$ and $d \in (\exists r_1^n. \top)^{\mathcal{I}}$, we also have $e \in (\exists r_1^n. \top)^{\mathcal{J}}$. Let now d' be an element reachable from d via an r -path of length n (which exists due to $d \in (\exists r^n. \top)^{\mathcal{I}}$). Since $r \sqsubseteq r_i$ for $i = 1, 2$, there are also arbitrary r_1/r_2 -paths of length n from d to d' . Since $\mathcal{I}, d \sim_{\mathcal{ALCH}, \Sigma} \mathcal{J}, e$, there are also arbitrary r_1/r_2 -paths of length n starting in e and whose end points are all $\mathcal{ALCH}(\pm)$ -bisimilar to d' and thus also mutually $\mathcal{ALCH}(\pm)$ -bisimilar. The concept name R will enforce that

- (*) the end point of any r_1/r_2 -path of length n starting in e carries a counter value that describes the path in a canonical way.

We can thus use these 2^n different, but bisimilar end points to start the infinite trees which mimick the computation tree of M as in the proof of Lemma 38. Along these we maintain the same two counters as there:

- one counter starting at 0 and counting modulo 2^n to divide the tree into configurations of length 2^n ;
- another counter starting at *the value of the counter on the leaf* and also counting modulo 2^n .

Formally, the ontology \mathcal{O} is constructed as follows. In order to realize (*) above, we use concept names $A_i, 0 \leq i < n$ realizing the counter and the following concept inclusions:

$$\begin{aligned}
R &\sqsubseteq R_0 \\
R_i &\sqsubseteq \forall r_1. (A_i \sqcap R_{i+1}) \sqcap \forall r_2. (\neg A_i \sqcap R_{i+1}) && i < n \\
R_i \sqcap A_j &\sqsubseteq \forall r_1. A_j \sqcap \forall r_2. A_j && 0 \leq j < i < n \\
R_i \sqcap \neg A_j &\sqsubseteq \forall r_1. \neg A_j \sqcap \forall r_2. \neg A_j && 0 \leq j < i < n \\
R_n &\sqsubseteq L_R
\end{aligned}$$

Using the concept name L_R , we start the s -trees with two counters, realized using concept names U_i and V_i , $0 \leq i < n$, and initialized to 0 and the value of the A -counter, respectively:

$$\begin{aligned} L_R &\sqsubseteq (U = 0) \\ L_R &\sqsubseteq A_j \leftrightarrow V_j && 0 \leq j < n \\ \top &\sqsubseteq \exists s. \top \end{aligned}$$

The structure of the computation tree, the initial configuration, and the coordination between consecutive configurations is done using the same concept inclusions as in the proof of Lemma 38, starting from inclusion (†) and replacing I_s with L_R . We can then prove the following very similarly to Lemma 39.

Lemma 43. *The following conditions are equivalent:*

1. M accepts w ;
2. there exist models \mathcal{I} and \mathcal{J} of \mathcal{O} such that $\mathcal{I}, d \sim_{\mathcal{ALCH}, \Sigma} \mathcal{J}, e$, for some $d \in (\exists r^n. \top)^{\mathcal{I}}$ and $e \notin (\exists r^n. \top)^{\mathcal{J}}$.

Proof. “1 \Rightarrow 2”. If M accepts w , there is a computation tree of M on w . We construct a single interpretation \mathcal{I} with $\mathcal{I}, d \sim_{\mathcal{ALCH}, \pm} \mathcal{I}, e$ for some d, e with $d \in (\exists r^n. \top)^{\mathcal{I}}$ and $e \notin (\exists r^n. \top)^{\mathcal{I}}$ as follows. Let $\hat{\mathcal{J}}$ be the infinite tree-shaped interpretation that represents the computation tree of M on w as described above, that is, configurations are represented by sequences of 2^n elements linked by role s and labeled by $B_{\forall}, B_{\exists}^1, B_{\exists}^2$ depending on whether the configuration is universal or existential, and in the latter case the superscript indicates which choice has been made for the existential state. Finally, the first element of the first successor configuration of a universal configuration is labeled with Z . Observe that $\hat{\mathcal{J}}$ interprets only the symbols in Σ as non-empty. Now, we obtain interpretations \mathcal{I}_k , $k < 2^n$ from $\hat{\mathcal{J}}$ by interpreting non- Σ -symbols as follows:

- the root of \mathcal{I}_k satisfies L_R ;
- the U -counter starts at 0 at the root and counts modulo 2^n along each s -path;
- the V -counter starts at k at the root and counts modulo 2^n along each s -path;
- the auxiliary concept names of the shape A_{σ}^i and A'_{σ} are interpreted in a minimal way so as to satisfy the concept inclusions that enforce the coordination between consecutive configurations (cf. the concept inclusions in proof of Lemma 38).

Now obtain \mathcal{I} from $\widehat{\mathcal{J}}$ and the \mathcal{I}_k as follows: First, create a path of length n from some element d so that consecutive elements are connected with r, r_1, r_2 , and identify the end point of the path with the root of $\widehat{\mathcal{J}}$. Then create a binary tree of depth n , rooted in e , in which left children are always r_1 -successors and right children are always r_2 -successors. Label the nodes of the tree with R_i and A_j as described above and identify the leaf having A -counter value k with the root of \mathcal{I}_k , for all $k < 2^n$. \mathcal{I} is as required since, by construction, $d \in (\exists r^n. \top)^{\mathcal{I}}$, $e \notin (\exists r^n. \top)^{\mathcal{I}}$, and the reflexive, transitive, and symmetric closure of

- all pairs (d', e') such that d' has distance $\ell \leq n$ from d and e' has distance ℓ from e , and
- all pairs (b, b') , with b in $\widehat{\mathcal{J}}$ and b' a copy of b in some tree \mathcal{I}_k

is an $\mathcal{ALCH}(\pm)$ -bisimulation S on \mathcal{I} with $(d, e) \in S$.

“2 \Rightarrow 1”. Assume that $\mathcal{I}, d \sim_{\mathcal{ALCH}, \pm} \mathcal{J}, e$ for models \mathcal{I}, \mathcal{J} of \mathcal{O} and some d, e with $d \in (\exists r^n. \top)^{\mathcal{I}}$ and $e \notin (\exists r^n. \top)^{\mathcal{J}}$. As argued above, there are r_1/r_2 -paths of length n whose end points carry all possible counters $< 2^n$ and are all $\mathcal{ALCH}(\pm)$ -bisimilar. In addition, all these end points root s -trees which are $\mathcal{ALCH}(\pm)$ -bisimilar. Since \mathcal{J} is a model of \mathcal{O} , all these s -trees are additionally labeled with some auxiliary concept names not in Σ , depending on the value of the A -counter of the corresponding leaf. Using the concept inclusions in \mathcal{O} and the arguments given in their description, it can be shown that all s -trees contain a computation tree of M on input w (which is solely represented with concept names in Σ). \square

The same proof works as well for \mathcal{ALCH}^\square as the relation S constructed in the direction “1 \Rightarrow 2” above is actually an $\mathcal{ALCH}^\square(\pm)$ -bisimulation. For $\mathcal{L} \in \{\mathcal{ALCHI}, \mathcal{ALCHI}^u\}$, we have to slightly adapt the model construction in “1 \Rightarrow 2”, following the idea provided in Example 24 (except that we do not need to take the union of $\mathcal{I}_1, \mathcal{I}_2$ here, since we construct a single interpretation $\mathcal{I} = \mathcal{I}_1 = \mathcal{I}_2$). Let d_0, \dots, d_n be the elements on the r -path that starts in d , that is, $d_0 = d$ and d_ℓ has distance ℓ from d . Recall that $(d_\ell, e') \in S$ for every element e' in level ℓ in the binary tree rooted at e . Observe that S is not an $\mathcal{L}(\Sigma)$ -bisimulation since, for $\ell > 0$, d_ℓ has both an r_1 and an r_2 -predecessor (both are $d_{\ell-1}$), but elements in the binary tree lack either an r_1 - or an r_2 -predecessor. To repair this, we add for every element e' in level $\ell > 0$ in the binary tree the following connections:

$$(d_{\ell-1}, e') \in r_1^{\mathcal{I}} \quad \text{and} \quad (d_{\ell-1}, e') \in r_2^{\mathcal{I}}.$$

It can be verified that the modified interpretation is still a model of \mathcal{O} , and that S is an $\mathcal{L}(\Sigma)$ -bisimulation as required.

We conclude the section by remarking that one can analyze the structure of the explicit $\mathcal{ALCH}(\pm)$ -definitions that arise in the proof of Lemma 42 along the lines of Section 7.2. In contrast to that section, the size of the FO-definition

$$\varphi(x_1) = \exists x_2 \dots \exists x_n. \bigwedge_{i=1}^{n-1} r_1(x_i, x_{i+1}) \wedge r_2(x_i, x_{i+1})$$

of $\exists r^n. \top$ under \mathcal{O} is not constant, but depends on n .

7.4 CI-interpolant Existence

We show the 2EXPTIME lower bound for CI-interpolant existence stated in Theorem 31. We employ the ontology \mathcal{O} , individual b , and signature Σ constructed in the proof of Lemma 39 and remind the reader that the claim of Lemma 39 holds also for \mathcal{ALCO}^u and \mathcal{ALCOI}^u . Let \mathcal{O}_1 be defined as \mathcal{O} without $\{b\} \sqsubseteq \exists r. \{b\}$ and with $\neg\{b\} \sqcap \exists r. \top \sqsubseteq I_s$ replaced by $\exists r. \top \sqsubseteq I_s$. Also define \mathcal{O}_2 as \mathcal{O} without $\{b\} \sqsubseteq \exists r. \{b\}$ and with all concept and role names not in Σ replaced by fresh symbols. Transform \mathcal{O}_2 into an equivalent ontology of the form $\{\top \sqsubseteq D\}$. Observe that \mathcal{O}_1 does not use b . In fact, the shared symbols of \mathcal{O}_1 and the CI $\forall u. D \sqcap \{b\} \sqsubseteq \neg \exists r. \{b\}$ are exactly the symbols in Σ . The 2EXPTIME lower bound now follows from the following lemma.

Lemma 44. *Let $\mathcal{L} \in \{\mathcal{ALCO}^u, \mathcal{ALCOI}^u\}$. Then the following conditions are equivalent:*

1. *Point 2 of Lemma 39 holds; that is, there exist models \mathcal{I} and \mathcal{J} of \mathcal{O} such that $\mathcal{I}, b^{\mathcal{I}} \sim_{\mathcal{L}, \Sigma} \mathcal{J}, d$, for some $d \neq b^{\mathcal{J}}$;*
2. *there does not exist an \mathcal{L} -CI interpolant for \mathcal{O}_1 and $\forall u. D \sqcap \{b\} \sqsubseteq \neg \exists r. \{b\}$.*

Proof. Assume Point (1) holds and take \mathcal{I}, \mathcal{J} , and d witnessing this. We may assume that $\mathcal{I} = \mathcal{J}$ is the interpretation constructed in the proof of “1 \Rightarrow 2” of Lemma 39. Assume for a proof by contradiction that \mathcal{O}' is an \mathcal{L} -CI interpolant for \mathcal{O}_1 and $\forall u. D \sqcap \{b\} \sqsubseteq \neg \exists r. \{b\}$. Let \mathcal{I}' denote the restriction of \mathcal{I} to elements that cannot be reached from b along a path following $r^{\mathcal{I}}$ or $s^{\mathcal{I}}$ and reinterpret b as an element of $\Delta^{\mathcal{I}'}$. Then \mathcal{I}' is a model of \mathcal{O}_1 by the definition of \mathcal{I} and since \mathcal{O}_1 does not contain any CIs with the individual b . Moreover, we have $\mathcal{I}, b^{\mathcal{I}} \sim_{\mathcal{L}, \Sigma} \mathcal{I}', d$ since $b \notin \Sigma$. Then, as \mathcal{L} admits the universal role, \mathcal{I} is a model of \mathcal{O}' . We now reinterpret in \mathcal{I} the fresh concept and role names in \mathcal{O}_2 in the same way

as the original ones in \mathcal{I} and obtain a model \mathcal{I}'' with $\Delta^{\mathcal{I}''} = D^{\mathcal{I}''}$ since \mathcal{I} is a model of \mathcal{O} . But then $\mathcal{I}'' \not\models \forall u.D \sqcap \{b\} \sqsubseteq \neg \exists r.\{b\}$ and so (as \mathcal{I}'' is still a model of \mathcal{O}' since $\text{sig}(\mathcal{O}') \subseteq \Sigma$) $\mathcal{O}' \not\models \forall u.D \sqcap \{b\} \sqsubseteq \neg \exists r.\{b\}$, a contradiction.

Conversely, assume there does not exist an \mathcal{L} -CI interpolant for \mathcal{O}_1 and $\forall u.D \sqcap \{b\} \sqsubseteq \neg \exists r.\{b\}$. As in the proof of Theorem 21 and using the fact that \mathcal{L} admits the universal role, we obtain a model \mathcal{J} of \mathcal{O}_1 , $d \in \Delta^{\mathcal{J}}$, and an interpretation \mathcal{I} with $\mathcal{I} \not\models \forall u.D \sqcap \{b\} \sqsubseteq \neg \exists r.\{b\}$ and $\mathcal{I}, b^{\mathcal{I}} \sim_{\mathcal{L}, \Sigma} \mathcal{J}, d$. We may assume that \mathcal{I} and \mathcal{J} are disjoint. Observe that \mathcal{J} satisfies all CIs in \mathcal{O} with the exception of $\{b\} \sqsubseteq \exists r.\{b\}$. By reinterpreting in \mathcal{I} the original concept and role names in \mathcal{O} in the same way as the fresh concept and role names in \mathcal{O}_2 , we obtain a model \mathcal{I}' of \mathcal{O} . Take the union $\mathcal{I}' \cup \mathcal{J}$ of \mathcal{I}' and \mathcal{J} with $b^{\mathcal{I}' \cup \mathcal{J}}$ defined as $b^{\mathcal{I}'}$. Then $\mathcal{I}' \cup \mathcal{J}$ is a model of \mathcal{O} such that $\mathcal{I}' \cup \mathcal{J}, b^{\mathcal{I}' \cup \mathcal{J}} \sim_{\mathcal{L}, \Sigma} \mathcal{I}' \cup \mathcal{J}, d$, for some $d \neq b^{\mathcal{J}}$, as required for Point (1). \square

8 Upper Bound Proofs without Ontology

The upper bound for Points 1 and 2 of Theorem 28 is a consequence of the respective upper bounds in Theorem 27. For showing the upper bounds of Points 3 and 4 in Theorem 28, we prove that joint consistency is in NEXPTIME and then apply Theorem 21. Indeed, the NEXPTIME upper bound follows directly from the following exponential size witness model property.

Lemma 45. *Let $\mathcal{L} \in DL_{nr}$ admit neither the universal role nor both inverse roles and nominals simultaneously. Let \mathcal{O} be a set of RIs, C_1, C_2 \mathcal{L} -concepts, and Σ a signature. If C_1 and C_2 are jointly consistent under \mathcal{O} modulo $\mathcal{L}(\Sigma)$ -bisimulations, then there exist pointed interpretations \mathcal{I}_1, d_1 and \mathcal{I}_2, d_2 with $\mathcal{I}_1, \mathcal{I}_2$ models of \mathcal{O} and of at most exponential size in $\|\mathcal{O}\| + \|C_1\| + \|C_2\|$ such that $d_1 \in C_1^{\mathcal{I}_1}$, $d_2 \in C_2^{\mathcal{I}_2}$, and $\mathcal{I}_1, d_1 \sim_{\mathcal{L}, \Sigma} \mathcal{I}_2, d_2$.*

Before we prove Lemma 45, we introduce some notation. The *depth* of a concept C is the number of nestings of existential restrictions in C . For instance, a concept name has depth 0 and $\exists r.\exists r.B$ has depth 2. Given the ontology \mathcal{O} , concepts C_1, C_2 , and the signature Σ , we use the notation introduced in Section 6. For instance, the set of concepts Ξ , Ξ -types t , and mosaics (T_1, T_2) are defined as in Section 6. While in Section 6 we used the relation \rightsquigarrow_r between mosaics to guide the construction of interpretations, here we use a relation between mosaics that is directly

induced by interpretations. Assume interpretations \mathcal{I}_1 and \mathcal{I}_2 are given. Consider mosaics $p = (T_1(d), T_2(d))$ and $q = (T_1(d'), T_2(d'))$ such that there exists a role name $r \in \Sigma$ with $(d, d') \in r^{\mathcal{I}_i}$, for some $i \in \{1, 2\}$. Then define, for every role name s and $i \in \{1, 2\}$, relations $R_{p,q}^{s,i} \subseteq T_i(d) \times T_i(d')$ by setting $(t, t') \in R_{p,q}^{s,i}$ if there exist e and e' realizing t and t' , respectively, with $(T_1(e), T_2(e)) = p$ and $(T_1(e'), T_2(e')) = q$, such that $(e, e') \in s^{\mathcal{I}_i}$.

Now assume that C_1 and C_2 are jointly consistent under \mathcal{O} modulo $\mathcal{L}(\Sigma)$ -bisimulations. By definition, there exist pointed models \mathcal{I}_1, d_1 and \mathcal{I}_2, d_2 of \mathcal{O} such that $d_1 \in C_1^{\mathcal{I}_1}$, $d_2 \in C_2^{\mathcal{I}_2}$, and $\mathcal{I}_1, d_1 \sim_{\mathcal{L}, \Sigma} \mathcal{I}_2, d_2$. Let k be the maximum depth of C_1, C_2 .

We start with the case involving nominals and without inverse roles. We construct exponential size $\mathcal{J}_1, \mathcal{J}_2$ with the same properties as $\mathcal{I}_1, \mathcal{I}_2$ above. Intuitively, \mathcal{J}_i is obtained via a suitable unraveling operation up to the depth k of the concepts C_1, C_2 ; during the unraveling, we take care of the nominals and, moreover, restrict the outdegree of the produced interpretation by keeping only necessary successors. Formally, let \mathcal{B} be some minimal set of mosaics defined by $\mathcal{I}_1, \mathcal{I}_2$ such that

- $(T_1(d_1), T_2(d_1)) \in \mathcal{B}$;
- \mathcal{B} contains every mosaic generated by some nominal, or formally, $(T_1(d), T_2(d)) \in \mathcal{B}$ for every $d \in \Delta^{\mathcal{I}_i}$ such that $d = a^{\mathcal{I}_i}$ for some nominal $a \in \text{sig}(C_i)$;
- for every type t realized in \mathcal{I}_i there exists $(T_1, T_2) \in \mathcal{B}$ with $t \in T_i$.

Intuitively, \mathcal{B} serves to describe the behavior of the root of the unraveling (first item), of the nominals (second item), and of potential witnesses for existential restrictions for non- Σ -roles (third item). Observe that the size of \mathcal{B} is at most exponential in the size of \mathcal{O}, C_1, C_2 . To restrict the outdegree, select, for any mosaic $p = (T_1, T_2)$ defined by $\mathcal{I}_1, \mathcal{I}_2$ and any $\exists s.C \in t \in T_i$ such that there exists $r \in \Sigma$ with $\mathcal{O} \models s \sqsubseteq r$, a mosaic $q = (T'_1, T'_2)$ such that $(t, t') \in R_{p,q}^{s,i}$ and $C \in t'$, and denote the resulting set by $\mathcal{S}(p)$. Form the set \mathcal{T} of sequences

$$\sigma = p_0 \cdots p_j = (T_1^0, T_2^0) \cdots (T_1^j, T_2^j)$$

with $j \leq k$, $p_0 \in \mathcal{B}$ and $p_{i+1} \in \mathcal{S}(p_i)$ for $i < j$. Let $\text{tail}(\sigma) = p_j$ and $\text{tail}_i(\sigma) = T_i^j$. We next define the domain of \mathcal{J}_1 and \mathcal{J}_2 as

$$\begin{aligned} \Delta^{\mathcal{J}_i} = \{ & (t, p) \mid t \in \text{tail}_i(p), p \in \mathcal{B} \} \cup \\ & \{ (t, \sigma) \mid \sigma \in \mathcal{T}, t \in \text{tail}_i(\sigma), |\sigma| > 1, t \text{ contains no nominal} \} \end{aligned}$$

and define the interpretation of individual, concept and role names in $\mathcal{J}_1, \mathcal{J}_2$ in the expected way:

- for any individual name a and $(T_1, T_2) \in \mathcal{B}$ with $\{a\} \in t \in T_i$, we set $a^{\mathcal{J}_i} = (t, (T_1, T_2))$;
- for any concept name A , $(t, \sigma) \in A^{\mathcal{J}_i}$ iff $A \in t$;
- for any role name r we let for $\sigma p \in \mathcal{T}$,
 - $((t, \sigma), (t', \sigma p)) \in r^{\mathcal{J}_i}$ if $(t, t') \in R_{\text{tail}(\sigma), p}^{r, i}$ and t' contains no nominal;
 - $((t, \sigma), (t', p)) \in r^{\mathcal{J}_i}$ if $(t, t') \in R_{\text{tail}(\sigma), p}^{r, i}$ and t' contains a nominal.

Next assume that $\text{tail}(\sigma) = (T_1, T_2)$ and σ has length k . If $\text{tail}(\sigma') = (T_1, T_2)$ for some $|\sigma'| < k$, then choose as r -successors of any element of the form (t, σ) exactly the r -successors of (t, σ') defined above. If no such σ' exists, then all elements of the form $(t, \text{tail}(\sigma))$ have distance exactly k from the roots (since no nominal occurs in any type in any mosaic in σ) and no successors are added.

It remains to take care of existential restrictions $\exists r.C$ for the role names r that do not entail any role name in Σ . If $\sigma \in \mathcal{T}$, $\exists r.C \in t \in T_i$ with $\text{tail}_i(\sigma) = T_i$ and $\mathcal{O} \not\models r \sqsubseteq s$ for any $s \in \Sigma$, we add $((t, \sigma), (t', p))$ to $r^{\mathcal{J}_i}$ (and all $s^{\mathcal{J}_i}$ with $\mathcal{O} \models r \sqsubseteq s$) for some $p = (T'_1, T'_2) \in \mathcal{B}$ and $t' \in T'_i$ with $C \in t'$ such that there are e, e' realizing t, t' in \mathcal{I}_i and $(e, e') \in r^{\mathcal{I}_i}$.

The following example illustrates the construction of $\mathcal{J}_1, \mathcal{J}_2$ using the interpretations $\mathcal{I}_1, \mathcal{I}_2$ introduced in Example 22.

Example 46. Let $t_0 = \text{tp}_{\Xi}(\mathcal{I}_1, a^{\mathcal{I}_1})$, $t_1 = \text{tp}_{\Xi}(\mathcal{I}_2, b^{\mathcal{I}_2})$, and $t_2 = \text{tp}_{\Xi}(\mathcal{I}_2, d)$. We ignore the types realized by $b^{\mathcal{I}_1}$ in \mathcal{I}_1 and by $a^{\mathcal{I}_2}$ in \mathcal{I}_2 as they are not relevant for understanding the construction. Then only the mosaic $p = (T_1, T_2)$ with $T_1 = \{t_0\}$ and $T_2 = \{t_1, t_2\}$ remains and \mathcal{J}_1 and \mathcal{J}_2 are depicted in Figure 8. \dashv




Fig. 8. Interpretations \mathcal{J}_1 and \mathcal{J}_2 illustrating Example 46.

We show that $\mathcal{J}_1, \mathcal{J}_2$ are as required. First, for $i \in \{1, 2\}$, $\mathcal{J}_i \models \mathcal{O}$ follows from the definition of \mathcal{J}_i and the fact that $\mathcal{I}_i \models \mathcal{O}$. Indeed, given $r \sqsubseteq s \in \mathcal{O}$, let $((t, \sigma), (t', \sigma')) \in r^{\mathcal{J}_i}$. This means that $(t, t') \in R_{\text{tail}(\sigma), \text{tail}(\sigma')}^{r, i}$, that is, there exist e, e' realizing t and t' , respectively, with $(T_1(e), T_2(e)) = \text{tail}(\sigma)$ and $(T_1(e'), T_2(e')) = \text{tail}(\sigma')$, such that $(e, e') \in r^{\mathcal{I}_i}$. Since $\mathcal{I}_i \models \mathcal{O}$, we obtain that $(e, e') \in s^{\mathcal{I}_i}$ as well, and thus $(t, t') \in R_{\text{tail}(\sigma), \text{tail}(\sigma')}^{s, i}$, meaning that $((t, \sigma), (t', \sigma')) \in s^{\mathcal{J}_i}$. Hence, $\mathcal{J}_i \models r \sqsubseteq s$.

We next prove that, for every $(t, \sigma) \in \Delta^{\mathcal{J}_i}$ and every concept $C \in \Xi$ of depth $\leq k - |\sigma|$,

$$(t, \sigma) \in C^{\mathcal{J}_i} \text{ iff } C \in t.$$

The proof is by induction on the structure of C . We consider the case $C = \exists r.D$, where D has depth $< k - |\sigma|$. We can assume that $|\sigma| < k$, since for $|\sigma| = k$ the claim holds trivially.

(\Rightarrow) Let $(t, \sigma) \in \exists r.D^{\mathcal{J}_i}$. Then $\exists r.D \in t$ follows by construction of $r^{\mathcal{J}_i}$ as we only have $((t, \sigma), (t', \sigma')) \in r^{\mathcal{J}_i}$ if there are e, e' realizing t, t' in \mathcal{I}_i such that $(e, e') \in r^{\mathcal{I}_i}$.

(\Leftarrow) Let $\text{tail}(\sigma) = p = (T_1, T_2)$ and suppose that $\exists r.D \in t \in T_i$. We distinguish two cases.

- There exists $s \in \Sigma$ such that $\mathcal{O} \models r \sqsubseteq s$. Then there exists $q = (T'_1, T'_2) \in \mathcal{S}(p)$ and $t' \in T'_i$ such that $(t, t') \in R_{p, q}^{r, i}$ and $D \in t'$. We distinguish two cases.
 - t' does not contain nominals. Then we have that $((t, \sigma), (t', \sigma q)) \in r^{\mathcal{J}_i}$. By inductive hypothesis, $(t', \sigma q) \in D^{\mathcal{J}_i}$, and thus $(t, \sigma) \in \exists r.D^{\mathcal{J}_i}$.
 - t' contains a nominal. Then we have that $((t, \sigma), (t', q)) \in r^{\mathcal{J}_i}$. By inductive hypothesis, $(t', q) \in D^{\mathcal{J}_i}$, hence $(t, \sigma) \in \exists r.D^{\mathcal{J}_i}$.
- For every $s \in \Sigma$, $\mathcal{O} \not\models r \sqsubseteq s$. By definition of \mathcal{J}_i , we have $((t, \sigma), (t', q)) \in r^{\mathcal{J}_i}$, for some $q = (T'_1, T'_2) \in \mathcal{B}$ and $t' \in T'_i$ such that $D \in t'$. By inductive hypothesis, $(t', q) \in D^{\mathcal{J}_i}$. Thus, $(t, \sigma) \in \exists r.D^{\mathcal{J}_i}$.

Next observe that the relation

$$S = \{((t, \sigma), (t', \sigma')) \in \Delta^{\mathcal{J}_1} \times \Delta^{\mathcal{J}_2} \mid \text{tail}(\sigma) = \text{tail}(\sigma')\}$$

is an $\mathcal{ALCHO}(\Sigma)$ -bisimulation. Indeed, for $((t, \sigma), (t', \sigma')) \in S$, we have the following.

- [AtomC] Let $(t, \sigma) \in A^{\mathcal{J}_1}$ and $A \in \Sigma$. By definition of \mathcal{J}_1 , we have that $(t, \sigma) \in A^{\mathcal{J}_1}$ iff $A \in t \in \text{tail}_1(\sigma)$, and thus $A \in t' \in \text{tail}_2(\sigma) = \text{tail}_2(\sigma')$, by definition of mosaics. But then $(t', \sigma') \in A^{\mathcal{J}_2}$. The converse direction is analogous.

[AtomI] Let $(t, \sigma) = a^{\mathcal{J}_1}$ and $a \in \Sigma$. By definition of \mathcal{J}_1 , $(t, \sigma) = a^{\mathcal{J}_1}$ iff $\{a\} \in t \in \text{tail}_1(\sigma)$, and thus $\{a\} \in t' \in \text{tail}_2(\sigma) = \text{tail}_2(\sigma')$, by definition of mosaics. But then $(t', \sigma') = a^{\mathcal{J}_2}$.

[Forth] Suppose that $((t, \sigma), (\hat{t}, \hat{\sigma})) \in r^{\mathcal{J}_1}$ with $r \in \Sigma$.

First, consider the case with $|\sigma|, |\sigma'| < k$. We have two possibilities.

– \hat{t} does not contain nominals. The following proof is illustrated in Figure 9. There is a mosaic p with $\hat{\sigma} = \sigma p$, and from $((t, \sigma), (\hat{t}, \sigma p)) \in r^{\mathcal{J}_1}$ we obtain $(t, \hat{t}) \in R_{\text{tail}(\sigma), p}^{r, 1}$. This means that there exist d, \hat{d} realizing t and \hat{t} , respectively, with $(T_1(d), T_2(d)) = \text{tail}(\sigma)$ and $(T_1(\hat{d}), T_2(\hat{d})) = p$, such that $(d, \hat{d}) \in r^{\mathcal{I}_1}$. As $t' \in T_2(d)$, there exists $e \in \Delta^{\mathcal{I}_2}$ with $\mathcal{I}_1, d \sim_{\mathcal{ALCH}\mathcal{O}, \Sigma} \mathcal{I}_2, e$ and e realizes t' . By the definition of bisimulations, there exists \hat{e} with $(e, \hat{e}) \in r^{\mathcal{I}_2}$ and $\mathcal{I}_1, \hat{d} \sim_{\mathcal{ALCH}\mathcal{O}, \Sigma} \mathcal{I}_2, \hat{e}$. Assume that \hat{e} realizes \hat{t}' . Then $\hat{t}' \in T_2(\hat{d})$ and $(\hat{t}, \hat{t}') \in R_{\text{tail}(\sigma'), p}^{r, 2}$. Now we consider again two possibilities.

- \hat{t}' does not contain nominals. Then from $(t', \hat{t}') \in R_{\text{tail}(\sigma'), p}^{r, 2}$ we obtain $((t', \sigma'), (\hat{t}', \sigma' p)) \in r^{\mathcal{J}_2}$. Since $\text{tail}(\sigma p) = \text{tail}(\sigma' p)$, we also obtain $((\hat{t}, \sigma p), (\hat{t}', \sigma' p)) \in S$.
- \hat{t}' contains nominals. Then from $(t', \hat{t}') \in R_{\text{tail}(\sigma'), p}^{r, 2}$ we obtain $((t', \sigma'), (\hat{t}', p)) \in r^{\mathcal{J}_2}$. Since $\text{tail}(\sigma p) = p$, we get that $((\hat{t}, \sigma p), (\hat{t}', p)) \in S$ as well.

In both cases, we obtain some $(\hat{t}', \hat{\sigma}')$ with $((t', \sigma'), (\hat{t}', \hat{\sigma}')) \in r^{\mathcal{J}_2}$ and $((\hat{t}, \hat{\sigma}), (\hat{t}', \hat{\sigma}')) \in S$, as required.

– \hat{t} contains nominals. In this case, $\hat{\sigma} = p$ for some mosaic p , and from $((t, \sigma), (\hat{t}, p)) \in r^{\mathcal{J}_1}$ we obtain $(t, \hat{t}) \in R_{\text{tail}(\sigma), p}^{r, 1}$. Now we can reason as above.

Now consider the case with $|\sigma| = k$ and $|\sigma'| < k$. As $\text{tail}(\sigma) = \text{tail}(\sigma')$, there exists σ'' such that $|\sigma''| < k$ and $\text{tail}(\sigma'') = \text{tail}(\sigma)$ and the r -successors of any node of the form (t, σ) are exactly the r -successors of (t, σ'') , and thus to show [Forth] one can proceed as above. The same argument applies if $|\sigma| < k$ and $|\sigma'| = k$ and if $|\sigma| = |\sigma'| = k$ and there exists σ'' with $\text{tail}(\sigma'') = \text{tail}(\sigma') = \text{tail}(\sigma)$ and $|\sigma''| < k$. Finally, if $|\sigma| = |\sigma'| = k$ but there does not exist any σ'' with $\text{tail}(\sigma'') = \text{tail}(\sigma') = \text{tail}(\sigma)$ and $|\sigma''| < k$, then there are no r -successors to consider.

[Back] Dual to [Forth].

Observe that the models \mathcal{J}_i , $i = 1, 2$, are at most exponential in the size of \mathcal{O}, C_1, C_2 . Moreover, we have $(T_1(d_1), (T_2(d_1))) \in \mathcal{B}$ and so $(\text{tp}_{\Xi}(\mathcal{I}_1, d_1), T_1(d_1)) \in C_1^{\mathcal{J}_1}$, $(\text{tp}_{\Xi}(\mathcal{I}_2, d_2), T_2(d_1)) \in C_2^{\mathcal{J}_2}$, and

$$((\text{tp}_{\Xi}(\mathcal{I}_1, d_1), T_1(d_1)), (\text{tp}_{\Xi}(\mathcal{I}_2, d_2), T_2(d_1))) \in S,$$




Fig. 9. Proof step to show that S satisfies [Forth], with $\hat{\sigma} = \sigma p$ and $\hat{\sigma}' = \sigma' p$ (if \hat{t}' does not contain nominals) and $\hat{\sigma}' = p$ (if \hat{t}' contains nominals).

as required.

We next consider the case with inverse roles, but without nominals. In this case, we let \mathcal{B} be some minimal set of mosaics defined by $\mathcal{I}_1, \mathcal{I}_2$ containing $(T_1(d_1), T_2(d_1))$ and such that for every type t realized in \mathcal{I}_i there exists $(T_1, T_2) \in \mathcal{B}$ with $t \in T_i$. We extend the relations $R_{p,q}^{s,i}$ defined previously to inverse roles s in the obvious way and select for any mosaic $p = (T_1, T_2)$ and any $\exists s.C \in t \in T_i$ such that there exists a Σ -role r with $\mathcal{O} \models s \sqsubseteq r$ a mosaic $q = (T'_1, T'_2)$ such that $(t, t') \in R_{p,q}^{s,i}$ and $C \in t'$ and denote the resulting set by $\mathcal{S}(p)$.

Form again the set \mathcal{T} of sequences

$$\sigma = p_0 \cdots p_j = (T_1^0, T_2^0) \cdots (T_1^j, T_2^j)$$

with $j \leq k$, $p_0 \in \mathcal{B}$ and $p_{i+1} \in \mathcal{S}(p_i)$ for $i < j$. Let $\text{tail}(\sigma) = p_j$ and $\text{tail}_i(\sigma) = T_i^j$. We next define the domain of \mathcal{J}_1 and \mathcal{J}_2 as

$$\Delta^{\mathcal{J}_i} = \{(t, \sigma) \mid \sigma \in \mathcal{T}, t \in \text{tail}_i(\sigma)\}$$

We define interpretations $\mathcal{J}_1, \mathcal{J}_2$ in the expected way.

- For any concept name A , $(t, \sigma) \in A^{\mathcal{J}_i}$ iff $A \in t$;
- Let r be a role name. Then we let for $\sigma p \in \mathcal{T}$,
 - $((t, \sigma), (t', \sigma p)) \in r^{\mathcal{J}_i}$ if $(t, t') \in R_{\text{tail}(\sigma), p}^{r,i}$;
 - $((t', \sigma p), (t, \sigma)) \in r^{\mathcal{J}_i}$ if $(t, t') \in R_{\text{tail}(\sigma), p}^{r-,i}$.
- We still have to take care of existential restrictions $\exists r.C$ with r a role that does not entail any Σ -role. If $\sigma \in \mathcal{T}$, $\exists r.C \in t \in T_i$ with $\text{tail}_i(\sigma) = T_i$ and $\mathcal{O} \not\models r \sqsubseteq s$ for any Σ -role s , we add $((t, \sigma), (t', p))$ to $r^{\mathcal{J}_i}$ (and all $s^{\mathcal{J}_i}$ with $\mathcal{O} \models r \sqsubseteq s$) for some $p = (T'_1, T'_2) \in \mathcal{B}$ and $t' \in T'_i$ with $C \in t'$ such that there are e, e' realizing t, t' in \mathcal{I}_i and $(e, e') \in r^{\mathcal{I}_i}$.

The fact that $\mathcal{J}_i \models \mathcal{O}$, for $i \in \{1, 2\}$, is proved similarly to the case with nominals. One can also prove again by induction on the structure of C that for every $(t, \sigma) \in \Delta^{\mathcal{J}_i}$ and every $C \in \Xi$ of depth $\leq k - |\sigma|$,

$$(t, \sigma) \in C^{\mathcal{J}_i} \text{ iff } C \in t.$$

Next we observe that the relation

$$S = \{((t, \sigma), (t', \sigma)) \in \Delta^{\mathcal{J}_1} \times \Delta^{\mathcal{J}_2} \mid \sigma \in \mathcal{T}\}$$

is an $\mathcal{ALCH}\mathcal{I}(\Sigma)$ -bisimulation. Indeed, it can be seen, similar to the case with nominals, that S satisfies [AtomC]. We now give a proof of [Forth]. We provide the proof for role names; the proof for inverse roles is similar.

[Forth] Let $((t, \sigma), (t', \sigma)) \in S$ and $((t, \sigma), (\hat{t}, \hat{\sigma})) \in r^{\mathcal{J}_1}$. We distinguish two cases. Assume first that there exists a mosaic p with $\hat{\sigma} = \sigma p$. Then $(t, \hat{t}) \in R_{\text{tail}(\sigma), p}^{r, 1}$. Thus, there exist d, \hat{d} realizing t, \hat{t} , respectively, such that $(T_1(d), T_2(d)) = \text{tail}(\sigma)$, $(T_1(\hat{d}), T_2(\hat{d})) = p$, and $(d, \hat{d}) \in r^{\mathcal{I}_1}$. Since $((t, \sigma), (t', \sigma)) \in S$, there exists e realizing t' such that $\mathcal{I}_1, d \sim_{\mathcal{ALCH}\mathcal{I}, \Sigma} \mathcal{I}_2, e$. As d and e are bisimilar, we also have some $\hat{e} \in \Delta^{\mathcal{I}_2}$ such that $(e, \hat{e}) \in r^{\mathcal{I}_2}$ and $\mathcal{I}_1, \hat{d} \sim_{\mathcal{ALCH}\mathcal{I}, \Sigma} \mathcal{I}_2, \hat{e}$, with \hat{e} realizing some \hat{t}' . Hence, $(t', \hat{t}') \in R_{\text{tail}(\sigma), p}^{r, 2}$, and it follows that $((t', \sigma), (\hat{t}', \sigma p)) \in r^{\mathcal{J}_2}$. Moreover, $((\hat{t}, \sigma p), (\hat{t}', \sigma p)) \in S$.

Assume now that $\sigma = \hat{\sigma} p$ for some mosaic p . Then $(\hat{t}, t) \in R_{\text{tail}(\hat{\sigma}), p}^{r^-, 1}$.

Thus, there exist \hat{d}, d realizing \hat{t}, t , respectively, such that $(T_1(\hat{d}), T_2(\hat{d})) = \text{tail}(\hat{\sigma})$, $(T_1(d), T_2(d)) = p$, and $(\hat{d}, d) \in (r^-)^{\mathcal{I}_1}$. Since $((t, \sigma), (t', \sigma)) \in S$, there exists e realizing t' such that $\mathcal{I}_1, d \sim_{\mathcal{ALCH}\mathcal{I}, \Sigma} \mathcal{I}_2, e$. As d and e are bisimilar, we also have some $\hat{e} \in \Delta^{\mathcal{I}_2}$ such that $(\hat{e}, e) \in (r^-)^{\mathcal{I}_2}$ and $\mathcal{I}_1, \hat{d} \sim_{\mathcal{ALCH}\mathcal{I}, \Sigma} \mathcal{I}_2, \hat{e}$, with \hat{e} realizing some \hat{t}' . Hence, $(\hat{t}', t') \in R_{\text{tail}(\hat{\sigma}), p}^{r^-, 2}$, and it follows that $((t', \sigma), (\hat{t}', \hat{\sigma})) \in r^{\mathcal{J}_2}$. Moreover, $((\hat{t}, \hat{\sigma}), (\hat{t}', \hat{\sigma})) \in S$.

Observe that again the models \mathcal{J}_i , $i = 1, 2$, are of at most exponential size in the size of \mathcal{O}, C_1, C_2 . We also have $(T_1(d_1), T_2(d_1)) \in \mathcal{B}$ and so $(\text{tp}_{\Xi}(\mathcal{I}_1, d_1), T_1(d_1)) \in C_1^{\mathcal{J}_1}$, $(\text{tp}_{\Xi}(\mathcal{I}_2, d_2), T_2(d_1)) \in C_2^{\mathcal{J}_2}$, and

$$((\text{tp}_{\Xi}(\mathcal{I}_1, d_1), T_1(d_1)), (\text{tp}_{\Xi}(\mathcal{I}_2, d_2), T_2(d_1))) \in S,$$

as required.

9 Lower Bound Proofs without Ontology

In this section, we first show (the hardness part of) Points 1 and 2 of Theorem 28 by a reduction of the case with ontologies, and then show

the nondeterministic exponential time lower bounds for Points 3 and 4 of that Theorem. Points 1 and 2 of Theorem 28 are a direct consequence of the following lemma.

Lemma 47. *Let $\mathcal{L} \in DL_{nr}$ admit the universal role or both inverse roles and nominals. Then the following holds:*

1. *if \mathcal{L} admits RIs, then projective \mathcal{L} -definition existence can be reduced in polynomial time to RI-ontology projective \mathcal{L} -definition existence;*
2. *if \mathcal{L} does not admit RIs, then projective \mathcal{L} -definition existence can be reduced in polynomial time to ontology-free projective \mathcal{L} -definition existence.*

Proof. Assume \mathcal{O} , C , C_0 , and Σ are given. We may assume that \mathcal{O} takes the form $\{\top \sqsubseteq D\} \cup \mathcal{O}'$ with \mathcal{O}' a set of RIs.

Assume first that \mathcal{L} admits the universal role. Then one can easily show that there exists an explicit $\mathcal{L}(\Sigma)$ -definition of C_0 under \mathcal{O} and C iff there exists an explicit $\mathcal{L}(\Sigma)$ -definition of C_0 under \mathcal{O}' and $C \sqcap \forall u.D$.

Now assume that \mathcal{L} admits inverse roles and nominals. We use the spy-point technique to encode the universal role [1]. Introduce a fresh individual a and a fresh role name r_0 and define U as the conjunction of the concepts

$$\{a\}, \quad \exists r_0.\{a\}, \quad \exists r_0.(\{b\} \sqcap \exists r_0.\{a\}), \quad \forall r_0^-. \forall s. \exists r_0.\{a\},$$

for all $s \in \{r, r^-\}$ with $r \in \text{sig}(\mathcal{O}, C, C_0)$ and $b \in \text{sig}(\mathcal{O}, C, C_0)$. Observe that if $d \in (U \sqcap \forall r_0^-. F)^{\mathcal{I}}$ for some interpretation \mathcal{I} and concept F , then $e \in F^{\mathcal{I}}$ holds for all elements e in $\Delta^{\mathcal{I}}$ that can be reached in \mathcal{I} from d or any $b^{\mathcal{I}}$ with $b \in \text{sig}(\mathcal{O}, C, C_0)$ along roles in $\text{sig}(\mathcal{O}, C, C_0)$. It follows that for any $\mathcal{L}(\Sigma)$ -concept E , we have

$$\mathcal{O} \models C \sqsubseteq (C_0 \leftrightarrow E) \quad \text{iff} \quad \mathcal{O}' \models (C \sqcap U \sqcap \forall r_0^-. D) \sqsubseteq (C_0 \leftrightarrow E).$$

Hence there exists an explicit $\mathcal{L}(\Sigma)$ -definition of C_0 under \mathcal{O} and C iff there exists an explicit $\mathcal{L}(\Sigma)$ -definition of C_0 under \mathcal{O}' and $C \sqcap U \sqcap \forall r_0^-. D$. \square

We show the lower bound for Theorem 28, Points 3 and 4, by proving NEXPTIME-hardness for the version of joint consistency formulated in Theorem 23. We reduce the exponential torus tiling problem. A *tiling system* is a triple $P = (T, H, V)$, where $T = \{0, \dots, k\}$ is a finite set of *tile types* and $H, V \subseteq T \times T$ are the *horizontal* and *vertical* matching conditions, respectively. An *initial condition* for P takes the form $c =$

$(c_0, \dots, c_{n-1}) \in T^n$. A mapping $\tau : \{0, \dots, 2^n - 1\} \times \{0, \dots, 2^n - 1\} \rightarrow T$ is a *solution for P and c* if $\tau(i, 0) = c_i$ for all $i < n$, and for all $i, j < 2^n$, the following conditions hold (where \oplus_k denotes addition modulo k):

- if $\tau(i, j) = t_1$ and $\tau(i \oplus_{2^n} 1, j) = t_2$, then $(t_1, t_2) \in H$;
- if $\tau(i, j) = t_1$ and $\tau(i, j \oplus_{2^n} 1) = t_2$, then $(t_1, t_2) \in V$.

It is well-known that the problem of deciding whether there is a solution for given P and c is NEXPTIME-hard [9, Section 5.2.2]. For the following constructions, assume a tiling system P and an initial condition c of length n .

For the reduction for \mathcal{ALCO} , we give concepts C, C_0 and a signature Σ such that here exist $\mathcal{I}_1, d_1 \sim_{\mathcal{ALCO}, \Sigma} \mathcal{I}_2, d_2$ with $d_1 \in (C \sqcap C_0)^{\mathcal{I}_1}$ and $d_2 \in (C \sqcap \neg C_0)^{\mathcal{I}_2}$ iff P has a solution given c . We start with setting

$$C_0 = \exists r^{2^n}. \{a\} \sqcap \forall r^{2^n}. \{a\}$$

with $a \notin \Sigma$ and $r \in \Sigma$. In addition to r , Σ contains concept names B_0, \dots, B_{2^n-1} that serve as bits in the binary representation of *grid positions* (i, j) with $0 \leq i, j \leq 2^n - 1$, where bits B_0, \dots, B_{n-1} represent the horizontal position i and B_n, \dots, B_{2^n-1} the vertical position j , and concept names T_0, \dots, T_k representing tile types. We also use the following concept names that are not in Σ : another four sets of concepts names A_0, \dots, A_{2^n-1} and V_0, \dots, V_{2^n-1} with $V \in \{X, Y, Z\}$ that also serve as bits in the binary representation of grid position (i, j) with $0 \leq i, j \leq 2^n - 1$, and concept names $R_0, \dots, R_{2^n}, M, M_1$, and M_2 . We now define the concept C as a conjunction of several concepts. The first conjunct is

$$\neg C_0 \sqcap \exists r^{2^n}. \top \rightarrow R_0.$$

Intuitively, R_0 generates a binary r -tree of depth $2n$ with R_i true at level i for $0 \leq i \leq 2^{2n}$ and each leaf represents a grid position (i, j) using the concept names A_i . To achieve this let C contain the following conjuncts for generating the binary tree:

$$\begin{aligned} & \prod_{0 \leq i < 2n} \forall r^i. (R_i \rightarrow (\exists r. (A_i \sqcap R_{i+1}) \sqcap \exists r. (\neg A_i \sqcap R_{i+1}))) \\ & \prod_{1 \leq i < 2n} \prod_{0 \leq j < i} \forall r^i. ((A_j \rightarrow \forall r. A_j) \sqcap (\neg A_j \rightarrow \forall r. \neg A_j)) \end{aligned}$$

As usual, $\forall r^i$ abbreviates a sequence of i times $\forall r$.

We next express using additional conjuncts of C that any leaf d representing (i, j) using A_i has the following properties (A) – (C):

- (A) d has an r -successor representing (i, j) using B_i with a tile type $T_{(i,j)}$ true in it; moreover, no r -successor of d representing (i, j) satisfies a tile type different from $T_{(i,j)}$. This is achieved using the marker M which holds in exactly those r -successors of d that represent (i, j) using B_i . The latter condition is expressed using the counter X_i which represents (i, j) on all r -successors of d . In detail, we add the following conjuncts to C :

$$\begin{aligned}
& \forall r^{2n}. \exists r. M \\
& \forall r^{2n}. \left(\prod_{i < 2n} (A_i \rightarrow \forall r. X_i) \cap (\neg A_i \rightarrow \forall r. \neg X_i) \right) \\
& \forall r^{2n+1}. (M \leftrightarrow \prod_{i < 2n} (X_i \leftrightarrow B_i) \cap (\neg X_i \leftrightarrow \neg B_i)) \\
& \forall r^{2n}. (\forall r. (M \rightarrow \bigsqcup_{i \leq k} T_i) \cap \prod_{i \leq k} \exists r. (M \cap T_i) \rightarrow \forall r. (M \rightarrow T_i)) \\
& \forall r^{2n+1}. \prod_{i \neq j} \neg (T_i \cap T_j)
\end{aligned}$$

- (B) d has an r -successor representing $(i \oplus_{2^n} 1, j)$ using B_i with a tile type $T_{(i,j)}^{\text{right}}$ true in it such that $(T_{(i,j)}, T_{(i,j)}^{\text{right}}) \in H$; moreover, no r -successor of d representing $(i \oplus_{2^n} 1, j)$ satisfies a tile type different from $T_{(i,j)}^{\text{right}}$. This is achieved in a similar way as (A) using the marker M_1 which holds in exactly those r -successors of d that represent $(i \oplus_{2^n} 1, j)$ using B_i . The latter condition is expressed using the counter Y_i which represents $(i \oplus_{2^n} 1, j)$ on all r -successors of d . The implementation of these conditions is similar to (A) and omitted.
- (C) d has an r -successor representing $(i, j \oplus_{2^n} 1)$ using B_i with a tile type $T_{(i,j)}^{\text{up}}$ true in it such that $(T_{(i,j)}, T_{(i,j)}^{\text{up}}) \in V$; moreover, no r -successor of d representing $(i, j \oplus_{2^n} 1)$ satisfies a tile type different from $T_{(i,j)}^{\text{up}}$. This is achieved in a similar way as (A) using the marker M_2 which holds in exactly those r -successors of d that represent $(i, j \oplus_{2^n} 1)$ using B_i . The latter condition is expressed using the counter Z_i which represents $(i, j \oplus_{2^n} 1)$ on all r -successors of d . The implementation is again similar to (A) and omitted.

Finally, we ensure that the initial condition holds, that is $T_{(i,0)} = c_i$ for $i < n$. To this end we add the conjuncts

$$\forall r^{2n}. (A = (i, 0) \rightarrow (\forall r. (M \rightarrow c_i)))$$

for $i < n$, where $A = (i, 0)$ stands for the representation of $(i, 0)$ using A_i ; for instance, $A = (0, 0)$ stands for $\prod_{0 \leq i < 2^n} \neg A_i$.

This finishes the definition of C, C_0 and we verify next that they are as required.

Claim. There exist $\mathcal{I}_1, d_1 \sim_{\mathcal{ALCO}, \Sigma} \mathcal{I}_2, d_2$ with $d_1 \in (C \sqcap C_0)^{\mathcal{I}_1}$ and $d_2 \in (C \sqcap \neg C_0)^{\mathcal{I}_2}$ iff P has a solution given c .

Proof of the Claim. Observe that if $\mathcal{I}_1, d_1 \sim_{\mathcal{ALCO}, \Sigma} \mathcal{I}_2, d_2$ with $d_1 \in (C \sqcap C_0)^{\mathcal{I}_1}$ and $d_2 \in (C \sqcap \neg C_0)^{\mathcal{I}_2}$, then there are elements $e_{(i,j)}$, $0 \leq i, j \leq 2^n - 1$ such that

$$\mathcal{I}_1, a^{\mathcal{I}_1} \sim_{\mathcal{ALCO}, \Sigma} \mathcal{I}_2, e_{(i,j)}$$

and $e_{(i,j)}$ has (at least) three r -successors satisfying Conditions (A) to (C) and the initial condition. By Σ -bisimilarity and since $r \in \Sigma$, all $e_{(i,j)}$ have r -successors satisfying the same concept names in Σ . Hence, since the concept names B_i and T_i are in Σ , for every grid position (i, j) every $e_{(i',j')}$ has an r -successor representing (i, j) using B_i and all r -successors representing (i, j) using B_i satisfy the same tile type $T_{(i,j)}$. Moreover, $T_{(i \oplus_{2^n} 1, j)} = T_{(i,j)}^{\text{right}}$ and $T_{(i, j \oplus_{2^n} 1)} = T_{(i,j)}^{\text{up}}$. It follows that the mapping τ defined by setting $\tau(i, j) = T_{(i,j)}$ is a solution of P given c .

Conversely, assume that P and c have a solution τ . The definition of an interpretation \mathcal{I} with elements d_1 and d_2 such that $\mathcal{I}, d_1 \sim_{\mathcal{ALCO}, \Sigma} \mathcal{I}, d_2$ with $d_1 \in (C \sqcap C_0)^{\mathcal{I}}$ and $d_2 \in (C \sqcap \neg C_0)^{\mathcal{I}}$ is rather straightforward. An abstract version is depicted in Figure 10. We omit the counters, and note that $a^{\mathcal{I}}$ and *all* elements at level R_{2^n} have, for all $0 \leq i, j < 2^n$, an r -successor representing (using concept names B_i) grid position (i, j) which satisfies the concept name $T_{\tau(i,j)}$. We show only the three special successors from Conditions (A)–(C). This finishes the proof of the Claim and thus the reduction for \mathcal{ALCO} .

We come to the lower bound for \mathcal{ALCH} and \mathcal{ALCHI} . Let

$$\mathcal{O} = \{r \sqsubseteq r_1, r \sqsubseteq r_2, r_1 \sqsubseteq v, r_2 \sqsubseteq v\},$$

and Σ contains r_1, r_2 but not r nor v . In addition to r_1 and r_2 , Σ contains exactly the same concept names as in the \mathcal{ALCO} proof and we also use the same concept names not in Σ . We aim to construct concepts C, C_0 such that there exist models $\mathcal{I}_1, \mathcal{I}_2$ of \mathcal{O} and $d_1 \in (C \sqcap C_0)^{\mathcal{I}_1}$ and $d_2 \in (C \sqcap \neg C_0)^{\mathcal{I}_2}$ with $\mathcal{I}_1, d_1 \sim_{\mathcal{ALCH}, \Sigma} \mathcal{I}_2, d_2$ iff P has a solution given c .

We set $C_0 = \exists r^{2^n} . \top$. The concept C is again a conjunction of several concepts; we start in a similar way as for \mathcal{ALCO} with

$$\neg C_0 \sqcap \exists v^{2^n} . \top \rightarrow R_0$$




Fig. 10. Interpretation \mathcal{I} with elements $d_1 \in (C \sqcap C_0)^{\mathcal{I}}$ and $d_2 \in (C \sqcap \neg C_0)^{\mathcal{I}}$ such that $\mathcal{I}, d_1 \sim_{\mathcal{ALCO}, \pm} \mathcal{I}, d_2$.

The concept name R_0 will enforce that

- (**) the end point of any r_1/r_2 -path of length $2n$ starting in an element satisfying R_0 carries a pair of counter values (i, j) represented by concept names A_i which describe the path in a canonical way.²

To achieve this, we include the following conjuncts in C :

$$\prod_{0 \leq i < 2n} \forall v^i. (R_i \rightarrow \forall r_1. (A_i \sqcap R_{i+1}) \sqcap \forall r_2. (\neg A_i \sqcap R_{i+1}))$$

$$\prod_{1 \leq i < 2n} \prod_{0 \leq j < i} \forall v^i. ((A_j \rightarrow \forall v. A_j) \sqcap (\neg A_j \rightarrow \forall v. \neg A_j))$$

Note that we can use the role name v to address all elements reachable along r_1/r_2 -paths of length i via $\forall v^i$. We continue the definition of C in exactly the same way as for \mathcal{ALCO} except that we use $\forall v^{2n}$ to reach the end points of the paths mentioned in (**) and r_1 -successors of the leaves to encode a solution of the tiling problem. One can then easily prove the following.

Claim. There exist $\mathcal{I}_1, d_1 \sim_{\mathcal{ALCH}, \Sigma} \mathcal{I}_2, d_2$ with $\mathcal{I}_1, \mathcal{I}_2$ models of \mathcal{O} , $d_1 \in (C \sqcap \exists r^{2n}. \top)^{\mathcal{I}_1}$, and $d_2 \in (C \sqcap \neg \exists r^{2n}. \top)^{\mathcal{I}_2}$ iff P has a solution given c .

Proof of the Claim. Observe that if $\mathcal{I}_1, d_1 \sim_{\mathcal{ALCH}, \Sigma} \mathcal{I}_2, d_2$ with $\mathcal{I}_1, \mathcal{I}_2$ models of \mathcal{O} , $d_1 \in (C \sqcap \exists r^{2n}. \top)^{\mathcal{I}_1}$, and $d_2 \in (C \sqcap \neg \exists r^{2n}. \top)^{\mathcal{I}_2}$, then there exists an element e reachable from d_1 along an r -path of length $2n$ in \mathcal{I}_1 .

² Notice the similarity with Property (*) from the proof of Lemma 42.

Since $d_2 \in R_0^{\mathcal{I}_2}$ and e is reachable via arbitrary r_1/r_2 -paths of length $2n$ from d_1 , Property (**) implies that there are elements $e_{(i,j)}$, $0 \leq i, j \leq 2^n - 1$, reachable from d_2 along a v -path of length $2n$ in \mathcal{I}_2 such that $\mathcal{I}_1, e \sim_{\mathcal{ALCH}\Sigma} \mathcal{I}_2, e_{(i,j)}$ and $e_{i,j}$ represents the pair (i, j) using the concept names A_i . The remaining proof is now essentially the same as for \mathcal{ALCO} .

The converse direction is rather straightforward and similar to the proof for \mathcal{ALCO} . The difference is that the binary tree over role r in the right side of interpretation \mathcal{I} depicted in Figure 10 is now a binary tree over roles r_1 (left successor) and r_2 (right successor). This finishes the proof of the Claim.

To prove the claim above for \mathcal{ALCHI} , we adapt the model construction in a similar way as in the case with ontologies (Section 7.3). More precisely, for each element e at level $\ell > 0$ in the binary tree below d_2 , add $(d, e) \in r_1^{\mathcal{I}}$ and $(d, e) \in r_2^{\mathcal{I}}$, where d is the element in distance $\ell - 1$ from d_1 . One can then verify that \mathcal{I} is as required, that is, $\mathcal{I}, d_1 \sim_{\mathcal{ALCHI}} \mathcal{I}, d_2$, $d_1 \in (C \sqcap C_0)^{\mathcal{I}}$, and $d_2 \in (C \sqcap \neg C_0)^{\mathcal{I}}$.

10 Computation Problem

In the previous sections, we have presented algorithms for *deciding* the existence of interpolants and explicit definitions, but these algorithms (and their correctness proofs) do not give immediately rise to a way of *computing* interpolants and explicit definitions in case they exist. Intuitively, this is due to the fact that compactness is used in the proof of the model-theoretic characterization of interpolant and explicit definition existence in terms of joint consistency modulo bisimulations which was provided in Theorems 21 and 23, respectively. In this section, we address the computation problem for logics in DL_{nr} that do not admit nominals, by showing that we can actually compute interpolants in case they exist. We use *DAG representation* for the interpolants; recall that in DAG representation common sub-formulas are stored only once, and that thus DAG representation is more succinct than formula representation. Our approach is inspired by a recent note on a type elimination based computation of interpolants in modal logic [88] which was originally provided for the guarded fragment [15].

Theorem 48. *Let $\mathcal{L} \in DL_{nr}$ not admit nominals, \mathcal{O} be an \mathcal{L} -ontology, C_1, C_2 be \mathcal{L} -concepts, and Σ be a signature. Then, if there is an $\mathcal{L}(\Sigma)$ -interpolant for $C_1 \sqsubseteq C_2$ under \mathcal{O} , we can compute the DAG representation of an $\mathcal{L}(\Sigma)$ -interpolant in time $2^{2^{p(n)}}$ where p is a polynomial and $n = \|\mathcal{O}\| + \|C_1\| + \|C_2\|$.*

Note that this implies that the DAG representation is also of double exponential size, and that a formula representation of the interpolant can be computed in triple exponential time. Moreover, this also allows us to compute explicit definitions since, given \mathcal{O} , C , and Σ , any $\mathcal{L}(\Sigma)$ -interpolant for $C_\Sigma \sqsubseteq C$ under $\mathcal{O} \cup \mathcal{O}_\Sigma$ is an explicit $\mathcal{L}(\Sigma)$ -definition of C under \mathcal{O} , where \mathcal{O}_Σ and C_Σ are obtained from \mathcal{O} and C by replacing all symbols not in Σ by fresh symbols. We conjecture that the triple exponential upper bound on formula size is actually tight, given the discussion on the explicit definitions that arise in the hardness proofs in Sections 7.2 and 7.3.

Let \mathcal{L} , \mathcal{O} , C_1, C_2 , and Σ be as in Theorem 48. By Theorem 21, the existence of an $\mathcal{L}(\Sigma)$ -interpolant for $C_1 \sqsubseteq C_2$ under \mathcal{O} is equivalent to joint consistency of C_1 and $\neg C_2$ under \mathcal{O} modulo $\mathcal{L}(\Sigma)$ -bisimulations. Recall that we have provided before Lemma 35 in Section 6 a mosaic elimination procedure for deciding the latter. In fact, the computation of the $\mathcal{L}(\Sigma)$ -interpolant relies on a finer analysis of that procedure. We need one more notion to formalize this analysis.

Let T be a set of \mathcal{E} -types. Let \mathcal{I} be an interpretation and, for each $t \in T$, let d_t be a domain element of \mathcal{I} . We say that \mathcal{I} and the elements d_t , $t \in T$ *jointly realize T modulo $\mathcal{L}(\Sigma)$ -bisimulations* if, for all $t, t' \in T$, we have that $\text{tp}_\Sigma(\mathcal{I}, d_t) = t$ and $\mathcal{I}, d_t \sim_{\mathcal{L}, \Sigma} \mathcal{I}, d_{t'}$. We call T *jointly realizable under \mathcal{O} modulo $\mathcal{L}(\Sigma)$ -bisimulations* if there is a model \mathcal{I} of \mathcal{O} and elements d_t for each $t \in T$ that jointly realize T modulo $\mathcal{L}(\Sigma)$ -bisimulations. In contrast to the notion of joint consistency, we require here a single model \mathcal{I} of \mathcal{O} . In what follows, let Real denote the set of all sets of types T which are jointly realizable under \mathcal{O} modulo $\mathcal{L}(\Sigma)$ -bisimulations. We can effectively determine Real since joint realizability of a set T can be decided in double exponential time, similar to joint consistency—we refrain from giving details.

In the (proof of the) following lemma we show how to compute a concept differentiating between T_1 and T_2 when (T_1, T_2) is eliminated for $T_1, T_2 \in \text{Real}$. In Lemma 50 below, we show how to assemble these differentiating concepts to an interpolant (in case it exists).

Lemma 49. *Let $T_1, T_2 \in \text{Real}$. If (T_1, T_2) is eliminated in the mosaic elimination procedure, then we can compute an $\mathcal{L}(\Sigma)$ -concept I_{T_1, T_2} such that*

1. *for all models \mathcal{I} of \mathcal{O} and elements d_t , for each $t \in T_1$, that jointly realize T_1 modulo $\mathcal{L}(\Sigma)$ -bisimulations, $d_t \in I_{T_1, T_2}^\mathcal{I}$ for some (equivalently: all) $t \in T_1$;*

2. for all models \mathcal{I} of \mathcal{O} and elements d_t , for each $t \in T_2$, that jointly realize T_2 modulo $\mathcal{L}(\Sigma)$ -bisimulations, $d_t \notin I_{T_1, T_2}^{\mathcal{I}}$ for some (equivalently: all) $t \in T_2$.

Moreover, a DAG representation of I_{T_1, T_2} can be computed in time $2^{2p(n)}$ for some polynomial p and $n = \|\mathcal{O}\| + \|C_1\| + \|C_2\|$.

Proof. We compute the I_{T_1, T_2} inductively in the order in which the (T_1, T_2) got eliminated in the elimination procedure. We distinguish cases why (T_1, T_2) got eliminated.

Suppose first that (T_1, T_2) was eliminated because of (failing) Σ -concept name coherence. Since T_1, T_2 are both jointly realizable, there are the following two cases.

- (a) There is a concept name $A \in \Sigma$ such that $A \in t$ for all $t \in T_1$, but $A \notin t$, for all $t \in T_2$. Then $I_{T_1, T_2} = A$.
- (b) There is a concept name $A \in \Sigma$ such that $A \in t$ for all $t \in T_2$, but $A \notin t$, for all $t \in T_1$. Then $I_{T_1, T_2} = \neg A$.

Clearly, in both cases, I_{T_1, T_2} satisfies Points (1) and (2) of Lemma 49.

Now, suppose that (T_1, T_2) was eliminated due to (failing) existential saturation from \mathcal{S}_i during the elimination procedure. Since T_1, T_2 are both jointly realizable under \mathcal{O} , there are the following two cases.

- (a) There exist $t \in T_1$, $\exists r.C \in t$, and a Σ -role s with $\mathcal{O} \models r \sqsubseteq s$, such that there is no $(T'_1, T'_2) \in \mathcal{S}_i$ such that (i) $(T_1, T_2) \rightsquigarrow_s (T'_1, T'_2)$ and (ii) there is $t' \in T'_1$ with $C \in t'$ and $t \rightsquigarrow_{r, \mathcal{O}} t'$. Then, take

$$I_{T_1, T_2} = \exists s. \left(\bigsqcup_{\substack{T'_1 \in \text{Real}, \\ T_1 \rightsquigarrow_s T'_1, t \rightsquigarrow_{r, \mathcal{O}} t', C \in t' \in T'_1}} \prod_{\substack{T'_2 \in \text{Real}, \\ T_2 \rightsquigarrow_s T'_2}} I_{T'_1, T'_2} \right)$$

- (b) There exist $t \in T_2$, $\exists r.C \in t$, and a Σ -role s with $\mathcal{O} \models r \sqsubseteq s$, such that there is no $(T'_1, T'_2) \in \mathcal{S}$ such that (i) $(T_1, T_2) \rightsquigarrow_s (T'_1, T'_2)$ and (ii) there is $t' \in T'_2$ with $C \in t'$ and $t \rightsquigarrow_{r, \mathcal{O}} t'$. Then, take

$$I_{T_1, T_2} = \forall s. \left(\bigsqcup_{\substack{T'_1 \in \text{Real}, \\ T_1 \rightsquigarrow_s T'_1}} \prod_{\substack{T'_2 \in \text{Real}, \\ T_2 \rightsquigarrow_s T'_2, t \rightsquigarrow_{r, \mathcal{O}} t', C \in t' \in T'_2}} I_{T'_1, T'_2} \right)$$

We show Points (1) and (2) of the lemma for Case (a); Case (b) is dual. So suppose Case (a) applies and fix $t \in T_1, \exists r.C \in t$, and a Σ -role s witnessing that.

To show Point (1) of the lemma, let \mathcal{I} be a model of \mathcal{O} and fix d_{t_1} for each $t_1 \in T_1$ such that \mathcal{I} and the d_{t_1} jointly realize T_1 modulo $\mathcal{L}(\Sigma)$ -bisimulations. It suffices to show that $d_t \in I_{T_1, T_2}^{\mathcal{I}}$ for the type t that was fixed in the application of Case (a). Since d_t realizes t and $\exists r.C \in t$, there is some $e \in C^{\mathcal{I}}$ with $(d_t, e) \in r^{\mathcal{I}}$. Since $\mathcal{O} \models r \sqsubseteq s$, also $(d_t, e) \in s^{\mathcal{I}}$. Since the $d_{t_1}, t_1 \in T_1$ are mutually $\mathcal{L}(\Sigma)$ -bisimilar and s is a Σ -role, we find elements $e_{t_1}, t_1 \in T_1$ such that:

- $e_{t_1}, t_1 \in T_1$ are mutually $\mathcal{L}(\Sigma)$ -bisimilar,
- $(d_{t_1}, e_{t_1}) \in s^{\mathcal{I}}$, for all $t_1 \in T_1$,
- $e_t = e$.

Let

$$T'_1 = \{\text{tp}_{\Xi}(\mathcal{I}, e_{t_1}) \mid t_1 \in T_1\},$$

and let further $T'_2 \in \mathbf{Real}$ be arbitrary with $T_2 \rightsquigarrow_s T'_2$. By definition of T'_1 , we have $T'_1 \in \mathbf{Real}$ and $T_1 \rightsquigarrow_s T'_1$. Thus, (T'_1, T'_2) has been eliminated before (T_1, T_2) : otherwise, Case (a) would not apply to the fixed $t, \exists r.C, s$. By induction, we can conclude that $e = e_t \in I_{T'_1, T'_2}^{\mathcal{I}}$, and hence $d \in I_{T_1, T_2}^{\mathcal{I}}$.

To show Point (2) of the lemma, let \mathcal{I} be a model of \mathcal{O} and fix d_{t_2} for each $t_2 \in T_2$ such that \mathcal{I} and the d_{t_2} jointly realize T_2 modulo $\mathcal{L}(\Sigma)$ -bisimulations. Suppose, to the contrary of what has to be shown, that $d_{\hat{t}} \in I_{T_1, T_2}^{\mathcal{I}}$ for some $\hat{t} \in T_2$. Then, there is an e with $(d_{\hat{t}}, e) \in s^{\mathcal{I}}$ and a $T'_1 \in \mathbf{Real}$ with $T_1 \rightsquigarrow_s T'_1$ and a type $t'_1 \in T_1$ with $t \rightsquigarrow_{r, \mathcal{O}} t'_1$ and $C \in t'_1$ such that

(*) $e \in I_{T'_1, T}^{\mathcal{I}}$ for all $T \in \mathbf{Real}$ with $T_2 \rightsquigarrow_s T$.

Since \mathcal{I} and the elements $d_{t_2}, t_2 \in T_2$ jointly realize T_2 modulo $\mathcal{L}(\Sigma)$ -bisimulations and s is a Σ -role, there are elements $e_{t_2}, t_2 \in T_2$ such that:

- $e_{t_2}, t_2 \in T_2$ are mutually $\mathcal{L}(\Sigma)$ -bisimilar,
- $(d_{t_2}, e_{t_2}) \in s^{\mathcal{I}}$, for all $t_2 \in T_2$,
- $e_{\hat{t}} = e$.

Let

$$T'_2 = \{\text{tp}_{\Xi}(\mathcal{I}, e_{t_2}) \mid t_2 \in T_2\}.$$

By definition of T'_2 , we have $T'_2 \in \mathbf{Real}$ and $T_2 \rightsquigarrow_s T'_2$. Thus, (T'_1, T'_2) has been eliminated before (T_1, T_2) : otherwise, Case (a) would not apply to the fixed $t, \exists r.C, s$. By induction, we obtain $e = e_{\hat{t}} \notin I_{T'_1, T'_2}^{\mathcal{I}}$, in contradiction to (*).

For the analysis of the DAG representation, observe that we can use a single node for every I_{T_1, T_2} . Moreover, I_{T_1, T_2} looks as follows:

- If (T_1, T_2) was eliminated due to failing Σ -concept name coherence, I_{T_1, T_2} is a single concept name A or its negation $\neg A$.
- Otherwise, it is a node labeled with $\exists s$ (resp., $\forall s$), which has a single successor labeled with \sqcup . This successor has then at most double exponentially many successor nodes, each labeled with \sqcap and each having at most double exponentially many successor nodes I_{T_1, T_2} .

Overall, we obtain double exponentially many nodes in the DAG and the DAG can be constructed in double exponential time (both in $p(\|\mathcal{O}\| + \|C_1\| + \|C_2\|)$). \square

Lemma 50. *Suppose the result \mathcal{S}^* of the mosaic elimination procedure does not contain a pair $(T_1, T_2) \in \text{Real} \times \text{Real}$ such that $C_1 \in t_1$ and $\neg C_2 \in t_2$ for some types $t_1 \in T_1$ and $t_2 \in T_2$. Then,*

$$C = \bigsqcup_{\substack{T_1 \in \text{Real}: \\ \text{there is } t_1 \in T_1 \text{ with } C_1 \in t_1}} \bigsqcap_{\substack{T_2 \in \text{Real}: \\ \text{there is } t_2 \in T_2 \text{ with } \neg C_2 \in t_2}} I_{T_1, T_2}$$

is an $\mathcal{L}(\Sigma)$ -interpolant for $C_1 \sqsubseteq C_2$ under \mathcal{O} . Moreover, a DAG representation of C can be computed in time $2^{2^{p(n)}}$, for some polynomial p and $n = \|\mathcal{O}\| + \|C_1\| + \|C_2\|$.

Proof. We have to show that $\mathcal{O} \models C_1 \sqsubseteq C$ and $\mathcal{O} \models C \sqsubseteq C_2$.

For $\mathcal{O} \models C_1 \sqsubseteq C$, let \mathcal{I} be a model of \mathcal{O} and suppose $d \in C_1^{\mathcal{I}}$. Let $T_1 = \{\text{tp}_{\sqsubseteq}(\mathcal{I}, d)\}$ consist of the single type of d . Clearly, $T_1 \in \text{Real}$. Let $T_2 \in \text{Real}$ be arbitrary such that $\neg C_2 \in t$, for some $t \in T_2$. By assumption of Lemma 50, (T_1, T_2) got eliminated in the elimination procedure. Point (1) of Lemma 49 implies $d \in I_{T_1, T_2}^{\mathcal{I}}$. Hence, $d \in C^{\mathcal{I}}$.

For $\mathcal{O} \models C \sqsubseteq C_2$, let \mathcal{I} be a model of \mathcal{O} and let $d \in (-C_2)^{\mathcal{I}}$. Now, let $T_1 \in \text{Real}$ be arbitrary such that $C_1 \in t$ for some $t \in T_1$, and set $T_2 = \{\text{tp}_{\sqsubseteq}(\mathcal{I}, d)\}$. Clearly, $T_2 \in \text{Real}$. By assumption of Lemma 50, (T_1, T_2) got eliminated in the elimination procedure. Point (2) of Lemma 49 implies $d \notin I_{T_1, T_2}^{\mathcal{I}}$. Hence, $d \notin C^{\mathcal{I}}$.

For the analysis of the DAG representation of C , it suffices to recall that the DAG representations of the I_{T_1, T_2} provided in Lemma 49 can be computed in time $2^{2^{p(n)}}$, and to observe that C adds only one \sqcup node and at most double exponentially many \sqcap -nodes. \square

To conclude the section, we give some intuition as to why the proof of Theorem 48 cannot be easily adapted to logics from DL_{nr} that admit nominals. Recall that in any two interpretations $\mathcal{I}_1, \mathcal{I}_2$, every nominal a is realized (modulo bisimulation) in exactly one mosaic. We addressed this by starting the elimination procedure for all possible choices of mosaics realizing the nominals. More specifically, in the proof of Lemma 37, we showed there is an interpolant for $C_1 \sqsubseteq C_2$ under \mathcal{O} iff, for all maximal sets \mathcal{U} of mosaics that are good for nominals, the mosaic elimination procedure started with \mathcal{U} leads to an \mathcal{S}^* which does not satisfy Condition 2 of Lemma 36, which is akin to the precondition of Lemma 50 above. It is, however, unclear how to combine these different runs of the elimination procedure in proving analogues of Lemmas 49 and 50. An alternative approach might be to derive the interpolants from a suitably constrained proof of $\mathcal{O} \models C \sqsubseteq D$ in an appropriate proof system, see e.g. [81].

11 Some Consequences for Modal and Hybrid Logics

In this section we formulate a few consequences of our results in terms of modal and hybrid logics. We focus on interpolant existence and do not discuss the transfer of results on explicit definition existence as they can be obtained in a similar way. We consider the local consequence relation and formulate results for standard hybrid modal languages without the backward modality but with any combination of nominals, the @-operator, and the universal modal modality. We also briefly discuss the reformulation of description logics with role inclusions into modal logic with inclusion conditions on the accessibility relations. For detailed introductions to (hybrid) modal logics we refer the reader to [2, 4].

Let $\text{ML}_{@}^u$ denote the modal hybrid language constructed using the rule

$$\varphi, \psi \quad := \quad p \mid \top \mid i \mid \neg\varphi \mid \varphi \wedge \psi \mid \Box\varphi \mid @_i\varphi \mid \Box_u\varphi,$$

where p ranges over a countably infinite set of *propositional variables*, i ranges over a countably infinite set of *nominals*, \Box ranges over an infinite set of modal operators \Box_0, \dots , and \Box_u denotes the *universal modality*. The fragment of $\text{ML}_{@}^u$ without the universal modality is denoted $\text{ML}_{@}$, the fragment of $\text{ML}_{@}$ without the operators $@_i$ is denoted ML_n , and the fragment of ML_n without nominals is the standard language of polymodal logic and denoted ML . By ML_n^u we denote the fragment of $\text{ML}_{@}^u$ without the operators $@_i$ and by ML^u the extension of ML with the universal modality.

The signature $\text{sig}(\varphi)$ of a formula φ is the set of propositional variables, nominals, and modal operators (without the universal role) occurring in it.

The language $ML_{@}^u$ and its fragments are interpreted in *Kripke models* $\mathfrak{M} = (W, (R_i)_{i < \omega}, V)$ with W a nonempty set of *worlds*, $R_i \subseteq W \times W$ accessibility relations, and V a valuation such that $V(p) \subseteq W$ for every propositional variable p , and $V(i) \subseteq W$ a singleton for every nominal i . Then the truth relation $\mathfrak{M}, w \models \varphi$ between *pointed models* \mathfrak{M}, w with $w \in W$ and formulas φ is defined inductively as follows:

$$\begin{array}{ll}
\mathfrak{M}, w \models \top, & \\
\mathfrak{M}, w \models p & \text{iff } w \in V(p), \\
\mathfrak{M}, w \models i & \text{iff } V(i) = \{w\}, \\
\mathfrak{M}, w \models \neg\psi & \text{iff } \mathfrak{M}, w \not\models \psi, \\
\mathfrak{M}, w \models \psi \wedge \chi & \text{iff } \mathfrak{M}, w \models \psi \text{ and } \mathfrak{M}, w \models \chi, \\
\mathfrak{M}, w \models \Box_n \psi & \text{iff } \mathfrak{M}, v \models \psi, \text{ for every } v \in W \text{ such that} \\
& (w, v) \in R_n, \\
\mathfrak{M}, w \models @_i \psi & \text{iff } \mathfrak{M}, v \models \psi, \text{ for the unique element } v \in V(i), \\
\mathfrak{M}, w \models \Box_u \psi & \text{iff } \mathfrak{M}, v \models \psi, \text{ for every } v \in W.
\end{array}$$

We set $\mathfrak{M} \models \varphi$ if $\mathfrak{M}, w \models \varphi$ for all $w \in W$. Observe that the $@$ -operator can be defined using the universal modality as $@_i \varphi = \Box_u(i \rightarrow \varphi)$ and so ML_n^u and $ML_{@}^u$ have the same expressive power.

There are two natural notions of consequence studied in modal and hybrid logics, local and global entailment, which also give rise to different notions of interpolants. We focus here on local entailment and briefly discuss global entailment at the end of this section. We say that φ *locally entails* ψ , in symbols $\varphi \models_{loc} \psi$, if for all pointed models \mathfrak{M}, w , if $\mathfrak{M}, w \models \varphi$ then $\mathfrak{M}, w \models \psi$. We note that deciding \models_{loc} is PSPACE-complete for any of the languages introduced above without the universal modality and EXPTIME-complete for any of the languages introduced above with the universal modality [4].

We formulate the interpolant existence problems for hybrid modal logics in the expected way. Call a formula χ an *interpolant for* φ, ψ if $\text{sig}(\chi) \subseteq \text{sig}(\varphi) \cap \text{sig}(\psi)$, $\varphi \models_{loc} \chi$ and $\chi \models_{loc} \psi$.

Definition 51. *Let \mathcal{L} be any of the languages introduced above. Then the interpolant existence problem for \mathcal{L} is the problem to decide for any $\varphi, \psi \in \mathcal{L}$ whether there exists an interpolant for φ, ψ in \mathcal{L} .*

Observe that since ML and ML^u enjoy the Craig interpolation property (if $\varphi \models_{loc} \psi$ then an interpolant for φ, ψ exists [37]), the interpolant existence problem reduces to checking $\varphi \models_{loc} \psi$ and is PSPACE-complete for ML and EXPTIME-complete for ML^u . The following tight complexity bounds for their extensions with nominals and the @-operator are the main result of this section.

Theorem 52. *1. Let $\mathcal{L} \in \{ML_n, ML_{@}\}$. Then the interpolant existence problem for \mathcal{L} is CONEXPTIME-complete.
2. The interpolant existence problem for ML_n^u is 2EXPTIME-complete.
These results also hold if one considers the language with a single modal operator only.*

Proof. (1) Let \cdot^m be the obvious bijection between \mathcal{ALCO} -concepts and ML_n -formulas and denote by \cdot^d its inverse. Then $\models C \sqsubseteq D$ iff $C^m \models_{loc} D^m$ for any \mathcal{ALCO} -concepts C, D . Hence the following conditions are equivalent, for all formulas $\varphi, \psi \in ML_n$:

- there exists an interpolant for φ, ψ in ML_n ;
- there exists an $\mathcal{ALCO}(\Sigma)$ -interpolant for φ^d, ψ^d , where $\Sigma = \text{sig}(\varphi^d) \cap \text{sig}(\psi^d)$.

The CONEXPTIME-completeness for interpolant existence for ML_n now follows from Point 3 of Theorem 28. We now come to $ML_{@}$. We did not consider the operator @ for DLs as it does not play a large role in description logic research.³ Note, however, that \mathcal{ALCO} can be extended to the DL $\mathcal{ALCO}_{@}$ with @ in a straightforward way by setting $@_a C := \forall u.(\{a\} \rightarrow C)$. The expressive power of $\mathcal{ALCO}_{@}$ -concepts is characterized by $\mathcal{ALCO}_{@}(\Sigma)$ -bisimulations, where an $\mathcal{ALCO}(\Sigma)$ -bisimulation S between interpretations \mathcal{I} and \mathcal{J} is an $\mathcal{ALCO}_{@}(\Sigma)$ -bisimulation if $(a^{\mathcal{I}}, a^{\mathcal{J}}) \in S$ for any $a \in \Sigma$. Then one can prove Lemma 3 also for $\mathcal{ALCO}_{@}$. Next one can prove the characterization (Theorem 21) for $\mathcal{ALCO}_{@}$ in exactly the same way as for \mathcal{ALCO} , and finally one can extend the NEXPTIME-upper bound proof for joint consistency modulo $\mathcal{ALCO}(\Sigma)$ -bisimulations to joint consistency modulo $\mathcal{ALCO}_{@}(\Sigma)$ -bisimulations (Lemma 45) by observing that for all nominal generated mosaics $(T_1(d), T_2(d))$ we now have that $T_i(d) \neq \emptyset$ for $i = 1, 2$. Hence $(a^{\mathcal{I}}, a^{\mathcal{J}}) \in S$ for any $a \in \Sigma$, for the bisimulation S constructed in the proof of Lemma 45.

The lower bound proof for Theorem 28, Point 3, provided in Section 9 still goes through as it does not use any nominal in the shared signature

³ An exception is the investigation of updates for description logic knowledge bases where the expressive power of the @-operator plays a significant role [66].

and so using @ does not make any difference. Note, moreover, that it uses only a single role name r which corresponds to using a single modal operator.

(2) can be proved in the same way as (1) by observing that there is a bijection \cdot^m between \mathcal{ALCO}^u -concepts and ML_n^u -formulas, that $\models C \sqsubseteq D$ iff $C^m \models_{loc} D^m$ for any \mathcal{ALCO}^u -concepts C, D , and then applying Point 1 of Theorem 28. Note that the lower bound holds for a single role, see Lemma 41, which again translates to a single modal operator (and the universal modality). \square

Description logics with RIs correspond to modal logics determined by Kripke models satisfying inclusions $R_i \subseteq R_j$ between accessibility relations R_i and R_j . For any finite set I of pairs (i, j) let \mathcal{M}_I denote the class of Kripke models satisfying $R_i \subseteq R_j$ for all $(i, j) \in I$. Define the consequence relation \models_{loc}^I in the usual way by setting $\varphi \models_{loc}^I \psi$ if for all pointed models \mathfrak{M}, w with $\mathfrak{M} \in \mathcal{M}_I$, if $\mathfrak{M}, w \models \varphi$ then $\mathfrak{M}, w \models \psi$. We then obtain the following complexity result directly from Points 4 and 2 of Theorem 28, respectively.

Theorem 53. *For all finite I , the interpolant existence problem for \models_{loc}^I in ML is in CONEXPTIME. There exists a finite I such that the interpolant existence problem for \models_{loc}^I in ML is CONEXPTIME-hard.*

For all finite I , the interpolant existence problem for \models_{loc}^I in ML^u is in 2EXPTIME. There exists a finite I such that the interpolant existence problem for \models_{loc}^I in ML^u is 2EXPTIME-hard.

We close this section with a brief discussion of interpolant existence for the global consequence relation. We say that φ *globally entails* ψ , in symbols $\varphi \models_{glo} \psi$, if for all models \mathfrak{M} from $\mathfrak{M} \models \varphi$ it follows that $\mathfrak{M} \models \psi$. Call a formula χ a *global interpolant* for φ, ψ if $\text{sig}(\chi) \subseteq \text{sig}(\varphi) \cap \text{sig}(\psi)$, $\varphi \models_{glo} \chi$ and $\chi \models_{glo} \psi$. The *global interpolant existence problem* for \mathcal{L} is the problem to decide for any $\varphi, \psi \in \mathcal{L}$ whether there exists a global interpolant for φ, ψ in \mathcal{L} . It is straightforward to show that global interpolant existence corresponds to CI-interpolant existence in DLs in the same way as interpolant existence for the local consequence relation corresponds to ontology-free interpolant existence in DLs. We therefore obtain 2EXPTIME-completeness of global interpolant existence for the language ML_n^u from Theorem 31. We conjecture that the same result holds for global interpolant existence for ML_n and $ML_{@}$ but leave the proofs for future work.

12 Conclusion

We have investigated the problem of deciding the existence of interpolants and explicit definitions for description and modal logics with nominals and role inclusions, and we also presented an algorithm computing them for logics with role inclusions. There are many challenging problems left for future work, for instance, an algorithm computing interpolants for logics with nominals and the design and implementation of practical algorithms that could be applied in supervised concept learning and referring expression generation. From a theoretical viewpoint it would be of interest to gain a better understanding of when the existence of interpolants is computationally harder than entailment, for logics that do not enjoy the CIP. Logics to consider include more expressive DLs with nominals such as those also admitting qualified number restrictions and/or transitive roles and extensions of the two-variable fragment of FO with counting and/or further constraints on relations [53]. Another class of interest are decidable fragments of first-order modal logics and products of modal logics which both often do not enjoy the CIP [32, 73]. Here it would be of interest to consider logics such as the one-variable or monodic fragments of K and S4 for which the complexity of interpolant existence was left open [61]. Finally, is it possible to prove general transfer results (for example, for families of normal modal logics) stating that decidable entailment implies decidability of interpolant existence?

Acknowledgements

Frank Wolter was supported by EPSRC grant EP/S032207/1. Ana Ozaki was supported by NFR grant 316022. We thank Agi Kurucz, Michael Zakharyashev, and two referees for helpful comments.

A Proofs for Section 4

For the proofs of Theorems 12 and 29, we require a few definitions and observations. Given an interpretation $\mathcal{I} = (\Delta^{\mathcal{I}}, \cdot^{\mathcal{I}})$ and non-empty set $V \subseteq \Delta^{\mathcal{I}}$, we define the *restriction of \mathcal{I} to V* as $\mathcal{I}|_V = (\Delta^{\mathcal{I}|_V}, \cdot^{\mathcal{I}|_V})$, where $\Delta^{\mathcal{I}|_V} = \Delta^{\mathcal{I}} \cap V$, $B^{\mathcal{I}|_V} = B^{\mathcal{I}} \cap V$, for every concept name B , $r^{\mathcal{I}|_V} = r^{\mathcal{I}} \cap (V \times V)$, for every role name r , and $a^{\mathcal{I}|_V} = a^{\mathcal{I}}$ if $a^{\mathcal{I}} \in V$, for every individual name a . Note that $\mathcal{I}|_V$ does not interpret individual names that are not interpreted in V . Hence, in our constructions of restrictions, we always make sure that all relevant individuals are interpreted in V .

The *relativization* $C|_A$ of a concept C to a concept name A describes, in any interpretation \mathcal{I} , the extension of C in the restriction of \mathcal{I} to $A^{\mathcal{I}}$. In detail, define $C|_A$ inductively by setting $\top|_A = A$, $B|_A = B \sqcap A$, $(\neg C)|_A = A \sqcap \neg C|_A$, $(C \sqcap D)|_A = C|_A \sqcap D|_A$, $\{a\}|_A = \{a\} \sqcap A$, and $(\exists r.C)|_A = A \sqcap \exists r.C|_A$. Then the following observations can be shown by induction over the construction of C . For any interpretation \mathcal{I} with $A^{\mathcal{I}} \neq \emptyset$ and $A \notin \text{sig}(C)$ such that all $a \in \text{sig}(C)$ are interpreted in $A^{\mathcal{I}}$, $d \in C|_A^{\mathcal{I}}$ iff $d \in C^{\mathcal{I}|_{A^{\mathcal{I}}}}$, for all $d \in \Delta^{\mathcal{I}}$. Given an ontology \mathcal{O} , we set $\mathcal{O}|_A = \{C|_A \sqsubseteq D|_A \mid C \sqsubseteq D \in \mathcal{O}\}$. Then $\mathcal{I} \models \mathcal{O}|_A$ iff $\mathcal{I}|_{A^{\mathcal{I}}} \models \mathcal{O}$ whenever all $a \in \text{sig}(\mathcal{O})$ are interpreted in $A^{\mathcal{I}}$ and $A^{\mathcal{I}} \neq \emptyset$.

We next introduce a generalization of cartesian products called bisimulation products. Consider pointed models \mathcal{I}_1, d_1 and \mathcal{I}_2, d_2 with $\mathcal{I}_1, d_1 \sim_{\mathcal{ALCCO}, \Sigma} \mathcal{I}_2, d_2$. Take a bisimulation S witnessing this. Then the *bisimulation product* \mathcal{I} induced by S is defined as follows: the domain $\Delta^{\mathcal{I}}$ of \mathcal{I} is the set of all pairs $(e_1, e_2) \in S$. The concept and role names in Σ are interpreted as in cartesian products: $(e_1, e_2) \in B^{\mathcal{I}}$ iff $(e_1, e_2) \in S$ and $e_i \in B^{\mathcal{I}_i}$ for $i = 1, 2$, and $((e_1, e_2), (e'_1, e'_2)) \in r^{\mathcal{I}}$ iff $(e_1, e_2), (e'_1, e'_2) \in S$ and $(e_i, e'_i) \in r^{\mathcal{I}_i}$ for $i = 1, 2$. A nominal a in Σ is interpreted as $(a^{\mathcal{I}_1}, a^{\mathcal{I}_2})$ if a is in the domain (equivalently, range) of S . Note that we have projection functions $f_i : S \rightarrow \Delta^{\mathcal{I}_i}$ with $f_i(e_1, e_2) = e_i$, for $i = 1, 2$. We denote by $f_i(S)$ the image of S under f_i in $\Delta^{\mathcal{I}_i}$ and set $f_i^{-1}(V) := \{(e_1, e_2) \in S \mid e_i \in V\}$, for any $V \subseteq \Delta^{\mathcal{I}_i}$. The following lemma is straightforward.

Lemma 54. *Let $\mathcal{L} \in \{\mathcal{ALCCO}, \mathcal{ALCCIO}\}$. If S is an $\mathcal{L}(\Sigma)$ -bisimulation, then f_i is an $\mathcal{L}(\Sigma)$ -bisimulation between \mathcal{I} and \mathcal{I}_i , for $i = 1, 2$.*

A subset V of $\Delta^{\mathcal{I}}$ is called *closed in \mathcal{I}* if $e' \in V$ whenever $(e, e') \in r^{\mathcal{I}}$ and $e \in V$, where r is a role name. V is called *fully closed in \mathcal{I}* if it is closed in \mathcal{I} and, moreover, $e \in V$ whenever $(e, e') \in r^{\mathcal{I}}$ and $e' \in V$, where r is a role name. Observe that \mathcal{ALCCO} -concepts are preserved under relativization to closed subsets in the sense that $d \in C^{\mathcal{I}}$ iff $d \in C^{\mathcal{I}|_V}$ for any closed subset V of $\Delta^{\mathcal{I}}$, $d \in V$, and \mathcal{ALCCO} -concept C . It is easy to see that \mathcal{ALCCIO} -concepts are not always preserved under relativization to closed subsets. They are, however, preserved under relativization to fully closed subsets. Concepts using the universal role are not always preserved under relativization to fully closed subsets.

For a pointed interpretation \mathcal{I}, d denote by $\Delta_{\downarrow d}^{\mathcal{I}}$ the smallest subset of $\Delta^{\mathcal{I}}$ such that $d \in \Delta_{\downarrow d}^{\mathcal{I}}$ and $\Delta_{\downarrow d}^{\mathcal{I}}$ is closed in \mathcal{I} . The restriction of \mathcal{I} to $\Delta_{\downarrow d}^{\mathcal{I}}$ is denoted $\mathcal{I}_{\downarrow d}$ and called the *interpretation generated by d in \mathcal{I}* . It follows from our observation about closed subsets above that $e \in D^{\mathcal{I}}$ iff $e \in D^{\mathcal{I}_{\downarrow d}}$

holds for all $e \in \Delta_{\downarrow d}^{\mathcal{I}}$ and all \mathcal{ALCCO} -concepts D . We are in a position now to prove Theorem 12.

Theorem 12. *The following statements hold.*

- (1) *No $\mathcal{L} \in \mathcal{DL}_{nr}$ enjoys the CIP nor the PBDP. The CIP and PBDP also do not hold for RI-ontologies and, if \mathcal{L} admits nominals, the empty ontology.*
- (2) *All $\mathcal{L} \in \mathcal{DL}_{nr} \setminus \{\mathcal{ALCCO}, \mathcal{ALCHO}\}$ enjoy the BDP. \mathcal{ALCCO} and \mathcal{ALCHO} do not enjoy the BDP.*
- (3) *All $\mathcal{L} \in \mathcal{DL}_{nr}$ enjoy the BDP for RI-ontologies and the BDP for the empty ontology.*

Proof. Point (1) follows from the proofs of our complexity lower bounds for explicit definition existence. More specifically, in the proofs of the lower bounds we present concepts that are implicitly definable, but not explicitly definable.

For Point (2), assume first that $\mathcal{L} \in \mathcal{DL}_{nr} \setminus \{\mathcal{ALCCO}, \mathcal{ALCHO}\}$ does not admit nominals. Then the BDP follows essentially from Theorem 2.5.4 in [87], see also [89]. Theorem 2.5.4 is formulated in terms of modal logic. Hence observe that modal logics are syntactic variants of descriptions logics (see Section 11 for details) and that inverse roles, role inclusions, and the universal role can be introduced as first-order definable conditions on frame classes that are preserved under generated subframes and bisimulation products of frames.

Next assume that $\mathcal{L} \in \mathcal{DL}_{nr} \setminus \{\mathcal{ALCCO}, \mathcal{ALCHO}\}$ admits nominals and the universal role. Then the @-operator from hybrid logic can be expressed in \mathcal{L} (see again Section 11) and so the BDP follows from Theorem 6.2.4 in [87], see also [89].

It remains to prove that \mathcal{ALCCOI} and \mathcal{ALCHOI} enjoy the BDP. This is done using bisimulation products. We also use the characterization of the existence of explicit definitions using bisimulations provided in Theorem 23. Consider an \mathcal{ALCHOI} -ontology \mathcal{O} , let A be a concept name, and let C be an \mathcal{ALCHOI} -concept. Let $\Sigma = \text{sig}(\mathcal{O}, C) \setminus \{A\}$. Assume A is not explicitly $\mathcal{ALCHOI}(\Sigma)$ -definable under \mathcal{O} and C . By Theorem 23, we find pointed models \mathcal{I}_1, d_1 and \mathcal{I}_2, d_2 such that \mathcal{I}_i is a model of \mathcal{O} and $d_i \in C^{\mathcal{I}_i}$ for $i = 1, 2$, $d_1 \in A^{\mathcal{I}_1}$, $d_2 \notin A^{\mathcal{I}_2}$, and $\mathcal{I}_1, d_1 \sim_{\mathcal{ALCCOI}, \Sigma} \mathcal{I}_2, d_2$. Take a bisimulation S witnessing this. Let \mathcal{I} be defined as the bisimulation product induced by S . By Lemma 54, the projection functions $f_i : S \rightarrow \Delta^{\mathcal{I}_i}$ are $\mathcal{ALCCOI}(\Sigma)$ -bisimulations between \mathcal{I} and \mathcal{I}_i . Moreover, as we have inverse

roles, the image of S under f_i is fully closed in \mathcal{I}_i . We now define interpretations \mathcal{J}_1 and \mathcal{J}_2 as the interpretation \mathcal{I} except that $A^{\mathcal{J}_i} = f_i^{-1}(A^{\mathcal{I}_i})$, for $i = 1, 2$. Then the f_i are $\mathcal{ALCCOI}(\Sigma \cup \{A\})$ -bisimulations between \mathcal{J}_i and \mathcal{I}_i , for $i = 1, 2$. Note, however, that the \mathcal{J}_i do not necessarily interpret all nominals, as for a nominal $\{a\}$, the element $a^{\mathcal{J}_1}$ might be in a different connected component than d_1 (equivalently: $a^{\mathcal{J}_2}$ is in a different connected component than d_2). To address this, let \mathcal{I}' be the restriction of \mathcal{I}_1 to $\Delta^{\mathcal{I}_1} \setminus f_1(S)$. Then, obtain \mathcal{J}'_i by taking the disjoint union of \mathcal{J}_i and \mathcal{I}' , $i = 1, 2$. It follows from the construction, the preservation properties of \mathcal{ALCCOI} -bisimulations, and the fact that the universal role is not used in \mathcal{O} nor \mathcal{C} that the following conditions hold:

- (a) \mathcal{J}'_1 and \mathcal{J}'_2 are models of \mathcal{O} , $(d_1, d_2) \in C^{\mathcal{J}'_1}$, and $(d_1, d_2) \in C^{\mathcal{J}'_2}$;
- (b) the Σ -reducts of \mathcal{J}'_1 and \mathcal{J}'_2 coincide;
- (c) $(d_1, d_2) \in A^{\mathcal{J}'_1}$ but $(d_1, d_2) \notin A^{\mathcal{J}'_2}$.

But then A is not implicitly Σ -definable under \mathcal{O} and \mathcal{C} , as required.

Finally, to show Point (2), we have to argue that \mathcal{ALCCO} and \mathcal{ALCHCO} do not enjoy the BDP. A counterexample to the BDP is given in [89]. It illustrates nicely the way in which the addition of inverse roles or the universal role to \mathcal{ALCHCO} restores the BDP. As this is relevant in the proof of Point (3) and also in the analysis of non-projective definitions later in Section B, we give the example here. Let

$$\mathcal{O} = \{A \sqsubseteq \{a\}, \{b\} \sqcap B \sqsubseteq \exists r.(\{a\} \sqcap A), \{b\} \sqcap \neg B \sqsubseteq \exists r.(\{a\} \sqcap \neg A)\}$$

and set $\Sigma = \{a, B, b, r\} = \text{sig}(\mathcal{O}) \setminus \{A\}$. Then $\mathcal{O} \models A \equiv \{a\} \sqcap \exists r. (B \sqcap \{b\})$ and so A is explicitly $\mathcal{ALCCOI}(\Sigma)$ -definable under \mathcal{O} . Also $\mathcal{O} \models A \equiv \{a\} \sqcap \exists u. (B \sqcap \{b\})$, and so A is also explicitly $\mathcal{ALCCO}^u(\Sigma)$ -definable under \mathcal{O} . Note, however, that A is not explicitly $\mathcal{ALCCO}(\Sigma)$ -definable under \mathcal{O} . Indeed, the models \mathcal{I}, \mathcal{J} of \mathcal{O} depicted in Figure 11 (where $a^{\mathcal{I}} = a$ in \mathcal{I} and $a^{\mathcal{J}} = a$ in \mathcal{J} , and similarly for b) show that $\mathcal{I}, a^{\mathcal{I}} \sim_{\mathcal{ALCCO}, \Sigma} \mathcal{J}, a^{\mathcal{J}}$, with $a^{\mathcal{I}} \in A^{\mathcal{I}}$ and $a^{\mathcal{J}} \notin A^{\mathcal{J}}$. By Lemma 3, we have $\mathcal{I}, a^{\mathcal{I}} \equiv_{\mathcal{ALCCO}, \Sigma} \mathcal{J}, a^{\mathcal{J}}$, and hence there cannot be any $\mathcal{ALCCO}(\Sigma)$ -concept C such that $\mathcal{O} \models A \equiv C$.

It follows that non explicit $\mathcal{ALCCO}(\Sigma)$ -definability of A is caused by the fact that one cannot reach b from a along a path following the role name r . One can reach b , however, using the universal role or inverse roles.

For Point (3), it remains to show that \mathcal{ALCHCO} enjoys the BDP for ontologies containing RIs only. The BDP for \mathcal{ALCCO} with empty ontology follows immediately. The proof is similar to the proof of Point (2) above for \mathcal{ALCHIO} with fully closed subsets replaced by point generated interpretations.




Fig. 11. Models \mathcal{I}, \mathcal{J} of \mathcal{O} used to show that A is not explicitly $\mathcal{ALCCO}(\Sigma)$ -definable under \mathcal{O} .

Assume \mathcal{O} contains RIs only, C is an \mathcal{ALCCO} -concept, and A is a concept name. Let $\Sigma = \text{sig}(\mathcal{O}, C) \setminus \{A\}$. Assume A is not explicitly $\mathcal{ALCCO}(\Sigma)$ -definable under \mathcal{O} and C . By Theorem 23, we find pointed models \mathcal{I}_1, d_1 and \mathcal{I}_2, d_2 such that \mathcal{I}_i is a model of \mathcal{O} and $d_i \in C^{\mathcal{I}_i}$ for $i = 1, 2$, $d_1 \in A^{\mathcal{I}_1}$, $d_2 \notin A^{\mathcal{I}_2}$, and $\mathcal{I}_1, d_1 \sim_{\mathcal{ALCH}\mathcal{O}, \Sigma} \mathcal{I}_2, d_2$. Take a bisimulation S witnessing this. As we do not have the universal role nor inverse roles, we may assume that S is a $\mathcal{ALCCO}(\Sigma)$ -bisimulation between the set $\Delta_{\downarrow d_1}^{\mathcal{I}_1}$ generated by d_1 in \mathcal{I}_1 and the set $\Delta_{\downarrow d_2}^{\mathcal{I}_2}$ generated by d_2 in \mathcal{I}_2 . Let \mathcal{I} be defined as the bisimulation product induced by S . By Lemma 54, the projection functions $f_i : S \rightarrow \Delta^{\mathcal{I}_i}$ are $\mathcal{ALCCO}(\Sigma)$ -bisimulations between \mathcal{I} and \mathcal{I}_i . Let \mathcal{J}_1 and \mathcal{J}_2 be again defined as the interpretation \mathcal{I} except that $A^{\mathcal{J}_i} = f_i^{-1}(A^{\mathcal{I}_i})$, for $i = 1, 2$. Then the f_i are $\mathcal{ALCCO}(\Sigma \cup \{A\})$ -bisimulations between \mathcal{J}_i and \mathcal{I}_i , for $i = 1, 2$. As in the proof above, \mathcal{I} does not necessarily interpret all nominals in C . As \mathcal{O} is an ontology using RIs only, this problem can be addressed in a straightforward manner. Define interpretations \mathcal{J}'_i as the disjoint union of \mathcal{J}_i and the singleton interpretation \mathcal{I}' with domain $\{d\}$ such that $a^{\mathcal{J}'_i} = d$ for all a not interpreted in \mathcal{I} and $B^{\mathcal{I}'} = r^{\mathcal{I}'} = \emptyset$ for all concept and role names B and r . Then \mathcal{J}'_1 and \mathcal{J}'_2 satisfy the conditions (a) to (c) above and show that A is not implicitly Σ -definable under \mathcal{O} and C . \square

B Proof of Theorem 29

We show Theorem 29 which states that for $\mathcal{L} \in \{\mathcal{ALCCO}, \mathcal{ALCH}\mathcal{O}\}$ non-projective \mathcal{L} -definition existence of concept names is EXPTIME-complete. The lower bound follows from the corresponding lower bound for subsumption and any upper bound for $\mathcal{ALCH}\mathcal{O}$ trivially implies the same upper bound for \mathcal{ALCCO} . We therefore focus on the upper bound for $\mathcal{ALCH}\mathcal{O}$ and show the following criterion for (the complement of) non-projective explicit definability of concept names.

Lemma 55. *Let \mathcal{O} be an $\mathcal{ALCH}\mathcal{O}$ -ontology, C an \mathcal{ALCCO} -concept, and A a concept name. Let $\Sigma = \text{sig}(\mathcal{O}, C) \setminus \{A\}$. Then A is not explicitly*

$\mathcal{ALCHO}(\Sigma)$ -definable under \mathcal{O} and C iff there are pointed interpretations \mathcal{I}_1, d and \mathcal{I}_2, d such that

- \mathcal{I}_i is a model of \mathcal{O} and $d \in C^{\mathcal{I}_i}$, for $i = 1, 2$;
- the Σ -reducts of $\mathcal{I}_{1 \downarrow d}$ and $\mathcal{I}_{2 \downarrow d}$ coincide;
- $d \in A^{\mathcal{I}_1}$ and $d \notin A^{\mathcal{I}_2}$.

Proof. Clearly, if the conditions of Lemma 55 hold, then A is not explicitly $\mathcal{ALCHO}(\Sigma)$ -definable under \mathcal{O} and C , by Theorem 23. Conversely, assume A is not explicitly $\mathcal{ALCHO}(\Sigma)$ -definable under \mathcal{O} and C . By Theorem 23, we find pointed models \mathcal{I}_1, d_1 and \mathcal{I}_2, d_2 such that \mathcal{I}_i is a model of \mathcal{O} and $d_i \in C^{\mathcal{I}_i}$ for $i = 1, 2$, $d_1 \in A^{\mathcal{I}_1}$, $d_2 \notin A^{\mathcal{I}_2}$, and $\mathcal{I}_1, d_1 \sim_{\mathcal{ALCHO}, \Sigma} \mathcal{I}_2, d_2$. Take a bisimulation S witnessing this. As we do not admit the universal role nor inverse roles, we may assume that S is a bisimulation between the set $\Delta_{\downarrow d_1}^{\mathcal{I}_1}$ generated by d_1 in \mathcal{I}_1 and the set $\Delta_{\downarrow d_2}^{\mathcal{I}_2}$ generated by d_2 in \mathcal{I}_2 . Let \mathcal{I} be the bisimulation product induced by S . By Lemma 54, the projection functions $f_i : S \rightarrow \Delta^{\mathcal{I}_i}$ are $\mathcal{ALCO}(\Sigma)$ -bisimulations between \mathcal{I} and \mathcal{I}_i . However, as in the proof of Theorem 12, \mathcal{I} does not necessarily interpret all nominals. We address this in the following. Let \mathcal{J}_i be the restriction of \mathcal{I}_i to $\Delta^{\mathcal{I}_i} \setminus \Delta_{\downarrow d_i}^{\mathcal{I}_i}$, for $i = 1, 2$. We now define interpretations \mathcal{J}'_1 and \mathcal{J}'_2 as follows: \mathcal{J}'_i is the disjoint union of \mathcal{I} and \mathcal{J}_i extended by

- adding to the interpretation of A all elements in $f_i^{-1}(A^{\mathcal{I}_i})$;
- adding $(e, (e_1, e_2))$ with $(e_1, e_2) \in S$ to the interpretation of a role name r if $e \in \Delta^{\mathcal{I}_i} \setminus \Delta_{\downarrow d_i}^{\mathcal{I}_i}$, $e_i \in \Delta_{\downarrow d_i}^{\mathcal{I}_i}$, and $(e, e_i) \in r^{\mathcal{I}_i}$.

Using the condition that \mathcal{O} and C do not use the universal role nor inverse roles, one can show that the interpretations $\mathcal{I}_i := \mathcal{J}'_i$ and $d := (d_1, d_2)$ satisfy the conditions of Lemma 55. \square

We now show that the conditions of Lemma 55 can be checked in EXP-TIME by providing a polynomial time reduction to checking non \mathcal{ALCHO} -subsumption. Let \mathcal{O} be an \mathcal{ALCHO} -ontology, C an \mathcal{ALCO} -concept, A a concept name, and $\Sigma = \text{sig}(\mathcal{O}, C) \setminus \{A\}$. Take

- a concept name D for the domain $\Delta^{\mathcal{I}_1 \downarrow d}$ of the interpretation $\mathcal{I}_{1 \downarrow d}$ generated by d ;
- concept names D_i for the domain $\Delta^{\mathcal{I}_i}$ of \mathcal{I}_i , $i = 1, 2$;
- a copy A' of A ;
- and copies a' of the individual names a in Σ .

We let \mathcal{O}^c denote the set of CIs in \mathcal{O} and \mathcal{O}^r denote the set of RIs in \mathcal{O} . Let \mathcal{O}^{cc} be the ontology obtained from \mathcal{O}^c by replacing A by A' and all

nominals a in \mathcal{O}^c by a' . Let $\mathcal{O}_{|D_1}^c$ be the relativization of \mathcal{O}^c to D_1 and let $\mathcal{O}_{|D_2}^{cc}$ be the relativization of \mathcal{O}^{cc} to D_2 . Now consider the following ontology encoding Points 1 and 2 of Lemma 55:

$$\begin{aligned} \mathcal{O}' = & \mathcal{O}^r \cup \mathcal{O}_{|D_1}^c \cup \mathcal{O}_{|D_2}^{cc} \cup \{D \sqsubseteq \forall r.D \mid r \in \Sigma\} \cup \{D \sqsubseteq D_1, D \sqsubseteq D_2\} \cup \\ & \{\{a\} \sqsubseteq D_1 \mid a \in \Sigma\} \cup \{\{a'\} \sqsubseteq D_2 \mid a \in \Sigma\} \cup \\ & \{D \sqcap \{a\} \sqsubseteq \{a'\} \mid a \in \Sigma\} \cup \{D \sqcap \{a'\} \sqsubseteq \{a\} \mid a \in \Sigma\} \end{aligned}$$

Observe that we have to treat the individual names in Σ differently from the concept names in Σ as we have to ensure that they are interpreted in D_1 and D_2 respectively. As their interpretation might be different outside D , we have to introduce copies of the individual names and then state that those that are interpreted in D are actually interpreted in the same way. It is now straightforward to show that the conditions of Lemma 55 hold iff $\mathcal{O}' \not\models A \sqcap D \sqsubseteq A'$.

References

1. Carlos Areces, Patrick Blackburn, and Maarten Marx. A road-map on complexity for hybrid logics. In *Proceedings of the 8th Annual Conference of the European Association for Computer Science Logic, CSL 1999*, pages 307–321. Springer, 1999.
2. Carlos Areces, Patrick Blackburn, and Maarten Marx. Hybrid logics: Characterization, interpolation and complexity. *J. Symb. Log.*, 66(3):977–1010, 2001.
3. Carlos Areces, Alexander Koller, and Kristina Striegnitz. Referring expressions as formulas of description logic. In *Proceedings of the 5th International Natural Language Generation Conference, INLG 2008*. The Association for Computer Linguistics, 2008.
4. Carlos Areces and Balder ten Cate. Hybrid logics. In *Handbook of Modal Logic*, volume 3, pages 821–868. Elsevier, 2007.
5. Alessandro Artale, Jean Christoph Jung, Andrea Mazzullo, Ana Ozaki, and Frank Wolter. Living without beth and craig: Explicit definitions and interpolants in description logics with nominals (extended abstract). In *Proceedings of the 33rd International Workshop on Description Logics (DL 2020) co-located with the 17th International Conference on Principles of Knowledge Representation and Reasoning (KR 2020), Online Event [Rhodes, Greece], September 12th to 14th, 2020*, 2020.
6. Alessandro Artale, Jean Christoph Jung, Andrea Mazzullo, Ana Ozaki, and Frank Wolter. Living without beth and craig: Definitions and interpolants in description logics with nominals and role inclusions. In *Proceedings of the 35th AAAI Conference on Artificial Intelligence, AAAI 2021*, pages 6193–6201. AAAI Press, 2021.
7. Alessandro Artale, Andrea Mazzullo, Ana Ozaki, and Frank Wolter. On free description logics with definite descriptions. In *Proceedings of the 18th International Conference on Principles of Knowledge Representation and Reasoning, KR 2021*, pages 63–73, 2021.
8. Franz Baader, Diego Calvanese, Deborah L. McGuinness, Daniele Nardi, and Peter F. Patel-Schneider, editors. *The Description Logic Handbook: Theory, Implementation, and Applications*. Cambridge University Press, 2003.

9. Franz Baader, Ian Horrocks, Carsten Lutz, and Ulrike Sattler. *An Introduction to Description Logic*. Cambridge University Press, 2017.
10. Vince Bárány, Michael Benedikt, and Balder ten Cate. Rewriting guarded negation queries. In *Proceedings of the 38th International Symposium on Mathematical Foundations of Computer Science, MFCS 2013*, pages 98–110. Springer, 2013.
11. Vince Bárány, Michael Benedikt, and Balder ten Cate. Some model theory of guarded negation. *J. Symb. Log.*, 83(4):1307–1344, 2018.
12. Michael Benedikt, Pierre Bourhis, and Michael Vanden Boom. Definability and interpolation within decidable fixpoint logics. *Log. Methods Comput. Sci.*, 15(3), 2019.
13. Michael Benedikt, Julien Leblay, Balder ten Cate, and Efthymia Tsamoura. *Generating Plans from Proofs: The Interpolation-based Approach to Query Reformulation*. Synthesis Lectures on Data Management. Morgan & Claypool Publishers, 2016.
14. Michael Benedikt, Balder ten Cate, and Michael Vanden Boom. Interpolation with decidable fixpoint logics. In *Proceedings of the 30th Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2015*, pages 378–389. IEEE Computer Society, 2015.
15. Michael Benedikt, Balder ten Cate, and Michael Vanden Boom. Effective interpolation and preservation in guarded logics. *ACM Trans. Comput. Log.*, 17(2):8:1–8:46, 2016.
16. Evert Willem Beth. On Padoa’s Method in the Theory of Definition. *J. Symb. Log.*, 21(2):194–195, 1956.
17. Alexander Borgida, David Toman, and Grant E. Weddell. On referring expressions in query answering over first order knowledge bases. In *Proceedings of the 15th International Conference on Principles of Knowledge Representation and Reasoning, KR 2016*, pages 319–328. AAAI Press, 2016.
18. Alexander Borgida, David Toman, and Grant E. Weddell. Concerning referring expressions in query answers. In *Proceedings of the 26th International Joint Conference on Artificial Intelligence, IJCAI 2017*, pages 4791–4795. ijcai.org, 2017.
19. Lorenz Bühmann, Jens Lehmann, and Patrick Westphal. DL-learner - A framework for inductive learning on the semantic web. *J. Web Sem.*, 39:15–24, 2016.
20. Lorenz Bühmann, Jens Lehmann, Patrick Westphal, and Simon Bin. DL-Learner Structured Machine Learning on Semantic Web Data. In *Companion Proceedings of The Web Conference 2018, WWW 2018*, pages 467–471. ACM, 2018.
21. Diego Calvanese, Silvio Ghilardi, Alessandro Gianola, Marco Montali, and Andrey Rivkin. Combined covers and beth definability. In *Proceedings of the 10th International Joint Conference on Automated Reasoning, Part I, IJCAR 2020*, pages 181–200. Springer, 2020.
22. Diego Calvanese, Silvio Ghilardi, Alessandro Gianola, Marco Montali, and Andrey Rivkin. Combination of uniform interpolants via beth definability. *J. Autom. Reason.*, 66(3):409–435, 2022.
23. Ashok K. Chandra, Dexter C. Kozen, and Larry J. Stockmeyer. Alternation. *J. ACM*, 28:114–133, 1981.
24. C.C. Chang and H. Jerome Keisler. *Model Theory*. Elsevier, 1998.
25. Alessandro Cimatti, Alberto Griggio, and Roberto Sebastiani. Interpolant generation for UTVPI. In *Proceedings of the 22nd International Conference on Automated Deduction, CADE 2022*, pages 167–182. Springer, 2009.
26. S. D. Comer. Classes without the amalgamation property. *Pacific J. Math.*, 28(2):309–318, 1969.

27. William Craig. Three uses of the herbrand-gentzen theorem in relating model theory and proof theory. *J. Symb. Log.*, 22(3):269–285, 1957.
28. Giovanna D’Agostino and Marco Hollenberg. Uniform interpolation, automata and the modal μ -calculus. In *Proceedings of the 1st Workshop on Advances in Modal Logic, AiML-1*, pages 73–84. CSLI Publications, 1996.
29. Razvan Diaconescu. Logical support for modularisation. *Logical environments*, 83:130, 1993.
30. Nicola Fanizzi, Claudia d’Amato, and Floriana Esposito. DL-FOIL concept learning in description logics. In *Proceedings of the 18th International Conference on Inductive Logic Programming, ILP 2008*, pages 107–121. Springer, 2008.
31. Nicola Fanizzi, Giuseppe Rizzo, Claudia d’Amato, and Floriana Esposito. Dlfoil: Class expression learning revisited. In *Proceedings of the 21st International Conference on Knowledge Engineering and Knowledge Management, EKAW 2018*, pages 98–113. Springer, 2018.
32. Kit Fine. Failures of the interpolation lemma in quantified modal logic. *J. Symb. Log.*, 44(2):201–206, 1979.
33. Marie Fortin, Boris Konev, and Frank Wolter. Interpolants and explicit definitions in extensions of the description logic \mathcal{EL} . In *Proceedings of the 19th International Conference on Principles of Knowledge Representation and Reasoning, KR 2022*, 2022.
34. Enrico Franconi and Volha Kerhet. Effective query answering with ontologies and dboxes. In *Description Logic, Theory Combination, and All That - Essays Dedicated to Franz Baader on the Occasion of His 60th Birthday*, pages 301–328. Springer, 2019.
35. Enrico Franconi, Volha Kerhet, and Nhung Ngo. Exact query reformulation over databases with first-order and description logics ontologies. *J. Artif. Intell. Res.*, 48:885–922, 2013.
36. Maurice Funk, Jean Christoph Jung, Carsten Lutz, Hadrien Pulcini, and Frank Wolter. Learning description logic concepts: When can positive and negative examples be separated? In *Proceedings of the 28th International Joint Conference on Artificial Intelligence, IJCAI 2019*, pages 1682–1688, 2019.
37. Dov M. Gabbay. Craig’s interpolation theorem for modal logics. In *Conference in Mathematical Logic - London 1970*, pages 111–127. Springer, 1972.
38. Dov M Gabbay, Agi Kurucz, Frank Wolter, and Michael Zakharyashev. Many-dimensional modal logics: theory and applications. 2003.
39. Amit Goel, Sava Krstic, and Cesare Tinelli. Ground interpolation for combined theories. In *Proceedings of the 22nd International Conference on Automated Deduction, CADE 2022*, pages 183–198. Springer, 2009.
40. Valentin Goranko and Martin Otto. Model theory of modal logic. In *Handbook of Modal Logic*, pages 249–329. Elsevier, 2007.
41. K. Henkell. Pointlike sets: the finest aperiodic cover of a finite semigroup. *J. Pure Appl. Algebra*, 55(1-2):85–126, 1988.
42. K. Henkell, J. Rhodes, , and B. Steinberg. Aperiodic pointlikes and beyond. *Internat. J. Algebra Comput.*, 20(2):287–305, 2010.
43. Eva Hoogland and Maarten Marx. Interpolation and definability in guarded fragments. *Studia Logica*, 70(3):373–409, 2002.
44. Eva Hoogland, Maarten Marx, and Martin Otto. Beth definability for the guarded fragment. In *Proceedings of the 6th International Conference on Logic Programming and Automated Reasoning, LPAR 1999*, pages 273–285. Springer, 1999.

45. Luigi Iannone, Ignazio Palmisano, and Nicola Fanizzi. An algorithm based on counterfactuals for concept learning in the semantic web. *Appl. Intell.*, 26(2):139–159, 2007.
46. Rosalie Iemhoff. Uniform interpolation and the existence of sequent calculi. *Annals of Pure and Applied Logic*, 170(11):102711, 2019.
47. Ernesto Jiménez-Ruiz, Terry R. Payne, Alessandro Solimando, and Valentina A. M. Tamma. Limiting logical violations in ontology alignment through negotiation. In *Proceedings of the 15th International Conference on Principles of Knowledge Representation and Reasoning, KR 2016*, pages 217–226, 2016.
48. Jean Christoph Jung, Carsten Lutz, Hadrien Pulcini, and Frank Wolter. Logical separability of incomplete data under ontologies. In *Proceedings of the 17th International Conference on Principles of Knowledge Representation and Reasoning, KR 2020*, pages 517–528, 2020.
49. Jean Christoph Jung, Carsten Lutz, Hadrien Pulcini, and Frank Wolter. Separating data examples by description logic concepts with restricted signatures. In *Proceedings of the 18th International Conference on Principles of Knowledge Representation and Reasoning, KR 2021*, pages 390–399, 2021.
50. Jean Christoph Jung, Carsten Lutz, Hadrien Pulcini, and Frank Wolter. Logical separability of labeled data examples under ontologies. *Artif. Intell.*, 313:103785, 2022.
51. Jean Christoph Jung, Andrea Mazzullo, and Frank Wolter. More on interpolants and explicit definitions for description logics with nominals and/or role inclusions. In *Proceedings of the 35th International Workshop on Description Logics, DL 2022*, 2022.
52. Jean Christoph Jung and Frank Wolter. Living without beth and craig: Definitions and interpolants in the guarded and two-variable fragments. In *Proceedings of the 36th Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2021*, pages 1–14. IEEE, 2021.
53. Emanuel Kieronski, Ian Pratt-Hartmann, and Lidia Tendera. Two-variable logics with counting and semantic constraints. *ACM SIGLOG News*, 5(3):22–43, 2018.
54. Boris Konev, Carsten Lutz, Denis K. Ponomaryov, and Frank Wolter. Decomposing description logic ontologies. In *Proceedings of the 12th International Conference on Principles of Knowledge Representation and Reasoning, KR 2010*. AAAI Press, 2010.
55. Boris Konev, Carsten Lutz, Dirk Walther, and Frank Wolter. Formal properties of modularisation. In *Modular Ontologies: Concepts, Theories and Techniques for Knowledge Modularization*, pages 25–66. Springer, 2009.
56. Boris Konev, Dirk Walther, and Frank Wolter. Forgetting and uniform interpolation in large-scale description logic terminologies. In *Proceedings of the 21st International Joint Conference on Artificial Intelligence, IJCAI 2009*, pages 830–835, 2009.
57. Patrick Koopmann and Renate A. Schmidt. Count and forget: Uniform interpolation of \mathcal{SHQ} -ontologies. In *Proceedings of the 7th International Joint Conference on Automated Reasoning, IJCAR 2014*, pages 434–448. Springer, 2014.
58. Patrick Koopmann and Renate A. Schmidt. Uniform interpolation and forgetting for \mathcal{ALC} ontologies with aboxes. In *Proceedings of the 29th AAAI Conference on Artificial Intelligence, AAAI 2015*, pages 175–181. AAAI Press, 2015.
59. Tomasz Kowalski and George Metcalfe. Uniform interpolation and coherence. *Annals of Pure and Applied Logic*, 170(7):825–841, 2019.

60. Emiel Kraahmer and Kees van Deemter. Computational generation of referring expressions: A survey. *Computational Linguistics*, 38(1):173–218, 2012.
61. Agi Kurucz, Frank Wolter, and Michael Zakharyashev. Definitions and (uniform) interpolants in first-order modal logic. *CoRR*, abs/2303.04598, 2023.
62. Jens Lehmann and Christoph Haase. Ideal downward refinement in the \mathcal{EL} description logic. In *Proceedings of the 19th International Conference on Inductive Logic Programming, ILP 2009*, pages 73–87. Springer, 2009.
63. Jens Lehmann and Pascal Hitzler. Concept learning in description logics using refinement operators. *Machine Learning*, 78:203–250, 2010.
64. Francesca A. Lisi. A formal characterization of concept learning in description logics. In *Proceedings of the 25th International Workshop on Description Logics, DL 2012*. CEUR-WS.org, 2012.
65. Francesca A. Lisi and Umberto Straccia. Learning in description logics with fuzzy concrete domains. *Fundamenta Informaticae*, 140(3-4):373–391, 2015.
66. Hongkai Liu, Carsten Lutz, Maja Milicic, and Frank Wolter. Foundations of instance level updates in expressive description logics. *Artif. Intell.*, 175(18):2170–2197, 2011.
67. Carsten Lutz, Robert Piro, and Frank Wolter. Description logic tboxes: Model-theoretic characterizations and rewritability. In *Proceedings of the 22nd International Joint Conference on Artificial Intelligence, IJCAI 2011*, pages 983–988. IJCAI/AAAI, 2011.
68. Carsten Lutz, Inanç Seylan, and Frank Wolter. An automata-theoretic approach to uniform interpolation and approximation in the description logic EL. In *Proceedings of the 13th International Conference on Principles of Knowledge Representation and Reasoning, KR 2012*. AAAI Press, 2012.
69. Carsten Lutz, Inanç Seylan, and Frank Wolter. The data complexity of ontology-mediated queries with closed predicates. *Logical Methods in Computer Science*, 15(3), 2019.
70. Carsten Lutz and Frank Wolter. Foundations for uniform interpolation and forgetting in expressive description logics. In *Proceedings of the 22nd International Joint Conference on Artificial Intelligence, IJCAI 2011*, pages 989–995. IJCAI/AAAI, 2011.
71. Larisa Maksimova and Dov Gabbay. *Interpolation and Definability in Modal and Intuitionistic Logics*. Clarendon Press, 2005.
72. Maarten Marx. Interpolation in modal logic. In *Proceedings of the 7th International Conference on Algebraic Methodology and Software Technology, AMAST 1998*, pages 154–163. Springer, 1998.
73. Maarten Marx and Carlos Areces. Failure of interpolation in combined modal logics. *Notre Dame J. Formal Log.*, 39(2):253–273, 1998.
74. Kenneth L. McMillan. Interpolation and sat-based model checking. In *Proceedings of the 15th International Conference on Computer Aided Verification, CAV 2003*, pages 1–13. Springer, 2003.
75. Nadeschda Nikitina and Sebastian Rudolph. (Non-)succinctness of uniform interpolants of general terminologies in the description logic \mathcal{EL} . *Artif. Intell.*, 215:120–140, 2014.
76. D. Pigozzi. Amalgamation, congruence-extension, and interpolation properties in algebras. *Algebra Univers.*, (1):269–349, 1971.
77. Andrew M. Pitts. On an interpretation of second order quantification in first order intuitionistic propositional logic. *J. Symb. Log.*, 57(1):33–52, 1992.

78. Thomas Place. Separating regular languages with two quantifier alternations. *Log. Methods Comput. Sci.*, 14(4), 2018.
79. Thomas Place and Marc Zeitoun. Separating regular languages with first-order logic. *Log. Methods Comput. Sci.*, 12(1), 2016.
80. Thomas Place and Marc Zeitoun. Adding successor: A transfer theorem for separation and covering. *ACM Trans. Comput. Log.*, 21(2):9:1–9:45, 2020.
81. Wolfgang Rautenberg. Modal tableau calculi and interpolation. *J. Philos. Log.*, 12(4):403–423, 1983.
82. Giuseppe Rizzo, Nicola Fanizzi, and Claudia d’Amato. Class expression induction as concept space exploration: From dl-foi to dl-focl. *Future Gener. Comput. Syst.*, 108:256–272, 2020.
83. Giuseppe Rizzo, Nicola Fanizzi, Claudia d’Amato, and Floriana Esposito. A framework for tackling myopia in concept learning on the web of data. In *Proceedings of the 21st International Conference on Knowledge Engineering and Knowledge Management, EKAW 2018*, pages 338–354. Springer, 2018.
84. Md. Kamruzzaman Sarker and Pascal Hitzler. Efficient concept induction for description logics. In *Proceedings of the 33rd AAAI Conference on Artificial Intelligence, AAAI 2019*, pages 3036–3043. AAAI Press, 2019.
85. Inanç Seylan, Enrico Franconi, and Jos de Bruijn. Effective query rewriting with ontologies over dboxes. In *Proceedings of the 21st International Joint Conference on Artificial Intelligence, IJCAI 2009*, pages 923–925, 2009.
86. Balder ten Cate. Interpolation for extended modal languages. *J. Symb. Log.*, 70(1):223–234, 2005.
87. Balder ten Cate. *Model theory for extended modal languages*. PhD thesis, University of Amsterdam, 2005. ILLC Dissertation Series DS-2005-01.
88. Balder ten Cate. Lyndon interpolation for modal logic via type elimination sequences. Technical report, ILLC, Amsterdam, 01 2022.
89. Balder ten Cate, Willem Conradie, Maarten Marx, and Yde Venema. Definitorially complete description logics. In *Proceedings of the 10th International Conference on Principles of Knowledge Representation and Reasoning, KR 2006*, pages 79–89. AAAI Press, 2006.
90. Balder ten Cate, Enrico Franconi, and Inanç Seylan. Beth definability in expressive description logics. *J. Artif. Intell. Res.*, 48:347–414, 2013.
91. David Toman and Grant E. Weddell. *Fundamentals of Physical Design and Query Compilation*. Synthesis Lectures on Data Management. Morgan & Claypool Publishers, 2011.
92. David Toman and Grant E. Weddell. First order rewritability for ontology mediated querying in horn-dlfd. In *Proceedings of the 33rd International Workshop on Description Logics, DL 2020*. CEUR-WS.org, 2020.
93. David Toman and Grant E. Weddell. FO Rewritability for OMQ using Beth Definability and Interpolation. In *Proceedings of the 34th International Workshop on Description Logics, DL 2021*. CEUR-WS.org, 2021.
94. An C. Tran, Jens Dietrich, Hans W. Guesgen, and Stephen Marsland. Parallel symmetric class expression learning. *J. Mach. Learn. Res.*, 18:64:1–64:34, 2017.
95. Johan Van Benthem. The many faces of interpolation. *Synthese*, pages 451–460, 2008.
96. Albert Visser et al. Uniform interpolation and layered bisimulation. In *Gödel’96: Logical foundations of mathematics, computer science and physics—Kurt Gödel’s legacy*, pages 139–164. Association for Symbolic Logic, 1996.

97. Yizheng Zhao and Renate A. Schmidt. Forgetting concept and role symbols in *ALCOH*-ontologies. In *Proceedings of the 25th International Joint Conference on Artificial Intelligence, IJCAI 2016*, pages 1345–1353. IJCAI/AAAI Press, 2016.