




Article

Association between Organochlorine Pesticides and Vitamin D in Female Subjects

Edwina Brennan ^{1,*}, Alexandra E. Butler ¹, Manjula Nandakumar ¹, Daniel S. Drage ^{2,3}, Thozhukat Sathyapalan ⁴ and Stephen L. Atkin ¹

¹ School of Medicine, Royal College of Surgeons in Ireland-Medical University of Bahrain, Busaiteen 15503, Bahrain; abutler@rcsi.com (A.E.B.); mnandakumar@rcsi.com (M.N.); satkin@rcsi.com (S.L.A.)

² School of Geography, Earth and Environmental Sciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK; d.s.drage@bham.ac.uk

³ Queensland Alliance for Environmental Health Sciences, The University of Queensland, 39 Kessels Road, Brisbane, QLD 4108, Australia

⁴ Hull York Medical School, University of Hull, Hull HU6 7RX, UK; thozhukat.sathyapalan@hyms.ac.uk

* Correspondence: ebrennan@rcsi.com

Abstract: In human population studies, organochlorine pesticides (OCPs) have been linked to vitamin D deficiency. Therefore, this study examined the association between OCPs, vitamin D₃ (cholecalciferol, 25(OH)D₃), and the active metabolite 1,25-dihydrovitamin D₃ (1,25(OH)₂D₃) in a cohort of non-obese women. The serum samples of 58 female participants (age—31.9 ± 4.6 years; body mass index (BMI)—25.7 ± 3.7 kg/m²) were screened for 10 indicator OCPs. 25(OH)D₃ and 1,25(OH)₂D₃ levels were determined via isotope dilution liquid chromatography tandem mass spectrometry. In this cohort, the 25(OH)D₃ and 1,25(OH)₂D₃ levels were 22.9 ± 11.2 ng/mL and 0.05 ± 0.02 ng/mL, respectively, with 28 participants classified as 25(OH)D₃-deficient (<50 nmol/L). In the study cohort, no correlations were found between individual or total OCPs (ΣOCPs) and 25(OH)D₃. *p,p'*-dichlorodiphenyldichloroethylene (DDE) and ΣOCPs correlated positively with 1,25(OH)₂D₃, with the latter being negatively correlated with estimated glomerular filtration rate (eGFR). In women with sufficient 25(OH)D₃ levels, *p,p'*-dichlorodiphenyltrichloroethane (DDT) was positively correlated with 1,25(OH)₂D₃, whilst in the deficient group, hexachlorobenzene (HCB) and *p,p'*-DDE were positively correlated with 1,25(OH)₂D₃, β-Hexachlorocyclohexane (HCH) was positively correlated with 25(OH)D₃, and none of the OCPs were associated with measures of renal function. Overall, OCPs and ΣOCPs were not associated with 25(OH)D₃, suggesting that they are unrelated to vitamin D deficiency, but *p,p'*-DDE and ΣOCPs correlated positively with active 1,25(OH)₂D₃, while ΣOCPs correlated negatively with eGFR, suggesting a possible renal effect. Analysis of vitamin D deficiency revealed an association between β-HCH and 25(OH)D₃, and between HCB and *p,p'*-DDE and 1,25(OH)₂D₃, suggesting that OCP effects may be enhanced in cases of vitamin D deficiency.

Keywords: Organochlorine pesticides (OCPs); organic pollutants; vitamin D; cholecalciferol



Citation: Brennan, E.; Butler, A.E.; Nandakumar, M.; Drage, D.S.; Sathyapalan, T.; Atkin, S.L. Association between Organochlorine Pesticides and Vitamin D in Female Subjects. *Biomedicines* **2023**, *11*, 1451. <https://doi.org/10.3390/biomedicines11051451>

Academic Editors: Giuseppe Murdaca and Sebastiano Gangemi

Received: 23 April 2023

Revised: 4 May 2023

Accepted: 10 May 2023

Published: 15 May 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Organochlorine pesticides (OCPs) are a group of chlorinated hydrocarbons that were once used extensively in agriculture and mosquito control and have since been termed persistent organic pollutants (POPs) due to their high toxicity, slow degradation, and bioaccumulation [1]. OCPs are highly persistent in the environment and can accumulate in the fatty tissues of animals and humans [2,3]. They can also be transported over great distances in the air and water, leading to the contamination of soil and water sources far from the OCPs' original application site [4]. Due to the potential health risks associated with OCPs, many countries have banned or severely restricted their use in recent years [5].

However, these chemicals can persist in the environment for decades [4] and may still be found in some food and water sources.

Exposure to OCPs has been linked to a variety of health issues. OCPs have been linked to an increased risk of breast [6] and prostate [7] cancer, and lymphoma [8]. OCPs have been linked to reproductive problems in both men and women, including decreased fertility [9,10] and miscarriages, particularly with respect to hexachlorocyclohexane (HCH) [11], and prenatal exposure to α -HCH, β -HCH, *o,p'*-dichlorodiphenyldichloroethane (DDD), and *p,p'*-DDD has been associated with an increased risk for neural tube birth defects [12]. OCPs have been linked to neurological problems, which they are thought to contribute to via their accumulation in the body over time, potentially leading to Parkinson's disease (HCH) [13] and Alzheimer's disease (*p,p'*-dichlorodiphenyldichloroethylene (DDE)) [14]. OCPs have been linked to immune system disorders, with dichlorodiphenyltrichloroethane (DDT) causing the dysregulation of interleukin 6 (IL-6), leading to chronic inflammation [15], and having been related to oxidative stress and immune suppression [16], thus increasing susceptibility to infections and other diseases. OCPs have been classed as endocrine disruptors that can cause developmental problems in children (DDT) [17] and reproductive problems in adults [9]. In addition, low doses of OCPs have been strongly linked to various chronic diseases including diabetes [18] and cardiovascular diseases [19].

OCPs may have several mechanisms of action that contribute to their reported diverse effects. OCPs function by binding to specific target sites in the body, including binding to and blocking the GABAA receptors of sodium channels in the brain [20]. By binding to these sites, they disrupt the normal functioning of the nervous system, leading to the overstimulation or inhibition of nerve impulses. OCPs can also interfere with the normal function of the endocrine system by disrupting the production, release, transport, and metabolism of hormones at several sites of action such as the hypothalamus (increasing gonadotropin-releasing hormone mRNA levels), pituitary glands (affecting luteinizing hormone release), and ovaries (decreased sex steroid hormone production) [20].

Vitamin D deficiency is associated with various health conditions such as osteoporosis, cancer, cardiovascular disease, autoimmune diseases, and increased mortality [21,22]. In addition, vitamin D deficiency is associated with an increased incidence of type 2 diabetes [23–25] and microvascular and macrovascular diabetes-related complications [26]. Vitamin D₃ (cholecalciferol, 25(OH)D₃) is endogenously produced via the UV-B irradiation of 7-dehydrocholesterol and is hydroxylated to 25(OH)D₃ by multiple 25-hydroxylases in the liver [27]. 25(OH)D₃ is transported to the kidneys and converted to active 1,25-dihydrovitamin D₃ (1,25(OH)₂D₃) by 1 alpha hydroxylase present in the renal tubular cells [28]. It has recently been reported that extrarenal tissues may also convert 25(OH)D₃ to 1,25(OH)₂D₃, although activation in renal and non-kidney tissues is regulated differently via the macrophage production of 1,25(OH)₂D₃ through the type 2 interferon (IFN) response [29].

There is some evidence suggesting that exposure to OCPs may be associated with lower levels of vitamin D in the body [19,30]. Population studies have suggested that OCPs, including *p,p'*-DDT, *p,p'*-DDE, and β -HCH, can cause vitamin D deficiency [31]. In one such study, Inuit people who were heavily exposed to these pesticides through their traditional diet of fish and marine mammals, causing accumulation of these pesticides, showed higher levels of OCPs in their blood and had lower levels of vitamin D [32]; however, chronobiological factors within this population are likely to play a role.

OCPs may contribute to vitamin D deficiency through their effects on vitamin D metabolism in the liver, the disruption of the conversion of 25(OH)D₃ to 1,25(OH)₂D₃, interference with the function of vitamin D receptors in the body, or the impairment of calcium homeostasis [33]. It is important to note that the evidence linking OCPs and vitamin D levels is still limited, and more research is needed to fully understand the relationship between these two factors. However, the evidence still highlights the potential health risks associated with exposure to these persistent environmental pollutants. Therefore, this study was undertaken to examine the association between OCPs (pentachlorobenzene (PeCB),

hexachlorobenzene (HCB), HCH, lindane, trans-chlordane, cis-chlordane, DDE, DDD, DDT, and dodecachlorooctahydro-1,3,4-metheno-1H-cyclobuta[cd]pentalene (Mirex)) and 25(OH)D₃ and its active metabolite 1,25(OH)₂D₃ in a group of non-obese healthy women prior to undergoing in vitro fertilization (IVF). In addition, we sought to determine if such an association may be precipitated by the impairment of calcium homeostasis, specifically with respect to its association with calcium/calmodulin-dependent (CaM) kinases (CaMK), CaMK kinase alpha (CaMKK α), and CaM phosphodiesterase (PDE) proteins.

2. Materials and Methods

2.1. Patient Recruitment

Subjects were recruited from the Hull IVF Unit, UK, following ethical approval from The Yorkshire and The Humber NRES ethical committee, UK (approval number 02/03/043). Venesection was performed on 58 fasting, non-obese Caucasian women; samples were taken at 09:00 on day 21 of the menstrual cycle prior to IVF downregulation. Venous blood samples were centrifuged at 3500 \times g for 15 min at 4 °C, and serum was stored at –80 °C within 1 h of collection for further analysis. No participant was on any prescribed or over-the-counter medication. All participants provided written informed consent.

2.2. OCP Measurement

Samples were screened for 10 indicator OCPs (PeCB, HCB, HCH, lindane, trans-chlordane, cis-chlordane, DDE, DDD, DDT, and Mirex). A sum-of-OCPs (Σ OCPs) variable was calculated by adding the molar concentrations of each OCP analyzed. After vortexing the samples for 1 min, 6 mL of acetonitrile, 3 mL of milliQ water, 5 g of anhydrous MgSO₄, and 1 g of NaCl were added along with a ceramic homogenizer. Samples were manually shaken for 1 min before being centrifuged at 4700 RPM for 8 min at 10 °C. The supernatant layer (approximately 6 mL of acetonitrile) was collected and transferred to 12 mL glass vials. An aliquot of 200 μ L was collected and cleaned for non-targeted analysis (not presented here). The remainder of the extract was transferred to a 15 mL glass tube, evaporated under a gentle stream of nitrogen to near-dryness, and reconstituted in approximately 1 mL of hexane. Approximately 1 mL of >98% concentrated sulfuric acid was added, and the sample was vortexed for at least 30 s. The aqueous and hexane layers were left to separate overnight at <4 °C. The supernatant (hexane) layer was transferred directly onto a silica solid-phase extraction (SPE) cartridge (Supelco LC-Si, 3 mL/500 mg) (preconditioned with 6 mL dichloromethane (DCM) followed by 6 mL of hexane). Target compounds were eluted into a 15 mL glass collection tube using 6 mL of hexane followed by 8 mL of DCM. Samples were evaporated to near-dryness and reconstituted in 50 μ L of methanol containing 2.5 ng ¹³C₁₂-PCB-141 as a recovery standard. Samples were transferred to autosampler vial inserts prior to analysis.

2.3. Instrumental Analysis

OCPs were identified in serum samples via high-resolution gas chromatography coupled with high-resolution mass spectrometry (HRGC/HRMS). A Thermofisher TRACE 1300 gas chromatograph was coupled to a Thermofisher DFS mass spectrometer. The injector was operated in splitless mode with separation achieved on an Agilent DB-5ms column (30 m length \times 0.25 mm in diameter \times 0.25 μ m film thickness). Experiments were conducted in multiple ion detection (MID) mode at a resolution level of 10,000 (10% valley definition). The inlet, transfer line, and source were maintained at 250 °C, 280 °C, and 280 °C, respectively. The flow rate was maintained at 1.0 mL/min. In the GC oven, an initial temperature of 80 °C was held for two minutes before being ramped up to 180 °C at 20 °C/min and held for 0.5 min. The temperature was then increased to 300 °C at 10 °C/min and held for 5 min.

2.4. Quality Assurance/Quality Control

The limit of reporting (LOR) was determined according to the quality of the peak shape and using a minimum signal-to-noise ratio of 10:1. Contamination was assessed via inclusion of procedural blanks (milli-Q water, $n = 6$) extracted in parallel with unknown samples. If a target compound was detected in a blank at below 5% of a sample, then no correction occurred. When blank concentrations were 5–25% of a respective sample, the sample was blank corrected. Linearity was established with a seven-point calibration across the range of 0.1 to 25 ng/mL for all target analytes. R^2 values > 0.99 were achieved for all compounds. Analytes were identified based on retention time ($< 2\%$ RSD from standard) and confirmed with reference to a second ion (HRGC/HRMS). Quantification was performed using isotope dilution mass spectrometry. All OCPs were quantified against $^{13}\text{C}_{12}$ -HCB. All data were validated by spiking 5 aliquots of bovine serum with 300 pg of target analytes. The concentrations of the spiked samples were measured, and the recoveries (measured concentration/spiked concentration $\times 100$) were calculated. Average recoveries were 80–120% with a relative standard deviation of $< 15\%$, thus conforming to the OECD guidelines of good laboratory practice (Supplementary Table S1).

2.5. Vitamin D₃ and Biochemical Parameters

Biochemical and hormonal parameters were measured as per a previously detailed method [34]. Briefly, serum 25(OH)D₃ and 1,25(OH)₂D₃ levels were quantified using isotope dilution liquid chromatography tandem mass spectrometry (LC-MS/MS). Vitamin D metabolites and labelled internal standards were extracted from 250 μL of serum using supportive liquid–liquid extraction and Diels–Alder derivatization prior to LC-MS/MS analysis. Chromatographic separations were achieved using Hypersil Gold C18 column (150 \times 2.1 mm; 1.9 μm) at a flow rate of 0.2 mL/min and operated in Electrospray Ionization (ESI) positive mode and analyzed using the multiple-reaction-monitoring (MRM) method. The limit of quantification (LOQ) for 1,25(OH)₂D₃ was 10 pg/mL and 25(OH)D₃ 1.2 nmol/L (0.5 ng/mL) [35]. Vitamin D deficiency was defined as a cutoff of 20 ng/mL (50 nmol/L) as per the clinical practice guidelines of the Endocrine Society Task Force on Vitamin D [36].

2.6. CaM Proteomic Measurement

Circulating levels of CaM-associated proteins (PDE1A, CaMK2A, CaMK2B, CaMK1D, CaMK2D, CaMK1, and CaMKK α) were determined via Slow Off-rate Modified Aptamer (SOMA)-scan plasma protein measurement (Somalogic, Boulder, CO, USA), the details of which have been previously reported [37]. Briefly, EDTA plasma samples were diluted, and the following assay steps were performed: (a) Binding: analytes and primer bead–SOMAmers (fully synthetic fluorophore-labelled SOMAmer coupled to a biotin moiety through a photocleavable linker) are equilibrated. (b) Catch I: all analyte–SOMAmer complexes are immobilized on a streptavidin-substituted support. Washing steps: proteins not stably bound to primer bead–SOMAmers are removed, and bound protein is biotinylated. (c) Cleave: long-wave ultraviolet light is applied to release analyte SOMAmer complexes into the solution. (d) Catch II: analyte–SOMAmer complexes are selectively immobilized on streptavidin support via the introduced analyte-borne biotinylation. Further washing is performed to select against non-specific analyte–SOMAmer complexes. (e) Elution: denaturation is induced to disrupt analyte–SOMAmer complexes. Released SOMAmers serve as surrogates for quantification of analyte concentrations. (f) Quantification: SOMAmer complementary oligonucleotides are hybridized to custom arrays. Normalization of raw intensities, hybridization, median signal, and calibration signal were performed based on the standard samples included on each plate, as previously described [38].

2.7. Statistics

No previous studies were available to enable the performance of a power analysis; therefore, this pilot study was designed according to the research conducted by Birkett and Day [39]. Descriptive data are presented as mean \pm standard deviation (SD) for

continuous data. Potential correlations between OCPs, 25(OH)D₃, and 1,25(OH)₂D₃ were examined using exploratory Spearman's rank order correlations. A *p*-value of <0.05 was considered to indicate statistical significance. Statistical analysis was carried out using Jamovi (version 2.0.0).

3. Results

3.1. Whole-Cohort Analysis

Demographic and biochemical data for this cohort are shown in Table 1. The participants had a mean age of 31.9 ± 4.6 years and a mean body mass index (BMI) of 25.7 ± 3.7 kg/m². Thyroid function and C-reactive protein (CRP) levels, constituting measures of underlying inflammation, were normal. The mean levels of 25(OH)D₃ and 1,25(OH)₂D₃ were 22.9 ± 11.2 ng/mL and 0.05 ± 0.02 ng/mL, respectively. Of the 58 women recruited, 28 had a 25(OH)D₃ level less than 20 ng/mL (50 nmol/L). The obtained levels of OCPs, 25(OH)D₃, and 1,25(OH)₂D₃ are shown in Table 1.

Table 1. Mean and standard deviation of OCPs in the population studied.

	Female Subjects (n = 58)	
	Mean	SD
Age (years)	31.9	4.6
BMI (kg/m ²)	25.7	3.7
CRP (mg/L)	2.6	2.5
TSH (mU/L)	2.4	2.2
Free-T3 (pmol/L)	4.8	0.7
Free-T4 (pmol/L)	11.3	1.8
25(OH)D ₃ (ng/mL)	23	11.2
1,25(OH) ₂ D ₃ (ng/mL)	0.05	0.02
Calcium (mmol/L)	2.3	0.07
Urea (nmol/L)	4.66	5.21
Creatinine (nmol/L)	66	8.86
eGFR (mL/min/1.73 m ²)	92.3	14.8
PDE1A (RFU)	789	1060
CAMK2A (RFU)	373	453
CaMK2B (RFU)	859	1423
CaMK1D (RFU)	1695	580
CaMK2D (RFU)	2379	3462
CaMK1 (RFU)	4681	1371
CaMKK α (RFU)	343	463
PeCB (ng/g Lipid)	9.12	9.1
α-HCH (ng/g Lipid)	<LOR	
β-HCH (ng/g Lipid)	4.63	4.92
HCB (ng/g Lipid)	9.81	6.5
Lindane (ng/g Lipid)	1.35	0.28
Trans-Chlordane (ng/g Lipid)	1.4	0.97
cis-Chlordane (ng/g Lipid)	1.59	0.58
p,p'-DDE (ng/g Lipid)	40.7	60.4
o,p'-DDE (ng/g Lipid)	2.23	0.32
o,p'-DDD (ng/g Lipid)	<LOR	
p,p'-DDD+o,p'-DDT (ng/g Lipid)	<LOR	
p,p'-DDT (ng/g Lipid)	4.99	2.3
Mirex (ng/g Lipid)	3.27	0.69
ΣOCPs (ng/g Lipid)	63.3	66.8

BMI: body mass index; CRP: C-reactive protein; TSH: thyroid-stimulating hormone; Free-T3: free-triiodothyronine; Free-T4: free-thyroxine; eGFR: estimated glomerular filtration rate; PDE: calcium/calmodulin-dependent phosphodiesterase; CaMK: calcium/calmodulin-dependent kinase; CaMKK α: calcium/calmodulin-dependent kinase alpha; PeCB: pentachlorobenzene; α: alpha; β: beta; HCH: hexachlorocyclohexane; HCB: hexachlorobenzene; p: para; o: ortho; DDE: dichlorodiphenyldichloroethylene; DDD: dichlorodiphenyldichloroethane; DDT: dichlorodiphenyltrichloroethane; Mirex: dodecachlorooctahydro-1,3,4-metheno-1H-cyclobuta[cd]pentalene; SD: standard deviation; LOR: limit of reporting; ΣOCPs: total organochlorine pesticides.

3.2. OCP Levels

The detection frequency of individual OCPs ranged from 100–12%. *p,p'*-DDE was detected in all serum samples, making it the most abundant OCP, followed by HCB (95%), *p,p'*-DDT (74%), β -HCH (67%), *cis*-chlordane (48%), Mirex (41%), PeCB (38%), *trans*-chlordane (24%), *o,p'*-DDE (22%), and lindane (12%). The OCPs α -HCH, *o,p'*-DDD, and *p,p'*-DDD+*o,p'*-DDT were not detected.

3.3. Whole-Group Correlations

In the study cohort, no correlations were found between Σ OCPs or individual OCPs and 25(OH) D_3 . The OCP *p,p'*-DDE and the Σ OCPs positively correlated with 1,25(OH) $_2D_3$, ($\rho = 0.41$; $p = 0.01$) and ($\rho = 0.34$; $p = 0.04$), respectively (Figure 1A,B), but no other correlations were found between individual OCPs and 1,25(OH) $_2D_3$ (Table 2). The Σ OCPs variable negatively correlated with eGFR ($\rho = -0.28$; $p = 0.04$) (Figure 1C). There were no correlations between *p,p'*-DDE or Σ OCPs with urea and creatinine.




Figure 1. Significant correlations of OCPs with 1,25(OH) $_2D_3$ and measures of renal function in whole study cohort. (A) 1,25(OH) $_2D_3$ and *p,p'*-DDE; (B) 1,25(OH) $_2D_3$ and Σ OCPs; (C) eGFR and Σ OCPs.


Table 2. Spearman’s rho (ρ) correlations of OCPs with 25(OH)D₃ and 1,25(OH)₂D₃. Data in bold indicate statistically significant correlations.

	PeCB	β -HCH	HCB	Lindane	trans-Chlordane	cis-Chlordane	o,p'-DDE	p,p'-DDE	p,p'-DDT	Mirex	Σ OCPs
25(OH)D ₃	0.289 (0.192)	0.174 (0.339)	0.101 (0.502)	0.393 (0.40)	−0.14 (0.648)	0.325 (0.091)	0.374 (0.258)	0.272 (0.061)	0.096 (0.595)	−0.204 (0.361)	0.284 (0.051)
1,25(OH) ₂ D ₃	−0.108 (0.66)	0.315 (0.14)	0.333 (0.05)	−0.2 (0.92)	−0.205 (0.74)	0.278 (0.263)	0.06 (0.89)	0.408 (0.01)	0.364 (0.09)	−0.03 (0.92)	0.34 (0.04)

3.4. Sufficient and Deficient 25(OH)D₃ Subgroup Correlations

The study cohort was divided into two subgroups based on the participants’ vitamin D levels: sufficient 25(OH)D₃ (>20 ng/mL; >50 nmol/L) women and deficient 25(OH)D₃ (<20 ng/mL; <50 nmol/L) women. In the sufficient 25(OH)D₃ women, p,p'-DDT strongly positively correlated with 1,25(OH)₂D₃ ($\rho = 0.70$; $p = 0.01$) (Figure 2A). There were no correlations found between OCPs and 25(OH)D₃ in the sufficient 25(OH)D₃ women subgroup. In deficient 25(OH)D₃ women, HCB and p,p'-DDE positively correlated with 1,25(OH)₂D₃ ($\rho = 0.55$; $p = 0.03$) and ($\rho = 0.52$; $p = 0.045$), respectively (Figure 2B,C)), while β -HCH strongly positively correlated with 25(OH)D₃ ($\rho = 0.85$; $p = 0.01$) (Figure 2D). Calcium levels did not differ between vitamin D sufficiency (2.3 ± 0.07 mmol/L) and vitamin D deficiency groups (2.3 ± 0.06 mmol/L).

We have found that 1,25(OH)₂D₃ levels may be modulated through CaMK1 by poly-brominated diethyl ether 153 (unpublished results); therefore, the CaM-associated proteins (PDE1A, CaMK2A, CaMK2B, CaMK1D, CaMK2D, CaMK1, and CaMKK α) (Table 1) that effect the rapid action of 1,25(OH)₂D₃ at the cell membrane leading to increasing intracellular calcium levels were investigated, but no associations with the individual or total OCPs, or in vitamin-D-sufficient or deficient states, were found.

**Figure 2.** Cont.




Figure 2. Significant correlations of OCPs with 25(OH)D₃ and 1,25(OH)₂D₃ in cohort subgroups. The study cohort was divided into two subgroups based on the participants' vitamin D levels: sufficient 25(OH)D₃ (>20 ng/mL; >50 nmol/L) women and deficient 25(OH)D₃ (<20 ng/mL; <50 nmol/L) women. Sufficient vitamin D subgroup—(A) 1,25(OH)₂D₃ and p,p'-DDT. Deficient vitamin D subgroup—(B) 1,25(OH)₂D₃ and HCB; (C) 1,25(OH)₂D₃ and p,p'-DDE; (D) 25(OH)D₃ and β-HCH.

4. Discussion

These data show that the 10 indicator OCPs and ΣOCPs analyzed do not appear to be associated with 25(OH)D₃ but that the OCP p,p'-DDE and ΣOCPs correlated positively with active 1,25(OH)₂D₃. Overall, the ΣOCPs were negatively associated with a reduced eGFR level, as noted by others [40], indicating that their action may be exerted on the kidneys, as suggested by the reported association of DDT burden with chronic kidney disease [41], but this was an observational study with no mechanistic insights. An inverse relationship of OCPs, including p,p'-DDT, p,p'-DDE, and β-HCH, with vitamin D has been reported [31], but this was not seen in the data presented herein; however, this difference is potentially due to the differing study populations, as those associations were seen amongst subjects of an older age, who were Caucasian, or who suffered from chronic diseases, whereas the population analyzed in this study was young and healthy.

When the study cohort was divided into two subgroups, namely, sufficient and deficient groups based on their vitamin D levels, in the vitamin-D-sufficient 25(OH)D₃ women, p,p'-DDT strongly positively correlated with 1,25(OH)₂D₃. There were no correlations found between OCPs and 25(OH)D₃ in the vitamin-D-sufficient 25(OH)D₃ subgroup of women. In the vitamin-D-deficient 25(OH)D₃ women, both p,p'-DDE and HCB positively correlated with 1,25(OH)₂D₃. In the vitamin-D-deficient group, β-HCH strongly positively correlated with 25(OH)D₃, suggesting that a reduction in vitamin D may signal a potential effect of both HCB and HCH, but it is unclear what those specific effects may be. The mechanism(s) behind the positive associations between OCPs and 1,25(OH)₂D₃ is unclear, but it has been reported that in cells exposed to p,p'-DDT and p,p'-DDE, this led to an increase in the vitamin D binding protein [42] that binds to 1,25(OH)₂D₃ for its activity, perhaps leading to an increase in 1,25(OH)₂D₃ levels.

1,25(OH)₂D₃ increases calcium absorption via vitamin D receptor (VDR) binding and exerts rapid action at the cell membrane, increasing intracellular calcium levels by activating protein kinase cascades that stimulate VDR-mediated transcription-enhancing steroid receptor coactivator (SRC) activity, with the overall effect of enhancing 1,25(OH)₂D₃-dependent interactions between VDR and SRC coactivators [43]. To investigate a possible mechanism by which OCPs may affect 1,25(OH)₂D₃ action, the determination of the plasma levels of CaM proteins (CaMK, CaMKK, and PDE), which may be indicative of 1,25(OH)₂D₃

action, was undertaken. No correlations with total or individual OCPs were found, suggesting that the OCPs impacting $1,25(\text{OH})_2\text{D}_3$ are not acting at the membrane level through calmodulin-associated proteins. Therefore, it is clear that $25(\text{OH})\text{D}_3$ is converted to its biologically active form $1,25(\text{OH})_2\text{D}_3$ in the presence of OCPs without affecting serum calcium levels, as has been described in other animal species [44], and it is unclear how OCPs may then impair the vitamin D biological response. That the OCPs appeared to be associated with vitamin D and vitamin D sufficiency or deficiency suggests that OCPs' effects may be mediated through vitamin D. Of particular concern, OCPs may potentiate vitamin D deficiency, which is a condition that affects over 50% of the global population [45]. Moreover, the diseases that are associated with vitamin D deficiency, such as diabetes and cardiovascular disease [21,22], are the same as those attributed to OCP exposure [19,30], suggesting that one may represent an epiphenomenon of the other. Another potential explanation is that the association between OCPs and vitamin D may be caused by indirect mechanisms. In vitro, OCPs induce inflammation and oxidative stress, with β -HCH, DDE, and Dieldrin causing the upregulation of pro-inflammatory cytokines such as tumor necrosis factor (TNF)- α , IL-1 β , IL-6, nuclear factor kappa B (NF- κ B), and cyclooxygenase (COX)-2 expression in OCP-treated human ovary surface epithelial cells [46]. These findings are in line with observations that DDT causes the dysregulation of IL-6 leading to chronic inflammation [15] and that DDT is related to oxidative stress and immune suppression [15,16]. This is also reflected in in vivo findings, where, in a systematic review, exposure to OCPs was positively associated with CRP, IL-1 β , IL-2, and IL-10, which is indicative of dysregulated inflammatory responses; however, the underlying mechanisms remain unclear [47]. Conversely, vitamin D prevents oxidative stress, while vitamin D deficiency induces the disruption of mitochondrial function [48] and is also associated with markers of inflammation [49]. Consequently, in vitamin D sufficiency, this may generally mitigate the inflammatory and oxidative stress effects of the OCPs; however, in vitamin D deficiency, these effects have already been uncovered and indeed may be exacerbated (Figure 3), as seen with results reporting that vitamin D deficiency promotes colonic inflammation [50] and, in an animal model of acetaminophen-induced liver injury, exacerbates induced oxidative stress [51].




Figure 3. OCPs analyzed in this study. The association of OCPs with various diseases in vitamin D sufficiency and deficiency were analyzed. Vitamin D sufficiency may generally mitigate the inflammatory and oxidative stress effects and associated disorders/diseases arising due to exposure to OCPs, but in vitamin D deficiency, those effects are unmasked and aggravated.

In addition, extrarenal tissues may also convert $25(\text{OH})\text{D}_3$ to $1,25(\text{OH})_2\text{D}_3$; however, notably, activation in renal and non-kidney tissues is regulated differently, with macrophage production of $1,25(\text{OH})_2\text{D}_3$ through the type 2 IFN response [29]. In animal models, OCPs do not appear to affect macrophages; therefore, their influence through an extrarenal effect on $1,25(\text{OH})_2\text{D}_3$ appears less likely [52]. However, DDT caused an increase in IFN γ secretion [53]; increasing IFN γ may have the effect of increasing the conversion of $25(\text{OH})\text{D}_3$ to $1,25(\text{OH})_2\text{D}_3$ in the extrarenal tissues.

The strengths of this study include its state-of-the-art measurement of OCPs and vitamin D ($25(\text{OH})\text{D}_3$ and $1,25(\text{OH})_2\text{D}_3$). Its limitations include the small number of subjects included, who were all Caucasian females, so these findings may not be generalizable to male subjects or those of different ethnicities. The low numbers compounded by the detection number may have resulted in a type-two statistical error (false negative); however, this study will potentiate the determinative power of a larger study focusing on vitamin D deficiency. OCPs have been associated with parathyroid hyperplasia leading to elevated parathyroid hormone (PTH) and calcium levels [54]; however, the calcium levels were in the normal reference range in these patients, thus obviating the need to measure PTH.

5. Conclusions

In conclusion, our findings show that, overall, OCPs and ΣOCPs were not associated with $25(\text{OH})\text{D}_3$, suggesting that they are unrelated to vitamin D deficiency, and that the OCP p,p'-DDE and the ΣOCPs correlated positively with active $1,25(\text{OH})_2\text{D}_3$ and negatively with eGFR, suggesting a renal effect. Vitamin D deficiency unmasked an association between $\beta\text{-HCH}$ and $25(\text{OH})\text{D}_3$ and between HCB and p,p'-DDE and $1,25(\text{OH})_2\text{D}_3$, suggesting that OCPs' effects may be enhanced in patients with vitamin D deficiency.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/biomedicines11051451/s1>. Supplementary Table S1. OCP validation data.

Author Contributions: Conceptualization, S.L.A. and T.S.; methodology, S.L.A. and T.S.; validation, S.L.A.; formal analysis, E.B. and A.E.B.; investigation, D.S.D.; resources, S.L.A.; data curation, S.L.A.; writing—original draft preparation, E.B., A.E.B. and S.L.A.; writing—review and editing, E.B., A.E.B., S.L.A., D.S.D., M.N. and T.S.; visualization, E.B., A.E.B., M.N. and S.L.A. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki, and approved by the Institutional Review Board (or Ethics Committee) of The Yorkshire and The Humber NRES ethical committee, UK (approval number 02/03/043).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: The data presented in this study are available on reasonable request from the corresponding author.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Jayaraj, R.; Megha, P.; Sreedev, P. Organochlorine pesticides, their toxic effects on living organisms and their fate in the environment. *Interdiscip. Toxicol.* **2016**, *9*, 90–100. [[CrossRef](#)] [[PubMed](#)]
2. Deshmukh, H.; Aylward, L.L.; Rose, M.; Fernandes, A.; Sedman, P.; Thatcher, N.J.; Atkin, S.L.; Sathyapalan, T. Association of endocrine active environmental compounds with body mass index and weight loss following bariatric surgery. *Clin. Endocrinol.* **2020**, *93*, 280–287. [[CrossRef](#)] [[PubMed](#)]
3. Buah-Kwofie, A.; Humphries, M.S.; Combrink, X.; Myburgh, J.G. Accumulation of organochlorine pesticides in fat tissue of wild Nile crocodiles (*Crocodylus niloticus*) from iSimangaliso Wetland Park, South Africa. *Chemosphere* **2018**, *195*, 463–471. [[CrossRef](#)] [[PubMed](#)]
4. Ma, Y.; Yun, X.; Ruan, Z.; Lu, C.; Shi, Y.; Qin, Q.; Men, Z.; Zou, D.; Du, X.; Xing, B.; et al. Review of hexachlorocyclohexane (HCH) and dichlorodiphenyltrichloroethane (DDT) contamination in Chinese soils. *Sci. Total Environ.* **2020**, *749*, 141212. [[CrossRef](#)]

5. Keswani, C.; Dilnashin, H.; Birla, H.; Roy, P.; Tyagi, R.K.; Singh, D.; Rajput, V.D.; Minkina, T.; Singh, S.P. Global footprints of organochlorine pesticides: A pan-global survey. *Environ. Geochem. Health* **2022**, *44*, 149–177. [[CrossRef](#)] [[PubMed](#)]
6. Miao, Y.; Rong, M.; Li, M.; He, H.; Zhang, L.; Zhang, S.; Liu, C.; Zhu, Y.; Deng, Y.L.; Chen, P.P.; et al. Serum concentrations of organochlorine pesticides, biomarkers of oxidative stress, and risk of breast cancer. *Environ. Pollut.* **2021**, *286*, 117386. [[CrossRef](#)]
7. Antignac, J.P.; Figiel, S.; Pinault, M.; Blanchet, P.; Bruyère, F.; Mathieu, R.; Lebdaï, S.; Fournier, G.; Rigaud, J.; Mahéo, K.; et al. Persistent organochlorine pesticides in periprostatic adipose tissue from men with prostate cancer: Ethno-geographic variations, association with disease aggressiveness. *Environ. Res.* **2023**, *216*, 114809. [[CrossRef](#)] [[PubMed](#)]
8. Luo, D.; Zhou, T.; Tao, Y.; Feng, Y.; Shen, X.; Mei, S. Exposure to organochlorine pesticides and non-Hodgkin lymphoma: A meta-analysis of observational studies. *Sci. Rep.* **2016**, *6*, 25768. [[CrossRef](#)]
9. Rattan, S.; Zhou, C.; Chiang, C.; Mahalingam, S.; Brehm, E.; Flaws, J.A. Exposure to endocrine disruptors during adulthood: Consequences for female fertility. *J. Endocrinol.* **2017**, *233*, R109–R129. [[CrossRef](#)]
10. Giulioni, C.; Maurizi, V.; Castellani, D.; Scarcella, S.; Skrami, E.; Balercia, G.; Galosi, A.B. The environmental and occupational influence of pesticides on male fertility: A systematic review of human studies. *Andrology* **2022**, *10*, 1250–1271. [[CrossRef](#)]
11. Pathak, R.; Mustafa, M.; Ahmed, R.S.; Tripathi, A.K.; Guleria, K.; Banerjee, B.D. Association between recurrent miscarriages and organochlorine pesticide levels. *Clin. Biochem.* **2010**, *43*, 131–135. [[CrossRef](#)] [[PubMed](#)]
12. Yin, S.; Sun, Y.; Yu, J.; Su, Z.; Tong, M.; Zhang, Y.; Liu, J.; Wang, L.; Li, Z.; Ren, A.; et al. Prenatal exposure to organochlorine pesticides is associated with increased risk for neural tube defects. *Sci. Total Environ.* **2021**, *770*, 145284. [[CrossRef](#)]
13. Xu, S.; Yang, X.; Qian, Y.; Luo, Q.; Song, Y.; Xiao, Q. Analysis of serum levels of organochlorine pesticides and related factors in Parkinson's disease. *Neurotoxicology* **2022**, *88*, 216–223. [[CrossRef](#)] [[PubMed](#)]
14. Medehouenou, T.C.M.; Ayotte, P.; Carmichael, P.H.; Kröger, E.; Verreault, R.; Lindsay, J.; Dewailly, É.; Tyas, S.L.; Bureau, A.; Laurin, D. Exposure to polychlorinated biphenyls and organochlorine pesticides and risk of dementia, Alzheimer's disease and cognitive decline in an older population: A prospective analysis from the Canadian Study of Health and Aging. *Environ. Health* **2019**, *18*, 57. [[CrossRef](#)] [[PubMed](#)]
15. Martin, T.J.; Gabure, S.; Maise, J.; Snipes, S.; Peete, M.; Whalen, M.M. The organochlorine pesticides pentachlorophenol and dichlorodiphenyltrichloroethane increase secretion and production of interleukin 6 by human immune cells. *Environ. Toxicol. Pharmacol.* **2019**, *72*, 103263. [[CrossRef](#)]
16. Koner, B.C.; Banerjee, B.D.; Ray, A. Organochlorine pesticide-induced oxidative stress and immune suppression in rats. *Indian J. Exp. Biol.* **1998**, *36*, 395–398.
17. Jeddy, Z.; Kordas, K.; Allen, K.; Taylor, E.V.; Northstone, K.; Dana Flanders, W.; Namulanda, G.; Sjodin, A.; Hartman, T.J. Prenatal exposure to organochlorine pesticides and early childhood communication development in British girls. *Neurotoxicology* **2018**, *69*, 121–129. [[CrossRef](#)]
18. Evangelou, E.; Ntritsos, G.; Chondrogiorgi, M.; Kavvoura, F.K.; Hernández, A.F.; Ntzani, E.E.; Tzoulaki, I. Exposure to pesticides and diabetes: A systematic review and meta-analysis. *Environ. Int.* **2016**, *91*, 60–68. [[CrossRef](#)]
19. Zago, A.M.; Faria, N.M.X.; Fávero, J.L.; Meucci, R.D.; Woskie, S.; Fassa, A.G. Pesticide exposure and risk of cardiovascular disease: A systematic review. *Glob. Public Health* **2022**, *17*, 3944–3966. [[CrossRef](#)]
20. Costa, L.G. The neurotoxicity of organochlorine and pyrethroid pesticides. *Handb. Clin. Neurol.* **2015**, *131*, 135–148. [[CrossRef](#)]
21. Bjelakovic, G.; Gluud, L.L.; Nikolova, D.; Whitfield, K.; Wetterslev, J.; Simonetti, R.G.; Bjelakovic, M.; Gluud, C. Vitamin D supplementation for prevention of mortality in adults. *Cochrane Database Syst. Rev.* **2014**, Cd007470. [[CrossRef](#)] [[PubMed](#)]
22. Osorio Landa, H.K.; Perez Diaz, I.; Laguna Barcanas, S.D.C.; Lopez Navarro, J.M.; Abella Roa, M.F.; Corral Orozco, M.; Mancilla Ortega, J.P.; Martinez Duarte, D.A.; Morales Montalvo, S.I.; Muzquiz Aguirre, S.; et al. Association of serum vitamin D levels with chronic disease and mortality. *Nutr. Hosp.* **2020**, *37*, 335–342. [[CrossRef](#)] [[PubMed](#)]
23. Nakashima, A.; Yokoyama, K.; Yokoo, T.; Urashima, M. Role of vitamin D in diabetes mellitus and chronic kidney disease. *World J. Diabetes* **2016**, *7*, 89–100. [[CrossRef](#)] [[PubMed](#)]
24. Husemoen, L.L.; Thuesen, B.H.; Fenger, M.; Jorgensen, T.; Glumer, C.; Svensson, J.; Ovesen, L.; Witte, D.R.; Linneberg, A. Serum 25(OH)D and type 2 diabetes association in a general population: A prospective study. *Diabetes Care* **2012**, *35*, 1695–1700. [[CrossRef](#)] [[PubMed](#)]
25. Mathieu, C. Vitamin D and diabetes: Where do we stand? *Diabetes Res. Clin. Pract.* **2015**, *108*, 201–209. [[CrossRef](#)]
26. Butler, A.E.; Dargham, S.R.; Latif, A.; Mokhtar, H.R.; Robay, A.; Chidiac, O.M.; Jayyousi, A.; Al Suwaidi, J.; Crystal, R.G.; Abi Khalil, C.; et al. Association of vitamin D3 and its metabolites in subjects with and without Type 2 diabetes and their relationship to diabetes complications. *Ther. Adv. Chronic Dis.* **2020**, *11*, 2040622320924159. [[CrossRef](#)]
27. Bikle, D.D. Vitamin D metabolism, mechanism of action, and clinical applications. *Chem. Biol.* **2014**, *21*, 319–329. [[CrossRef](#)]
28. Christakos, S.; Dhawan, P.; Verstuyf, A.; Verlinden, L.; Carmeliet, G. Vitamin D: Metabolism, Molecular Mechanism of Action, and Pleiotropic Effects. *Physiol. Rev.* **2016**, *96*, 365–408. [[CrossRef](#)]
29. Adams, J.S.; Rafison, B.; Witzel, S.; Reyes, R.E.; Shieh, A.; Chun, R.; Zavala, K.; Hewison, M.; Liu, P.T. Regulation of the extrarenal CYP27B1-hydroxylase. *J. Steroid Biochem. Mol. Biol.* **2014**, *144 Pt A*, 22–27. [[CrossRef](#)]
30. Han, X.; Zhang, F.; Meng, L.; Xu, Y.; Li, Y.; Li, A.; Turyk, M.E.; Yang, R.; Wang, P.; Zhang, J.; et al. Exposure to organochlorine pesticides and the risk of type 2 diabetes in the population of East China. *Ecotoxicol. Environ. Saf.* **2020**, *190*, 110125. [[CrossRef](#)]
31. Yang, J.H.; Lee, Y.M.; Bae, S.G.; Jacobs, D.R., Jr.; Lee, D.H. Associations between organochlorine pesticides and vitamin D deficiency in the U.S. population. *PLoS ONE* **2012**, *7*, e30093. [[CrossRef](#)] [[PubMed](#)]

32. Schæbel, L.K.; Bonefeld-Jørgensen, E.C.; Vestergaard, H.; Andersen, S. The influence of persistent organic pollutants in the traditional Inuit diet on markers of inflammation. *PLoS ONE* **2017**, *12*, e0177781. [[CrossRef](#)] [[PubMed](#)]
33. Andrews, J.E.; Courtney, K.D.; Donaldson, W.E. Impairment of calcium homeostasis by hexachlorobenzene (HCB) exposure in Fischer 344 rats. *J. Toxicol. Environ. Health* **1988**, *23*, 311–320. [[CrossRef](#)] [[PubMed](#)]
34. Brennan, E.; Kumar, N.; Drage, D.S.; Cunningham, T.K.; Sathyapalan, T.; Mueller, J.F.; Atkin, S.L. A case-control study of polychlorinated biphenyl association with metabolic and hormonal outcomes in polycystic ovary syndrome. *J. Environ. Sci. Health C Toxicol. Carcinog.* **2022**, *40*, 86–105. [[CrossRef](#)] [[PubMed](#)]
35. Javed, Z.; Papageorgiou, M.; Deshmukh, H.; Kilpatrick, E.S.; Mann, V.; Corless, L.; Abouda, G.; Rigby, A.S.; Atkin, S.L.; Sathyapalan, T. A Randomized, Controlled Trial of Vitamin D Supplementation on Cardiovascular Risk Factors, Hormones, and Liver Markers in Women with Polycystic Ovary Syndrome. *Nutrients* **2019**, *11*, 188. [[CrossRef](#)]
36. Holick, M.F.; Binkley, N.C.; Bischoff-Ferrari, H.A.; Gordon, C.M.; Hanley, D.A.; Heaney, R.P.; Murad, M.H.; Weaver, C.M. Evaluation, treatment, and prevention of vitamin D deficiency: An Endocrine Society clinical practice guideline. *J. Clin. Endocrinol. Metab.* **2011**, *96*, 1911–1930. [[CrossRef](#)]
37. Moin, A.S.M.; Sathyapalan, T.; Butler, A.E.; Atkin, S.L. Vitamin D association with coagulation factors in polycystic ovary syndrome is dependent upon body mass index. *J. Transl. Med.* **2021**, *19*, 239. [[CrossRef](#)]
38. Moin, A.S.M.; Sathyapalan, T.; Butler, A.E.; Atkin, S.L. Coagulation factor dysregulation in polycystic ovary syndrome is an epiphenomenon of obesity. *Clin. Endocrinol.* **2023**, *98*, 796–802. [[CrossRef](#)]
39. Birkett, M.A.; Day, S.J. Internal pilot studies for estimating sample size. *Stat. Med.* **1994**, *13*, 2455–2463. [[CrossRef](#)]
40. Siddharth, M.; Datta, S.K.; Bansal, S.; Mustafa, M.; Banerjee, B.D.; Kalra, O.P.; Tripathi, A.K. Study on organochlorine pesticide levels in chronic kidney disease patients: Association with estimated glomerular filtration rate and oxidative stress. *J. Biochem. Mol. Toxicol.* **2012**, *26*, 241–247. [[CrossRef](#)]
41. Lv, J.; Guo, L.; Gu, Y.; Xu, Y.; Xue, Q.; Yang, X.; Wang, Q.N.; Meng, X.M.; Xu, D.X.; Pan, X.F.; et al. National temporal trend for organophosphate pesticide DDT exposure and associations with chronic kidney disease using age-adapted eGFR model. *Environ. Int.* **2022**, *169*, 107499. [[CrossRef](#)] [[PubMed](#)]
42. Pavlíková, N.; Daniel, P.; Šrámek, J.; Jelínek, M.; Šrámková, V.; Němcová, V.; Balušíková, K.; Halada, P.; Kovář, J. Upregulation of vitamin D-binding protein is associated with changes in insulin production in pancreatic beta-cells exposed to p,p'-DDT and p,p'-DDE. *Sci. Rep.* **2019**, *9*, 18026. [[CrossRef](#)]
43. Ellison, T.I.; Dowd, D.R.; MacDonald, P.N. Calmodulin-dependent kinase IV stimulates vitamin D receptor-mediated transcription. *Mol. Endocrinol.* **2005**, *19*, 2309–2319. [[CrossRef](#)] [[PubMed](#)]
44. Nowicki, H.G.; Myrtle, J.F.; Norman, A.W. Effects of organochlorine insecticides on metabolism of cholecalciferol (vitamin D₃) in Rachitic cockerel. *J. Agric. Food Chem.* **1972**, *20*, 380–384. [[CrossRef](#)] [[PubMed](#)]
45. Siddiquee, M.H.; Bhattacharjee, B.; Siddiqi, U.R.; MeshbahurRahman, M. High prevalence of vitamin D deficiency among the South Asian adults: A systematic review and meta-analysis. *BMC Public Health* **2021**, *21*, 1823. [[CrossRef](#)]
46. Shah, H.K.; Sharma, T.; Banerjee, B.D. Organochlorine pesticides induce inflammation, ROS production, and DNA damage in human epithelial ovary cells: An in vitro study. *Chemosphere* **2020**, *246*, 125691. [[CrossRef](#)]
47. Liu, Z.; Lu, Y.; Zhong, K.; Wang, C.; Xu, X. The associations between endocrine disrupting chemicals and markers of inflammation and immune responses: A systematic review and meta-analysis. *Ecotoxicol. Environ. Saf.* **2022**, *234*, 113382. [[CrossRef](#)]
48. Dzik, K.P.; Kaczor, J.J. Mechanisms of vitamin D on skeletal muscle function: Oxidative stress, energy metabolism and anabolic state. *Eur. J. Appl. Physiol.* **2019**, *119*, 825–839. [[CrossRef](#)]
49. Querfeld, U. Vitamin D and inflammation. *Pediatr. Nephrol.* **2013**, *28*, 605–610. [[CrossRef](#)]
50. Wei, X.; Li, X.; Du, J.; Ge, X.; Sun, Y.; Li, X.; Xun, Z.; Liu, W.; Wang, Z.Y.; Li, Y.C. Vitamin D Deficiency Exacerbates Colonic Inflammation Due to Activation of the Local Renin-Angiotensin System in the Colon. *Dig. Dis. Sci.* **2021**, *66*, 3813–3821. [[CrossRef](#)]
51. Wang, Y.Q.; Geng, X.P.; Wang, M.W.; Wang, H.Q.; Zhang, C.; He, X.; Liang, S.M.; Xu, D.X.; Chen, X. Vitamin D deficiency exacerbates hepatic oxidative stress and inflammation during acetaminophen-induced acute liver injury in mice. *Int. Immunopharmacol.* **2021**, *97*, 107716. [[CrossRef](#)] [[PubMed](#)]
52. Glinski, A.; Liebel, S.; Pelletier, È.; Voigt, C.L.; Randi, M.A.; Campos, S.X.; Oliveira Ribeiro, C.A.; Filipak Neto, F. Toxicological interactions of silver nanoparticles and organochlorine pesticides in mouse peritoneal macrophages. *Toxicol. Mech. Methods* **2016**, *26*, 251–259. [[CrossRef](#)] [[PubMed](#)]
53. Massawe, R.; Drabo, L.; Whalen, M. Effects of pentachlorophenol and dichlorodiphenyltrichloroethane on secretion of interferon gamma (IFN γ) and tumor necrosis factor alpha (TNF α) from human immune cells. *Toxicol. Mech. Methods* **2017**, *27*, 223–235. [[CrossRef](#)] [[PubMed](#)]
54. McMullin, J.L.; Codner, J.; Patel, S.G.; Sharma, J.; Hu, X.; Jones, D.P.; Weber, C.J.; Saunders, N.D. Environmental Chemicals and their Association with Hyperparathyroidism. *World J. Surg.* **2023**, *47*, 296–303. [[CrossRef](#)] [[PubMed](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.