



Dermatology Reports

<https://www.pagepress.org/journals/index.php/dr/inde>

X

eISSN 2036-7406



Publisher's Disclaimer. E-publishing ahead of print is increasingly important for the rapid dissemination of science. **Dermatology Reports** is, therefore, E-publishing PDF files of an early version of manuscripts that undergone a regular peer review and have been accepted for publication, but have not been through the copyediting, typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one. The final version of the manuscript will then appear on a regular issue of the journal. E-publishing of this PDF file has been approved by the authors.

Please cite this article as: Mortato E, Baratta S, Rubino L, et al. **Cost analysis in the management of moderate-to-severe psoriasis: comparison between conventional and biological systemic therapies.** *Dermatol Rep* 2023 [Epub Ahead of Print] doi: 10.4081/dr.2023.9755

 © the Author(s), 2023
Licensee PAGEPress, Italy

Submitted: 25/05/2023 – Accepted 06/06/2023

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Cost analysis in the management of moderate-to-severe psoriasis: comparison between conventional and biological systemic therapies

Edoardo Mortato, Silvia Baratta, Luca Rubino, Anna Paola De Caro, Francesco Loconsole

Clinica Dermatologica Azienda Ospedaliera Universitaria Policlinico Consorziale Bari, Italy

Correspondence: Francesco Loconsole, Clinica Dermatologica Azienda Ospedaliera Universitaria Policlinico Consorziale Bari Piazza Giulio Cesare, 11, 70124, Bari, Italy.

Tel.: +39.0805591111.

E-mail: franciscus59@gmail.com

Key words: antirheumatic agents, biological therapy, costs and cost analysis, psoriasis.

Acknowledgments: editorial support, provided by Health Publishing and Services Srl, was funded by Novartis Farma SpA.

Contributions: all authors contributed equally.

Conflict of interest: the authors declare no potential conflict of interest.

Funding: none.

Ethical approval and consent to participate: not applicable.

Data availability: the data presented in this study are available on reasonable request from the corresponding author.

Abstract

Background. Psoriasis is a chronic inflammatory skin disease with an important socio-economic burden. Available therapies include conventional systemic drugs and biological drugs, such as Tumor Necrosis Factor (TNF)- α inhibitors, that are characterized by high costs.

Aim. Perform a cost-estimation analysis of conventional treatment vs therapy with biosimilar TNF- α inhibitor between January 2021 and January 2022, according to the Apulia regional cost list.

Methods. The average annual expenditure per patient on conventional treatment (cyclosporine and methotrexate) vs therapy with biosimilar TNF- α inhibitor was compared. The 'cost per responder' was determined by analyzing the percentages of 'responders' (patients achieving PASI 75 and PASI 90) and 'non-responders' (PASI <75) after one year of treatment.

Results. The annual per capita expenditure with cyclosporin was €3,515.35, with methotrexate was €1,048.87, while for treatment with TNF- α biosimilar inhibitor drug was €3,030.11. The "cost per responder" analysis showed a value of €8,573 for cyclosporine, €2,834 for methotrexate, and €3,564 for TNF- α biosimilar inhibitor.

Conclusions. Conventional drugs have a greater impact on healthcare expenditure than TNF- α biosimilar inhibitors.

To the Editor:

Psoriasis is a chronic inflammatory skin disease with an estimated worldwide prevalence of 0.5% to 11.4% in adults and up to 1.4% in children.¹

It is a disease characterized by erythematous, infiltrated, itchy and often painful skin lesions. Recent advances in the scientific community have enabled a greater understanding of the predisposing genetic factors, pathophysiology, comorbidities and treatment of psoriasis. Currently available therapies for moderate-severe psoriasis [Psoriasis Area Severity Index (PASI) greater than 10] include Disease-modifying antirheumatic drugs (DMARDs)/conventional systemic drugs [cyclosporine, methotrexate (mtx) and acitretin] and systemic biological drugs, such as Tumor Necrosis Factor (TNF)- α inhibitors and Interleukin (IL) inhibitors. According to the Official Bulletin of the Apulia Region no. 149 of 20-11-2018 about the rationalization of pharmaceutical expenditure, interventions aimed at increasing the prescriptive appropriateness of high-cost systemic immunosuppressive drugs for authorized uses in moderate-to-severe plaque psoriasis were outlined, on the basis of the update guidelines in literature. DMARDs/conventional systemic drugs (cyclosporine, mtx, acitretin) were chosen as first line of treatment, systemic biological TNF- α inhibitors drugs as second line of treatment, and systemic biological IL inhibitors drugs as third line.

We performed a cost estimation according to the regional cost list² and we compared the average annual expenditure per patient on conventional DMARDs treatment vs. therapy with biosimilar TNF- α inhibitor, considering the costs of outpatient controls, pre-treatment and follow-up haematochemical/instrumental examinations, and lost working days compared to the different therapies. In the group of conventional DMARDs, we only considered cyclosporine and mtx. Acitretin, due to its limited therapeutic indications, was not relevant to our study.

We also divided the first 40 consecutive naive patients admitted between January 2021 and January 2022 into 2 groups similar for age (21-73) and gender (M:F). The first group consisted of 20 patients treated with conventional systemic therapy (12 on cyclosporine therapy and 8 on mtx therapy); the second group of 20 patients was treated with biosimilar TNF- α inhibitor, of which 8 were previously treated with conventional DMARDs and 12 naive due to contraindications/intolerance to conventional systemic therapies. Within each group, we analyzed the percentage of 'responders' (patients achieving PASI 75 and PASI 90) and 'non-responders' (patients achieving PASI <75) after one year of treatment. We calculated the 'cost per responder', *i.e.*, the measure of the effectiveness of healthcare technology, by dividing the total expenditure of each drug (cyclosporine, mtx, TNF- α inhibitor) by the percentage number of 'responders' patients.

We also analyzed the Number Needed to Treat (NNT), which is an absolute effect measure representing the number of patients who need to be treated to obtain a therapeutic benefit (responders).

The NNT corresponds to the reciprocal of the absolute risk reduction (ARR), where the ARR coincides with the difference in the incidence of an event between the experimental group [experimental event rate (EER) in our analysis corresponds to the population on treatment with anti-TNF- α biosimilar] and the control group (control event rate, CER - in our case represented by the population of patients on DMARDs therapy), as shown in Table 1.

In general, the NNT represents the number of patients who need to be treated with drug A (in our analysis corresponding to anti-TNF- α biosimilar) versus drug B (in our analysis corresponding to DMARDs) in order to achieve a therapeutic benefit (in our analysis: attainment of PASI 75 and PASI 90). It is worth mentioning that the lower the NNT is, the greater the efficacy of the intervention is, compared to the selected comparators³.

According to our cost estimate, the annual per capita expenditure for treatment with cyclosporin was €3,515.35; the one for treatment with mtx was €1,048.87. Annual per capita expenditure for treatment with systemic biological TNF- α biosimilar inhibitor drug was €3,030.11 (Table 2).

We observed 7 "non-responders" (60%) among the 12 patients treated with cyclosporine, 5 "non-responders" (62.5%) among the 8 patients treated with mtx, and 3 "non-responders" (15%) among

the 20 patients treated with anti-TNF- α biosimilar. Considering the total of 20 patients treated with DMARDs, the number of 'non-responders' becomes 12 (Table 3). The analysis of "cost per responder" showed a value of €8,573 for cyclosporine, €2,834 for mtx, and €3,564 for TNF- α biosimilar inhibitor.

The NNT between TNF- α biosimilar inhibitor and DMARDs showed a value of 6.6 for "responder-PASI 75", while reported a value of 3.3 for "responder-PASI 90 (Table 1).

We conclude that despite conventional DMARDs being among the first-line therapies for moderate-severe psoriasis, mainly because of their economic implications, our cost estimation shows that they have a greater impact on healthcare expenditure than TNF- α biosimilar inhibitors, which are more manageable, more effective and even more cost-effective than their predecessors. The major limitations of this study are the lack of randomization and the small number of patients included. Prospective studies on larger series will be needed in order to validate the results shown in this analysis.

References

1. Michalek IM, Loring B, John SM. A systematic review of worldwide epidemiology of psoriasis. *J Eur Acad Dermatol Venereol* 2017;31:205-12.
2. Delibera giunta regionale n°951, 13 Maggio 2013.
3. Ravasio RCA, Antonelli S, Maiorino A, Losi S. Number needed to treat for interleukin inhibitors approved for the treatment of moderate-to-severe plaque psoriasis in Italy. *Grhta* 2021;15:8:53-7.

Table 1. Analysis of the Number Needed to Treat (NNT) between TNF- α biosimilar inhibitor and DMARDs.

PASI75	$NNT = \frac{1}{ARR} = \frac{1}{EER - CER} = \frac{1}{PASI\ 75A - PASI\ 75B}$ $NNT = \frac{1}{PASI\ 75A - PASI\ 75B} = \frac{1}{0.4 - 0.25} = \frac{1}{0.15}$ $= 6.6$
---------------	---

PASI90	$NNT = \frac{1}{ARR} = \frac{1}{EER - CER} = \frac{1}{PASI\ 90A - PASI\ 90B}$ $NNT = \frac{1}{PASI\ 90A - PASI\ 75B} = \frac{1}{0.45 - 0.15} = \frac{1}{0.3}$ $= 3.3$

Determination of NNT value to obtain PASI 75 or PASI 90 in patients treated with either drug A or drug B; NNT, Number Needed to Treat; ARR, absolute risk reduction; CER, experimental event rate; EER, control event rate; PASI, Psoriasis Area Severity Index; A, TNF-α biosimilar inhibitor; B=DMARDs.

Table 2. Per capita annual treatment cost per drug.

Drug	Per capita annual treatment cost
Cyclosporine	€3,515.35
Mtx	€1,048.87
TNF-α biosimilar inhibitor	€3,030.11

Cost per capita for annual treatment with Cyclosporine, Metotrexate (Mtx) and a TNF-α biosimilar inhibitor.

Table 3. Percentage of responders and non-responders per drug.

Drug (N patients)	PASI 90 (%)	PASI 75 (%)	Non responders (%)
Cyclosporine (12)	8	32	60
Mtx (8)	25	12.5	62.5
TNF-α biosimilar inhibitor (20)	45	40	15

DMARDs (20)	15	25	60
-------------	----	----	----

Percentage of PASI 90, PASI 75 responders and non-responders in the treated groups. PASI, Psoriasis Area Severity Index; DMARDs, Disease-modifying antirheumatic drugs; Mtx, Metotrexate.