



OPEN ACCESS

EDITED BY

Ranadip Chowdhury,
Society for Applied Studies, India

REVIEWED BY

Sitanshi Sharma,
Center for Health Research and Development,
Society for Applied Studies, India
Rukman Manapurath,
Center for Health Research and Development,
Society for Applied Studies, India

*CORRESPONDENCE

Diana Sunardi

✉ diana_sunardi@yahoo.com

RECEIVED 20 February 2023

ACCEPTED 09 June 2023

PUBLISHED 26 June 2023

CITATION

Sunardi D, Wibowo Y, Mak TN and Wang D (2023) Micronutrients intake inadequacy in different types of milk consumers in Indonesian children 1-5 years: dietary modeling with young child milk improved.

Front. Nutr. 10:1169904.

doi: 10.3389/fnut.2023.1169904

COPYRIGHT

© 2023 Sunardi, Wibowo, Mak and Dantong. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Micronutrients intake inadequacy in different types of milk consumers in Indonesian children 1-5 years: dietary modeling with young child milk improved

Diana Sunardi^{1*}, Yulianti Wibowo², Tsz Ning Mak³ and Dantong Wang³

¹Department of Nutrition, Faculty of Medicine, Universitas Indonesia – Cipto Mangunkusumo Hospital, Jakarta, Indonesia, ²Medical Nutrition Services, Nestle Indonesia, Jakarta, Indonesia, ³Nestle Institute of Health Sciences, Singapore Hub, Nestle Research, Singapore, Singapore

Background: Indonesian children under-five have a high prevalence of micronutrient deficiencies. Improving young child feeding practices may be the solution. Increasing the consumption of appropriate milk products could help to reduce nutrient inadequacy.

Methods: The objective of this study was to assess nutrient inadequacy in Indonesian children to evaluate the potential improvement using dietary modeling analysis. Data from children aged 1–5 years from the Indonesian Individual Dietary Consumption Survey in 2014 were used in this analysis ($n=11,020$). Diet modeling was conducted in two scenarios, substitution volume to volume and calories to calories.

Results: The proportion of children consuming young child milk (YCM) was the highest compared to other milk types across all age groups, followed by condensed milk and cow's milk. YCM, also called "Growing-Up Milk" (GUM), are marketed as a product specifically formulated for the nutritional needs of young children. YCM consumers had lower prevalence of inadequate intakes in iron, zinc, vitamins A, C and D across age groups when compared to condensed milk consumers. The prevalence of inadequate intakes of nutrients in condensed milk consumers was Vitamin A (67, 64%), folate (92, 91%), Vitamin D (87, 84%), iron (84, 76%), and zinc (76, 76%) in 1–2y and 3–4y, respectively. The substitution of condensed milk with a YCM reduced the prevalence of inadequate intakes of micronutrients, such as Vitamin A, vitamin D, folate, iron and zinc, which are important for immune function. YCM reduced the prevalence of inadequate intakes of micronutrients by 20–40% (Vit A and folate) and 40–50% (Vit D and zinc). The reduction of prevalence of inadequate iron intake was 31% in 1–2y and 63% in 3–4y.

Conclusion: The prevalence of inadequate micronutrient intakes was high among children aged 1–5 years old in Indonesia. YCM consumers had better nutrient intake. The substitution of condensed milk with a YCM reduced the prevalence of inadequate of micronutrient intake. Thus, nutrient intakes could be improved by YCM consumption in 1–5 years old children in Indonesia, along with nutrition education on feeding practices.

KEYWORDS

nutrient intake, children under-five, young child milk, dietary modeling, inadequacy

1. Background

Indonesia has achieved several milestones in its developmental journey to a middle-income country. Decreases in children's deaths and significant increases in primary school enrolment are among them. However, improvement in children's nutritional status, has not been observed. Millions of Indonesian children and adolescents remain threatened by staggering rates of stunting and wasting and the 'double burden' of malnutrition where undernutrition and overnutrition co-exist. While poverty contributes to malnutrition, inadequate knowledge, and practices of childcaring and child feeding also sustain the high rates of malnutrition (1). Indonesia has made some progress toward achieving the target for stunting and wasting; however, 30.8 and 10.2% of children under 5 years of age were affected (2). Better dietary intake is one of the key aspects to improve the situation. A systematic review showed that in children under-5 years in Indonesia, micronutrient intake varied, for example iron ranged from 2.40 to 12.8 mg/d, zinc ranged from 1.20 to 5.53 mg/d, calcium ranged from 83 to 707.9 mg/d, vitamin A ranged from 53 to 1,335 ug/d, and vitamin D ranged 2.75–14.10 ug/d (3). Prevalence of micronutrient deficiency in children under-five was also reported in a national survey, such as anemia (urban 17.6% vs. rural 18.5%), iron deficiency (urban 4.6% vs. rural 8.8%), and vitamin D deficiency (urban 43% vs. rural 44.2%) (4). Although several good practices have been put in place, some critical ones are not fully implemented. These are related to dietary diversity, healthy feeding and hygiene practices. Poor feeding practices result in suboptimal dietary intake. The available data suggest that children's consumption of nutrient rich foods was limited (5) and feeding practices are not optimal among children aged 6 months and above (6). Improving feeding practices is therefore crucial to tackle malnutrition among Indonesian young children.

Milk and dairy products are nutrient-dense foods, supplying energy and high-quality protein with a range of essential micronutrients (especially calcium, magnesium, potassium, zinc, and phosphorus) in an easily absorbed form for normal growth and development in children (7). Increasing the consumption of milk products among children may help to reduce nutrient inadequacy. To test this hypothesis, dietary modeling is a useful tool for assessing the theoretical nutritional impact of increasing dairy consumption in the study population. Dietary modeling is a statistical technique that has been increasingly applied to inform the development of dietary guidelines worldwide. In 2020, Dietary Guidelines for Americans Advisory Committee defined it as a method to understand how changes to the amounts or types of foods and beverages in a diet might impact meeting nutrient needs (8). Using dietary modeling, a study in Filipino children 1–5 years old suggested an improvement of macro- and micronutrient intakes by adding a serving of a milk product to the diets of children who did not meet their dairy intake recommendations (9). In the present study, we analyzed the nutrient intakes in Indonesian children aged 1–5 years who consumed different types of milk and applied dietary modeling approach to investigate the potential impact on nutrient intakes. This is, to our knowledge, the first such report for Indonesia.

2. Methods

The objective of this study was to assess nutrient inadequacy in children 1–5 years old in Indonesia and to evaluate the potential

improvement in nutrient adequacy by dietary modeling analysis. Data of children aged 1–5 years from the Indonesian Individual Food Consumption Survey (IFCS) in 2014 ($n=11,020$) were used for this analysis. This Survey was part of the Indonesian Total Dietary Intake Study (SDT 2014), providing quantitative information on food intake by households and individuals (10). Inclusion criteria is all children aged 12–60 months old, with the exclusion criteria, if the demographic and dietary intake were not complete.

The energy and nutrient intakes of each child were estimated based on a 1 day 24-h dietary recall using Nutri-Survey software (EBISpro, Germany). Milk consumers were defined based on the consumption of different types of milk, including young child milk (YCM), condensed milk or cow's milk, by children that were recorded in the 24 h food recall in the SDT survey. The energy and nutrient intakes of children were compared among the different milk consumer groups defined above. The SDT 2014 classification approach of energy and protein intake status, and WHO estimated average requirements (EAR) for micronutrients were applied to define inadequate intakes in this study. A substitution approach was applied in diet modeling, that is, by substituting condensed milk with equal volume and isocaloric YCM. YCM also called "Growing-Up Milk" (GUM), are marketed as products specifically formulated for the nutritional needs of young children (11). Dancow 1+ and 3+ composition was used in the diet modeling as YCM examples for children aged 12–35 months and 36–60 months, respectively. Statistical analysis was done using SPSS software version 20. To investigate the potential difference in nutrient intake across age, the analyses were stratified by age group 12–23, 24–35, 36–47, and 48–60 months for nutrient adequacy analysis, and 12–35 and 36–60 months for dietary modeling, respectively. Differences in subject's characteristics (gender, living area, SES, and mother's education) were also tested. The Chi-square test was performed to test the statistical significance of characteristics by type of milk consumption group, Fisher's exact test was used to compare the percentages of inadequate intake in YCM consumers to those of condensed milk and cow's milk consumers, as well as to compare the prevalence of baselines to that of the modeling scenarios. The Wilcoxon signed-rank test was applied to compare the nutrient intake distribution (12).

3. Results

3.1. Subjects

This study assessed the dietary intakes of 11,020 children aged 1–5 years from the SDT survey. We found a higher proportion of children living in rural areas (57.1%) than in urban areas (42.9%), while the distribution of children by gender and age was similar. The detailed distributions of the subject's characteristics distribution are shown in Table 1.

3.2. Subjects and households' characteristics of consumers of different milk beverages

Based on the dietary recall, YCM, condensed milk and cow's milk were the top three of dairy foods consumed by the study population.

TABLE 1 Subject characteristics (n=11,020).

| Variable | Frequencies | % |
|----------|-------------|------|
| Region | | |
| Urban | 4,731 | 42.9 |
| Rural | 6,289 | 57.1 |
| Gender | | |
| Boys | 5,694 | 51.7 |
| Girl | 5,326 | 48.3 |
| Age | | |
| Months | | |
| 12–23 | 2,529 | 22.9 |
| 24–35 | 2,701 | 24.5 |
| 36–47 | 2,847 | 25.8 |
| 48–59 | 2,943 | 26.7 |

Hence, in this study, subjects were classified into four groups namely non-milk consumers ($n=6,283$), YCM consumers ($n=3,138$), condensed milk consumers ($n=991$), and cow's milk (pasteurized milk) consumers ($n=608$; Table 2). Approximately half of the children (46–57%) did not consume any milk product between 12 and 35 months and 48–60 months. However, only 18% of the children did not consume milk products between 36 and 47 months. Among milk consumers, the proportion of children who consumed YCM was the highest across age groups, followed by condensed milk and cow's milk. Children aged 24–35 and 36–47 months had, respectively, the lowest (33.4%) and highest percentage of YCM consumption (57.7%), respectively. The proportion of children consuming condensed milk showed an increasing trend from 12–23 to 36–47 months, and slightly decreased between 48 and 60 months. The proportion of children consuming cow's milk was doubled in children aged 24–35 months compared to 12–23 months and remained stable at around 8%. A similar trend was observed for condensed milk intake (Table 2).

Overall, 67% of the non-milk consumers were lived in rural areas. Compared to rural areas, a higher proportion of children from urban areas consumed YCM (60%) and cow's milk (54%), whereas condensed milk was mostly consumed by children in rural areas (56%). The type of milk consumption differed by socioeconomic status (SES). In this study, we found that a higher proportion of non-milk consumers were from lower SES families, and that a higher proportion of children from higher SES families consumed YCM. Mothers of children in the non-milk consumer group tended to have lower education levels than those of the milk consumer groups. Conversely, mothers of children in the YCM consumer group tended to have higher educational levels than those in other milk consumer groups (Table 2).

3.3. Nutrient intake according to types of milk consumption

More than 80% of the children were below the recommended energy intake in this population, and no difference was found among children who consumed different types of milk (not shown). YCM consumers had a lower prevalence of inadequate intake of iron, zinc, vitamins A, C, and D across age groups when compared to non-milk

consumers, and condensed milk consumers. When compared to cow's milk consumers, the prevalence of inadequate intake of iron, zinc, folate, vitamins C and D was lower between 12 and 23 months, but no significant difference was found in the older groups (Figure 1).

3.4. Diet modeling

Condensed milk was the second-most consumed milk in the study population. To investigate the potential impact of YCM consumption compared to condensed milk consumption, a diet modeling analysis was performed using a substitution approach. Two scenarios were evaluated among condensed milk consumers. First, substitution of condensed milk with the YCM, on an equal volume basis. Second, substitution of condensed milk with isocaloric YCM. For children aged 12–35 months and 36–60 months, Dancow 1+ and Dancow 3+ compositions were used in the analysis, respectively, as described in the Method. Tables 3, 4 showed that in both scenarios, the substitution of condensed milk with YCM reduced the prevalence of inadequate intakes of micronutrients, such as Vit A, iron, folate and zinc that are important for the maintenance of normal immune function.

4. Discussion

There are studies on children's nutrient intakes in Indonesia, but an adequate summary of nutrient inadequacies was challenging for Arini and colleagues to provide, due to the heterogeneity in subject characteristics among studies (3). In this study, data were derived from Indonesian 2014 Food Intake Survey, a national representative survey. This is an appropriate dataset to assess the nutrient intakes in Indonesia with 11,020 children aged 1–5 years, with representative distribution in location, gender, and age groups (10).

Meeting nutritional requirements is often a challenge, especially in the period in which children's diets become diversified due to the introduction of solid foods (13). In general, milk and milk products should be important parts of children's diet as they are rich in macronutrients and micronutrient such as protein, calcium, magnesium, phosphorus, zinc, potassium, vitamin D, vitamin A, riboflavin, and vitamin B-12. The SEANUTS study (The Southeast Asian Nutrition Survey), conducted in children 1–12 years of age, found that dairy consumption was positively associated with nutritional status and consumers were less likely to be deficient in vitamin A or insufficient in vitamin D (14).

However, there are various types of milk available in the market and their nutritional value varies considerably (15). Condensed milk is consumed by people at different ages and can be found easily in many parts of the world, including Indonesia. Condensed milk is a dairy product in the form of a viscous liquid obtained by removing some part of the water from a the mixture of milk and sugar or by other processes so that it reaches a certain level of concentration. Sugar was then added to prevent damage to the product. This product is usually used as a topping for various desserts or is mixed with beverages. Because of the high content of sugar used as a preservative (estimated close to 50% of the total calories), normally condensed milk should not be offered to children <12 months and older as a substitute for breast milk, YCM or cow's milk. However, improper

TABLE 2 Subjects and household characteristic by type of milk consumption.

| Characteristic | | Non-milk consumer | YCM | Condensed milk | Cow's milk |
|------------------|--------------------------|-----------------------------|-----------------------------|--|--|
| | | <i>n</i> =6,283 | <i>n</i> =3,138 | <i>n</i> =991 | <i>n</i> =608 |
| Gender | Boys | 51.7% | 50.7% | 53.4% | 52.6% |
| | Girl | 48.3% | 49.3% | 46.6% | 47.4% |
| | <i>p</i> -value | <i>P</i> _b >0.05 | <i>P</i> _a >0.05 | <i>P</i> _a >0.05, <i>P</i> _b >0.05 | <i>P</i> _a >0.05, <i>P</i> _b >0.05 |
| Age | 12–23 m | 54.80% | 35.3% | 6.5% | 3.4% |
| | 24–35 m | 45.80% | 33.4% | 13.5% | 7.3% |
| | 36–47 m | 18.50% | 57.7% | 15.5% | 8.3% |
| | 47–59 m | 56.80% | 21.9% | 13.3% | 8.0% |
| | <i>P</i> -value | <i>P</i> _b <0.01 | <i>P</i> _a <0.01 | <i>P</i> _a <0.01, <i>P</i> _b <0.01 | <i>P</i> _a <0.01, <i>P</i> _b <0.01 |
| Location | Urban | 33% | 60% | 44% | 54% |
| | Rural | 67% | 40% | 56% | 46% |
| | <i>P</i> -value | <i>P</i> _b <0.01 | <i>P</i> _a <0.01 | <i>P</i> _a <0.01, <i>P</i> _b <0.01 | <i>P</i> _a <0.01, <i>P</i> _b <0.01 |
| SES | Low | 69.6% | 40.0% | 60.9% | 48.4% |
| | Middle | 27.2% | 46.7% | 34.8% | 43.2% |
| | High | 3.3% | 14.4% | 4.3% | 8.4% |
| | <i>P</i> -value | <i>P</i> _b <0.01 | <i>P</i> _a <0.01 | <i>P</i> _a <0.01, <i>P</i> _b <0.01 | <i>P</i> _a <0.01, <i>P</i> _b <0.01 |
| Mother education | No education | 6.2% | 3.1% | 4.1% | 3.1% |
| | Not graduated elementary | 12.4% | 8.2% | 9.2% | 8.2% |
| | Elementary | 34.0% | 25.5% | 34.7% | 33.7% |
| | Junior high | 20.6% | 20.4% | 23.5% | 21.4% |
| | Senior high | 20.6% | 30.6% | 21.4% | 25.5% |
| | Diploma | 3.1% | 5.1% | 4.1% | 4.1% |
| | University | 3.1% | 7.1% | 4.1% | 4.1% |
| | <i>P</i> -value | <i>P</i> _b <0.01 | <i>P</i> _a <0.01 | <i>P</i> _a <0.01, <i>P</i> _b <0.01 | <i>P</i> _a <0.01, <i>P</i> _b <0.01 |

Chi-square test was performed to test the statistical significance among the groups. *P*_a, non-milk consumer group was used as the reference group; *P*_b, YCM group was used as the reference group.

utilization of condensed milk has been reported previously in Indonesia (15–19). A study in Yogyakarta (a province in Indonesia) showed that, although milk consumption among preschool children was dominated by YCM (48.6%), there was still a significant proportion of children consuming condensed milk (29.3%) (14). Our results were similar to those of the above study, in that the highest proportion of milk consumers in 1–5 years old children consumed YCM, followed by condensed milk and cow's milk.

The ESPGHAN Committee on Nutrition suggested that YCM can be considered as an option to increase nutrient intakes (20). A study in 300 healthy children aged 1–3 years in Jakarta observed that a daily consumption of two services YCM (≥ 300 mL/day) could be considered as a potential preventive measure for stunting in children (adjusted OR=0.28, 95%CI 0.13–0.63) (21). The YCM was reported as a beneficial contributor of bioavailable calcium and potassium, and had a specific stimulating effect on linear growth and IGF-1 (insulin-like growth factor 1) levels in the children (22, 23). The study in Yogyakarta also reported the benefit of YCM consumption in a population where the prevalence of iron (low ferritin), zinc, and iodine deficiencies, vitamin D insufficiency and deficiency were still high, and the lowest prevalence of nutrient deficiencies was found in YCM consuming group (15). The result of the present analysis support the Yogyakarta study. We also found that YCM consumers had lower prevalence of inadequate intakes of iron, zinc, vitamins A, C and D across age groups

when compared to condensed milk consumers. Substitution of condensed milk with YCM in our modeling analysis resulted in a significant increase of nutrient intake, showing that low nutrient intake could be significantly improved by changing the low-nutrient dense milk (condensed milk) with YCM. The results of this modeling were aligned with those of studies in the UK, China and the Philippines (12, 23, 24). The simulation scenarios study in the UK suggested more specifically that the consumption of YCM, instead of whole cow's milk, might lead to improvements in nutrient intake, particularly for protein, fat quality, vitamin D and iron (24). The study conducted in China showed that YCM contributed significantly to the nutrient intakes of its consumers and the YCM consumers had higher nutrient intakes than non-consumers in many nutrients, such as iron, calcium, vitamin C, vitamin D, vitamin E, vitamin B6, and folate, all of which are important for growth and immune function (12). The dietary modeling in children aged 1–5 years in the Philippines, showed an improvement in minerals and vitamin intakes especially when a serving of YCM was added to child total diet (25). Finally, a randomized controlled trial study in New Zealand showed that, compare to unfortified cow's milk consumers, YCM consuming group had lower prevalence of iron and vitamin D deficiencies in children aged 1–2 years (26). Results of these studies all point to the same direction, i.e., diet quality can be improved by YCM consumption. These studies indicate the importance of choosing the right milk, suggesting potential measures to ameliorate




FIGURE 1
Percent of children with inadequate micronutrient intakes by types of dairy products in each age group. YCM group was used as the reference group. Fisher exact test was performed to compare the percentages of inadequate intakes in YCM consumers to that of condensed milk and cow's milk consumers. * $p < 0.05$ after applying Bonferroni correction for multiple comparison.

TABLE 3 Modeling for nutrient intakes and inadequacy among condensed milk consumers (1–2y, $N=507$).

| | | Mean | SD | P25 | P50 | P75 | p1 | Reference values (12–35m) | % Children below reference value | p2 |
|---------|------------|-------|-------|-------|-------|-------|-------|---------------------------|----------------------------------|-------|
| Vit. A | Baseline | 319.5 | 671.2 | 17.6 | 123.2 | 401.2 | | 286 | 67% | |
| | Scenario 1 | 458.1 | 687.6 | 161.7 | 277.1 | 528.1 | <0.01 | 286 | 52% | <0.01 |
| | Scenario 2 | 467.9 | 710.8 | 168.5 | 283.4 | 532.8 | <0.01 | 286 | 50% | <0.01 |
| Folate | Baseline | 46.5 | 66.4 | 9.6 | 26.1 | 54.7 | | 120 | 92% | |
| | Scenario 1 | 104.5 | 95.6 | 48.3 | 76.4 | 125.9 | <0.01 | 120 | 73% | <0.01 |
| | Scenario 2 | 106 | 98.2 | 49 | 76.8 | 127.8 | <0.01 | 120 | 73% | <0.01 |
| Vit. D | Baseline | 2.1 | 4.9 | 0 | 0.3 | 2.2 | | 5 | 87% | |
| | Scenario 1 | 8.4 | 9.6 | 3.4 | 5.7 | 10.6 | <0.01 | 5 | 47% | <0.01 |
| | Scenario 2 | 8.7 | 9.9 | 3.6 | 5.8 | 11 | <0.01 | 5 | 47% | <0.01 |
| Calcium | Baseline | 335.3 | 417.3 | 146.4 | 242.7 | 402 | | 417 | 77% | |
| | Scenario 1 | 530.9 | 497 | 267.3 | 420.2 | 647.2 | <0.01 | 417 | 49% | <0.01 |
| | Scenario 2 | 544.9 | 509.3 | 275.3 | 433.9 | 653.1 | <0.01 | 417 | 49% | <0.01 |
| Iron | Baseline | 17.8 | 107.9 | 1.7 | 2.8 | 4.5 | | 5.8 | 84% | |
| | Scenario 1 | 20.6 | 107.8 | 3.6 | 5.2 | 8 | <0.01 | 5.8 | 58% | <0.01 |
| | Scenario 2 | 20.7 | 107.8 | 3.6 | 5.3 | 8.1 | <0.01 | 5.8 | 58% | <0.01 |
| Zinc | Baseline | 2.6 | 2.3 | 1.3 | 2.2 | 3.3 | | 3.4 | 76% | |
| | Scenario 1 | 4.4 | 3.2 | 2.6 | 3.7 | 5.3 | <0.01 | 3.4 | 44% | <0.01 |
| | Scenario 2 | 4.5 | 3.3 | 2.6 | 3.7 | 5.4 | <0.01 | 3.4 | 44% | <0.01 |

The intakes of different nutrients were compared to the baseline levels after substitution. Scenario 1, Substituted condensed milk with the same volume of YCM (volume to volume); Scenario 2, Substituted condensed milk with isocaloric YCM (kcal to kcal); SD, standard deviation; p1, p -value of Wilcoxon signed-rank test; p2, p -value of Fisher's exact test.

TABLE 4 Modeling for Nutrient intakes and inadequacy among condensed milk consumers (3–4y, N=797).

| | | Mean | SD | P25 | P50 | P75 | p1 | Reference values (36–47m) | Reference values (48–60m) | % Children below reference value | p2 |
|---------|------------|-------|--------|-------|-------|-------|-------|---------------------------|---------------------------|----------------------------------|-------|
| Vit. A | Baseline | 370.2 | 733.1 | 23.5 | 167.5 | 448.4 | | 286 | 321 | 63.70% | |
| | Scenario 1 | 638.7 | 1846.2 | 199.8 | 375.3 | 614.9 | <0.01 | 286 | 321 | 39.60% | <0.01 |
| | Scenario 2 | 569.2 | 868 | 196.2 | 361.8 | 605.8 | <0.01 | 286 | 321 | 40.30% | <0.01 |
| Folate | Baseline | 59.3 | 82.2 | 14.8 | 33.8 | 68.2 | | 120 | 160 | 90.70% | |
| | Scenario 1 | 155.1 | 458 | 64.4 | 101.5 | 155 | <0.01 | 120 | 160 | 68.10% | <0.01 |
| | Scenario 2 | 136.9 | 159.2 | 63.2 | 99.5 | 152.3 | <0.01 | 120 | 160 | 69.00% | <0.01 |
| Vit. D | Baseline | 2.4 | 4.3 | 0 | 0.5 | 3.2 | | 5 | 5 | 84.10% | |
| | Scenario 1 | 13.1 | 52.5 | 4.4 | 7.8 | 12.2 | <0.01 | 5 | 5 | 26.90% | <0.01 |
| | Scenario 2 | 10.9 | 16.5 | 4.2 | 7.7 | 12.1 | <0.01 | 5 | 5 | 32.00% | <0.01 |
| Calcium | Baseline | 389.9 | 481.8 | 182.8 | 278.8 | 433.2 | | 417 | 500 | 76.80% | |
| | Scenario 1 | 711.9 | 2170.1 | 328.5 | 489 | 720.2 | <0.01 | 417 | 500 | 43.40% | <0.01 |
| | Scenario 2 | 624.2 | 711.2 | 322.7 | 481.2 | 704.6 | <0.01 | 417 | 500 | 45.40% | <0.01 |
| Iron | Baseline | 24.2 | 160.5 | 2.4 | 3.7 | 5.9 | | 5.8 | 6.3 | 75.80% | |
| | Scenario 1 | 30.9 | 163.2 | 5.8 | 8.4 | 12 | <0.01 | 5.8 | 6.3 | 27.40% | <0.01 |
| | Scenario 2 | 29.6 | 160.5 | 5.7 | 8.2 | 11.9 | <0.01 | 5.8 | 6.3 | 28.60% | <0.01 |
| Zinc | Baseline | 3.1 | 3.9 | 1.6 | 2.5 | 3.9 | | 4 | 4 | 76.20% | |
| | Scenario 1 | 6.8 | 19.1 | 3.5 | 5 | 7 | <0.01 | 4 | 4 | 32.70% | <0.01 |
| | Scenario 2 | 6.1 | 6.8 | 3.4 | 4.9 | 6.8 | <0.01 | 4 | 4 | 34.90% | <0.01 |

The intakes of different nutrients were compared to the baseline levels after substitution. Scenario 1, Substituted condensed milk with the same volume of YCM (volume to volume); Scenario 2, Substituted condensed milk with isocaloric YCM (kcal to kcal); SD, standard deviation; p1, *p*-value of Wilcoxon signed-rank test; p2, *p*-value of Fisher's exact test.

micronutrient deficiency through consumption the proper type of milk and establishing healthy dietary habits.

The present study is the first to analyze the impact of different milk consumptions on nutrient intake using data from a national representative survey in Indonesia. The dietary modeling approach was also applied for the first time and the results highlighted the advantage of YCM consumption in the study population. The study had some limitations. First, this was a cross sectional study, and no causal relationship could be inferred. Second, we found differences among different milk consumer groups in dwelling location, SES, and mother education levels. A multivariate analysis is required to further reveal the influence of different milk consumption. Third, the modeling approach described a potential theoretical impact; other aspects, such as cost and eating habits, need to be considered in future studies.

5. Conclusion

In this study, we found that in children 1–5y in Indonesia, the most consumed milk type was YCM, followed by condensed milk and cow's milk, which could be related to SES. The prevalence of inadequate micronutrient intake was still high in the study population. Compared to condensed milk and cow's milk consumers, YCM consumers had improved micronutrient intake, including vit A, iron, and zinc. Substitution of condensed milk with YCM in dietary modeling showed a reduction in the prevalence of inadequate intakes of micronutrients, such as Vit A,

Vit D, iron, folate, and zinc, which are important for the maintenance of normal immune function. Thus, nutrient intake could be improved by YCM consumption in 1–5 years old children in Indonesia, along with nutrition education on food diversity.

Data availability statement

The original contributions presented in the study are included in the article/[Supplementary material](#), further inquiries can be directed to the corresponding author.

Ethics statement

Ethical approval number LB.02.01/5.2/KE.189/2014 were received from the Ethical Commission Health Research, Health Research and Development Commission in Indonesian Ministry of Health. Written informed consent to participate in this study was provided by the participants' legal guardian/next of kin.

Author contributions

DS, YW, TM, and DW participated in the proposal development and the manuscript writing. DS conducted the data analysis. All authors contributed to the article and approved the submitted version.

Funding

This study was funded by the Nestle Research. Nestle Research Center Team was involved in the study design and the development of statistical analysis plan, however, it was not involved in data analysis or result reporting.

Acknowledgments

We would like to thank the Research and Development of Health Division of the Indonesian Ministry of Health for sharing us data from Indonesian the Individual Dietary Consumption Survey in 2014 of children aged 1–5 years.

Conflict of interest

DS is a lecturer at the Faculty of Medicine, Universitas Indonesia, and a Physician Clinical Nutrition Specialist at Cipto

Mangunkusumo Hospital. YW, TM, and DW are part of Nestle Scientific team.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fnut.2023.1169904/full#supplementary-material>

References

- Unicef Indonesia. *Nutrition: tackling the 'double burden' of malnutrition in Indonesia*. (2022) Available at: [unicef.org](https://www.unicef.org).
- Global Nutrition Report: Indonesia. (2022). *The burden of malnutrition at a glance*. Available at: [globalnutritionreport.org](https://www.globalnutritionreport.org).
- Arini HRB, Hadju V, Thomas P, Ferguson M. Nutrient and food intake of Indonesian children under 5 years of age: a systematic review. *Asia Pac J Public Health*. (2022) 34:25–35. doi: 10.1177/10105395211041001
- Sandjaja S, Budiman B, Harahap H, Ernawati F, Soekatri M, Widodo Y, et al. Food consumption and nutritional and biochemical status of 0-5-12-year-old Indonesian children: the SEANUTS study. *Br J Nutr*. (2013) 110:S11–20. doi: 10.1017/S0007114513002109
- Southeast Asian Ministers of Education Organization (SEAMEO) - Tropical Medicine and Public Health Network (TROPMED) Regional Centre for Community Nutrition (RCCN) University of Indonesia. *Health and nutritional status among children under-five in selected sub-districts of Mentawai and Nias Islands: a baseline survey for community-based health program by SurfAid international*. Jakarta: SEAMEO-TROPMED RCCN, University of Indonesia (2007).
- Blaney S, Februhartanty J, Sukotjo S. Feeding practices among Indonesian children above six months of age: a literature review on their magnitude and quality (part 1). *Asia Pac J Clin Nutr*. (2015) 24:16–27. doi: 10.6133/apjcn.2015.24.1.13
- Górska-Warsewicz H, Rejman K, Laskowski W, Czacotko M. Milk and dairy products and their nutritional contribution to the average polish diet. *Nutrients*. (2019) 11:1771. doi: 10.3390/nu11081771
- Davis KA, Esslinger K, Elvidge-Munene LA, St-Pierre S. International approaches to developing healthy eating patterns for national dietary guidelines. *Nutr Rev*. (2019) 77:388–403. doi: 10.1093/nutrit/nyy058
- Mak T-N, Angeles-Agdeppa I, Tassy M, Capanzana MV, Offord EA. Contribution of milk beverages to nutrient adequacy of young children and preschool children in the Philippines. *Nutrients*. (2020) 12:392. doi: 10.3390/nu12020392
- DDK S, Permaesih D, Lamid A, Prihatini S, Hermina MK, Puspitasari DS, et al. In: A Trihono, AB Jahari and D Kartono, editors. *Studi Diet Total: survei konsumsi makanan individu Indonesia 2014*: Lembaga Penerbitan Badan Litbangkes (2014).
- Chouraqui JP, Turck D, Tavoularis G, Ferry C, Dupont C. The role of young child formula in ensuring a balanced diet in young children (1–3 years old). *Nutrients*. (2019) 11:2213. doi: 10.3390/nu11092213
- Zhang J, Wang D, Zhang Y. Patterns of the consumption of young children formula in Chinese children aged 1–3 years and implications for nutrient intake. *Nutrients*. (2020) 12:1672. doi: 10.3390/nu12061672
- Verduci E, Di Profio E, Corsello A, Scatigno L, Fiore G, Bosetti A, et al. Which milk during the second year of life: a personalized choice for a healthy future? *Nutrients*. (2021) 13:3412. doi: 10.3390/nu13103412
- Bao KLN, Sandjaja S, Poh BK, Rojroongwasinkul N, Huu CN, Sumedi E, et al. The consumption of dairy and its association with nutritional status in the South East Asian Nutrition Surveys (SEANUTS). *Nutrients*. (2018) 10:759. doi: 10.3390/nu10060759
- Prawirohartono EP, Lestari SK, Nurani N, Sitaresmi MN. Difference in nutrient biomarkers concentration by habitual intake of milk among preschool children in an urban area of Indonesia. *J Hum Nutr Food Sci*. (2015) 3:1055.
- Hess HJ. *Condensed milk*. Reproduced from: Encyclopaedia of Food Science, Food Technology and Nutrition, Copyright 1993, Academic Press. Vevey: Nestec Ltd. (2003).
- UNICEF. *Celebrating the Innocenti declaration on the protection, promotion and support of breastfeeding: past achievements, present challenges and the way forward for infant and young child feeding* UNICEF Innocenti Research Centre Piazza SS Annunziata. Florence, Italy. (2005).
- Martha E, Amelia T, Myranti T. Toddlers' eating behavior in slum urban and semi urban communities: study in Kampung Melayu and Bantul, Indonesia. In The 1st International Conference on Global Health, KNe Life Sciences. Jakarta, Indonesia (2017); 1–7.
- Sugito FS, Djoharnas H, Darwita RR. Breastfeeding and early childhood caries (ECC) severity of children under three years old in DKI Jakarta. *Makara Kesehatan*. (2008) 12:86–91. doi: 10.7454/msk.v12i2.310
- Hojsak I, Bronsky J, Campoy C, Domellöf M, Embleton N, Fidler Mis N, et al. Young child formula: a position paper by the ESPGHAN committee on nutrition. *J Pediatr Gastroenterol Nutr*. (2018) 66:177–85. doi: 10.1097/MPG.0000000000001821
- Sjarif DR, Yuliarti K, Iskandar WJ. Daily consumption of growing-up milk is associated with less stunting among Indonesian toddlers. *Med J Indones*. (2019) 28:70–6. doi: 10.13181/mji.v28i1.2607
- FAO/WHO. *Protein quality evaluation in human diets*. Food and Agriculture Organization, Food and Nutrition Paper 51. Rome: FAO/WHO (1991).
- Michaelsen KE. Cow's milk in the prevention and treatment of stunting and wasting. *Food Nutr Bull*. (2013) 34:249–51. doi: 10.1177/156482651303400219
- Eussen SRBM, Pean J, Olivier L, Delaere F, Lluich A. Theoretical impact of replacing whole cow's milk by young-child formula on nutrient intake of UK young children: results of a simulation study. *Ann Nutr Metab*. (2015) 67:247–56. doi: 10.1159/000440682
- Tn M, Agdeppa IM, Tassy M, Capanzana MV, Offord EA. The nutritional impact of Milk beverages in reducing nutrient inadequacy among children aged one to five years in the Philippines: a dietary modelling study. *Nutrients*. (2020) 12. doi: 10.3390/nu12113330
- Lovell AL, Davies PSW, Hill RJ, Milne T, Matsuyama M, Jiang Y, et al. Compared with cow milk, a growing-up milk increases vitamin D and Iron status in healthy children at 2 years of age: the growing-up milk-lite (GUMLi) randomized controlled trial. *J Nutr*. (2018) 148:1570–9. doi: 10.1093/jn/nxy167