



EVALUATION OF POTENTIAL OF HYDROGEN (PH) OF SODIUM HYPOCHLORITE AT THREE DIFFERENT MOMENTS BEFORE ITS EXPIRATION DATE

AVALIAÇÃO DO POTENCIAL HIDROGÊNIO (PH) DO HIPOCLORITO DE SÓDIO EM TRÊS DIFERENTES MOMENTOS ANTES DO SEU VENCIMENTO

EVALUACIÓN DEL POTENCIAL DE HIDRÓGENO (PH) DEL HIPOCLORITO DE SODIO EN TRES MOMENTOS DIFERENTES ANTES DE SU FECHA DE CADUCIDAD

Sarah Regina dos Santos Pereira¹, Thaís Cristine Pereira², Vanessa Marques Meccatti², Lara Steffany de Carvalho², Enrico Coser Bridi¹, Luciane Dias de Oliveira², Amjad Abu Hasna³

e453110

<https://doi.org/10.47820/recima21.v4i5.3110>

RECEIVED: 03/19/2023

APPROVED: 04/19/2023

PUBLISHED: 05/03/2023

ABSTRACT

Sodium hypochlorite (NaOCl) is the most common used irrigant for root canal treatment, it was introduced into endodontics in the last half of XVIII century. the aim of this study was to evaluate the pH stability of sodium hypochlorite (of different concentrations) and saline solution at three different moments. The pH of the irrigants was measured in triplicate using a pH meter Mettler Toledo SevenEasy S20 (Mettler-Toledo AG. Analytical Instruments. Schwerzenbach, Switzerland) at three moments: S1: upon unpacking the solution bottle; S2: one month later; and S3: four months later. Data were submitted to normality tests. Then were analyzed by One-way Anova and Tukey tests by GraphPad Prism 6 (La Jolla, CA, USA) with a significance level of 5%. It was found that there was a statistically significant difference among the Sodium hypochlorite 1% and 2.5% in comparison with the saline solution and distilled water groups in S1, S2 and S3. Besides, it was found that sodium hypochlorite 1% loses its alkaline pH since the first month. However, sodium hypochlorite 2.5% was more stable after 4 months. It was concluded that The storage time has an effect on 1% sodium hypochlorite pH stability.

KEYWORDS: Sodium hypochlorite. Saline solution. Hydrogen-Ion Concentration.

RESUMO

O hipoclorito de sódio (NaOCl) é o irrigante mais utilizado no tratamento endodôntico, tendo sido introduzido na endodontia na última metade do século XVIII. O objetivo deste estudo foi avaliar a estabilidade do pH do hipoclorito de sódio (de diferentes concentrações) e solução salina em três momentos diferentes. O pH dos irrigantes foi medido em triplicata usando um medidor de pH Mettler Toledo SevenEasy S20 (Mettler-Toledo AG. Analytical Instruments. Schwerzenbach, Suíça) em três momentos: S1: ao desembalar o frasco da solução; S2: um mês depois; e S3: quatro meses depois. Os dados foram submetidos a testes de normalidade. Em seguida, foram analisados pelos testes One-way Anova e Tukey pelo GraphPad Prism 6 (La Jolla, CA, EUA) com nível de significância de 5%. Verificou-se que houve diferença estatisticamente significativa entre os grupos Hipoclorito de Sódio 1% e 2,5% em comparação com os grupos soro fisiológico e água destilada em S1, S2 e S3. Além disso, constatou-se que o hipoclorito de sódio 1% perde seu pH alcalino desde o primeiro mês. No entanto, o hipoclorito de sódio 2,5% foi mais estável após 4 meses. Concluiu-se que o tempo de armazenamento tem efeito na estabilidade do pH do hipoclorito de sódio 1%.

PALAVRAS-CHAVE Hipoclorito de sódio. Solução salina. Concentração de íons de hidrogênio.

¹ Instituto Nacional de Ensino Superior e Pós-Graduação Padre Gervásio (INAPÓS).

² Department of Biosciences and Oral Diagnosis, Institute of Science and Technology, São Paulo State University (ICT-UNESP), São José dos Campos.

³ São Paulo State University.



RECIMA21 - REVISTA CIENTÍFICA MULTIDISCIPLINAR ISSN 2675-6218

EVALUATION OF POTENTIAL OF HYDROGEN (PH) OF SODIUM HYPOCHLORITE AT THREE DIFFERENT MOMENTS BEFORE ITS EXPIRATION DATE
Sarah Regina dos Santos Pereira, Thais Cristine Pereira, Vanessa Marques Meccatti, Lara Steffany de Carvalho, Enrico Coser Bridi, Luciane Dias de Oliveira, Amjad Abu Hasna

RESUMEN

El hipoclorito de sodio (NaOCl) es el irrigante más utilizado en el tratamiento de endodoncia, habiéndose introducido en la endodoncia en la última mitad del siglo XVIII. El objetivo de este estudio fue evaluar la estabilidad del pH del hipoclorito de sodio (de diferentes concentraciones) y solución salina en tres tiempos diferentes. El pH de los irrigantes se midió por triplicado utilizando un pHmetro Mettler Toledo SevenEasy S20 (Mettler-Toledo AG. Analytical Instruments. Schwerzenbach, Suiza) en tres momentos: S1: al desempacar el frasco de solución; S2: un mes después; y S3: cuatro meses después. Los datos fueron sometidos a pruebas de normalidad. Luego, se analizaron mediante pruebas One-way Anova y Tukey por GraphPad Prism 6 (La Jolla, CA, EE. UU.) con un nivel de significancia del 5%. Se encontró que hubo una diferencia estadísticamente significativa entre los grupos de Hipoclorito de Sodio 1% y 2,5% en comparación con los grupos de solución salina y agua destilada en S1, S2 y S3. Además, se encontró que el hipoclorito de sodio al 1% pierde su pH alcalino desde el primer mes. Sin embargo, el hipoclorito de sodio al 2,5% fue más estable después de 4 meses. Se concluyó que el tiempo de almacenamiento tiene efecto en la estabilidad del pH del hipoclorito de sodio al 1%.

PALABRAS CLAVE: Hipoclorito de sódio. Solución salina. Concentración de iones de hidrogeno.

INTRODUCTION

The root canal treatment is carried-out to disinfect/prevent the endodontic infection, in which the microorganisms' role is essential. For this, cleaning and shaping (1) aim to, mechanically and chemically, eliminate the microorganisms and their subproducts (2–4). However, despite all the technological evolution in the area of endodontics, it is clear that an adequate decontamination of root canals demands the use of auxiliary chemical substances, because of the anatomy complexity and the microorganisms resistance (5–7).

Sodium hypochlorite (NaOCl) is the most common used irrigant for root canal treatment, it was introduced into endodontics in the last half of XVIII century (8). It has a wide bactericidal effect against variety of Gram-positive and Gram-negative microorganisms, (9–11) and their byproducts. like Lipopolysaccharides (LPS) and lipoteichoic acid (LTA) (10,12). It has the ability to dissolve organic matter from the root canal (13). Its effectiveness is directly related to its concentration, becoming greater as its concentration increases, being found in concentrations at 0.5%, 1%, 2.5%, 5.25%. However, the higher it is, the greater its toxic effect on periapical tissues, and may also cause incidents during its use (14).

Its major disadvantage is the storage instability (15). Thus, its effectiveness can be reduced upon increased temperature, exposure to light, transport, and long period storage. Its chemical effectiveness depends on the active chlorine content, which is influenced by concentration, exposure time, pH and temperature in which the alkaline pH maintains the stability of NaOCl solutions, and have a greater bactericidal effect when pH is close to neutral or basic (16).

Therefore, the aim of this study was to evaluate the pH stability of sodium hypochlorite (of different concentrations) and saline solution at three different moments. The null hypothesis was that the storage time have no effect on the pH stability of the irrigant.



RECIMA21 - REVISTA CIENTÍFICA MULTIDISCIPLINAR ISSN 2675-6218

EVALUATION OF POTENTIAL OF HYDROGEN (PH) OF SODIUM HYPOCHLORITE AT THREE DIFFERENT MOMENTS BEFORE ITS EXPIRATION DATE
Sarah Regina dos Santos Pereira, Thais Cristine Pereira, Vanessa Marques Meccatti, Lara Steffany de Carvalho, Enrico Coser Bridi, Luciane Dias de Oliveira, Amjad Abu Hasna

MATERIAL AND METHODS

Ph measurement

Sodium hypochlorite of 1% and 2.5 % (Asfer, São Caetano do Sul, SP, Brazil), sterilized saline solution (Eurofarma laboratórios, São Paulo, SP, Brazil), and purified water (prepared at our laboratory) were used in this study as found in table 1.

Table 1 the evaluated endodontic irrigants

Irrigant	Manufacturer
Sodium hypochlorite 1%	Asfer, São Caetano do Sul, SP, Brazil
Sodium hypochlorite 2.5 %	Asfer, São Caetano do Sul, SP, Brazil
Sterilized saline solution	Eurofarma laboratórios, São Paulo, SP, Brazil
Distilled water	Institute of science and technology, São Paulo State University.

The pH of the irrigants was measured in triplicate using a pH meter Mettler Toledo SevenEasy S20 (Mettler-Toledo AG. Analytical Instruments. Schwerzenbach, Switzerland) at three moments:

1. S1: upon unpacking the solution bottle;
2. S2: one month later;
3. S3: four months later.

The pH meter was calibrated, then was immersed in a Falcon-tube containing the irrigant. All the irrigants were stored at refrigerator at 9 °C between the intervals of measurement.

STATISTICAL ANALYSIS

Data were submitted to normality tests. Then were analyzed by One-way Anova and Tukey tests by GraphPad Prism 6 (La Jolla, CA, USA) with a significance level of 5%.

RESULTS

Inter-group comparisons

It was found that there was a statistically significant difference among the Sodium hypochlorite 1% and 2.5% in comparison with the saline solution and distilled water groups in S1, S2 and S3. (Table 1) and (Figure 1).




RECIMA21 - REVISTA CIENTÍFICA MULTIDISCIPLINAR ISSN 2675-6218

EVALUATION OF POTENTIAL OF HYDROGEN (PH) OF SODIUM HYPOCHLORITE AT THREE DIFFERENT MOMENTS BEFORE ITS EXPIRATION DATE
Sarah Regina dos Santos Pereira, Thais Cristine Pereira, Vanessa Marques Meccatti, Lara Steffany de Carvalho, Enrico Coser Bridi, Luciane Dias de Oliveira, Amjad Abu Hasna

Table 1. Mean values and statistical difference among the experimental groups at three moments of pH measurement (S1, S2 and S3) (inter-groups comparisons).

Samples	Saline solution	Distilled water	NaOCl 1%	NaOCl 2.5%
S1	5.84 (A)	6.12 (A)	12.18 (B)	12.96 (B)
S2	6.20 (A)	7.30 (A)	9.11 (B)	12.84 (C)
S3	6.80 (A)	7.21 (A)	9.00 (B)	12.89 (C)

Figure 1. pH values and statistical difference among the experimental groups at three moments of pH measurement (S1, S2 and S3). Different uppercase letters mean statistically significant difference.



INTRA-GROUPS COMPARISONS

It was found that sodium hypochlorite 1% loses its alkaline pH since the first month. However, sodium hypochlorite 2.5% was more stable after 4 months (Table 2) and (Figure 2).

Table 2. Mean values and statistical difference among the three moments of pH measurement (S1, S2 and S3) of experimental groups (intra-groups comparisons).


	S1	S2	S3
Saline Solution	5.84 (A)	6.20 (A)	6.80 (A)
Distilled water	6.12 (A)	7.30 (B)	7.21 (B)
NaOCl 1%	12.18 (A)	9.11 (B)	9.00 (B)
NaOCl 2.5%	12.96 (A)	12.84 (A)	12.89 (A)



RECIMA21 - REVISTA CIENTÍFICA MULTIDISCIPLINAR ISSN 2675-6218

EVALUATION OF POTENTIAL OF HYDROGEN (PH) OF SODIUM HYPOCHLORITE AT THREE DIFFERENT MOMENTS BEFORE ITS EXPIRATION DATE
Sarah Regina dos Santos Pereira, Thais Cristine Pereira, Vanessa Marques Meccatti, Lara Steffany de Carvalho, Enrico Coser Bridi, Luciane Dias de Oliveira, Amjad Abu Hasna

Figure 2. pH values and statistical difference among three moments of pH measurement (S1, S2 and S3) of the experimental groups. Different uppercase letters mean statistically significant difference.



DISCUSSION

It is known that the purpose of endodontic treatment is to remove organic tissues, microorganisms and their by-products presents in the root canal system (2,6,13,17). Among the irrigation protocols, the use of sodium hypochlorite (NaOCl) is clinically predominant (2,4), however its chemical instability puts the effectiveness of disinfection and tissue dissolution to the test. The present study consisted of a longitudinal analysis to evaluate the pH stability of NaOCl at different concentrations (1% and 2.5%) during the storage period, the null hypothesis being rejected.

The pH of the NaOCl solution has a direct influence on the biological effect, whether for tissue dissolution capacity or antimicrobial effect. In alkaline solutions, the hypochlorite ion (OCI^-) prevails, which exhibits greater capacity for tissue dissolution. In acidic solutions, hypochlorous acid (HOCl) is more abundant, presenting an antimicrobial effect (18,19). In the study by Nicoletti et al. (2005), it was verified that factors such as sun exposure and temperature promote the chemical instability of



RECIMA21 - REVISTA CIENTÍFICA MULTIDISCIPLINAR ISSN 2675-6218

EVALUATION OF POTENTIAL OF HYDROGEN (PH) OF SODIUM HYPOCHLORITE AT THREE DIFFERENT MOMENTS BEFORE ITS EXPIRATION DATE
Sarah Regina dos Santos Pereira, Thais Cristine Pereira, Vanessa Marques Meccatti, Lara Steffany de Carvalho, Enrico Coser Bridi, Luciane Dias de Oliveira, Amjad Abu Hasna

chlorinated solutions. Care during storage of the solutions influences the maintenance of properties and stability (15). Even more, it was reported that there is a relation between the pH, antimicrobial effect, detergent and dissolving capacity of these solutions when exposed to sunlight or high temperatures. According to Siqueira (2000) sodium hypochlorite solutions have a greater bactericidal effect when at a pH close to neutral or basic, and even if kept under refrigeration they can remain stable for 120 days (20).

According to Camargo et. Al (2008), it was found that the NaOCl solutions showed pH stability for 30 days (21). In the present study, it was found that the 1% NaOCl solution underwent a significant change in pH, with an initial value of 12.18 (when opening the package) and 9.00 at the end of 4 months. Such results raise questions regarding the effectiveness of these solutions used for cleaning and disinfecting root canals, since, at first, the pH conditions of the solution were more favorable to the dissolution of the organic tissue and, after 4 months, its conditions of pH were no longer the same, which could influence their biological actions and consequently their therapeutic efficacy. Regarding the highest concentration evaluated, NaOCl 2.5%, it was observed that there were no significant changes in the pH of the solution. This remained stable until the fourth month of analysis. Such results are in line with findings in the literature, as it is known that NaOCl instability occurs due to the decomposition of hypochlorite ions into chlorate and chlorine ions and this process directly depends on the concentration of the solution (16).

By associating physicochemical factors in the search for greater elimination of microorganisms, 2.5% NaOCl at pH 12 was able to significantly decrease the cell viability of the mixed bacterial biofilm (formed in situ), but when this solution (NaOCl 2, 5%) had an acidic pH, a more significant reduction was observed than that found at basic pH (19).

Finally, NaOCl has a great effect on the success of the root canal treatment (2,4,5,13). However, due to the fact that the present study has a longitudinal character, in which the same sample was analyzed at different times, with the objective of evaluating the influence of time, it was possible to verify the instability of the 1% NaOCl solution, and this fact is of great relevance clinic, since the biological properties of this solution may be compromised.

CONCLUSION

The storage time has an effect on 1% sodium hypochlorite pH stability.

REFERENCES

1. Abu Hasna A, khoury RD, Toia CC, Gonçalves GB, De Andrade FB, Talge Carvalho CA, et al. In vitro evaluation of the antimicrobial effect of n-acetylcysteine and photodynamic therapy on root canals infected with enterococcus faecalis. *Iran Endod J.* 2020;15(4):236–45.
2. Abu Hasna A, Ungaro DM De T, De Melo AAP, Yui KCK, Da Silva EG, Martinho FC, et al. nonsurgical endodontic management of dens invaginatus: a report of two cases. [version 1; peer review: 2 approved]. *F1000RES.* 2019 dec 2;8:2039.



RECIMA21 - REVISTA CIENTÍFICA MULTIDISCIPLINAR ISSN 2675-6218

EVALUATION OF POTENTIAL OF HYDROGEN (PH) OF SODIUM HYPOCHLORITE AT THREE DIFFERENT MOMENTS BEFORE ITS EXPIRATION DATE
Sarah Regina dos Santos Pereira, Thais Cristine Pereira, Vanessa Marques Meccatti, Lara Steffany de Carvalho, Enrico Coser Bridi, Luciane Dias de Oliveira, Amjad Abu Hasna

3. Abu Hasna A, Pereira Da Silva L, Pelegrini FC, Ferreira CLR, De Oliveira LD, Carvalho CAT. Effect of sodium hypochlorite solution and gel with/without passive ultrasonic irrigation on enterococcus faecalis, escherichia coli and their endotoxins. F1000RES. 2020 jun 24;9:642.
4. Abu Hasna A, Monteiro JB, Abreu RT, Camillo W, Nogueira Matuda AG, De Oliveira LD, et al. Effect of passive ultrasonic irrigation over organic tissue of simulated internal root resorption. Int J Dent. 2021 oct 29;2021:3130813.
5. Ali A, Bhosale A, Pawar S, kakti A, Bichpuriya A, Agwan MA. Current trends in root canal irrigation. Cureus. 2022 may 8;14(5):E24833.
6. Camargo SEA, Blanco TM, Lima RY, Rode SM, Camargo CHR. Avaliação do ph das soluções de hipoclorito de sódio 1% e 2,5% e digluconato de clorexidina 2% em função do tempo. Odonto. 2008 jun 30;16(31):85–91.
7. Carvalho CAT, Hasna AA, Carvalho AS, Vilela P das GF, Ramos L de P, Valera MC, et al. Clinical study of sodium hypochlorite, polymyxin b and limewater effect on MMP-3,-8,-9 in apical periodontitis. Braz dent j. 2020 APR;31(2):116–21.
8. Del Carpio-Perochena A, Bramante CM, De Andrade FB, Maliza AGA, Cavenago BC, Marciano MA, et al. Antibacterial and dissolution ability of sodium hypochlorite in different phs on multi-species biofilms. Clin Oral Investig. 2015 nov;19(8):2067–73.
9. Domingues N, Ramos L de P, Pereira LM, Do Rosário Estevam Dos Santos PB, Scorzoni L, Pereira TC, et al. Antimicrobial action of four herbal plants over mixed-species biofilms of candida albicans with four different microorganisms. Aust Endod J. 2022 sep 4.
10. Leonardo R de T, Leonardo MR. Aspectos atuais do tratamento da infecção endodôntica. revista da associacao paulista de cirurgioes dentistas. 2012 sep.
11. Macedo RG, Herrero NP, Wesselink P, Versluis M, Van Der Sluis L. Influence of the dentinal wall on the ph of sodium hypochlorite during root canal irrigation. J Endod. 2014 jul;40(7):1005–8.
12. Maekawa LE, Valera MC, Oliveira LD de, Carvalho CAT, Koga-Ito CY, Jorge AOC. In vitro evaluation of the action of irrigating solutions associated with intracanal medications on escherichia coli and its endotoxin in root canals. J Appl Oral Sci. 2011 apr;19(2):106–12.
13. Nicoletti MA, Siqueira EL, Bombana AC, Oliveira GG de. Shelf-life of a 2.5% sodium hypochlorite solution as determined by arrhenius equation. Braz Dent J. 2009;20(1):27–31.
14. Oliveira LD de, Oliveira FE de, Hatje BA, Valera MC, Carvalho CAT, Hasna AA. Detoxification of Ita by intracanal medication: analysis by macrophages proinflammatory cytokines production. Braz Dent J. 2022;33(6):36–43.
15. Oliveira LD de, Oliveira FE de, Hatje BA, Valera MC, Carvalho CAT, Hasna AA.. Detoxification of Ita by intracanal medication: analysis by macrophages proinflammatory cytokines production. Braz Dent J. 2022;33(6):36–43.
16. Orozco EIF, Toia CC, Cavalli D, Khoury RD, Cardoso FG da R, Bresciani E, et al. Effect of passive ultrasonic activation on microorganisms in primary root canal infection: a randomized clinical trial. J Appl Oral Sci. 2020;28:E20190100.
17. Pradhan MS, Gunwal M, Shenoi P, Sonarkar S, Bhattacharya S, Badole G. Evaluation of ph and chlorine content of a novel herbal sodium hypochlorite for root canal disinfection: an experimental in vitro study. Contemp Clin Dent. 2018 jun;9(Suppl 1):S74–8.



RECIMA21 - REVISTA CIENTÍFICA MULTIDISCIPLINAR
ISSN 2675-6218

EVALUATION OF POTENTIAL OF HYDROGEN (PH) OF SODIUM HYPOCHLORITE AT THREE
DIFFERENT MOMENTS BEFORE ITS EXPIRATION DATE
Sarah Regina dos Santos Pereira, Thais Cristine Pereira, Vanessa Marques Meccatti,
Lara Steffany de Carvalho, Enrico Coser Bridi, Luciane Dias de Oliveira, Amjad Abu Hasna

- 18.Schilder H. Cleaning and shaping the root canal. Dent Clin North Am. 1974 apr;18(2):269–96.
- 19.Sedgley C. Root canal irrigation--a historical perspective. J Hist Dent. 2004 jul;52(2):61–5.
- 20.Valera MC, Silva KCG da, Maekawa LE, Carvalho CAT, Koga-Ito CY, Camargo CHR, Et al. Antimicrobial activity of sodium hypochlorite associated with intracanal medication for candida albicans and enterococcus faecalis inoculated in root canals. J Appl Oral Sci. 2009 dec;17(6):555–9.
- 21.Valera MC, Maekawa LE, Chung A, Cardoso FGR, Oliveira LD de, Oliveira CL de, et al. The effect of sodium hypochlorite and ginger extract on microorganisms and endotoxins in endodontic treatment of infected root canals. Gen Dent. 2014 jun;62(3):25–9.