



The paleoenvironmental and thermal histories of the Permian Irati Formation shale in the Paraná Basin, Brazil: An integrated approach based on mineralogical and organic imprints

Haline V. Rocha^{a,*}, Lucy G. Sant'Anna^a, Cristina F.A. Rodrigues^b, Márcia Mendes^c, Zélia Pereira^c, Gilda Lopes^{d,e}, Paulo Fernandes^d, Egberto Pereira^f, Colombo C.G. Tassinari^a, Manuel J. Lemos de Sousa^b

^a IEE-USP, Institute of Energy and Environment, University of São Paulo, Av. Prof. Luciano Gualberto 1289, São Paulo, SP, 05508-080, Brazil

^b I3ID, Fundação Fernando Pessoa, Praça de 9 de Abril 349, 4249-004, Porto, Portugal

^c LNEG, Rua da Amieira, 4465-965, S. Mamede de Infesta, Portugal

^d Plants, Photosynthesis and Soil, School of Biosciences, University of Sheffield, Western Bank, Sheffield, S10 2TN, United Kingdom

^e CIMIA - Centro de Investigação Marinha e Ambiental, Universidade Do Algarve, Campus de Gambelas, 8005-130, Faro, Portugal

^f Department of Stratigraphy and Paleontology, Rio de Janeiro State University, São Francisco Xavier Street, 524 - 2^o floor-Maracanã, Rio de Janeiro, RJ, 20550-900, Brazil

ARTICLE INFO

Keywords:

Authigenesis
Clay coating
Intracratonic basin
Organofacies
Marine shale
Serra geral magmatism
Shale gas
Shale oil
Thermal maturity

ABSTRACT

Mineralogical assemblages and organofacies are important sources of information to recover the paleoenvironmental and thermal histories of shale deposits. In this study, a detailed qualitative and quantitative characterization of the Permian Irati Formation (Assistência Member) shale is based on mineralogical (XRD and SEM-EDS) and organic components (TOC, Rock-Eval pyrolysis, organofacies, TAI, fluorescence and vitrinite reflectance measurements) and provides integrated data about sediment provenance, depositional environment, diagenesis, and thermal history, while supporting interpretations on the Paraná Basin (PB), Brazil, paleogeography and its correlation to the southwest Gondwana. The results revealed a prevailing type I/II kerogen, with type III kerogen being also present but mainly confined along the paleoshoreline of the PB. The dominance of fluorescent amorphous organic matter (AOM) combined with framboidal pyrite suggests microbial activity in an anoxic-dysoxic neritic-marine paleoenvironment. Additionally, common to abundant well-preserved phytoclasts, as well as the occurrence of *Botryococcus braunii*, indicates freshwater influx in a brackish marine depositional setting. Immature to early-oil window thermal maturities prevail across the PB, according to the organic maturation indicators. The combined analysis between the organic matter evolution with clay mineralogy, such as the occurrence of interstratified clays (e.g., I/S) and its positive correlation with depth suggest that burial diagenesis reached the transition to early catagenesis on the north, southeast, and south of the basin, attributing a shale oil potential for the Irati Formation on a regional scale. Local scale imprints of the Early Cretaceous Paraná-Etendeka Large Igneous Province (LIP), and its thermal effect in the Irati Formation shale components, are recorded as clay authigenesis (e.g., smectite webby texture and clay coating development), crystallization of minerals by low to high-grade of thermal alteration (e.g., corrensite, talc, lizardite and diopside), and by local scale gas-window maturities. Such thermal alteration, identified in the proximity to intruded sills and dykes, led to a heterogeneous organic maturation pattern with implications on shale gas and shale oil potential of the Irati Formation shale, demonstrating that these subjects in the Paraná Basin should be assessed locally.

1. Introduction

Clay minerals and organic matter hold imprints on the paleohistory

of sedimentary basins, such as their relative ages, sediment provenance, depositional environments, as well as on post-depositional events (e.g., Chamley, 1989; Tyson, 1995; Merriman, 2005; Bjørlykke, 2014).

* Corresponding author.

E-mail address: haline@usp.br (H.V. Rocha).

<https://doi.org/10.1016/j.marpetgeo.2023.106328>

Received 31 January 2023; Received in revised form 6 May 2023; Accepted 17 May 2023

Available online 19 May 2023

0264-8172/© 2023 Elsevier Ltd. All rights reserved.