

The University of Notre Dame Australia ResearchOnline@ND

Theses

2023

Charting and Navigating the "Scylla and Charybdis" Conundrum of our Ageing Hearts - Heart Failure & Atrial Fibrillation

Simon Stewart
The University of Notre Dame Australia

Follow this and additional works at: https://researchonline.nd.edu.au/theses



COMMONWEALTH OF AUSTRALIA Copyright Regulations 1969

WARNING

The material in this communication may be subject to copyright under the Act. Any further copying or communication of this material by you may be the subject of copyright protection under the Act.

Do not remove this notice.

Publication Details

Stewart, S. (2023). Charting and Navigating the "Scylla and Charybdis" Conundrum of our Ageing Hearts - Heart Failure & Atrial Fibrillation (Doctor of Medical Science). University of Notre Dame Australia. https://researchonline.nd.edu.au/theses/367

This dissertation/thesis is brought to you by ResearchOnline@ND. It has been accepted for inclusion in Theses by an authorized administrator of ResearchOnline@ND. For more information, please contact researchonline@nd.edu.au.



CHARTING AND NAVIGATING THE "SCYLLA AND CHARYBDIS" CONUNDRUM OF OUR AGEING HEARTS — HEART FAILURE & ATRIAL FIBRILLATION



Henry Fuseli's painting of Odysseus choosing the unpalatable choice between Scylla and Charybdis, circa 1794/6 (Public Domain)

SIMON STEWART

RN (ITUCERT), BA, BN, GRAD DIP AD ED, PHD, FESC, FAHA, FHFA

Submitted in fulfilment of the requirements for the **Doctor of Medical Science**



School of Medicine, Sydney, NSW, Australia

March 2023

DECLARATION

To the best of the candidate's knowledge, this thesis contains no material previously published by another person, except where due acknowledgement has been made. This includes the portfolio of published, peer-reviewed manuscripts that form the basis for the thesis where the candidate is listed as the first and/or senior corresponding author.

This thesis is the candidate's own work and contains no material which has been accepted for the award of any other degree or diploma in any institution.

HUMAN ETHICS

The research projects presented and reported in this thesis were conducted in accordance with the National Health and Medical Research Council National Statement on Ethical Conduct in Human Research (2007, updated 2018). Ethics approval was granted by all relevant Human Research Ethics Committees in Australia, UK, USA, Sweden, and South Africa.

SIMON STEWART

30TH MARCH 2023

ABSTRACT

Residing on opposite sides of the Strait of Messina between Sicily and Calabria, *Scylla* and *Charybdis* were mythical sea monsters (one a fearsome creature, the other a deadly whirlpool) described by Homer. For anyone seeking to navigate that narrow and treacherous passage, avoid one of these lethal threats and you would be devoured by the other. For many populations around the world, the conundrum of successfully preventing and/or treating heart failure (**HF**) to increase longevity, only to increase the probability that it will increase the number of people living with and dying from atrial fibrillation (**AF**), represents a conundrum of *Scylla* and *Charybdis* dimensions. So how did this happen? In the late 20th Century, heart disease was characterised by a predominance of middle-aged men suffering often fatal acute coronary events. However, this pattern began to change due to the combination of two key factors. Firstly, in high-income countries at least, there were increasingly successful attempts to prevent and treat acute coronary events (thereby reducing premature mortality rates). Secondly, the vanguard of the *Post-War Baby Boomer* generation had reached their sixth decade of life, with more to come. Thus, more people were living longer with an ageing/damaged heart and those numbers would inevitably rise.

The first manifestation of this phenomenon was an increasing number of HF cases (well before a noticeable increase in AF cases). This mandated increasing recognition that HF was a discrete physical entity/syndrome that needed to be more widely recognised. This trend (rising HF cases) meant that the main burden of heart disease was shifting from the fifth/sixth decade of life with a male predominance, to the seventh/eighth decade of life affecting both men and women. Unless urgently addressed, it seemed clear that HF cases would overwhelm future health care services. At this point in time, AF attracted far less scrutiny because case numbers remained low. However, fuelled by the inevitable wave of ageing <code>Baby Boomers</code>, combined with successful attempts to prolong the lives of those affected by "earlier" forms of heart disease (including HF), it was inevitable, therefore, that more people would be successfully reaching their eight/ninth decade of life. Thus, the perfect "recipe" for a rising tide of AF had emerged.

So does every individual face a *Scylla* and *Charybdis* choice between HF and AF – of course not! However, as a society in successfully recognising and treating a rising epidemic of HF, we <u>inevitably</u> fuelled (as some of us were predicting) a consequential epidemic of AF. In simple terms, unless we could "cure" heart disease – we had no choice in a classical *Scylla* and *Charybdis* conundrum, by solving/avoiding one problem and then creating/running into, another. Thus, as with HF 10-20 years earlier, AF has since challenged health care systems to cope with the demands it places on all healthcare services. Concurrent to the rise of AF, HF now represents a sustained threat to the heart health of successfully ageing populations worldwide (creating a "twin epidemic"), with AF and HF often occurring in the same person.

It is within this context, that this thesis describes two closely related portfolios of research (comprising >100 primary and topic-related, original reports) that were instrumental in -1) Describing the evolving burden of disease imposed by HF and AF, 2) Developing targeted, multidisciplinary management programs with the capacity to simultaneously reduce the risk of recurrent hospitalisation and prolong the lives of vulnerable people affected by one or both of these deadly/disabling conditions, and 3) Considering their broader impact in vulnerable/disadvantaged communities and regions of the world (from Central Australia to Sub-Saharan Africa).

In addition to describing how this portfolio of research represents a *cogent investigation* of two critical aspects of HF and AF (i.e., what is the pattern of disease and how can its impact from an individual to societal perspective be attenuated?), this thesis outlines the concrete impact these research reports have made in our <u>collective</u> awareness, understanding and response to them.

My own (small) contribution to this field of research includes a series of important studies generating new knowledge (e.g., the prognostic impact and economic burden of HF and AF) and trials of HF and AF management that have directly influenced the provision and design of new health programs and services. Consistent with this impact, the 33 original research studies presented in this thesis have attracted >100 individual citations in Expert Clinical Guidelines/Consensus Reports over the last 20 years.

FOREWORD

Fulfilling the criteria for the award of a *Doctor of Medical Science* administered by the *University of Notre Dame Australia*, this thesis provides a critical appraisal of a body of research undertaken and led by me over a 20-year period. Following the completion of my PhD within the Department of Medicine at the University of Adelaide ("*Optimising therapeutic efficacy in acute and chronic cardiac disease states*") under the wonderful supervision of Professor John Horowitz (Head/Professor of Cardiology at the Queen Elizabeth Hospital) and Professor John Marley (Head/Professor of General Practice), I was awarded the 1999 *National Heart Foundation of Australia Overseas Ralph Reader Fellowship*.

With great fortune, the Ralph Reader Fellowship led me to the University of Glasgow, Scotland under the mentorship/tutelage of Professor John McMurray (now one of the world's most influential and respected Cardiologists). During my Fellowship I also worked with other luminaries in the fields of cardiology/public health – including Professors Simon Capewell, Jill Pell, David Hole, and Caroline Morrison.

My time at the University of Glasgow (noting that I proudly remain an active collaborator and Visiting Professor with Professor McMurray and his team) represents the starting point for the body of work presented in this thesis. When I landed in Glasgow in 1999, I had already had a strong interest in the clinical management of *heart failure*. By the time I returned to Professor John Horowitz's group in 2002, I had developed a further (intense) interest in the burden of disease imposed by *heart failure*. As a prospective component of my Fellowship (prompted by Professor Horowitz), I also extended my interest in developing and adapting clinical management programs to the emerging problem of *atrial fibrillation*.

When comparing notes with the parallel experience of the late, great Professor Henry Krum (who also returned home to Australia after working with Professor Milton Packer in the USA), I realise how much better it would have been for my career if I had stayed in the UK (for the recognition and opportunities it would have generated). Until I joined the *Baker Heart Institute* as Head of Preventative Cardiology in 2007, I struggled to gain any traction (or research project funding) back home in Australia. Nevertheless, through the support of an *Inaugural NHMRC Career Development Award* and then successively more senior NHMRC Fellowships (culminating in a *Senior Principal Research Fellowship* 15 years later), I persisted in developing my own research program/team focussed on *heart failure* and *atrial fibrillation*.

Not surprisingly (given its genesis/origin), the portfolio of research presented in this thesis has many international components – stretching from Sweden in Northern Europe to Soweto in South Africa. I've been extremely fortunate to collaborate with amazing people like Professors Karen Sliwa (Director of the Cape Heart Institute/Immediate Past-President of the World Heart Federation) and Ana Mocumbi (Non-Communicable Disease Lead, Instituto Nacional de Saúde, Mozambique Ministry of Health).

Given the above context (good fortune, geographically disperse research projects and a myriad of collaborations and opportunities not necessarily conducive to creating a linear narrative), I've deliberately sought to present a set of primary manuscripts with three key characteristics -1) I was the 1st or last/senior author, 2) When combined (even if out-of-synch in terms of timing) they form a cogent narrative focussed on *heart failure* and *atrial fibrillation*, and 3) Each report has had a proven impact on our wider understanding of these conditions and/or changed clinical practice.

No research is perfect. Thus, in linking each manuscript, I have been mindful of the imperative to present a balanced and critical account of each study and how it built on previous research (either my own or someone else's). If there is just one lesson, I've learned in my research journey, it is that the goal of truly understanding and responding to complex phenomena such as *heart failure* and *atrial fibrillation* will only end when we've eliminated disease and created the "eternal" heart. I hope this lesson is reflected in this thesis! Ironically, therefore, after fretting early in my career whether I'd still have anything to research once heart disease had been "cured" (remembering the early hype around stem cells, regeneration and near immortality!), I can confidently predict that conditions like *heart failure* and *atrial fibrillation* will continue to confound and challenge us for many more years to come.

Simon Stewart March 2023

ACKNOWLEDGEMENTS

I hope it is entirely clear from the Foreword, that I recognise and owe much of my research career success to the senior people (many of whom are named above) who have mentored and supported me. I've also been blessed with amazing research teams. Some members of these teams, like the late Ms. Ana Nadonza, Ms. Nerolie Stickland, and Dr Kai Chan, are deserving as much credit as me for the work we've collectively produced. I've also been blessed with "visionary" PhD students like Professor Geoff Strange who've created their own career path and taken me on the journey with them. I'm eternally grateful for everyone who has collaborated and supported me along the way.

It would be rare for anyone at my career stage to be good at listening and taking advice. This is especially true for someone writing a thesis about their beloved research – and I'm no exception! For their patience and forbearance of my peculiar writing style and penchant for long sentences, I'd like to warmly thank Professors Jim Codde and David Playford for helping me to compile this thesis. Both have been instrumental in me being able to complete this labour of love.

Unfortunately, my father Tom Stewart (the reason why I chose Glasgow over Boston, San Francisco, or Gothenburg for my Post-Doctoral Fellowship), never witnessed me gain a PhD or, hopefully, this higher degree. It strikes me that his witty twist on a simple label ("Simple Simon") into "It's not so Simple Simon" neatly describes my complex clinical and academic career. Nevertheless, his memory and legacy (never to be forgotten) is, and always will be, a central motivation to my research endeavours. I can only honour him properly by striving to make a difference and supporting others who seek to do the same – regardless of where they come or their circumstances. Despite all her health challenges, my mother, Jo Stewart, was still here to inspire me with her fierce intellect and support for her "boys". She played an integral role in my achieving a PhD in the first instance. Subsequently, I was grateful for the opportunity to explain how I had written and completed my thesis ready for evaluation. Thus, providing her with further concrete evidence that her sacrifices to give us a new life in Australia were still being honoured. Sadly, she died before I could share the good news about my being awarded the degree. Nevertheless - "top of the world Ma!".

"This one is important!". I can't count how many times I said this to my wife Tania as I burned through yet another evening and night finishing one of the manuscripts included in this thesis. There is really no excuse for my single-mindedness and journeying so far away from home T. However, I sincerely hope you understand why I'm so driven towards answering the next question (and the one beyond)! Regardless, I couldn't have achieved anything without your love and support and this thesis (dedicated to you and the girls) is just as much your creation as mine.

By the same token, I never had the opportunity to truly explain to my three daughters why I was absent (physically or mentally distracted) so often when they were growing up. Laura, Amy and Sarah, the fact that you grew-up to be such amazingly talented, accomplished, and caring young women is more credit to yourselves and your mum than anything I ever did. Regardless, I'm so very proud of you and the career choices you've made. I hope this thesis at least allows you to understand me a little better. If ever I worry, that I'm not working hard enough or not doing something for the right reason, I have you (and your grandfather) to thank for driving me on. I'm truly inspired and honoured to be your father.

LIST OF THESIS PUBLICATIONS

- 1. McMurray JJ, <u>Stewart S</u>. Epidemiology, aetiology, and prognosis of heart failure. *Heart*. 2000;83:596-602
- 2. <u>Stewart S</u>, MacIntyre K, Hole DJ, Capewell S and McMurray JJ. **More 'malignant' than cancer? Five-year survival following a first admission for heart failure**. *Eur J Heart Fail*. 2001;3:315-22.
- 3. <u>Stewart S</u>, MacIntyre K, MacLeod MM, Bailey AE, Capewell S, McMurray JJ. **Trends in hospitalization** for heart failure in Scotland, 1990-1996. An epidemic that has reached its peak? *Eur Heart J*. 2001;22:209-17.
- 4. <u>Stewart S</u>, Jenkins A, Buchan S, McGuire A, Capewell S, McMurray JJ. The current cost of heart failure to the National Health Service in the UK. *Eur J Heart Fail*. 2002;4:361-71.
- 5. **Stewart S**, MacIntyre K, Capewell S and McMurray JJ. **Heart failure and the aging population: an increasing burden in the 21st century?** *Heart***. 2003;89:49-53.**
- 6. Clark RA, McLennan S, Dawson A, Wilkinson D, **Stewart S**. **Uncovering a hidden epidemic: a study of the current burden of heart failure in Australia.** *Heart Lung Circ*. 2004;13:266-73.
- 7. <u>Stewart S</u>, McIntyre K, Capewell S, McMurray JJ. Heart failure in a cold climate. Seasonal variation in heart failure-related morbidity and mortality. *J Am Coll Cardiol*. 2002;39:760-6.
- 8. <u>Stewart S</u>, Demers C, Murdoch DR, McIntyre K, MacLeod ME, Kendrick S, Capewell S, McMurray JJ. Substantial between-hospital variation in outcome following first emergency admission for heart failure. *Eur Heart J*. 2002;23:650-7.
- 9. Clark RA, McLennan S, Eckert K, Dawson A, Wilkinson D, <u>Stewart S</u>. Chronic heart failure beyond city limits. *Rural Remote Health*. 2005;5:443.
- 10. <u>STEWART S</u>, Ekman I, Ekman T, Oden A, Rosengren A. <u>Population impact of heart failure and the most common forms of cancer: a study of 1 162 309 hospital cases in Sweden (1988 to 2004). *Circ Cardiovasc Qual Outcomes*. 2010;3:573-80.</u>
- 11. Clark RA, Driscoll A, Nottage J, McLennan S, Coombe DM, Bamford EJ, Wilkinson D. <u>Stewart S.</u> Inequitable provision of optimal services for patients with chronic heart failure: a national geomapping study. *Med J Aust.* 2007;186:169-73.
- 12. <u>STEWART S</u>, Playford D, Scalia GM, Currie P, Celermajer DS, Prior D, Codde J, Strange G (on behalf of the NEDA Investigators). <u>Ejection fraction and mortality</u>: a nationwide register-based cohort study of 499 153 women and men. *Eur J Heart Fail*. 2021;23:406-416.
- 13. Sliwa K, Wilkinson D, Hansen C, Ntyintyane L, Tibazarwa K, Becker A, <u>Stewart S</u>. Spectrum of heart disease and risk factors in a black urban population in South Africa (The Heart of Soweto Study): a cohort study. *Lancet*. 2008;371:915-22.
- 14. <u>Stewart S</u>, Wilkinson D, Hansen C, Vaghela V, Mvungi R, McMurray J, Sliwa K. <u>Predominance of heart failure in the Heart of Soweto Study cohort: emerging challenges for urban African communities. *Circulation*. 2008;118:2360-7.</u>
- 15. <u>Stewart S</u>, Horowitz JD. Home-based intervention in congestive heart failure: long-term implications on readmission and survival. *Circulation*. 2002;105:2861-6.
- 16. <u>Stewart S</u>, Blue L, Walker A, Morrison C, McMurray JJ. **An economic analysis of specialist heart failure** nurse management in the UK; can we afford not to implement it? *Eur Heart J*. 2002;23:1369-78.
- 17. McLennan SN, Pearson SA, Cameron J, <u>Stewart S</u>. Prognostic importance of cognitive impairment in chronic heart failure patients: does specialist management make a difference? *Eur J Heart Fail*. 2006;8:494-501.
- 18. Clark RA, Inglis SC, McAlister FA, Cleland JG, <u>Stewart S</u>. Telemonitoring or structured telephone support programmes for patients with chronic heart failure: systematic review and meta-analysis. *BMJ*. 2007;334:942.
- 19. Sochalski J, Jaarsma T, Krumholz HM, Laramee A, McMurray JJ, Naylor MD, Rich MW, Riegel B, Stewart S. What works in chronic care management: the case of heart failure. Health Aff (Millwood). 2009;28:179-89.

- 20. <u>STEWART S Carrington MJ</u>, Marwick TH, Davidson PM, Macdonald P, Horowitz JD, Krum H, Newton PJ, Reid C, Chan YK, Scuffham PA. Impact of home versus clinic-based management of chronic heart failure: the WHICH? (Which Heart Failure Intervention Is Most Cost-Effective & Consumer Friendly in Reducing Hospital Care) multicenter, randomized trial. *J Am Coll Cardiol*. 2012;60:1239-48.
- 21. <u>Stewart S</u>, Chan YK, Wong C, Jennings G, Scuffham P, Esterman A, Carrington M (on behalf of the NIL-CHF Investigators). Impact of a nurse-led home and clinic-based secondary prevention programme to prevent progressive cardiac dysfunction in high-risk individuals: the Nurse-led Intervention for Less Chronic Heart Failure (NIL-CHF) randomized controlled study. *Eur J Heart Fail*. 2015;17:620-30.
- 22. <u>Stewart S</u>, Wiley JF, Ball J, Chan YK, Ahamed Y, Thompson DR, Carrington MJ. <u>Impact of Nurse-Led, Multidisciplinary Home-Based Intervention on Event-Free Survival Across the Spectrum of Chronic Heart Disease: Composite Analysis of Health Outcomes in 1226 Patients From 3 Randomized Trials. *Circulation*. 2016;133:1867-77.</u>
- 23. Scuffham PA, Ball J, Horowitz JD, Wong C, Newton PJ, Macdonald P, McVeigh J, Rischbieth A, Emanuele N, Carrington MJ, Reid CM, Chan YK, <u>Stewart S</u> (on behalf of the WHICH? II Trial Investigators). Standard vs. intensified management of heart failure to reduce healthcare costs: results of a multicentre, randomized controlled trial. *Eur Heart J*. 2017;38:2340-2348.
- 24. <u>Stewart S</u>, Hart CL, Hole DJ, McMurray JJ. <u>Population prevalence, incidence, and predictors of atrial fibrillation in the Renfrew/Paisley study</u>. *Heart*. 2001;86:516-21.
- 25. <u>Stewart S</u>, Hart CL, Hole DJ, McMurray JJ. A population-based study of the long-term risks associated with atrial fibrillation: 20-year follow-up of the Renfrew/Paisley study. *Am J Med*. 2002;113:359-64.
- 26. <u>STEWART S</u>, MacIntyre K, MacLeod MM, Bailey AE, Capewell S, McMurray JJ. Trends in hospital activity, morbidity and case fatality related to atrial fibrillation in Scotland, 1986--1996. *Eur Heart J.* 2001;22:693-701.
- 27. <u>STEWART S</u>, MacIntyre K, Chalmers JW, Boyd J, Finlayson A, Redpath A, Pell JP, Capewell S, McMurray JJ. Trends in case-fatality in 22968 patients admitted for the first time with atrial fibrillation in Scotland, 1986-1995. *Int J Cardiol*. 2002;82:229-36.
- 28. <u>Stewart S</u>, Murphy NF, Walker A, McGuire A. McMurray JJ. **Cost of an emerging epidemic: an economic analysis of atrial fibrillation in the UK**. *Heart*. 2004;90:286-92.
- 29. Sliwa K, Carrington MJ, Klug E, Opie L, Lee G, Ball J, <u>Stewart S</u>. **Predisposing factors and incidence of newly diagnosed atrial fibrillation in an urban African community: insights from the Heart of Soweto Study**. *Heart*. 2010;96:1878-82.
- 30. Ball J, Carrington MJ, McMurray JJ, <u>Stewart S</u>. Atrial fibrillation: profile and burden of an evolving epidemic in the 21st century. *Int J Cardiol*. 2013;167:1807-24.
- 31. Ball J, Thompson DR, Ski CF, Carrington MJ, Gerber T, <u>Stewart S</u>. Estimating the current and future prevalence of atrial fibrillation in the Australian adult population. *Med J Aust*. 2015;202:32-5.
- 32. Ball J, Carrington MJ, <u>Stewart S</u> (on behalf of the SAFETY Trial investigators). **Mild cognitive** impairment in high-risk patients with chronic atrial fibrillation: a forgotten component of clinical management? *Heart*. 2013;99:542-7.
- 33. Ball J, Carrington MJ, Wood KA, <u>Stewart S</u> (on behalf of the SAFETY Trial investigators). **Women versus** men with chronic atrial fibrillation: insights from the Standard versus Atrial Fibrillation spEcific management study (SAFETY). *PLoS One*. 2013;8:e65795.
- 34. <u>Stewart S</u>, Ball J, Horowitz JD, Marwick TH, Mahadevan G, Wong C, Abhayaratna WP, Chan YK, Esterman A, Thompson DR, Scuffham PA, Carrington MJ. <u>Standard versus atrial fibrillation-specific management strategy (SAFETY) to reduce recurrent admission and prolong survival: pragmatic, multicentre, randomised controlled trial. *Lancet*. 2015;385:775-84.</u>

Intellectual Contribution & Leadership

APPENDIX I provides the details of the 12 peer-reviewed funding schemes (from prestigious career Fellowships to substantive program of research grants) that directly relate to these 33 selected manuscripts.

A key component of a good proportion of this portfolio, is the co-authorship of PhD candidates under my principal/co- supervision. Many were supported by their own prestigious post-graduate scholarships. I am fortunate to have navigated 20 such candidates to completion in the past 20 years. Many of these early career researchers were initially funded by the National Heart Foundation of Australia and/or National Health & Medical Research Council of Australia. Moreover, many now hold senior academic appointments (at last count at least 5 are full Professors) through which they lead their own research teams and undertake impactful research programs.

When selecting the 33 primary/original research manuscripts to compile this thesis and survive the scrutiny of expert peer-review, I consciously decided to apply a combination of the following criteria, in respect to <u>unequivocally demonstrating</u> my intellectual contribution/leadership of each output -

- **A.** Being listed as the first/primary author or last/senior corresponding author of the manuscript.
- **B.** Conducting the described research/project as the primary recipient/ Principal Investigator of the peer-reviewed funding that supported the research being described.
- **C.** Demonstrated leadership of large multidisciplinary research teams such as the *WHICH? Trial* and *SAFETY Trial* and *Heart of Soweto Study* Investigators, to successfully complete (and report on) large-scale disease surveillance and pragmatic health service intervention studies.

APPENDIX II contains all 34 of the primary manuscripts that have been carefully selected to form the core of this portfolio of research. They comprise a key review of the HF literature co-authored by me with the inspirational Professor John McMurray and published in *Heart* in 2000 plus <u>33 original research papers</u> spanning 20 years thereafter. A further <u>78 co-authored manuscripts</u> that support the results and interpretation of study findings of these primary reports have been added as references and are highlighted in the **References**.

PLACING THIS PORTFOLIO OF RESEARCH INTO MY CAREER CONTEXT

During my research career as a clinician/scientist/health services researcher who has spanned multiple disciplines (basic science, nursing, cardiology, medicine, clinical trials, public health, epidemiology, and health economics), I've been privileged to publish more than 400 research articles, more than 1000 research abstracts and authored/edited (all as senior author/editor) ten books.

My current *Scopus h-index (March 2023) stands at 69 from 420 listed papers and ~22,000 citations*. I am proud that this important metric is not inflated by guidelines or "group" research papers where I have minimal intellectual control or input. Thus, as reflected in the portfolio of research informing this thesis, most of my highly cited, original research reports were led and written by me.

RESEARCH IMPACT

As was succinctly commented to me when compiling this portfolio of research — "it is one thing to be highly cited, it's another to be impactful and make a difference". Indeed, the conditions for this award are clear in that the published body of research must —

- **a.** Represent a significant advance in knowledge (in this case, our understanding of the evolving epidemiology/burden of disease and optimal management of heart failure and atrial fibrillation).
- **b.** Given rise to significant debate amongst recognised scholars (in this case, generating numerous editorials and high citations from subsequent research projects and reviews of the literature).
- c. Have directly changed the direction of research or practice in a newer generation of scholars (in this case, contributing to an expanded focus on heart failure as a truly "malignant" syndrome and pioneering dedicated health services, identifying atrial fibrillation as a rising public health threat requiring dedicated care programs AND highlighting the increasing intersection between these two deadly and disabling conditions.

In framing the clinical impact of my portfolio of research TABLE 1 and TABLE 2, specifically identify the >50 individual guideline/expert consensus reports that have referenced the 33 original research reports that form the core of this thesis. To date, my research has informed >200 expert documents. This includes current clinical guidelines published by the European Society of Cardiology, American Heart Association and Cardiac Society of Australia & New Zealand.

TABLE OF CONTENTS

SECTION	Page
Introduction	14
HEART FAILURE – THE SCYLLA OF CHRONIC HEART DISEASE IN THE LATE 20^{TH} & Early 21^{ST}	17
CENTURY	
Prognostic Impact of HF	17-21
 A RISING TIDE OF HF ADMISSIONS 	22-24
COUNTING THE COST OF HF	24-25
THE FUTURE OF HF WITHIN THE AGEING POPULATIONS OF HIGH-INCOME COUNTRIES	25-26
 CLOSING THE KNOWLEDGE GAPS AROUND AN EVOLVING EPIDEMIC OF HF 	26-32
Consolidating the Evidence Around Multidisciplinary HF Management	32
 Are the Initial Benefits of Nurse-Led, Home-Based Management 	
Prolonged?	33-34
 Is There an Economic Case for Supporting Nurse-led HF Services 	34-35
 Nurse-Led Management Should Be Applied — But Which Model? 	35-40
 Is it Possible to Prevent HF in High-Risk Patients? 	40-42
 Does Nurse-Led HF Management Work Every Time? 	42-44
New Directions in HF Management	44-45
Translational Impact – Greater HF Awareness & Dedicated Management Programs	46-49
ATRIAL FIBRILLATION – THE CHARYBDIS OF CHRONIC HEART DISEASE IN THE 21 ST CENTURY	50
	50-52
CHARACTERISING THE POPULATION PROFILE & IMPACT OF AF PENSALING THE PROGNESTIC MARKET OF AF	52-53
REVEALING THE PROGNOSTIC IMPACT OF AF AF DELATED LICENSEL FORCES & OUTSIANSE OUTSIANS ASSOCIATED ASSOCI	53-54
AF-RELATED HOSPITAL EPISODES & OUTCOMES	55
COUNTING THE COST OF AN EMERGING EPIDEMIC Province Cost of An Emerging Epidemic A Forest A Fore	55
Bringing Order to Chaos – Global Burden & Impact of AF in the 21 st Course. Course.	55-60
CENTURY	60
PIONEERING ATRIAL FIBRILLATION-SPECIFIC DISEASE MANAGEMENT PROGRAMS	60-66
A New Solution to an Old Problem Table Transport Control Average Special Transport Control Trans	00 00
Translational Impact – Greater Awareness & Dedicated Management Programs for	66-68
ATRIAL FIBRILLATION	
Thesis Conclusions	69-70
HEART FAILURE BURDEN PORTFOLIO REFERENCES	71
HEART FAILURE MANAGEMENT REFERENCES	71-72
ATRIAL FIBRILLATION BURDEN REFERENCES	72
ATRIAL FIBRILLATION MANAGEMENT REFERENCES	72
Additional References	73-84
APPENDIX I - COMPETITIVE FUNDING SUPPORTING THIS PORTFOLIO OF RESEARCH	85-88
Appendix II - Original Thesis Publications	-

LIST OF TABLES

	PAGE
TABLE 1 – A PORTFOLIO OF HIGHLY INFLUENTIAL RESEARCH REPORTS ON THE EVOLVING BURDEN	
& Management of HF	47-49
Table 2 – A Portfolio of Highly Influential Research Reports on the Evolving Burden	
& MANAGEMENT OF AF	67-68

LIST OF FIGURES (THIS THESIS PRESENTS SOURCE FIGURES AS THEY WERE ORIGINALLY PUBLISHED)

	PAGE
FIGURE 1 – PORTFOLIO OF HEART FAILURE RESEARCH	15
FIGURE 2 – PORTFOLIO OF ATRIAL FIBRILLATION RESEARCH	16
FIGURE 3 – HF IS MORE MALIGNANT THAN CANCER	17
FIGURE 4 – IMPROVING HF SURVIVAL RATES IN SWEDEN	18
FIGURE 5 – SEX-SPECIFIC PATTERNS OF MORTALITY ACCORDING TO LVFEF	19
FIGURE 6 – PROGNOSTIC IMPACT OF CHANGE IN LVEF	20
FIGURE 7 – SEASONAL PATTERNS OF HF MORTALITY	21
FIGURE 8 – HF SEASONALITY IS A GLOBAL PHENOMENON	21
FIGURE 9 – RISING HF ADMISSIONS IN THE LATE 20 TH CENTURY	22
FIGURE 10 – A PLATEAU OF PRIMARY HF ADMISSIONS IN SCOTLAND	23
FIGURE 11 – AN AGE-ADJUSTED DECLINE IN HF ADMISSIONS IN SCOTLAND	23
FIGURE 12 – INTER-HOSPITAL VARIATIONS IN HF RE-ADMISSIONS IN SCOTLAND	24
FIGURE 13 – THE RISING COST OF HEART FAILURE	25
FIGURE 14 – THE FUTURE COST-BURDEN OF HEART FAILURE	26
FIGURE 15 – ESTIMATED BURDEN OF HEART FAILURE IN AUSTRALIA (CIRCA 2000)	27
FIGURE 16 – ESTIMATED BURDEN OF HEART FAILURE IN AUSTRALIA (CIRCA 2016)	28
FIGURE 17 – HEART FAILURE BEYOND CITY LIMITS	28
FIGURE 18 – MAPPING THE TYRANNY OF DISTANCE – CARDIAC ARIA MAP	29
FIGURE 19 – THE HEART OF SOWETO	30
FIGURE 20 – HEART FAILURE IN SOWETO	30
FIGURE 21 – SOWETO IS SITTING ON THE CROSSROADS OF PAST & FUTURE HEART DISEASE	31
FIGURE 22 – HEART FAILURE IN THE "HEART" OF AUSTRALIA	32
FIGURE 23 – IMPACT OF A NURSE-LED, HOME-BASED INTERVENTION IN HEART FAILURE ON	33
Mortality	
FIGURE 24 – LONG-TERM IMPACT OF A NURSE-LED, HOME-BASED INTERVENTION IN HEART	34
FAILURE ON READMISSION & MORTALITY	
FIGURE 25 – ECONOMIC BLUE-PRINT FOR A HEART FAILURE SERVICE IN THE UK	35
FIGURE 26 – META-ANALYSIS OF HEART FAILURE MANAGEMENT & MORTALITY	36
Figure 27 – Meta-Analysis of Remote Heart Failure Management & Readmissions	37
Figure 28 – Individual Meta-Analysis of Heart Failure Management Modality &	38
Readmissions	
FIGURE 29 – HOME- VERSUS CLINIC-BASED MANAGEMENT OF HEART FAILURE – THE WHICH? I TRIAL	38
FIGURE 30 – PROLONGED IMPACT OF HOME- VERSUS CLINIC-BASED MANAGEMENT OF HEART FAILURE – THE WHICH I TRIAL	39
FIGURE 31 – PRIMARY ENDPOINT OUTCOMES – THE WHICH? II TRIAL	40
FIGURE 32 – TRAJECTORY OF CARDIAC DYSFUNCTION IN THE NIL-CHF TRIAL COHORT	41
FIGURE 33 – CARDIAC TRAJECTORY & INCIDENT HF OR DEATH IN THE NIL-CHF TRIAL COHORT	41
FIGURE 34 – IMPACT OF MILD COGNITIVE IMPAIRMENT ON HEART FAILURE MANAGEMENT	42
FIGURE 35 – DIFFERENTIAL IMPACT OF HOME-BASED HEART FAILURE MANAGEMENT	43
ACCORDING TO THE AGE & CLINICAL COMPLEXITY OF INDIVIDUALS	-
FIGURE 36 – THE ARISE-HF FRAMEWORK FOR ASSESSING MULTIMORBIDITY	44
FIGURE 37 – A BIO-BEHAVIOURAL MODEL OF SEASONAL VULNERABILITY IN HEART FAILURE	45
FIGURE 38 – POPULATION PREVALENCE OF ATRIAL FIBRILLATION (RENFREW-PAISLEY COHORT)	51
FIGURE 39 — POPULATION INCIDENCE OF ATRIAL FIBRILLATION (RENFREW-PAISLEY COHORT)	51
FIGURE 40 – PROGNOSTIC IMPORTANCE OF ATRIAL FIBRILLATION TO LONG-TERM MORTALITY	52
FIGURE 41 – ATRIAL-FIBRILLATION RELATED HOSPITAL TRENDS (ALL CASES)	53
FIGURE 42 – DE NOVO ATRIAL FIBRILLATION RELATED HOSPITAL TRENDS	54

54
56
57
58
58
59
59
61
62
63
63
64
64
64
69
70

Introduction

In the late 20th Century/early 21st Century there was a growing realisation among clinicians and epidemiologists alike that the syndrome heart failure (HF) was placing an increasing burden on health care systems worldwide.¹ However, in many jurisdictions, including the UK, the clinical entity "heart failure" was poorly defined and documented.¹ This was exacerbated by the fact it could not be placed on a death certificate as a contributory cause of death.¹ Moreover, the definition of HF was evolving in tandem with the growing armoury of neurohormonal agents being successfully (and unsuccessfully) trialled to improve health outcomes.¹


As summarised in Figure 1, arising from an expert review paper¹ that was published in the 12 months following the completion of my PhD in collaboration with the host of my National Heart Foundation of Australia Overseas Ralph Reader Fellowship (Professor John McMurray, University of Glasgow, Scotland, UK), over nearly two decades I completed and published an integrated portfolio of 22 original research manuscripts that focussed on —

- 1. Understanding the evolving burden and impact of all forms heart failure and,
- 2. Further developing clinical management strategies that would attenuate the high rates of morbidity and mortality among those hospitalised with this deadly and disabling syndrome.¹⁻²³

Although HF was already the focus of considerable interest (both in terms of its burden and optimal treatment/management), this was clearly not the case for atrial fibrillation (AF). It is notable that in early clinical trials focussing on HF, AF was barely mentioned. Moreover, pre-existing cases were being treated with "blunt" therapeutics such as warfarin and sotalol. However, as shown in Figure 2, after considering the key dynamics of how HF had developed into a substantive health issue (due to the progressive ageing of high-income populations within which an increasing component of individuals were living with cardiac risk factors and/or survived an acute coronary event), I/we published a closely related portfolio of 10 original research manuscripts focussed on —


- 1. Charting the emerging threat and impact of *all forms of atrial fibrillation* and,
- 2. Developing new clinical strategies that would specifically attenuate the high-rates of morbidity and mortality among those hospitalised with chronic forms of AF.²⁴⁻³⁴

As with many complex research programs shaped by data availability, the formation of opportunistic collaborations and, critically, funding to conduct key studies, both portfolios of research are not completely linear in their progression of ideas, concepts, and findings. Critically, however, nearly every discrete project and publication described below has been highly cited in the literature (as of March 2022, collectively they have attracted >10,000 Scopus Citations) — thereby shaping the literature in return.



Testament to the international collaborations and partnerships (as represented by the image on the left) underlying this research, the portfolios of research summarised in Figures 1 and 2 have been cited close to 100 times in expert guidelines/peer reports on a global basis (see Tables 1 and 2). This includes clinical guidelines for the contemporary management of HF and ΑF as currently recommended by the European Society of Cardiology (ESC)³⁵, American Heart Association (AHA)³⁶ and Cardiac Society of Australia and New Zealand (CSANZ).37


FIGURE 1 – A PORTFOLIO OF ORIGINAL RESEARCH REPORTS EXAMINING AND RESPONDING TO THE EVOLVING BURDEN OF HEART FAILURE IN THE 21ST CENTURY.



Within the thesis, six broad sections (comprising 22 key publications) relating to the epidemiological profile, burden and management of HF are presented. Each section/reference is colour coded.

LEGEND

FIGURE 2 – A PARALLEL PORTFOLIO OF ORIGINAL RESEARCH REPORTS IDENTIFYING AND RESPONDING TO THE EVOLVING BURDEN OF AF IN THE 21ST CENTURY.



LEGEND Within the thesis, six broad sections (comprising 10 key publications) relating to AF are presented. Each section/reference is colour coded.

Material not shown due to convinibility

CONCLUSIONS: Over a 20-year period, we conducted a range of pragmatic trials and analyses focussed on the benefits (and potential adverse effects) of multidisciplinary HF management. Pragmatic RCTs were conducted in the UK¹⁷⁰, Hong Kong ¹⁷¹ and Australia.²³ Similarly, international collaborations collated and synthesised outcome data from across the globe. A key focus of my research program (ongoing) is the translation of evidence into cost-effective and sustainable health services. It also involves monitoring health systems to determine if the evidence around HF management is being appropriately applied. In parallel to our evolving knowledge around the detection and treatment of HF (the definition and clinical scope of which is continually changing), we have re-tested the contemporary efficacy of our original approaches to HF management, including our own. Recent findings suggest that our old models of care need to be updated to accommodate a more challenging HF patient population and the environment in which they live. ¹⁶⁹ Consequently, new, more innovative approaches to care are being developed and tested in a new generation of appropriately powered RCTs (from Africa to Australia).

TRANSLATIONAL IMPACT – GREATER HF AWARENESS & DEDICATED MANAGEMENT PROGRAMS

Of the <u>22 research manuscripts</u> presented in this portfolio of research in HF, 16 (73%) were published in highly prestigious journals with an impact factor >10. This includes the *British Medical Journal, The Lancet, Circulation, Journal of the American College of Cardiology* and *European Heart Journal* and the *European Journal of Heart Failure*. Two manuscripts are the 1st and 3rd most cited research articles in the journal's history.

Consistent with the overall significance of the research being presented in these papers, combined they have accumulated >4,500 Scopus citations (~180 cites per paper) with a mean field-weighted citation index of 6.27 (topic rather than discipline profile). Critically, the importance of an early report focussing on seasonal patterns of HF events took many years to be recognised (hence it's low FWCI calculated over the first 3 years of publication). By the same token, due to the predominant reliance on external guidelines, the African-specific reports (whilst highly influential) are unlikely to be cited until the framework of African research expands and clinical guidelines based on African-led findings are published.

As outlined in Table 1*, nearly all papers have specifically informed the expert clinical guidelines (28 individual citations), expert consensus statements (10 individual citations) and expert position statements/reports (18 individual citations). These cover the following regions:

- Sub-Saharan Africa Pan-African Society of Cardiology [PASCAR]),
- Australia National Heart Foundation of Australia [NHFA]/Cardiac Society of Australia and New Zealand (CSANZ),
- Europe European Society of Cardiology (ESC) and its Heart Failure Association [HFA]),
- Latin America and North America American Heart Association (AHA), American College of Cardiology (ACC) and Heart Failure Society of America [HFSA])
- Global Lancet Commission reports

Overall, these papers have attracted **66** individual citations in these highly influential documents (by mid-2022). At the most fundamental level, my program of research has contributed to the development of a new role for nurses (the Specialist HF Nurse) and new Multidisciplinary HF Management programs delivering care to those affected by HF worldwide – the challenge being to extend such coverage to disadvantaged/low-resource regions (e.g., Sub-Saharan Africa) and ensure these programs are adapted to remain cost-effective.

TABLE 1 summarises the bibliometric profile and inclusion in expert guidelines of each manuscript groups into the categories described in **Figure 1**.

TABLE 1 – A PORTFOLIO OF RESEARCH REPORTS ON THE EVOLVING BURDEN & MANAGEMENT OF HF

Title	Journal [Year]	SS	Scopu s Cites	FWCI/ Centile	Significance of the report to HF awareness/management	Guidelines/Expert Consensus & Positions
DEFINING THE PROGNOS	TIC IMPLICATION	NS & BI	JRDEN O	F HEART FA	ILURE	
More malignant than cancer? Five-year survival following a first admission for HF	European Journal of Heart Failure [2001]	1st	838	11.22/ 99 th	Report on the prognostic impact of HF relative to the most common forms of cancer. It remains the most cited research paper in the history of the highly prestigious European Journal of Heart Failure.	 2019 Italian Society of Cardiology Consensus Statement 2017 Austrian Cardiology Society Position Statement 2015 EHRA Consensus Statement 2014 Polish Clinical Forum on Cardiac Imaging 2012 ESC HF guidelines 2011 Brazilian Cardio- oncology guidelines 2010 ESC HF guidelines 2008 Italian Cardiology Society guidelines 2008 ESC HF guidelines 2008 ESC HF guidelines 2006 HFSA guidelines 2005 ESC Position Paper
HF in a cold climate: Seasonal variation in HF- related morbidity & mortality	Journal of the American College of Cardiology [2002]	1st	159	1.54/ 81 st *	Early, report on the seasonal impact on HF-related events – most citations have appeared in the last decade. It now forms the initial rationale for a MRFF-funded RCT - the RESILIENCE Trial.	 2020 Family Practice (US) guidelines 2011 ESC guidelines
Population impact of HF & the most common forms of cancer: A study 1 162 309 hospital cases in Sweden (1988 to 2004)	Circulation: CV Quality & Outcomes [2010]	1st	158	3.31/ 81	One of the largest population studies of HF-related mortality (versus cancer) ever reported. It quantified (for the first-time) the number of life-years-lost due to HF among men and women within a whole population.	 2017 Austrian Cardiology Society 2011 ESC guideline
Ejection fraction & mortality: a nationwide register-based cohort study of 499 153 women & men	European Journal of Heart Failure [2021]	1st	11	5.50/ 99 th	Largest-ever patient cohort study of its kind to date. It has contributed to a paradigm change in the interpretation of ejection fraction levels and the risk of mortality on a sex-specific basis with calls to revisit sex-specific thresholds of risk according to VEF levels and a reinterpretation of clinical trial outcomes	2021 ESC guidelines
EXAMINING THE EVOLVI	NG PATTERN OF	HEART	FAILURE-	-RELATED H	OSPITAL ADMISSIONS & OUTCOMES	
Trends in hospitalization for HF in Scotland, 1990-1996: An epidemic that has reached its peak?	European Heart Journal [2001]	1st	195	7.94/ 98th	A controversial report on the peaking of admissions with a primary diagnosis of HF (but not secondary diagnosis) in Scotland - subsequently replicated by equivalent studies worldwide	
Substantial between-hospital variation in outcome following first emergency admission for HF	European Heart Journal [2002]	1st	29	0.92 / 71 st	Highly controversial and debated study within Scotland/UK that framed calls for the application of multidisciplinary HF management programs to address wide variations in HF-related outcomes	2005 Latin American guidelines
•					H CARE SYSTEMS OF HIGH-INCOME COUNTRIES	
The current cost of heart failure to the National Health Service in the UK	European Journal of Heart Failure [2002]	1st	450	6.71/ 98 th	One of the top 5 most cited research papers in the history of the highly prestigious European Journal of Heart Failure, this was the first study compare & contrast the cost of HF within the same health service a decade apart. Findings showed the increasing economic burden of HF -subsequently replicated by other studies worldwide	 2017 Position Statement Austrian Cardiology Society 2012 ESC HF guidelines 2010 ESC HF guidelines 2009 Italian Cardiology Society HF guidelines 2008 ESC HF guidelines
					LVING EPIDEMIC OF HEART FAILURE	
Heart failure and the aging population: An increasing burden in the 21st Century	Heart [2003]	1st	296	5.50/ 96 th	A study that challenged pre- existing/alarming warnings of an exponential rise in HF cases & the associated burden of disease. Study findings suggesting a more modest (~25%	 2012 German National HF guidelines 2008 AHA guidelines 2006 ESC guidelines

					increase) and plateauing burden of disease were subsequently proven more accurate.	
CLOSING KEY KNOWLEDG	GE GAPS ON THE	PATTE	RN OF HI	F: FROM AU	STRALIA TO SUB-SAHARAN AFRICA	
Uncovering a hidden epidemic: A study of the current burden of heart failure in Australia	Heart Lung & Circulation [2004]	Last	50	1.40/ 79 th	Australia relies on very poor & often misleading (e.g., asking the public if they have swollen ankles or HF). This study was the first quantify the burden of HF using the best available data at the time.	2010 NHFA/CSANZ guidelines 2010 AIS Position Statement 2006 NHFA/CSANZ guidelines
Chronic heart failure beyond city limits	Rural & Remote Health [2005]	Last	16	1.05/ 73 rd	Contemporary national health surveys in Australia focussed on urban populations. This study was the first to demonstrate a disproportionate burden of HF cases in rural Australia (due to socio-economic factors) - subsequently confirmed by other studies.	
Inequitable provision of optimal services for patients with chronic heart failure: A national geo-mapping study	Medical Journal of Australia [2007]	Last	53	3.11/ 92 nd	A geo-mapping study that built on the findings of the above report — demonstrating a mismatch between HF cases and available clinical services/expert care. It also led to the innovative CARDIAC-ARIA tool	2014 NHFA/CSANZ Consensus Statement 2010 NHFA/CSANZ guidelines
Cardiometabolic risk and disease in Indigenous Australians: The Heart of the Heart Study	International Journal of Cardiology [2014]	Last	30	1.49/ 81 st *	This was a study of the antecedents and epidemiological profile (reported in a separate manuscript as part of a PhD program) of HF and other forms of chronic heart disease within the Indigenous communities of Central Australia	
Spectrum of heart disease and risk factors in a black urban population in South Africa (the Heart of Soweto Study): a cohort study	The Lancet [2008]	Last	282	15.53/ 99 th *	This paradigm-changing report challenged the conventional wisdom that heart disease and, specifically HF, was almost non-existent in Sub-Saharan Africa.	2020 ESC/HFSA Position Paper 2020 PASCAR Position Paper 2019 AHA Heart diseases & stroke statistics 2018 AHA Heart diseases & stroke statistics 2017 AHA Heart diseases & stroke statistics 2017 AHA Heart diseases & stroke statistics 2016 AHA Heart diseases & stroke statistics
Predominance of heart failure in the heart of Soweto study cohort: Emerging challenges for urban African communities	Circulation [2008]	1st	127	2.39/ 89 th *	Published as the same time as the primary Heart of Soweto primary report, this study characterised the large volume of HF cases (many with hypertension as well as right-sided aetiology) in a large urban African community in epidemiological transition.	 2021 Lancet Commission on Women & Cardiovascular Disease 2020 Lancet Commission on Poverty & Non-Communicable Diseases
ESTABLISHING AND THEN	NEXPLORING TH	E BENE	FITS & PI	TFALLS OF H	EART FAILURE MANAGEMENT PROGRA	AMS
An economic analysis of specialist heart failure nurse management in the UK: Can we afford not to implement it?	European Heart Journal [2002]	1st	82	2.58/ 90 th	Arising from my trials of HF management (PhD program) & the world-first Glasgow Heart Failure Nurse Liaison Service I helped to establish, this paper modelled the cost-benefits of applying a UK-wide HF service – the results of which were then implemented by, the British Heart Foundation.	 2019 German Cardiac Society Position Paper 2017 ESC/HFSA Position Paper 2011 ESC Position Paper
Home-based intervention in congestive heart failure: long-term implications on readmission and survival	Circulation [2002]	1st	250	7.08/ 98 th	Arising from my trials of HF management (PhD program) this report demonstrated that the initial benefits of this kind of program (previously shown to be effective in the short-term) were sustained over the longer-term.	 2017 Austrian Society of Cardiology Consensus 2016 Spanish Society of Cardiology Consensus 2009 German Society of Cardiology Guidelines 2008 HFSA Consensus 2006 HFSA Guidelines 2005 Latin American HF Guidelines

Prognostic importance of cognitive impairment in chronic heart failure patients: Does specialist management make a difference?	European Journal of Heart Failure [2010]	Last	76	4.41/ 96 th	Despite growing support for HF management programs world-wide, this paper (for the first-time) identified the pitfalls associated with implementing such programs among those with cognitive impairment (including those with HF & chronic pulmonary disease).	 2015 American HF Association Position Paper 2011 ESC/HFSA Guidelines
Telemonitoring or structured telephone support programmes for patients with chronic heart failure: Systematic review and metaanalysis	British Medical Journal [2007]	1st	485	37.67/ 99 th	Following the publication of this meta- analysis of HF programs (of which I was the 2 nd author), this report focused on the evidence supporting the remote management of HF. Along with the original meta-analysis [R] and a series of related reports derived from this analysis, this has shaped the application of HF programs worldwide.	 2013 ACCF/AHA HF guidelines 2009 ACCF/AHA HF guidelines 2008 ESC guidelines
What works in chronic care management: The case of heart failure	Health Affairs [2009]	Last	185	13.99/ 99 th	Addressing the inherent limitations of meta-analysis of scientific reports, this was a global collaboration that combined individual RCT profiling and outcome data from studies of HF management led by worldwide experts – producing a service-defining recommendation that in-person, multidisciplinary HF teams produced the best health outcomes.	 2015 EHRA Consensus Statement 2013 ACC/AHA HF Guidelines 2012 ESC guidelines
Impact of home versus clinic- based management of chronic heart failure: The WHICH? (Which heart failure intervention is most cost- effective & consumer friendly in reducing hospital care) multicentre, randomized trial	Journal of the American College of Cardiology [2012]	1st	103	3.25/ 93 rd	This RCT (funded by the NHMRC of Australia), was the first to compare two forms of HF management head-to-head. Initially underpowered to detect program differences, longer-term follow-up showed the superiority of the homebased intervention, whilst patients also preferred home-based management.	 2014 ESC Position Statement 2016 NHFA/CSANZ HF guidelines
Impact of a nurse-led home and clinic-based secondary prevention programme to prevent progressive cardiac dysfunction in high-risk individuals: The Nurse-led Intervention for Less Chronic Heart Failure (NIL-CHF) randomized controlled study	European Journal of Heart Failure [2015]	1st	18	1.45/ 80 th	Funded by the NHMRC of Australia, this was <u>first study of its kind</u> (complementing one other trial in primary care) to determine if it's possible to prevent HF in high-risk individuals via multidisciplinary care & surveillance. Although initially negative, the original and subsequent long-term follow-up identified the potential to prevent disease progression in high-risk individuals.	■ 2021 ESC/HFSA Position Paper
Impact of nurse-led multidisciplinary home-based intervention on event-free survival across the spectrum of chronic heart disease	Circulation [2016]	1st	24	1.62/ 83 rd	Funded by the NHMRC of Australia, this was a composite analysis of 4 RCTs of home-based disease management (including patients with HF). This study demonstrated, for the first-time, the potential harmful effects of this type of intervention among younger and older patients with chronic heart disease (often with malignant multimorbidity) requiring new perspectives on how care should be delivered to such individuals.	2018 American Practitioner Nurse Association Consensus Statement
Standard versus intensified management of heart failure to reduce healthcare costs: Results of a multicentre, randomized controlled trial	European Heart Journal [2016]	Last	16	1.17/ 75 th	Funded by the NHMRC of Australia, this multicentre RCT compared an intensified versus standardised approach to HF management of those living in both urban and rural settings – no difference between the two groups being found.	2021 ESC/HFSA Position Paper

Material not shown due to convinibility

This new clinical reality mandates a re-evaluation of many longstanding hospital avoidance programs to determine if they are still "fit for purpose". There is a cogent argument that they should be modified to suit the needs of modern-day patient populations. It also mandates the development of new approaches to care. It is for this reason we are conducting the MRFF-funded RESILIENCE Trial to reduce external provocations to the cardiopulmonary health of high-risk patients affected by a combination of CAD, AF and/or HF – rather than focussing on AF alone. 166, 167

CONCLUSIONS: Despite a wealth of clinical to public health data (some of which has been presented in this portfolio of research) to characterise the evolving characteristics and burden of AF, the evidence in support of AF-specific management programs remains under-developed. This might be rectified with the publication of further RCTs (such as the HELP-AF Study) and the potential to then generate meta-analyses from multiple RCTs with a large volume of endpoints/outcomes. At the very least, ongoing efforts to address "seasonal" vulnerability (as described above) will automatically identify individuals living with chronic AF at high-risk of recurrently hospitalisation and premature mortality. This approach (individual rather than disease specific) is the better new/old approach to provide incremental health benefits within a growing patient population presenting to hospital with a combination of AF and/or HF and other forms of multimorbidity.

TRANSLATIONAL IMPACT — GREATER AWARENESS & DEDICATED MANAGEMENT PROGRAMS FOR ATRIAL FIBRILLATION

In stark contrast to the HF portfolio, of the 10 AF-focussed manuscripts described above (all presenting primary research outputs), only 2 (20%) were published in journals with an impact factor >10. Despite this, reflective of the <u>latent/delayed recognition</u> of these reports, combined they have accumulated ~2,900 **Scopus citations** (~290 cites per paper) with a **mean field-weighted citation index** of **6.19** (topic rather than discipline profile).

Overall, all 10 of these papers have specifically informed the expert clinical guidelines (23 individual citations), expert consensus statements (11 individual citations) and expert position statements/reports (5 individual citations). These cover the following regions:

- Sub-Saharan Africa Pan-African Society of Cardiology [PASCAR]),
- Australia/Asia-Pacific National Heart Foundation of Australia [NHFA]/Cardiac Society of Australia and New Zealand (CSANZ), Pan-Pacific Heart Rhythm Society
- Europe European Society of Cardiology (ESC) and its Heart Failure Association [HFA]),
- Latin America and North America American Heart Association (AHA), American College of Cardiology (ACC) and Heart Failure Society of America [HFSA])
- Global Lancet Commission reports

Overall, these papers have attracted *44 individual citations in these documents* (by mid-2022). At the most fundamental level, my program of research, whilst having a lesser (to date) impact in respect to triggering coordinated AF management, still played an important role in highlighting the rising individual to economic burden of AF – high health expenditure being one the main drivers of health service and commercial innovations.

TABLE 2 (PAGES 61-62) summarises the bibliometric profile and inclusion in expert guidelines of each manuscript groups into the categories described in **FIGURE 2**.

TABLE 2 – A PORTFOLIO OF RESEARCH REPORTS ON THE EVOLVING BURDEN & MANAGEMENT OF AF

Title	Journal [Year]	SS	Scopus Cites	FWCI/ Centile	Significance of the report to HF awareness/management	Guidelines/Expert Consensus & Positions
POPULATION PROFILE OF A	\F					
Population prevalence, incidence, and predictors of atrial fibrillation in the Renfrew/Paisley Study	Heart [2001]	1st	502	2.92/ 92 nd	It took many years for the importance of this report to be recognised – it represents one of the first-ever population studies of the incidence and prevalence of AF.	 2017 EHRA guidelines 2013 French Society of Geriatrics and Gerontology guidelines 2010 ESC guidelines 2004 Swiss Cardiology Society guidelines
PROGNOSTIC IMPACT OF A	F					
A population-based study of the long-term risks associated with atrial fibrillation: 20-Year follow-up of the Renfrew/Paisley Study	American Journal of Medicine [2002]	1st	1,183	8.66/ 98 th	This is a paradigm-changing population-based longitudinal research report that redefined the prognostic implications of AF. Despite winning a major population research prize at the American Heart Association's Scientific meeting, as with the "more malignant than cancer" HF report, there was a reluctance by the major medical journals to publish anything that hadn't been derived from the Framingham Heart Study. Its importance was finally recognised when for two iterations of the major AF guidelines, it was first reference quoted to underline the importance of recognising the public health threat and need for expert clinical management of AF. This is the most cited paper published by the American Journal of Medicine in 2002.	2020 ESC guidelines 2020 EHRA/APHRS guidelines 2019 ACC/AAT/AHA guidelines 2018 European Society of Hypertension Position Paper 2016 ESC guidelines 2013 French Society of Geriatrics and Gerontology Consensus Statement 2012 HRS/EHRA Consensus Statement 2012 Polish Cardiac Society Consensus Statement 2011 ACC/ACC guidelines 2010 ESC guidelines 2009 Canadian Hypertension Education guidelines 2009 AFNET-EHRA Consensus Statement 2009 Japanese Hypertension Society guidelines 2007 HRS/EHRA Consensus Statement 2007 HRS/EHRA Consensus Statement 2007 HRS/EHRA Consensus Statement 2006 ACC/AHA/ESC guidelines
TRENDS IN AF-RELATED H	OSPITAL ADMISSIO	NS				
Trends in hospital activity, morbidity and case fatality related to atrial fibrillation in Scotland, 1986-1996	European Heart Journal [2001]	1st	129	3.03/ 91 st	This report was the first-ever analysis of long-term trends in AF-related hospital activity and outcomes. Like those studies focussing on HF, it led to several studies focussing on seasonal patterns of admission and the future burden of disease.	 2011 ACC/AHA guidelines 2006 ACC/AHA guidelines
Trends in case-fatality in 22,938 patients admitted for the first time with atrial fibrillation in Scotland, 1986-1995	International Journal of Cardiology	1 st	37	1.44/ 80 th	This follow-up report (to that above), provided a granular age-and sex-specific analysis of de novo/incident admissions with AF in Scotland – showing that the future burden of disease would increase as the population progressively aged.	2012 Polish Cardiac Society guidelines
COST-BURDEN OF AN EME		F AF				
Cost of an emerging epidemic: An economic analysis of atrial fibrillation in the UK	Heart [2004]	1st	437	11.22/ 99 th	This report represents the first-ever analysis of the economic burden of AF. Up until its publication there was little appreciation of the growing health burden of AF. Many similar reports have now been generated from around the world. It is the most cited paper published by the Heart in 2004	2022 Asia Pacific HRS guidelines 2020 Canadian CV Society guidelines 2018 NHFA guidelines 2017 WHF AF Roadmap 2016 ESC guidelines 2011 AIAC guidelines 2011 ACCF/AHA guidelines 2006 ACC/AHA/ESC guidelines

EPIDEMIOLOGICAL FEATUR	EPIDEMIOLOGICAL FEATURES OF AF – A GLOBAL PERSPECTIVE							
Predisposing factors and incidence of newly diagnosed atrial fibrillation in an urban African community: Insights from the Heart of Soweto Study	Heart [2010]	Last	51	1.80/ 84 th *	Arising from the paradigm-changing Heart of Soweto Study, that redefined the modern spectrum of heart disease in the region, this represents an early and therefore important study of AF in Sub-Saharan Africa.	2021 Lancet Commission on Women & Cardiovascular Disease 2020 Lancet Commission on Poverty & Non- Communicable Diseases 2020 PASCAR Position Paper		
Atrial fibrillation: Profile and burden of an evolving epidemic in the 21 st century	International Journal of Cardiology [2013]	Last	394	10.19/ 99 th	This report represents one of the most comprehensive systematic reviews and meta-analyses of AF ever reported. It provides pooled analyses of every aspect of the population characteristics and health burden of AF in high-income countries, whilst highlighting the paucity of data in low-income regions of the world.	2017 EHRA Consensus Statement 2015 EHRA Consensus Statement		
Estimating the current and future prevalence of atrial fibrillation in the Australian adult population	Medical Journal of Australia [2015]	Last	61	11.22/ 99 th	This is one of the first report in Australia to examine the current and future burden of AF.	 2019 Italian Society of Cardiology Consensus Statement 2017 Austrian Cardiology Society Position Statement 2015 EHRA Consensus Statement 		
ESTABLISHING THE CASE FO	OR ATRIAL FIBRILLAT	TION-SI	PECIFIC MA	NAGEMEN	T PROGRAMS			
Mild cognitive impairment in high-risk patients with chronic atrial fibrillation: A forgotten component of clinical management?	Heart [2013]	Last	60	2.25/ 88 th	An insightful clinical report that led to recommendation that all inpatients chronic AF should be screened for cognitive impairment, with consequential implications on the education and post-discharge management of these cases.	2016 ESC guidelines2016 EIPAAHA Consensus Statement		
Women Versus Men with Chronic Atrial Fibrillation: Insights from the Standard Versus Atrial Fibrillation spEcific managemenT studY (SAFETY)	PLoS ONE [2013]	Last	52	0.90/ 68 th	Arising from the SAFETY Trial this study generated important data on the clinical differences between men and women affected by AF.	 2020 Canadian CV Society guidelines 2016 ESC guidelines 2016 EIPAAHA Consensus Statement 		
Standard versus atrial fibrillation-specific management strategy (SAFETY) to reduce recurrent admission and prolong survival: Pragmatic, multicentre, randomised controlled trial	The Lancet [2015]	1 st	87	9.67/ 99 th	Analogous to the RCT on HF management (published in <i>The Lancet</i> as part of my PhD), this study provided concrete evidence to support nurse-led multidisciplinary AF-specific management to simultaneously reduce hospital admissions and prolong survival in those patients with AF but not HF – hence its simultaneous presentation at the 2015 AHA Scientific Meeting and simultaneous publication in <i>The Lancet</i> .	 2020 ESC guidelines 2018 NHFA/CSANZ guidelines 2016 ESC guidelines 2016 EHRA Consensus Statement 2015 EHRA Consensus Statement 		

THESIS CONCLUSIONS

When the first of the 33 manuscripts comprising this portfolio of research was published, it would have been difficult to justify labelling HF and AF as the "Scylla and Charybdis" of chronic heart disease. Even with poetic license (noting the need for such a strategy when applying perfectly elegant mythologies and metaphors to the imperfect world of public health/clinical care), one might argue that this label is still not justified – particularly when an increasing number of individuals in high-income countries are achieving relatively heart healthy longevity. However, all the data presented in this manuscript suggest that –

- 1. Even if the rate of *de novo* HF is declining (there is no real data to support this at a time when its definition is being expanded), due to better management of common antecedents such as acute coronary syndromes, paradoxically the overall volume of people entering old age at risk of developing the syndrome will remain high. Critically, antecedent risk in the form of sedentary behaviours, metabolic diseases, and hypertension, along with improving management of acute coronary syndromes remain at historical highs. *Thus, the modern-day epidemic of HF that emerged in the late* 20th Century, will be sustained well into the foreseeable future.
- 2. Following on from the initial rise of HF, AF has since emerged as the new cardiac epidemic in the 21st Century. *There is no sign (yet) that it has reached its peak.* Like HF, there is a paradoxical twist to the dynamics of this epidemic given that a) The prevalence of AF naturally rises with advancing age (due to the "ageing heart" effect) in the absence of any substantive disease and, b) It is far more likely to occur in those individuals who reach older age with pre-existing risk factors and/or established forms of heart disease (the most notable one being HF).

In considering the first point/conclusion, it's worth considering the temporal pattern of cases presenting with an LVEF <45% and their subsequent 1-year mortality captured by NEDA - FIGURE 57. Over a prolonged period, the survival profile of such cases has improved. However, dynamics such as the age, sex, diagnostic and investigative thresholds, new therapeutics and even variability in outcomes (depending on which institutions contributed to NEDA cases over that period) confound simple interpretation of these data. The only thing we can say with (some) certainty, is that despite some gains in survival overall, there appears to be a "steady state" of individuals being investigated with impaired systolic function. Such data will provide a rich line of enquiry as NEDA expands to include more cases (>1 million from >40 Australian centres), hospital episodes and therapeutic data (including new treatment strategies).




FIGURE 57 - TEMPORAL TRENDS IN 1-YEAR ALL-CAUSE MORTALITY IN NEDA CASES WITH LVEF ≤45%

Legend: The insert boxes show 1-year mortality averaged over 3 years (2000-02 to 2015-17)

Similarly, when considering the second point/conclusion, it's worth considering the temporal pattern of cases presenting in non-sinus rhythm (an imperfect surrogate for individuals in AF during their echocardiogram) and their subsequent 1-year mortality captured by NEDA - FIGURE 58. Once again, without more detailed analyses/accurate identification of AF cases (something that will be addressed once NEDA integrates linked data to hospital episodes and treatment data), these data are difficult to interpret. Once again however, despite a myriad of potential confounders and positive survival trends, the volume of individuals presenting with potential AF in Australia remains high. Thus, the underlying tension to optimally prevent and manage AF will remain high for the foreseeable future.




FIGURE 58 - TEMPORAL TRENDS IN 1-YEAR ALL-CAUSE MORTALITY IN NEDA CASES NOT IN SINUS RHYTHM

Legend: The insert boxes show 1-year mortality averaged over 3 years (2000-02 to 2015-17)

Thus, no matter how we might seek to safely *navigate* our increasingly at-risk populations to old age without provoking the rise of costly, debilitating, and deadly forms of heart dysfunction at an early age, we are inevitably placing them at risk of the <u>twin threats</u> HF and AF. Optimal disease management programs based on the highest levels of evidence (including appropriately powered RCTs and robust meta-analyses) do offer some hope – particularly those designed to optimise the management of high-risk individuals hospitalised with HF and/or AF. New pharmacological agents typically require real-world health service capacity to replicate what was achieved in the trials that proved their worth. This typically includes more cautious uptitration and safety monitoring than first anticipated. However, generating additional gains within an increasingly older and more complex patient cardiac population with HF and AF is increasingly challenging.

Final Conclusions. Paradoxically, we are the victims of our own success and relative wealth. We've largely overcome the dangers of middle-age - although the recent spate of high-profile sudden cardiac deaths/events in men aged in their 50's is a sobering reminder that it still exists (especially in middle-aged men who chose to smoke!). Many more people are now entering old age. However, they are typically entering it with antecedent risk for developing chronic forms of heart disease. Thus, HF and AF represent the **Scylla & Charybdis** of this new world. Having emerged in the clinical to public consciousness separately (around 10 years apart to reflect their different natural history/disease trajectory), they've now merged and pose a twin threat to millions of people globally. This mandates more nuanced thinking about the triggers of events and how we might prolong the lives of those affected. This will include the challenge of maintaining good quality of life for those affected whilst not provoking more debilitating and costly hospital episodes.

HEART FAILURE BURDEN PORTFOLIO REFERENCES

- 1. McMurray JJ, <u>Stewart S</u>. Epidemiology, aetiology, and prognosis of heart failure. *Heart*. 2000;83:596-602.
- 2. <u>Stewart S</u>, MacIntyre K, Hole DJ, Capewell S and McMurray JJ. **More 'malignant' than cancer? Five-year survival following a first admission for heart failure**. *Eur J Heart Fail*. 2001;3:315-22.
- 3. <u>Stewart S</u>, MacIntyre K, MacLeod MM, Bailey AE, Capewell S, McMurray JJ. **Trends in hospitalization for heart failure in Scotland, 1990-1996. An epidemic that has reached its peak?** *Eur Heart J*. 2001;22:209-17.
- 4. <u>Stewart S</u>, Jenkins A, Buchan S, McGuire A, Capewell S, McMurray JJ. **The current cost of heart failure to the National Health Service in the UK**. *Eur J Heart Fail*. 2002;4:361-71.
- 5. <u>Stewart S</u>, MacIntyre K, Capewell S and McMurray JJ. Heart failure and the aging population: an increasing burden in the 21st century? *Heart*. 2003;89:49-53.
- 6. Clark RA, McLennan S, Dawson A, Wilkinson D, Stewart S. Uncovering a hidden epidemic: a study of the current burden of heart failure in Australia. *Heart Lung Circ*. 2004;13:266-73.
- 7. <u>Stewart S</u>, McIntyre K, Capewell S, McMurray JJ. **Heart failure in a cold climate. Seasonal variation** in heart failure-related morbidity and mortality. *J Am Coll Cardiol*. 2002;39:760-6.
- 8. <u>Stewart S</u>, Demers C, Murdoch DR, McIntyre K, MacLeod ME, Kendrick S, Capewell S, McMurray JJ. Substantial between-hospital variation in outcome following first emergency admission for heart failure. *Eur Heart J.* 2002;23:650-7.
- 9. Clark RA, McLennan S, Eckert K, Dawson A, Wilkinson D, <u>Stewart S</u>. Chronic heart failure beyond city limits. *Rural Remote Health*. 2005;5:443.
- 10. <u>Stewart S</u>, Ekman I, Ekman T, Oden A, Rosengren A. Population impact of heart failure and the most common forms of cancer: a study of 1 162 309 hospital cases in Sweden (1988 to 2004). *Circ Cardiovasc Qual Outcomes*. 2010;3:573-80.
- 11. Clark RA, Driscoll A, Nottage J, McLennan S, Coombe DM, Bamford EJ, Wilkinson D. <u>Stewart S.</u> Inequitable provision of optimal services for patients with chronic heart failure: a national geomapping study. *Med J Aust*. 2007;186:169-73.
- 12. <u>Stewart S</u>, Playford D, Scalia GM, Currie P, Celermajer DS, Prior D, Codde J, Strange G (on behalf of the NEDA Investigators). **Ejection fraction and mortality: a nationwide register-based cohort study of 499 153 women and men.** *Eur J Heart Fail*. 2021;23:406-416.
- 13. Sliwa K, Wilkinson D, Hansen C, Ntyintyane L, Tibazarwa K, Becker A, <u>Stewart S</u>. Spectrum of heart disease and risk factors in a black urban population in South Africa (The Heart of Soweto Study): a cohort study. *Lancet*. 2008;371:915-22.
- 14. <u>Stewart S</u>, Wilkinson D, Hansen C, Vaghela V, Mvungi R, McMurray J, Sliwa K. **Predominance of heart failure in the Heart of Soweto Study cohort: emerging challenges for urban African communities.** *Circulation*. 2008;118:2360-7.

HEART FAILURE MANAGEMENT REFERENCES

- 15. <u>Stewart S</u>, Horowitz JD. Home-based intervention in congestive heart failure: long-term implications on readmission and survival. *Circulation*. 2002;105:2861-6.
- 16. <u>Stewart S</u>, Blue L, Walker A, Morrison C,McMurray JJ. **An economic analysis of specialist heart failure** nurse management in the UK; can we afford not to implement it? *Eur Heart J*. 2002;23:1369-78.
- 17. McLennan SN, Pearson SA, Cameron J, <u>Stewart S</u>. Prognostic importance of cognitive impairment in chronic heart failure patients: does specialist management make a difference? *Eur J Heart Fail*. 2006;8:494-501.
- 18. Clark RA, Inglis SC, McAlister FA, Cleland JG, <u>Stewart S</u>. Telemonitoring or structured telephone support programmes for patients with chronic heart failure: systematic review and meta-analysis. *BMJ*. 2007;334:942.
- 19. Sochalski J, Jaarsma T, Krumholz HM, Laramee A, McMurray JJ, Naylor MD, Rich MW, Riegel B, Stewart S. What works in chronic care management: the case of heart failure. Health Aff (Millwood). 2009;28:179-89.

- 20. <u>Stewart S</u> Carrington MJ, Marwick TH, Davidson PM, Macdonald P, Horowitz JD, Krum H, Newton PJ, Reid C, Chan YK, Scuffham PA. Impact of home versus clinic-based management of chronic heart failure: the WHICH? (Which Heart Failure Intervention Is Most Cost-Effective & Consumer Friendly in Reducing Hospital Care) multicenter, randomized trial. *J Am Coll Cardiol*. 2012;60:1239-48.
- 21. <u>Stewart S</u>, Chan YK, Wong C, Jennings G, Scuffham P, Esterman A, Carrington M (on behalf of the NIL-CHF Investigators). Impact of a nurse-led home and clinic-based secondary prevention programme to prevent progressive cardiac dysfunction in high-risk individuals: the Nurse-led Intervention for Less Chronic Heart Failure (NIL-CHF) randomized controlled study. Eur J Heart Fail. 2015;17:620-30.
- 22. <u>Stewart S</u>, Wiley JF, Ball J, Chan YK, Ahamed Y, Thompson DR, Carrington MJ. Impact of Nurse-Led, Multidisciplinary Home-Based Intervention on Event-Free Survival Across the Spectrum of Chronic Heart Disease: Composite Analysis of Health Outcomes in 1226 Patients From 3 Randomized Trials. *Circulation*. 2016;133:1867-77.
- 23. Scuffham PA, Ball J, Horowitz JD, Wong C, Newton PJ, Macdonald P, McVeigh J, Rischbieth A, Emanuele N, Carrington MJ, Reid CM, Chan YK, <u>Stewart S</u> (on behalf of the WHICH? II Trial Investigators). Standard vs. intensified management of heart failure to reduce healthcare costs: results of a multicentre, randomized controlled trial. *Eur Heart J*. 2017;38:2340-2348.

ATRIAL FIBRILLATION BURDEN REFERENCES

- 24. <u>Stewart S</u>, Hart CL, Hole DJ, McMurray JJ. **Population prevalence, incidence, and predictors of atrial fibrillation in the Renfrew/Paisley study**. *Heart*. 2001;86:516-21.
- 25. <u>Stewart S</u>, Hart CL, Hole DJ, McMurray JJ. **A population-based study of the long-term risks associated** with atrial fibrillation: **20-year follow-up of the Renfrew/Paisley study**. *Am J Med*. 2002;113:359-64.
- 26. <u>Stewart S</u>, MacIntyre K, MacLeod MM, Bailey AE, Capewell S, McMurray JJ. **Trends in hospital** activity, morbidity and case fatality related to atrial fibrillation in Scotland, 1986--1996. *Eur Heart J.* 2001;22:693-701.
- 27. <u>Stewart S</u>, MacIntyre K, Chalmers JW, Boyd J, Finlayson A, Redpath A, Pell JP, Capewell S, McMurray JJ. Trends in case-fatality in 22968 patients admitted for the first time with atrial fibrillation in Scotland, 1986-1995. *Int J Cardiol*. 2002;82:229-36.
- 28. <u>Stewart S</u>, Murphy NF, Walker A, McGuire A. McMurray JJ. **Cost of an emerging epidemic: an economic analysis of atrial fibrillation in the UK**. *Heart*. 2004;90:286-92.
- 29. Sliwa K, Carrington MJ, Klug E, Opie L, Lee G, Ball J, <u>Stewart S</u>. Predisposing factors and incidence of newly diagnosed atrial fibrillation in an urban African community: insights from the Heart of Soweto Study. *Heart*. 2010;96:1878-82.
- 30. Ball J, Carrington MJ, McMurray JJ, <u>Stewart S</u>. Atrial fibrillation: profile and burden of an evolving epidemic in the 21st century. *Int J Cardiol*. 2013;167:1807-24.
- 31. Ball J, Thompson DR, Ski CF, Carrington MJ, Gerber T, <u>Stewart S</u>. Estimating the current and future prevalence of atrial fibrillation in the Australian adult population. *Med J Aust*. 2015;202:32-5.

ATRIAL FIBRILLATION MANAGEMENT REFERENCES

- 32. Ball J, Carrington MJ, <u>Stewart S</u> (on behalf of the SAFETY Trial investigators). **Mild cognitive** impairment in high-risk patients with chronic atrial fibrillation: a forgotten component of clinical management? *Heart*. 2013;99:542-7.
- 33. Ball J, Carrington MJ, Wood KA, <u>Stewart S</u> (on behalf of the SAFETY Trial investigators). Women versus men with chronic atrial fibrillation: insights from the Standard versus Atrial Fibrillation spEcific management study (SAFETY). *PLoS One*. 2013;8:e65795.
- 34. <u>Stewart S</u>, Ball J, Horowitz JD, Marwick TH, Mahadevan G, Wong C, Abhayaratna WP, Chan YK, Esterman A, Thompson DR, Scuffham PA, Carrington MJ. **Standard versus atrial fibrillation-specific management strategy (SAFETY) to reduce recurrent admission and prolong survival: pragmatic, multicentre, randomised controlled trial.** *Lancet***. 2015;385:775-84.**

ADDITIONAL REFERENCES - *SS (CO-)AUTHOR

- 35. McDonagh TA, Metra M, Adamo M, Gardner RS, Baumbach A, Bohm M, Burri H, Butler J, Celutkiene J, Chioncel O, Cleland JGF, Coats AJS, Crespo-Leiro MG, Farmakis D, Gilard M, Heymans S, Hoes AW, Jaarsma T, Jankowska EA, Lainscak M, Lam CSP, Lyon AR, McMurray JJV, Mebazaa A, Mindham R, Muneretto C, Francesco Piepoli M, Price S, Rosano GMC, Ruschitzka F, Kathrine Skibelund A. 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J*. 2021;42:3599-3726.
- 36. Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM, Deswal A, Drazner MH, Dunlay SM, Evers LR, Fang JC, Fedson SE, Fonarow GC, Hayek SS, Hernandez AF, Khazanie P, Kittleson MM, Lee CS, Link MS, Milano CA, Nnacheta LC, Sandhu AT, Stevenson LW, Vardeny O, Vest AR, Yancy CW. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2022;79:e263-e421.
- 37. Atherton JJ, Sindone A, De Pasquale CG, Driscoll A, MacDonald PS, Hopper I, Kistler P, Briffa TG, Wong J, Abhayaratna WP, Thomas L, Audehm R, Newton PJ, O'Loughlin J, Connell C, Branagan M. National Heart Foundation of Australia and Cardiac Society of Australia and New Zealand: Australian clinical guidelines for the management of heart failure 2018. *Med J Aust*. 2018;209:363-369.
- 38. Ho KK, Anderson KM, Kannel WB, Grossman, Levy D. Survival after the onset of congestive heart failure in Framingham Heart Study subjects. *Circulation*. 1993;88:107-15.
- 39. McKee PA, Castelli WP, McNamara PM, Kannel WB. The natural history of congestive heart failure: the Framingham study. *N Engl J Med*. 1971;285:1441-6.
- 40. Braunschweig F. Fighting the malignancy of recurring heart failure events. *J Am Coll Cardiol*. 2011;58:738-9.
- 41. Liang HY, Lo YC, Chiang HY, Chen MF, Kuo CC. Validation and Comparison of the 2003 and 2016 Diastolic Functional Assessments for Cardiovascular Mortality in a Large Single-Center Cohort. *J Am Soc Echocardiogr*. 2020;33:469-480.
- 42. Lip GY, Rasmussen LH, Skjoth F, Overvad K, Larsen TB. Stroke and mortality in patients with incident heart failure: the Diet, Cancer and Health (DCH) cohort study. *BMJ Open*. 2012;2.
- 43. Guha A, Dey AK, Armanious M, Dodd K, Bonsu J, Jneid H, Abraham W, Fradley MG, Addison D. Health care utilization and mortality associated with heart failure-related admissions among cancer patients. *ESC Heart Fail*. 2019;6:733-746.
- 44. Aimo A, Senni M, Barison A, Panichella G, Passino C, Bayes-Genis A, Emdin M. Management of heart failure with preserved ejection fraction: from neurohormonal antagonists to empagliflozin. *Heart Fail Rev.* 2022.
- 45. Tsuyuki RT, Yusuf S, Rouleau JL, Maggioni AP, McKelvie RS, Wiecek EM, Wang Y, Pogue J, Teo KK, White M, Avezum A, Jr., Latini R, Held P, Lindgren E. Probstfield J. Combination neurohormonal blockade with ACE inhibitors, angiotensin II antagonists and beta-blockers in patients with congestive heart failure: design of the Randomized Evaluation of Strategies for Left Ventricular Dysfunction (RESOLVD) Pilot Study. *Can J Cardiol*. 1997;13:1166-74.
- 46. Williams RE. The effect of neurohormonal antagonists in reducing heart failure hospitalizations. *Curr Med Res Opin*. 2006;22:139-50.
- 47. Taylor CJ, Ordonez-Mena JM, Roalfe AK, Lay-Flurrie S, Jones NR, Marshall T, Hobbs FDR. Trends in survival after a diagnosis of heart failure in the United Kingdom 2000-2017: population based cohort study. *BMJ*. 2019;364:l223.
- 48. *Jhund PS, Macintyre K, Simpson CR, Lewsey JD, <u>Stewart S</u>, Redpath A, Chalmers JW, Capewell S, McMurray JJ. Long-term trends in first hospitalization for heart failure and subsequent survival between 1986 and 2003: a population study of 5.1 million people. *Circulation*. 2009;119:515-23.
- *****Dudas K, Lappas G, **Stewart S**, Rosengren A. Trends in out-of-hospital deaths due to coronary heart disease in Sweden (1991 to 2006). *Circulation*. 2011;123:46-52.

- *Teng TH, Hung J, Knuiman M, <u>Stewart S</u>, Arnolda L, Jacobs I, Hobbs M, Sanfilippo F, Geelhoed E, Finn J. Trends in long-term cardiovascular mortality and morbidity in men and women with heart failure of ischemic versus non-ischemic aetiology in Western Australia between 1990 and 2005. *Int J Cardiol*. 2012;158:405-10.
- 51. Dewan P, Rorth R, Raparelli V, Campbell RT, Shen L, Jhund PS, Petrie MC, Anand IS, Carson PE, Desai AS, Granger CB, Kober L, Komajda M, McKelvie RS, O'Meara E, Pfeffer MA, Pitt B, Solomon SD, Swedberg K, Zile MR, McMurray JJV. Sex-Related Differences in Heart Failure With Preserved Ejection Fraction. *Circ Heart Fail*. 2019;12:e006539.
- 52. Ibrahim NE, Pina IL, Camacho A, Bapat D, Felker GM, Maisel AS, Butler J, Prescott MF, Abbas CA, Solomon SD, Januzzi JL, Jr., Prospective Study of Biomarkers SI and Ventricular Remodeling During Entresto Therapy for Heart Failure Study I. Sex-based differences in biomarkers, health status, and reverse cardiac remodelling in patients with heart failure with reduced ejection fraction treated with sacubitril/valsartan. *Eur J Heart Fail*. 2020;22:2018-2025.
- 53. Lam CSP, Arnott C, Beale AL, Chandramouli C, Hilfiker-Kleiner D, Kaye DM, Ky B, Santema BT, Sliwa K, Voors AA. Sex differences in heart failure. *Eur Heart J*. 2019;40:3859-3868c.
- 54. Merrill M, Sweitzer NK, Lindenfeld J, Kao DP. Sex Differences in Outcomes and Responses to Spironolactone in Heart Failure With Preserved Ejection Fraction: A Secondary Analysis of TOPCAT Trial. *JACC Heart Fail*. 2019;7:228-238.
- 55. Wilcox JE, Fang JC, Margulies KB, Mann DL. Heart Failure With Recovered Left Ventricular Ejection Fraction: JACC Scientific Expert Panel. *J Am Coll Cardiol*. 2020;76:719-734.
- 56. Solomon SD, Vaduganathan M, B LC, Packer M, Zile M, Swedberg K, Rouleau J, M AP, Desai A, Lund LH, Kober L, Anand I, Sweitzer N, Linssen G, Merkely B, Luis Arango J, Vinereanu D, Chen CH, Senni M, Sibulo A, Boytsov S, Shi V, Rizkala A, Lefkowitz M, McMurray JJV. Sacubitril/Valsartan Across the Spectrum of Ejection Fraction in Heart Failure. *Circulation*. 2020;141:352-361.
- 57. Wehner GJ, Jing L, Haggerty CM, Suever JD, Leader JB, Hartzel DN, Kirchner HL, Manus JNA, James N, Ayar Z, Gladding P, Good CW, Cleland JGF, Fornwalt BK. Routinely reported ejection fraction and mortality in clinical practice: where does the nadir of risk lie? *Eur Heart J*. 2020;41:1249-1257.
- *Strange G, Celermajer DS, Marwick T, Prior D, Ilton M, Codde J, Scalia GM, <u>Stewart S</u>, Bulsara M, Gabbay E, Playford D. The National Echocardiography Database Australia (NEDA): Rationale and methodology. *Am Heart J*. 2018;204:186-189.
- 59. McMurray JJV, Jackson AM, Lam CSP, Redfield MM, Anand IS, Ge J, Lefkowitz MP, Maggioni AP, Martinez F, Packer M, Pfeffer MA, Pieske B, Rizkala AR, Sabarwal SV, Shah AM, Shah SJ, Shi VC, van Veldhuisen DJ, Zannad F, Zile MR, Cikes M, Goncalvesova E, Katova T, Kosztin A, Lelonek M, Sweitzer N, Vardeny O, Claggett B, Jhund PS, Solomon SD. Effects of Sacubitril-Valsartan Versus Valsartan in Women Compared With Men With Heart Failure and Preserved Ejection Fraction: Insights From PARAGON-HF. *Circulation*. 2020;141:338-351.
- *Strange G, Playford D, Scalia GM, Celermajer DS, Prior D, Codde J, Chan YK, Bulsara MK, <u>Stewart S</u> (on behalf of the NEDA Investigators). Change in ejection fraction and long-term mortality in adults referred for echocardiography. *Eur J Heart Fail*. 2021;23:555-563.
- 61. Boulay F, Berthier F, Sisteron O, Gendreike Y, Gibelin P. Seasonal variation in chronic heart failure hospitalizations and mortality in France. *Circulation*. 1999;100:280-6.
- ***Stewart S**, Keates AK, Redfern A, McMurray JJV. Seasonal variations in cardiovascular disease. *Nat Rev Cardiol*. 2017;14:654-664.
- 63. Zannad F. Have we jugulated the epidemic of heart failure? One swallow does not make a summer. *Eur Heart J.* 2001;22:188-90.
- 64. Fonarow GC, Heywood JT, Heidenreich PA, Lopatin M, Yancy CW. Temporal trends in clinical characteristics, treatments, and outcomes for heart failure hospitalizations, 2002 to 2004: findings from Acute Decompensated Heart Failure National Registry (ADHERE). *Am Heart J.* 2007;153:1021-8.
- 65. Levy D, Kenchaiah S, Larson MG, Benjamin EJ, Kupka MJ, Ho KK, Murabito JM, Vasan RS. Long-term trends in the incidence of and survival with heart failure. *N Engl J Med*. 2002;347:1397-402.

- 66. Ng TP, Niti M. Trends and ethnic differences in hospital admissions and mortality for congestive heart failure in the elderly in Singapore, 1991 to 1998. *Heart*. 2003;89:865-70.
- 67. Senni M, De Maria R, Gregori D, Gonzini L, Gorini M, Cacciatore G, Gavazzi A, Pulignano G, Porcu M, Maggioni AP. Temporal trends in survival and hospitalizations in outpatients with chronic systolic heart failure in 1995 and 1999. *J Card Fail*. 2005;11:270-8.
- 68. Shafazand M, Rosengren A, Lappas G, Swedberg K, Schaufelberger M. Decreasing trends in the incidence of heart failure after acute myocardial infarction from 1993-2004: a study of 175,216 patients with a first acute myocardial infarction in Sweden. *Eur J Heart Fail*. 2011;13:135-41.
- 69. Shahar E, Lee S. Historical trends in survival of hospitalized heart failure patients: 2000 versus 1995. BMC Cardiovasc Disord. 2007;7:2.
- 70. Yeung DF, Boom NK, Guo H, Lee DS, Schultz SE, Tu JV. Trends in the incidence and outcomes of heart failure in Ontario, Canada: 1997 to 2007. *CMAJ*. 2012;184:E765-73.
- ***STEWART S**. Atrial fibrillation in the 21st century: the new cardiac "Cinderella" and new horizons for cardiovascular nursing? *Eur J Cardiovasc Nurs*. 2002;1:115-21.
- 72. Lampe FC, Morris RW, Whincup PH, Walker M, Ebrahim S, Shaper AG. Is the prevalence of coronary heart disease falling in British men? *Heart*. 2001;86:499-505.
- 73. Tunstall-Pedoe H, Kuulasmaa K, Mahonen M, Tolonen H, Ruokokoski E, Amouyel P. Contribution of trends in survival and coronary-event rates to changes in coronary heart disease mortality: 10-year results from 37 WHO MONICA project populations. Monitoring trends and determinants in cardiovascular disease. *Lancet*. 1999;353:1547-57.
- 74. Bonneux L, Barendregt JJ, Meeter K, Bonsel GJ, van der Maas PJ. Estimating clinical morbidity due to ischemic heart disease and congestive heart failure: the future rise of heart failure. *Am J Public Health*. 1994;84:20-8.
- 75. Kelly DT. Paul Dudley White International Lecture. Our future society. A global challenge. *Circulation*. 1997;95:2459-64.
- *****Macintyre K, **STEWART S**, Chalmers J, Pell J, Finlayson A, Boyd J, Redpath A, McMurray J, Capewell S. Relation between socioeconomic deprivation and death from a first myocardial infarction in Scotland: population based analysis. *BMJ*. 2001;322:1152-3.
- *Pell JP, MacIntyre K, Walsh D, Capewell S, McMurray JJ, Chalmers JW, Boyd JH, Finlayson AR, **Stewart S**, Redpath AD. Time trends in survival and readmission following coronary artery bypass grafting in Scotland, 1981-96: retrospective observational study. *BMJ*. 2002;324:201-2.
- *McAlister FA, Murphy NF, Simpson CR, <u>Stewart S</u>, MacIntyre K, Kirkpatrick M, Chalmers J, Redpath A, Capewell S, McMurray JJ. Influence of socioeconomic deprivation on the primary care burden and treatment of patients with a diagnosis of heart failure in general practice in Scotland: population based study. *BMJ*. 2004;328:1110.
- *Murphy NF, MacIntyre K, Capewell S, <u>Stewart S</u>, Pell J, Chalmers J, Redpath A, Frame S, Boyd J, McMurray JJ. Hospital discharge rates for suspected acute coronary syndromes between 1990 and 2000: population based analysis. *BMJ*. 2004;328:1413-4.
- 80. *Murphy NF, Simpson CR, McAlister FA, <u>Stewart S</u>, MacIntyre K, Kirkpatrick M, Chalmers J, Redpath A, Capewell S, McMurray JJ. National survey of the prevalence, incidence, primary care burden, and treatment of heart failure in Scotland. *Heart*. 2004;90:1129-36.
- 81. Bukhman G, Mocumbi AO, Atun R, Becker AE, Bhutta Z, Binagwaho A, Clinton C, Coates MM, Dain K, Ezzati M, Gottlieb G, Gupta I, Gupta N, Hyder AA, Jain Y, Kruk ME, Makani J, Marx A, Miranda JJ, Norheim OF, Nugent R, Roy N, Stefan C, Wallis L, Mayosi B (on behalf of the Lancet NPCSG). The Lancet NCDI Poverty Commission: bridging a gap in universal health coverage for the poorest billion. *Lancet*. 2020;396:991-1044.
- 82. Kwan GF, Mayosi BM, Mocumbi AO, Miranda JJ, Ezzati M, Jain Y, Robles G, Benjamin EJ, Subramanian SV and Bukhman G. Endemic Cardiovascular Diseases of the Poorest Billion. *Circulation*. 2016;133:2561-75.

- ***Stewart S**, Sliwa K. Preventing CVD in resource-poor areas: perspectives from the 'real-world'. *Nat Rev Cardiol*. 2009;6:489-92.
- 84. Abhayaratna WP, Smith WT, Becker NG, Marwick TH, Jeffery IM, McGill DA. Prevalence of heart failure and systolic ventricular dysfunction in older Australians: the Canberra Heart Study. *Med J Aust*. 2006;184:151-4.
- 85. Barr EL, Zimmet PZ, Welborn TA, Jolley D, Magliano DJ, Dunstan DW, Cameron AJ, Dwyer T, Taylor HR, Tonkin AM, Wong TY, McNeil J, Shaw JE. Risk of cardiovascular and all-cause mortality in individuals with diabetes mellitus, impaired fasting glucose, and impaired glucose tolerance: the Australian Diabetes, Obesity, and Lifestyle Study (AusDiab). *Circulation*. 2007;116:151-7.
- 86. Cameron AJ, Welborn TA, Zimmet PZ, Dunstan DW, Owen N, Salmon J, Dalton M, Jolley D, Shaw JE. Overweight and obesity in Australia: the 1999-2000 Australian Diabetes, Obesity and Lifestyle Study (AusDiab). *Med J Aust*. 2003;178:427-32.
- 87. Dunstan DW, Zimmet PZ, Welborn TA, Cameron AJ, Shaw J, de Courten M, Jolley D, McCarty DJ, Australian Diabetes O and Lifestyle S. The Australian Diabetes, Obesity and Lifestyle Study (AusDiab)-methods and response rates. *Diabetes Res Clin Pract*. 2002;57:119-29.
- *Chan YK, Tuttle C, Ball J, Teng TK, Ahamed Y, Carrington MJ, <u>Stewart S.</u> Current and projected burden of heart failure in the Australian adult population: a substantive but still ill-defined major health issue. BMC Health Serv Res. 2016;16:501.
- *Strange G, Scalia GM, Playford D, <u>Stewart S</u>. Uncovering the treatable burden of severe aortic stenosis in Australia: current and future projections within an ageing population. *BMC Health Serv Res*. 2021;21:790.
- 90. *Clark RA, Eckert KA, <u>Stewart S</u>, Phillips SM, Yallop JJ, Tonkin AM, Krum H. Rural and urban differentials in primary care management of chronic heart failure: new data from the CASE study. *Med J Aust*. 2007;186:441-5.
- *Carrington MJ, Jennings GL, Clark RA, <u>Stewart S</u>. Assessing cardiovascular risk in regional areas: the Healthy Hearts Beyond City Limits program. *BMC Health Serv Res*. 2012;12:296.
- 92. *Carrington MJ, <u>Stewart S</u>. Cardiovascular disease prevention via a nurse-facilitated intervention clinic in a regional setting: the Protecting Healthy Hearts Program. *Eur J Cardiovasc Nurs*. 2015;14:352-61.
- *Carrington MJ, Jennings GL, <u>Stewart S</u>. Pattern of blood pressure in Australian adults: results from a national blood pressure screening day of 13,825 adults. *Int J Cardiol*. 2010;145:461-7.
- *Clark RA, Coffee N, Turner D, Eckert KA, van Gaans D, Wilkinson D, **STEWART S**, Tonkin AM (Cardiac Aria & Remoteness Index for Australia Investigators). Application of geographic modeling techniques to quantify spatial access to health services before and after an acute cardiac event: the Cardiac Accessibility and Remoteness Index for Australia (ARIA) project. *Circulation*. 2012;125:2006-14.
- 95. *Pretorius S, Sliwa K, Ruf V, Walker K, <u>Stewart S</u>. Feeding the emergence of advanced heart disease in Soweto: a nutritional survey of black African patients with heart failure. *Cardiovasc J Afr.* 2012;23:245-51.
- ***Sliwa K, Carrington M, Mayosi BM, Zigiriadis E, Mvungi R, <u>Stewart S</u>. Incidence and characteristics of newly diagnosed rheumatic heart disease in urban African adults: insights from the heart of Soweto study.** *Eur Heart J.* **2010;31:719-27.**
- 97. *Sliwa K, Carrington MJ, Becker A, Thienemann F, Ntsekhe M, <u>Stewart S</u>. Contribution of the human immunodeficiency virus/acquired immunodeficiency syndrome epidemic to de novo presentations of heart disease in the Heart of Soweto Study cohort. *Eur Heart J.* 2012;33:866-74.
- 98. *Stewart S. Population screening for heart disease in vulnerable populations: lessons from the Heart of Soweto Study. *Heart Lung Circ*. 2009;18:104-6.
- 99. *Stewart S, Carrington M, Pretorius S, Methusi P, Sliwa K. Standing at the crossroads between new and historically prevalent heart disease: effects of migration and socio-economic factors in the Heart of Soweto cohort study. *Eur Heart J.* 2011;32:492-9.

- 100. *Stewart S, Libhaber E, Carrington M, Damasceno A, Abbasi H, Hansen C, Wilkinson D, Sliwa K. The clinical consequences and challenges of hypertension in urban-dwelling black Africans: insights from the Heart of Soweto Study. *Int J Cardiol*. 2011;146:22-7.
- 101. *Stewart S, Mocumbi AO, Carrington MJ, Pretorius S, Burton R, Sliwa K. A not-so-rare form of heart failure in urban black Africans: pathways to right heart failure in the Heart of Soweto Study cohort. *Eur J Heart Fail*. 2011;13:1070-7.
- 102. *Stewart S, Wilkinson D, Becker A, Askew D, Ntyintyane L, McMurray JJ, Sliwa K. Mapping the emergence of heart disease in a black, urban population in Africa: the Heart of Soweto Study. *Int J Cardiol*. 2006;108:101-8.
- 103. *Tibazarwa K, Ntyintyane L, Sliwa K, Gerntholtz T, Carrington M, Wilkinson D, <u>Stewart S</u>. A time bomb of cardiovascular risk factors in South Africa: results from the Heart of Soweto Study "Heart Awareness Days". *Int J Cardiol*. 2009;132:233-9.
- 104. *Stewart S, Carrington MJ, Pretorius S, Ogah OS, Blauwet L, Antras-Ferry J, Sliwa K. Elevated risk factors but low burden of heart disease in urban African primary care patients: a fundamental role for primary prevention. *Int J Cardiol*. 2012;158:205-10.
- *Ogah OS, Sliwa K, Akinyemi JO, Falase AO, <u>Stewart S</u>. Hypertensive heart failure in Nigerian Africans: insights from the Abeokuta Heart Failure Registry. *J Clin Hypertens (Greenwich)*. 2015;17:263-72.
- 106. *Ogah OS, <u>Stewart S</u>, Falase AO, Akinyemi JO, Adegbite GD, Alabi AA, Ajani AA, Adesina JO, Durodola A, Sliwa K. Contemporary profile of acute heart failure in Southern Nigeria: data from the Abeokuta Heart Failure Clinical Registry. *JACC Heart Fail*. 2014;2:250-9.
- 107. *Ogah OS, **STEWART S**, Onwujekwe OE, Falase AO, Adebayo SO, Olunuga T, Sliwa K. Economic burden of heart failure: investigating outpatient and inpatient costs in Abeokuta, Southwest Nigeria. *PLoS One*. 2014;9:e113032.
- *Ojji D, <u>Stewart S</u>, Ajayi S, Manmak M, Sliwa K. A predominance of hypertensive heart failure in the Abuja Heart Study cohort of urban Nigerians: a prospective clinical registry of 1515 de novo cases. *Eur J Heart Fail*. 2013;15:835-42.
- 109. *Thienemann F, Dzudie A, Mocumbi AO, Blauwet L, Sani MU, Karaye KM, Ogah OS, Mbanze I, Mbakwem A, Udo P, Tibazarwa K, Damasceno A, Keates AK, <u>Stewart S</u>, Sliwa K. The causes, treatment, and outcome of pulmonary hypertension in Africa: Insights from the Pan African Pulmonary Hypertension Cohort (PAPUCO) Registry. *Int J Cardiol*. 2016;221:205-11.
- *Thienemann F, Dzudie A, Mocumbi AO, Blauwet L, Sani MU, Karaye KM, Ogah OS, Mbanze I, Mbakwem A, Udo P, Tibazarwa K, Ibrahim AS, Burton R, Damasceno A, <u>Stewart S</u>, Sliwa K. Rationale and design of the Pan African Pulmonary hypertension Cohort (PAPUCO) study: implementing a contemporary registry on pulmonary hypertension in Africa. *BMJ Open*. 2014;4:e005950.
- *Brown A, Carrington MJ, McGrady M, Lee G, Zeitz C, Krum H, Rowley K, <u>Stewart S</u>. Cardiometabolic risk and disease in Indigenous Australians: the heart of the heart study. *Int J Cardiol*. 2014;171:377-83.
- *Tuttle C, Reeves M, Zhong Hu TC, Keates AK, Brady S, Maguire G, <u>Stewart S</u>. Pattern and Outcome of Heart Failure-Related Hospitalization Over 5 Years in a Remote Australian Population: A Retrospective Administrative Data Cohort of 617 Indigenous and non-Indigenous Cases. *J Card Fail*. 2017;23:729-738.
- *Tuttle CS, Carrington MJ, <u>Stewart S</u>, Brown A. Overcoming the tyranny of distance: An analysis of outreach visits to optimise secondary prevention of cardiovascular disease in high-risk individuals living in Central Australia. *Aust J Rural Health*. 2016;24:99-105.
- 114. Etches V, Frank J, Di Ruggiero E, Manuel D. Measuring population health: a review of indicators. *Annu Rev Public Health*. 2006;27:29-55.
- ***Stewart S**, Blue, L. *Improving outcomes in chronic heart failure: Specialist nurse intervention from research to practice (2nd Edition).* 2nd ed. London: BMJ Books; 2004.

- 116. Blue L, Lang E, McMurray JJ, Davie AP, McDonagh TA, Murdoch DR, Petrie MC, Connolly E, Norrie J, Round CE, Ford I and Morrison CE. Randomised controlled trial of specialist nurse intervention in heart failure. *BMJ*. 2001;323:715-8.
- ***Stewart S**, Pearson S, Horowitz JD. Effects of a home-based intervention among patients with congestive heart failure discharged from acute hospital care. *Arch Intern Med.* 1998;158:1067-72.
- 118. *Stewart S, Vandenbroek AJ, Pearson S, Horowitz JD. Prolonged beneficial effects of a home-based intervention on unplanned readmissions and mortality among patients with congestive heart failure. *Arch Intern Med.* 1999;159:257-61.
- 119. *Stewart S, Marley JE, Horowitz JD. Effects of a multidisciplinary, home-based intervention on unplanned readmissions and survival among patients with chronic congestive heart failure: a randomised controlled study. *Lancet*. 1999;354:1077-83.
- *Chan YK, David AM, Mainland C, Chen L, <u>Stewart S</u>. Applying Heart Failure Management to Improve Health Outcomes: But WHICH One? *Card Fail Rev.* 2017;3:113-115.
- *Stewart S. Home is where the heart is when it comes to transitional care in heart failure, but is it the only way to improve health outcomes? *Eur J Heart Fail*. 2017;19:1444-1446.
- *Stewart S, Riegel B, Boyd C, Ahamed Y, Thompson DR, Burrell LM, Carrington MJ, Coats A, Granger BB, Hides J, Weintraub WS, Moser DK, Dickson VV, McDermott CJ, Keates AK, Rich MW. Establishing a pragmatic framework to optimise health outcomes in heart failure and multimorbidity (ARISE-HF): A multidisciplinary position statement. *Int J Cardiol*. 2016;212:1-10.
- 123. Chen L, Chan YK, Busija L, Norekval TM, Riegel B, <u>Stewart S</u>. Malignant and Benign Phenotypes of Multimorbidity in Heart Failure: Implications for Clinical Practice. *J Cardiovasc Nurs*. 2019;34:258-266.
- ***Stewart S**, Pearson S, Luke CG, Horowitz JD. Effects of home-based intervention on unplanned readmissions and out-of-hospital deaths. *J Am Geriatr Soc.* 1998;46:174-80.
- 125. Rich MW, Beckham V, Wittenberg C, Leven CL, Freedland KE, Carney RM. A multidisciplinary intervention to prevent the readmission of elderly patients with congestive heart failure. *N Engl J Med*. 1995;333:1190-5.
- **126.** *McAlister FA, <u>Stewart S</u>, Ferrua S, McMurray JJ. Multidisciplinary strategies for the management of heart failure patients at high risk for admission: a systematic review of randomized trials. *J Am Coll Cardiol*. 2004;44:810-9.
- 127. Clark AM, Thompson DR. The future of management programmes for heart failure. *Lancet*. 2008;372:784-6.
- 128. The cascade effect in the clinical care of patients. N Engl J Med. 1986;315:319-20.
- 129. Mold JW, Stein HF. The cascade effect in the clinical care of patients. *N Engl J Med*. 1986;314:512-4.
- 130. Dickstein K, Cohen-Solal A, Filippatos G, McMurray JJ, Ponikowski P, Poole-Wilson PA, Stromberg A, van Veldhuisen DJ, Atar D, Hoes AW, Keren A, Mebazaa A, Nieminen M, Priori SG, Swedberg K. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2008: the Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2008 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association of the ESC (HFA) and endorsed by the European Society of Intensive Care Medicine (ESICM). *Eur Heart J.* 2008;29:2388-442.
- 131. McMurray JJ, Adamopoulos S, Anker SD, Auricchio A, Bohm M, Dickstein K, Falk V, Filippatos G, Fonseca C, Gomez-Sanchez MA, Jaarsma T, Kober L, Lip GY, Maggioni AP, Parkhomenko A, Pieske BM, Popescu BA, Ronnevik PK, Rutten FH, Schwitter J, Seferovic P, Stepinska J, Trindade PT, Voors AA, Zannad F, Zeiher A. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: The Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2012 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association (HFA) of the ESC. *Eur Heart J.* 2012;33:1787-847.
- 132. Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JGF, Coats AJS, Falk V, Gonzalez-Juanatey JR, Harjola VP, Jankowska EA, Jessup M, Linde C, Nihoyannopoulos P, Parissis JT, Pieske B, Riley JP,

Rosano GMC, Ruilope LM, Ruschitzka F, Rutten FH, van der Meer P. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC)Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur Heart J.* 2016;37:2129-2200.

- ***Stewart S**, Pearson S. Uncovering a multitude of sins: medication management in the home post acute hospitalisation among the chronically ill. *Aust N Z J Med*. 1999;29:220-7.
- *Stewart S, Horowitz JD. Detecting early clinical deterioration in chronic heart failure patients post-acute hospitalisation-a critical component of multidisciplinary, home-based intervention? *Eur J Heart Fail*. 2002;4:345-51.
- 135. Steering Committee of the Queensland Heart Failure Service. Standard for Heart Failure Support Services. 2022.
- *****Driscoll A, Tonkin A, Stewart A, Worrall-Carter L, Thompson DR, Riegel B, Hare DL, Davidson PM, Mulvany C, **STEWART S**. Complexity of management and health outcomes in a prospective cohort study of 573 heart failure patients in Australia: does more equal less? *J Clin Nurs*. 2013;22:1629-38.
- *Driscoll A, Worrall-Carter L, McLennan S, Dawson A, O'Reilly J, **Stewart S**. Heterogeneity of heart failure management programs in Australia. *Eur J Cardiovasc Nurs*. 2006;5:75-82.
- *Driscoll A, Worrall-Carter L, **Stewart S**. Rationale and design of the National Benchmarking and Evidence-based National Clinical Guidelines for Chronic Heart Failure Management Programs Study. *J Cardiovasc Nurs*. 2006;21:276-82.
- 139. Jaarsma T, Stromberg A. Heart failure clinics in Europe. *Prog Cardiovasc Nurs*. 2000;15:67-8.
- 140. Dickstein K, Jaarsma T. The challenge of implementing heart failure management programmes throughout Europe. *Eur J Cardiovasc Nurs*. 2006;5:188-9.
- **141.** *Jaarsma T, Stromberg A, De Geest S, Fridlund B, Heikkila J, Martensson J, Moons P, Scholte op Reimer W, Smith K, **Stewart S**, Thompson DR. Heart failure management programmes in Europe. *Eur J Cardiovasc Nurs*. 2006;5:197-205.
- 142. Davidson PM, Paull G, Introna K, Cockburn J, Davis JM, Rees D, Gorman D, Magann L, Lafferty M, Dracup K. Integrated, collaborative palliative care in heart failure: the St. George Heart Failure Service experience 1999-2002. *J Cardiovasc Nurs*. 2004;19:68-75.
- 143. Doughty RN, Wright SP, Pearl A, Walsh HJ, Muncaster S, Whalley GA, Gamble G, Sharpe N. Randomized, controlled trial of integrated heart failure management: The Auckland Heart Failure Management Study. *Eur Heart J.* 2002;23:139-46.
- 144. Inglis SC, Naismith C, White K, Hendriks JM, Bray J, Hickman LD, Aldridge C, Bardsley K, Cameron J, Candelaria D, Cartledge S, Du H, Ferguson C, Martin L, Selkow T, Xu X, Wynne R, Driscoll A, Gallagher R, Clark R, Davidson PM. CSANZ COVID-19 Cardiovascular Nursing Care Consensus Statement: Executive Summary. *Heart Lung Circ*. 2020;29:1263-1267.
- 145. Gill J. Implantable cardiovascular devices: current and emerging technologies for remote heart failure monitoring. *Cardiol Rev.* 2022.
- 146. Hajduczok AG, Muallem SN, Nudy MS, DeWaters AL and Boehmer JP. Remote monitoring for heart failure using implantable devices: a systematic review, meta-analysis, and meta-regression of randomized controlled trials. *Heart Fail Rev.* 2021.
- **147.** *Whitty JA, Carrington MJ, <u>Stewart S</u>, Holliday J, Marwick TH, Scuffham PA. Patient preferences for the delivery of disease management in chronic heart failure: a qualitative study. *J Cardiovasc Nurs*. 2012;27:201-7.
- *Whitty JA, <u>Stewart S</u>, Carrington MJ, Calderone A, Marwick T, Horowitz JD, Krum H, Davidson PM, Macdonald PS, Reid C,Scuffham PA. Patient preferences and willingness-to-pay for a home or clinic based program of chronic heart failure management: findings from the Which? trial. *PLoS One*. 2013;8:e58347.
- 149. *Stewart S, Carrington MJ, Horowitz JD, Marwick TH, Newton PJ, Davidson PM, Macdonald P, Thompson DR, Chan YK, Krum H, Reid C and Scuffham PA. Prolonged impact of home versus clinic-

- based management of chronic heart failure: extended follow-up of a pragmatic, multicentre randomized trial cohort. *Int J Cardiol*. 2014;174:600-10.
- 150. Ariti CA, Cleland JG, Pocock SJ, Pfeffer MA, Swedberg K, Granger CB, McMurray JJ, Michelson EL, Ostergren J, Yusuf S. Days alive and out of hospital and the patient journey in patients with heart failure: Insights from the candesartan in heart failure: assessment of reduction in mortality and morbidity (CHARM) program. *Am Heart J*. 2011;162:900-6.
- 151. Chen Y, Lawrence J, Stockbridge N. Days alive out of hospital in heart failure: Insights from the PARADIGM-HF and CHARM trials. *Am Heart J*. 2021;241:108-119.
- 152. Faggioni M, Fanaroff AC. Seeking patient-centered trial outcomes: The case for days alive out of hospital. *Am Heart J.* 2022;248:172-174.
- **153.** *Carrington MJ, Kok S, Jansen K, <u>Stewart S</u>. The Green, Amber, Red Delineation of Risk and Need (GARDIAN) management system: a pragmatic approach to optimizing heart health from primary prevention to chronic disease management. *Eur J Cardiovasc Nurs*. 2013;12:337-45.
- *Piepoli MF, Adamo M, Barison A, Bestetti RB, Biegus J, Bohm M, Butler J, Carapetis J, Ceconi C, Chioncel O, Coats A, Crespo-Leiro MG, de Simone G, Drexel H, Emdin M, Farmakis D, Halle M, Heymans S, Jaarsma T, Jankowska E, Lainscak M, Lam CSP, Lochen ML, Lopatin Y, Maggioni A, Matrone B, Metra M, Noonan K, Pina I, Prescott E, Rosano G, Seferovic PM, Sliwa K, Stewart S, Uijl A, Vaartjes I, Vermeulen R, Monique Verschuren WM, Volterrani M, von Heahling S, Hoes A. Preventing heart failure: a position paper of the Heart Failure Association in collaboration with the European Association of Preventive Cardiology. Eur J Heart Fail. 2022;24:143-168.
- *Piepoli MF, Adamo M, Barison A, Bestetti RB, Biegus J, Bohm M, Butler J, Carapetis J, Ceconi C, Chioncel O, Coats A, Crespo-Leiro MG, de Simone G, Drexel H, Emdin M, Farmakis D, Halle M, Heymans S, Jaarsma T, Jankowska E, Lainscak M, Lam CSP, Lochen ML, Lopatin Y, Maggioni A, Matrone B, Metra M, Noonan K, Pina I, Prescott E, Rosano G, Seferovic PM, Sliwa K, Stewart S, Uijl A, Vaartjes I, Vermeulen R, Verschuren WM, Volterrani M, Von Haehling S, Hoes A. Preventing heart failure: a position paper of the Heart Failure Association in collaboration with the European Association of Preventive Cardiology. Eur J Prev Cardiol. 2022;29:275-300.
- *Carrington MJ, <u>Stewart S</u> (on behalf of the NIL-CHF Investigators). Bridging the gap in heart failure prevention: rationale and design of the Nurse-led Intervention for Less Chronic Heart Failure (NIL-CHF) Study. *Eur J Heart Fail*. 2010;12:82-8.
- 157. Jaarsma T, Hill L, Bayes-Genis A, La Rocca HB, Castiello T, Celutkiene J, Marques-Sule E, Plymen CM, Piper SE, Riegel B, Rutten FH, Ben Gal T, Bauersachs J, Coats AJS, Chioncel O, Lopatin Y, Lund LH, Lainscak M, Moura B, Mullens W, Piepoli MF, Rosano G, Seferovic P, Stromberg A. Self-care of heart failure patients: practical management recommendations from the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail*. 2021;23:157-174.
- 158. Riegel B, Dunbar SB, Fitzsimons D, Freedland KE, Lee CS, Middleton S, Stromberg A, Vellone E, Webber DE, Jaarsma T. Self-care research: Where are we now? Where are we going? *Int J Nurs Stud*. 2021;116:103402.
- 159. Riegel B, Jaarsma T. Recognizing the Complexity of Self-Care. Int J Nurs Stud. 2021;116:103908.
- 160. Riegel B, Westland H, Iovino P, Barelds I, Bruins Slot J, Stawnychy MA, Osokpo O, Tarbi E, Trappenburg JCA, Vellone E, Stromberg A, Jaarsma T. Characteristics of self-care interventions for patients with a chronic condition: A scoping review. *Int J Nurs Stud.* 2021;116:103713.
- 161. Nasreddine ZS, Phillips NA, Bedirian V, Charbonneau S, Whitehead V, Collin I, Cummings JL, Chertkow H. The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *J Am Geriatr Soc.* 2005;53:695-9.
- **162.** *Mocumbi AO, Cebola B, Muloliwa A, Sebastiao F, Sitefane SJ, Manafe N, Dobe I, Lumbandali N, Keates A, Stickland N, Chan YK, <u>Stewart S</u>. Differential patterns of disease and injury in Mozambique: New perspectives from a pragmatic, multicenter, surveillance study of 7809 emergency presentations. *PLoS One*. 2019;14:e0219273.

- ***Stewart S**, Chan YK, Playford D, Strange GA (on behalf of the NEDA Investigators). Mild pulmonary hypertension and premature mortality among 154 956 men and women undergoing routine echocardiography. *Eur Respir J.* 2022;59.
- *Mocumbi AO, <u>Stewart S</u>, Patel S, Al-Delaimy WK. Cardiovascular Effects of Indoor Air Pollution from Solid Fuel: Relevance to Sub-Saharan Africa. *Curr Environ Health Rep.* 2019;6:116-126.
- 165. *Stewart S, Al-Delaimy W, Sliwa K, Yacoub M and Mocumbi A. Clinical algorithm to screen for cardiopulmonary disease in low-income settings. *Nat Rev Cardiol*. 2019;16:639-641.
- 166. *Stewart S, Moholdt TT, Burrell LM, Sliwa K, Mocumbi AO, McMurray JJ, Keates AK, Hawley JA. Winter Peaks in Heart Failure: An Inevitable or Preventable Consequence of Seasonal Vulnerability? Card Fail Rev. 2019;5:83-85.
- *Loader J, Chan YK, Hawley JA, Moholdt T, McDonald CF, Jhund P, Petrie MC, McMurray JJ, Scuffham PA, Ramchand J, Burrell LM, <u>Stewart S</u>. Prevalence and profile of "seasonal frequent flyers" with chronic heart disease: Analysis of 1598 patients and 4588 patient-years follow-up. *Int J Cardiol*. 2019;279:126-132.
- *Inglis SC, Clark RA, Shakib S, Wong DT, Molaee P, Wilkinson D, <u>Stewart S</u>. Hot summers and heart failure: seasonal variations in morbidity and mortality in Australian heart failure patients (1994-2005). *Eur J Heart Fail*. 2008;10:540-9.
- 169. *Stewart S, Stewart AR, Waite L, Beilby J. Out with the Old and In with the New: Primary Care Management of Heart Failure with Preserved Ejection Fraction. *Card Fail Rev.* 2022;8:e04.
- 170. *Thompson DR, Roebuck A, <u>Stewart S</u>. Effects of a nurse-led, clinic and home-based intervention on recurrent hospital use in chronic heart failure. *Eur J Heart Fail*. 2005;7:377-84.
- 171. *Yu DS, Lee DT, **STEWART S**, Thompson DR, Choi KC, Yu CM. Effect of Nurse-Implemented Transitional Care for Chinese Individuals with Chronic Heart Failure in Hong Kong: A Randomized Controlled Trial. *J Am Geriatr Soc.* 2015;63:1583-93.
- 172. Brieger D, Amerena J, Attia JR, Bajorek B, Chan KH, Connell C, Freedman B, Ferguson C, Hall T, Haqqani HM, Hendriks J, Hespe CM, Hung J, Kalman JM, Sanders P, Worthington J, Yan T, Zwar NA. NHFA & CSANZ: Australian clinical guidelines for the diagnosis and management of atrial fibrillation 2018. *Med J Aust.* 2018;209:356-362.
- 173. Camm AJ, Lip GY, De Caterina R, Savelieva I, Atar D, Hohnloser SH, Hindricks G, Kirchhof P. 2012 focused update of the ESC Guidelines for the management of atrial fibrillation: an update of the 2010 ESC Guidelines for the management of atrial fibrillation--developed with the special contribution of the European Heart Rhythm Association. *Europace*. 2012;14:1385-413.
- 174. Frendl G, Sodickson AC, Chung MK, Waldo AL, Gersh BJ, Tisdale JE, Calkins H, Aranki S, Kaneko T, Cassivi S, Smith SC, Jr., Darbar D, Wee JO, Waddell TK, Amar D, Adler D and American Association for Thoracic S. 2014 AATS guidelines for the prevention and management of perioperative atrial fibrillation and flutter for thoracic surgical procedures. *J Thorac Cardiovasc Surg*. 2014;148:e153-93.
- 175. Fuster V, Ryden LE, Cannom DS, Crijns HJ, Curtis AB, Ellenbogen KA, Halperin JL, Kay GN, Le Huezey JY, Lowe JE, Olsson SB, Prystowsky EN, Tamargo JL, Wann LS, Smith SC, Jr., Priori SG, Estes NA, 3rd, Ezekowitz MD, Jackman WM, January CT, Lowe JE, Page RL, Slotwiner DJ, Stevenson WG, Tracy CM, Jacobs AK, Anderson JL, Albert N, Buller CE, Creager MA, Ettinger SM, Guyton RA, Halperin JL, Hochman JS, Kushner FG, Ohman EM, Stevenson WG, Tarkington LG, Yancy CW (American College of Cardiology Foundation/American Heart Association Task Force). 2011 ACCF/AHA/HRS focused updates incorporated into the ACC/AHA/ESC 2006 guidelines for the management of patients with atrial fibrillation: a report of the American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines. *Circulation*. 2011;123:e269-367.
- 176. January CT, Wann LS, Alpert JS, Calkins H, Cigarroa JE, Cleveland JC, Jr., Conti JB, Ellinor PT, Ezekowitz MD, Field ME, Murray KT, Sacco RL, Stevenson WG, Tchou PJ, Tracy CM, Yancy CW. 2014 AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: executive summary: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines and the Heart Rhythm Society. *Circulation*. 2014;130:2071-104.

- 177. Kirchhof P. Integrated care of patients with atrial fibrillation: the 2016 ESC atrial fibrillation guidelines. *Heart*. 2017;103:729-731.
- 178. Kirchhof P, Benussi S, Kotecha D, Ahlsson A, Atar D, Casadei B, Castella M, Diener HC, Heidbuchel H, Hendriks J, Hindricks G, Manolis AS, Oldgren J, Popescu BA, Schotten U, Van Putte B, Vardas P, Agewall S, Camm J, Baron Esquivias G, Budts W, Carerj S, Casselman F, Coca A, De Caterina R, Deftereos S, Dobrev D, Ferro JM, Filippatos G, Fitzsimons D, Gorenek B, Guenoun M, Hohnloser SH, Kolh P, Lip GY, Manolis A, McMurray J, Ponikowski P, Rosenhek R, Ruschitzka F, Savelieva I, Sharma S, Suwalski P, Tamargo JL, Taylor CJ, Van Gelder IC, Voors AA, Windecker S, Zamorano JL, Zeppenfeld K. 2016 ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS. *Europace*. 2016;18:1609-1678.
- 179. Lip GYH, Halperin JL, Tse HF. The 2010 European Society of Cardiology Guidelines on the management of atrial fibrillation: an evolution or revolution? *Chest*. 2011;139:738-741.
- 180. Lip GY. Atrial fibrillation in general practice. *Practitioner*. 1999;243:746-51.
- 181. Lip GY. Assessment of risk of thromboembolism in atrial fibrillation: which patients should be anticoagulated? *Eur Heart J.* 1999;20:1757-8.
- 182. Lip GY. Atrial fibrillation and mortality. *Eur Heart J.* 1999;20:1525-7.
- 183. Lip GY. Thromboprophylaxis for atrial fibrillation. *Lancet*. 1999;353:1620-1.
- 184. Ferreira JP, Cleland JG, Lam CSP, Anker SD, Mehra MR, van Veldhuisen DJ, Byra WM, LaPolice DA, Greenberg B, Zannad F. New-onset atrial fibrillation in patients with worsening heart failure and coronary artery disease: the COMMANDER-HF trial. *Clin Res Cardiol*. 2022;111:50-59.
- 185. Kolben Y, Kessler A, Puris G, Nachman D, Alvarez P, Briasoulis A, Asleh R. Management of heart failure with reduced ejection fraction: challenges in patients with atrial fibrillation, renal disease and in the elderly. *Rev Cardiovasc Med*. 2022;23:16.
- 186. Mesubi OO, Anderson ME. Heart Failure and Atrial Fibrillation-Chicken or Egg? *Circ Res*. 2022;130:1011-1013.
- 187. Pabel S, Sossalla S. Atrial fibrillation and heart failure: novel insights into the chicken and egg dilemma. *Eur Heart J.* 2022.
- ***Stewart S**. Bringing order to chaos: the case for integrated management of atrial fibrillation. *Heart*. 2017;103:1928-1929.
- 189. Davey Smith G, Hart C, Ferrell C, Upton M, Hole D, Hawthorne V, Watt G. Birth weight of offspring and mortality in the Renfrew and Paisley study: prospective observational study. *BMJ*. 1997;315:1189-93.
- 190. Hart CL, Hole DJ, Smith GD. Risk factors and 20-year stroke mortality in men and women in the Renfrew/Paisley study in Scotland. *Stroke*. 1999;30:1999-2007.
- 191. Hart CL, Hole DJ, Smith GD. The contribution of risk factors to stroke differentials, by socioeconomic position in adulthood: the Renfrew/Paisley Study. *Am J Public Health*. 2000;90:1788-91.
- 192. Hart CL, Hole DJ, Smith GD. Comparison of risk factors for stroke incidence and stroke mortality in 20 years of follow-up in men and women in the Renfrew/Paisley Study in Scotland. *Stroke*. 2000;31:1893-6.
- 193. Hawthorne VM, Watt GC, Hart CL, Hole DJ, Smith GD, Gillis CR. Cardiorespiratory disease in men and women in urban Scotland: baseline characteristics of the Renfrew/Paisley (midspan) study population. *Scott Med J.* 1995;40:102-7.
- 194. Hole DJ, Watt GC, Davey-Smith G, Hart CL, Gillis CR, Hawthorne VM. Impaired lung function and mortality risk in men and women: findings from the Renfrew and Paisley prospective population study. *BMJ*. 1996;313:711-5; discussion 715-6.
- 195. McCarron P, Hart CL, Hole D, Smith GD. The relation between adult height and haemorrhagic and ischaemic stroke in the Renfrew/Paisley study. *J Epidemiol Community Health*. 2001;55:404-5.
- 196. Smith GD, Hart C, Watt G, Hole D, Hawthorne V. Individual social class, area-based deprivation, cardiovascular disease risk factors, and mortality: the Renfrew and Paisley Study. *J Epidemiol Community Health*. 1998;52:399-405.

- 197. Davenport RJ, Dennis MS, Warlow CP. The accuracy of Scottish Morbidity Record (SMR1) data for identifying hospitalised stroke patients. *Health Bull (Edinb)*. 1996;54:402-5.
- 198. Harley K, Jones C. Quality of Scottish Morbidity Record (SMR) data. *Health Bull (Edinb)*. 1996;54:410-7.
- 199. Kohli HS, Knill-Jones RP. How accurate are SMR1 (Scottish Morbidity Record 1) data? *Health Bull* (*Edinb*). 1992;50:14-23; discussion 29-31.
- 200. Wolf PA, Benjamin EJ, Belanger AJ, Kannel WB, Levy D, D'Agostino RB. Secular trends in the prevalence of atrial fibrillation: The Framingham Study. *Am Heart J.* 1996;131:790-5.
- 201. Frost L, Engholm G, Moller H and Husted. Decrease in mortality in patients with a hospital diagnosis of atrial fibrillation in Denmark during the period 1980-1993. *Eur Heart J*. 1999;20:1592-9.
- 202. Frost L, Vestergaard P, Mosekilde L, Mortensen LS. Trends in incidence and mortality in the hospital diagnosis of atrial fibrillation or flutter in Denmark, 1980-1999. *Int J Cardiol*. 2005;103:78-84.
- 203. Kim D, Yang PS, Jang E, Yu HT, Kim TH, Uhm JS, Kim JY, Pak HN, Lee MH, Joung B, Lip GYH. Increasing trends in hospital care burden of atrial fibrillation in Korea, 2006 through 2015. *Heart*. 2018;104:2010-2017.
- 204. Nisar MU, Munir MB, Sharbaugh MS, Thoma FW, Althouse AD, Saba S. Trends in atrial fibrillation hospitalizations in the United States: A report using data from the National Hospital Discharge Survey. *Indian Pacing Electrophysiol J.* 2018;18:6-12.
- 205. Reinhardt SW, Chouairi F, Miller PE, Clark KAA, Kay B, Fuery M, Guha A, Freeman JV, Ahmad T, Desai NR, Friedman DJ. National Trends in the Burden of Atrial Fibrillation During Hospital Admissions for Heart Failure. *J Am Heart Assoc.* 2021;10:e019412.
- *Olsson LG, Swedberg K, Lappas G, **STEWART S**, Rosengren A. Trends in stroke incidence after hospitalization for atrial fibrillation in Sweden 1987 to 2006. *Int J Cardiol*. 2013;167:733-8.
- 207. *Damasceno A, Gomes J, Azevedo A, Carrilho C, Lobo V, Lopes H, Madede T, Pravinrai P, Silva-Matos C, Jalla S, **Stewart S**, Lunet N. An epidemiological study of stroke hospitalizations in Maputo, Mozambique: a high burden of disease in a resource-poor country. *Stroke*. 2010;41:2463-9.
- 208. *Keates AK, Mocumbi AO, Ntsekhe M, Sliwa K, <u>Stewart S</u>. Cardiovascular disease in Africa: epidemiological profile and challenges. *Nat Rev Cardiol*. 2017;14:273-293.
- 209. Agbor VN, Aminde LN, Tianyi FL, Mbanga CM, Petnga SN, Ditah C, Noubiap JJ. Atrial fibrillation among adults with heart failure in sub-Saharan Africa prevalence, incidence and all-cause mortality: a systematic review and meta-analysis protocol. *BMJ Open*. 2019;9:e022320.
- 210. Jacobs MS, Adeoye AM, Owolabi MO, Tieleman RG, Postma MJ, Van Hulst M. Screening for Atrial Fibrillation in Sub-Saharan Africa: A Health Economic Evaluation to Assess the Feasibility in Nigeria. *Glob Heart*. 2021;16:80.
- 211. Noubiap JJ, Nyaga UF. A review of the epidemiology of atrial fibrillation in sub-Saharan Africa. *J Cardiovasc Electrophysiol*. 2019;30:3006-3016.
- 212. Stambler BS, Ngunga LM. Atrial fibrillation in Sub-Saharan Africa: epidemiology, unmet needs, and treatment options. *Int J Gen Med*. 2015;8:231-42.
- **213.** *Diouf I, Magliano DJ, Carrington MJ, <u>Stewart S</u>, Shaw JE. Prevalence, incidence, risk factors and treatment of atrial fibrillation in Australia: The Australian Diabetes, Obesity and Lifestyle (AusDiab) longitudinal, population cohort study. *Int J Cardiol*. 2016;205:127-132.
- 214. Gallagher C, Hendriks JM, Giles L, Karnon J, Pham C, Elliott AD, Middeldorp ME, Mahajan R, Lau DH, Sanders P, Wong CX. Increasing trends in hospitalisations due to atrial fibrillation in Australia from 1993 to 2013. *Heart*. 2019;105:1358-1363.
- 215. Chugh SS, Roth GA, Gillum RF, Mensah GA. Global burden of atrial fibrillation in developed and developing nations. *Glob Heart*. 2014;9:113-9.
- 216. Lip GYH, Brechin CM, Lane DA. The global burden of atrial fibrillation and stroke: a systematic review of the epidemiology of atrial fibrillation in regions outside North America and Europe. *Chest*. 2012;142:1489-1498.

- 217. Conroy SP, Stevens T, Parker SG, Gladman JR. A systematic review of comprehensive geriatric assessment to improve outcomes for frail older people being rapidly discharged from acute hospital: 'interface geriatrics'. *Age Ageing*. 2011;40:436-43.
- 218. Garrard JW, Cox NJ, Dodds RM, Roberts HC, Sayer AA. Comprehensive geriatric assessment in primary care: a systematic review. *Aging Clin Exp Res.* 2020;32:197-205.
- 219. Sum G, Nicholas SO, Nai ZL, Ding YY, Tan WS. Health outcomes and implementation barriers and facilitators of comprehensive geriatric assessment in community settings: a systematic integrative review [PROSPERO registration no.: CRD42021229953]. *BMC Geriatr.* 2022;22:379.
- 220. Coleman K, Austin BT, Brach C, Wagner EH. Evidence on the Chronic Care Model in the new millennium. *Health Aff (Millwood)*. 2009;28:75-85.
- **221.** *Inglis S, McLennan S, Dawson A, Birchmore L, Horowitz JD, Wilkinson D, **STEWART S**. A new solution for an old problem? Effects of a nurse-led, multidisciplinary, home-based intervention on readmission and mortality in patients with chronic atrial fibrillation. *J Cardiovasc Nurs*. 2004;19:118-27.
- *Carrington MJ, Ball J, Horowitz JD, Marwick TH, Mahadevan G, Wong C, Abhayaratna WP, Haluska B, Thompson DR, Scuffham PA, <u>Stewart S</u>. Navigating the fine line between benefit and risk in chronic atrial fibrillation: rationale and design of the Standard versus Atrial Fibrillation spEcific managemenT study (SAFETY). *Int J Cardiol*. 2013;166:359-65.
- 223. Hendriks JM, de Wit R, Crijns HJ, Vrijhoef HJ, Prins MH, Pisters R, Pison LA, Blaauw Y, Tieleman RG. Nurse-led care vs. usual care for patients with atrial fibrillation: results of a randomized trial of integrated chronic care vs. routine clinical care in ambulatory patients with atrial fibrillation. *Eur Heart J*. 2012;33:2692-9.
- 224. Hendriks JM, de Wit R, Vrijhoef HJ, Tieleman RG, Crijns HJ. An integrated chronic care program for patients with atrial fibrillation: study protocol and methodology for an ongoing prospective randomised controlled trial. *Int J Nurs Stud.* 2010;47:1310-6.
- 225. Hendriks J, Tomini F, van Asselt T, Crijns H, Vrijhoef H. Cost-effectiveness of a specialized atrial fibrillation clinic vs. usual care in patients with atrial fibrillation. *Europace*. 2013;15:1128-35.
- 226. Pathak RK, Evans M, Middeldorp ME, Mahajan R, Mehta AB, Meredith M, Twomey D, Wong CX, Hendriks JML, Abhayaratna WP, Kalman JM, Lau DH, Sanders P. Cost-Effectiveness and Clinical Effectiveness of the Risk Factor Management Clinic in Atrial Fibrillation: The CENT Study. *JACC Clin Electrophysiol*. 2017;3:436-447.
- 227. Hendriks JM, Brooks AG, Rowett D, Moss JR, Gallagher C, Nyfort-Hansen K, Simmons S, Middeldorp ME, Jones T, Thomas G, Lau DH, Sanders P. Home-Based Education and Learning Program for Atrial Fibrillation: Rationale and Design of the HELP-AF Study. *Can J Cardiol*. 2019;35:846-854.
- *Maru S, Byrnes J, Carrington MJ, Chan YK, Thompson DR, <u>Stewart S</u>, Scuffham PA (on behalf of the WHICH? II Trial Investigators). Cost-effectiveness of home versus clinic-based management of chronic heart failure: Extended follow-up of a pragmatic, multicentre randomized trial cohort The WHICH? study (Which Heart Failure Intervention Is Most Cost-Effective & Consumer Friendly in Reducing Hospital Care). *Int J Cardiol*. 2015;201:368-75.
- 229. *Byrnes J, Carrington M, Chan YK, Pollicino C, Dubrowin N, <u>Stewart S</u>, Scuffham PA. Cost-Effectiveness of a Home Based Intervention for Secondary Prevention of Readmission with Chronic Heart Disease. *PLoS One*. 2015;10:e0144545.
- *Gao L, Scuffham P, Ball J, <u>Stewart S</u>, Byrnes J. Long-term cost-effectiveness of a disease management program for patients with atrial fibrillation compared to standard care a multi-state survival model based on a randomized controlled trial. *J Med Econ*. 2021;24:87-95.
- *Byrnes J, Ball J, Gao L, Kai Chan Y, Kularatna S, <u>Stewart S</u>, Scuffham PA. Within trial cost-utility analysis of disease management program for patients hospitalized with atrial fibrillation: results from the SAFETY trial. *J Med Econ.* 2019;22:945-952.
- 232. Gallagher C, Orchard J, Nyfort-Hansen K, Sanders P, Neubeck L, Hendriks JM. NursE led Atrial Fibrillation Management: The NEAT Study: A Randomized Controlled Trial. *J Cardiovasc Nurs*. 2020;35:456-467.

APPENDIX I – COMPETITIVE FUNDING SUPPORTING THIS PORTFOLIO OF RESEARCH

TIMEFRAME	RESEARCH FUNDING	PORTFOLIO OUTPUT
1999-2002	SIMON STEWART 1998 NATIONAL HEART FOUNDATION RALPH READER POST-DOCTORAL OVERSEAS MEDICAL RESEARCH FELLOWSHIP (University of Glasgow, Scotland)	 McMurray JJ, Stewart S. Epidemiology, aetiology, and prognosis of heart failure. Heart 2000; 83(5): 596-602. STEWART S, MacIntyre K, Hole DJ, Capewell S, McMurray JJ. More 'malignant' than cancer? Five-year survival following a first admission for heart failure. Eur J Heart Fail 2001; 3(3): 315-22. STEWART S, MacIntyre K, MacLeod MM, Bailey AE, Capewell S, McMurray JJ. Trends in hospitalization for heart failure in Scotland, 1990-1996. An epidemic that has reached its peak? Eur Heart J 2001; 22(3): 209-17. STEWART S, Hart CL, Hole DJ, McMurray JJ. Population prevalence, incidence, and predictors of atrial fibrillation in the Renfrew/Paisley study. Heart 2001; 86(5): 516-21. STEWART S, Jenkins A, Buchan S, McGuire A, Capewell S, McMurray JJ. The current cost of heart failure to the National Health Service in the UK. Eur J Heart Fail 2002; 4(3): 361-71. STEWART S, MacIntyre K, Capewell S, McMurray JJ. Heart failure and the aging population: an increasing burden in the 21st century? Heart 2003; 89(1): 49-53. STEWART S, McIntyre K, Capewell S, McMurray JJ. Heart failure in a cold climate. Seasonal variation in heart failure-related morbidity and mortality. J Am Coll Cardiol 2002; 39(5): 760-6. STEWART S, Demers C, Murdoch DR, et al. Substantial between-hospital variation in outcome following first emergency admission for heart failure. Eur Heart J 2002; 23(8): 650-7. STEWART S, Hart CL, Hole DJ, McMurray JJ. A population-based study of the long-term risks associated with atrial fibrillation: 20-year follow-up of the Renfrew/Paisley study. Am J Med 2002; 113(5): 359-64. STEWART S, MacIntyre K, Chalmers JW, et al. Trends in case-fatality in 22968
		patients admitted for the first time with atrial fibrillation in Scotland, 1986-1995. <i>Int J Cardiol</i> 2002; 82(3): 229-36.
2002-2006	SIMON STEWART NATIONAL HEALTH & MEDICAL RESEARCH COUNCIL OF AUSTRALIA CAREER DEVELOPMENT RESEARCH AWARD (University of South Australia/University of Queensland, Australia)	 STEWART S, Murphy NF, Walker A, McGuire A, McMurray JJ. Cost of an emerging epidemic: an economic analysis of atrial fibrillation in the UK. Heart 2004; 90(3): 286-92. Clark RA, McLennan S, Dawson A, Wilkinson D, STEWART S. Uncovering a hidden epidemic: a study of the current burden of heart failure in Australia.

		*	Clark RA, McLennan S, Eckert K, Dawson A, Wilkinson D, STEWART S. Chronic heart failure beyond city limits. <i>Rural Remote Health</i> 2005; 5 (4): 443. McLennan SN, Pearson SA, Cameron J, STEWART S. Prognostic importance of cognitive impairment in chronic heart failure patients: does specialist management make a difference? <i>Eur J Heart Fail</i> 2006; 8 (5): 494-501.
2007-2011	SIMON STEWART, J HOROWITZ, T MARWICK, P SCUFFHAM, P DAVIDSON, P MACDONALD, H KRUM, C REID NATIONAL HEALTH & MEDICAL RESEARCH COUNCIL OF AUSTRALIA HEALTH SERVICES RESEARCH GRANT FOR THE WHICH INTERVENTION IS MOST COST- EFFECTIVE& CONSUMER FRIENDLY IN HEART FAILURE (WHICH?) TRIAL (Baker Heart Research Institute, Australia)	*	STEWART S, Carrington MJ, Marwick TH, et al. Impact of home versus clinic-based management of chronic heart failure: the WHICH? (Which Heart Failure Intervention Is Most Cost-Effective & Consumer Friendly in Reducing Hospital Care) multicenter, randomized trial. <i>J Am Coll Cardiol</i> 2012; 60 (14): 1239-48.
2007-2009	SIMON STEWART, A TONKIN, D WILKINSON, R CLARK, J MILLIGAN, P ASTLES AUSTRALIAN RESEARCH COUNCIL OF AUSTRALIA LINKAGE GRANT FOR THE CARDIAC-ARIA PROJECT (University of South Australia/University of Adelaide/AlphaPharm Australia Pty Ltd, Australia)	*	Clark RA, Driscoll A, Nottage J, et al, STEWART S. Inequitable provision of optimal services for patients with chronic heart failure: a national geomapping study. <i>Med J Aust</i> 2007; 186 (4): 169-73. Clark RA, Inglis SC, McAlister FA, Cleland JG, STEWART S. Telemonitoring or structured telephone support programmes for patients with chronic heart failure: systematic review and meta-analysis. <i>BMJ</i> 2007; 34 (7600): 942.
2008-2012	SIMON STEWART NATIONAL HEALTH & MEDICAL RESEARCH COUNCIL OF AUSTRALIA SENIOR RESEARCH FELLOWSHIP (Baker Heart Research Institute, Australia)	*	Sliwa K, Wilkinson D, Hansen C, et al, STEWART S. Spectrum of heart disease and risk factors in a black urban population in South Africa (the Heart of Soweto Study): a cohort study. Lancet 2008; 371(9616): 915-22. STEWART S, Wilkinson D, Hansen C, et al. Predominance of heart failure in the Heart of Soweto Study cohort: emerging challenges for urban African communities. Circulation 2008; 118(23): 2360-67 Sochalski J, Jaarsma T, Krumholz HM, et al, STEWART S. What works in chronic care management: the case of heart failure. Health Aff (Millwood) 2009; 28(1): 179-89. STEWART S, Carrington M, Pretorius S, Methusi P, Sliwa K. Standing at the crossroads between new and historically prevalent heart disease: effects of migration and socio-economic factors in the Heart of Soweto cohort study. Eur Heart J 2011; 32(4): 492-9.
2008-2012	SIMON STEWART, G JENNINGS, P BERGIN, B DE COURTEN, G LEE, M CARRINGTON, C WONG NATIONAL HEALTH & MEDICAL RESEARCH COUNCIL OF AUSTRALIA PROJECT RESEARCH GRANT FOR THE NURSE-LED INTERVENTION FOR LESS CHRONIC HEART FAILURE (NIL-CHF) STUDY (Baker Heart Research Institute, Australia)	*	STEWART S, Chan YK, Wong C, et al. Impact of a nurse-led home and clinic-based secondary prevention programme to prevent progressive cardiac dysfunction in high-risk individuals: the Nurse-led Intervention for Less Chronic Heart Failure (NIL-CHF) randomized controlled study. <i>Eur J Heart Fail</i> 2015; 17 (6): 620-30.

2009-2013	SIMON STEWART, D THOMPSON, W ABHAYARATNA NATIONAL HEALTH & MEDICAL RESEARCH COUNCIL OF AUSTRALIA - PROGRAM OF RESEARCH GRANT FOR "INTEGRATION OF RISK EVALUATION IN CARDIOVASCULAR DISEASE MANAGEMENT PROGRAMS" (Baker Heart Research Institute, Australia)	 Ball J, Carrington MJ, McMurray JJ, STEWART S. Atrial fibrillation: profile and burden of an evolving epidemic in the 21st century. Int J Cardiol 2013; 167(5): 1807-24. Ball J, Thompson DR, Ski CF, Carrington MJ, Gerber T, STEWART S. Estimating the current and future prevalence of atrial fibrillation in the Australian adult population. Med J Aust 2015; 202(1): 32-5. Ball J, Carrington MJ, STEWART S. Mild cognitive impairment in high-risk patients with chronic atrial fibrillation: a forgotten component of clinical management? Heart 2013; 99(8): 542-7. Ball J, Carrington MJ, Wood KA, STEWART S, Investigators S. Women versus men with chronic atrial fibrillation: insights from the Standard versus Atrial Fibrillation spEcific managemenT studY (SAFETY). PLoS One 2013; 8(5): e65795. STEWART S, Ball J, Horowitz JD, et al. Standard versus atrial fibrillation-specific management strategy (SAFETY) to reduce recurrent admission and prolong survival: pragmatic, multicentre, randomised controlled trial. Lancet 2015; 385(9970): 775-84.
2012-2016	SIMON STEWART, J HOROWITZ, M CARRINGTON, P SCUFFHAM, C WONG, P NEWTON, A RISCHBIETH NATIONAL HEALTH & MEDICAL RESEARCH COUNCIL OF AUSTRALIA HEALTH SERVICES PROJECT GRANT FOR THE WHICH INTERVENTION IS MOST COST- EFFECTIVE IN HEART FAILURE (WHICH 2?) TRIAL (Baker Heart Research Institute, Australia)	Scuffham PA, Ball J, Horowitz JD, et al, STEWART S. Standard vs. intensified management of heart failure to reduce healthcare costs: results of a multicentre, randomized controlled trial. Eur Heart J 2017; 38(30): 2340-8.
2012-2017	SIMON STEWART, A BROWN, DR THOMPSON, N STOCKS, P SCUFFHAM, M CARRINGTON, K SLIWA, G MAGUIRE NATIONAL HEALTH & MEDICAL RESEARCH COUNCIL OF AUSTRALIA — CENTRE OF RESEARCH EXCELLENCE TO REDUCE INEQUALITY IN HEART DISEASE (Baker Heart & Diabetes Research Institute/Mary MacKillop Health Services Research Institute, Australia)	 STEWART S, Chan YK, Wong C, et al. Impact of a nurse-led home and clinic-based secondary prevention programme to prevent progressive cardiac dysfunction in high-risk individuals: the Nurse-led Intervention for Less Chronic Heart Failure (NIL-CHF) randomized controlled study. Eur J Heart Fail 2015; 17(6): 620-30. STEWART S, Wiley JF, Ball J, et al. Impact of Nurse-Led, Multidisciplinary Home-Based Intervention on Event-Free Survival Across the Spectrum of Chronic Heart Disease: Composite Analysis of Health Outcomes in 1226 Patients From 3 Randomized Trials. Circulation 2016; 133(19): 1867-77. STEWART S, Ball J, Horowitz JD, et al. Standard versus atrial fibrillation-specific management strategy (SAFETY) to reduce recurrent admission and prolong survival: pragmatic, multicentre, randomised controlled trial. Lancet 2015; 385(9970): 775-84.
2013-2017	SIMON STEWART	 Ball J, Carrington MJ, McMurray JJ, STEWART S. Atrial fibrillation: profile and burden of an evolving epidemic in the 21st century. Int J Cardiol 2013; 167(5): 1807-24.

	NATIONAL HEALTH & MEDICAL RESEARCH COUNCIL OF AUSTRALIA PRINCIPAL RESEARCH FELLOWSHIP (Baker Heart & Diabetes Research Institute/Mary MacKillop Health Services Research Institute, Australia)	 Ball J, Thompson DR, Ski CF, Carrington MJ, Gerber T, STEWART S. Estimating the current and future prevalence of atrial fibrillation in the Australian adult population. Med J Aust 2015; 202(1): 32-5. Ball J, Carrington MJ, STEWART S. Mild cognitive impairment in high-risk patients with chronic atrial fibrillation: a forgotten component of clinical management? Heart 2013; 99(8): 542-7. Ball J, Carrington MJ, Wood KA, STEWART S. Women versus men with chronic atrial fibrillation: insights from the Standard versus Atrial Fibrillation spEcific managemenT studY (SAFETY). PLoS One 2013; 8(5): e65795. STEWART S, Wiley JF, Ball J, et al. Impact of Nurse-Led, Multidisciplinary Home-Based Intervention on Event-Free Survival Across the Spectrum of Chronic Heart Disease: Composite Analysis of Health Outcomes in 1226 Patients From 3 Randomized Trials. Circulation 2016; 133(19): 1867-77.
2014-2018	SIMON STEWART, D THOMPSON, L BURRELL NATIONAL HEALTH & MEDICAL RESEARCH COUNCIL OF AUSTRALIA - PROGRAM OF RESEARCH GRANT FOR "OPTIMISING HEART DISEASE PREVENTION & MANAGEMENT" (Baker Heart & Diabetes Research Institute/Mary MacKillop Health Services Research Institute, Australia)	 STEWART S, Ball J, Horowitz JD, et al. Standard versus atrial fibrillation-specific management strategy (SAFETY) to reduce recurrent admission and prolong survival: pragmatic, multicentre, randomised controlled trial. Lancet 2015; 385(9970): 775-84. STEWART S, Chan YK, Wong C, et al. Impact of a nurse-led home and clinic-based secondary prevention programme to prevent progressive cardiac dysfunction in high-risk individuals: the Nurse-led Intervention for Less Chronic Heart Failure (NIL-CHF) randomized controlled study. Eur J Heart Fail 2015; 17(6): 620-30. STEWART S, Wiley JF, Ball J, et al. Impact of Nurse-Led, Multidisciplinary Home-Based Intervention on Event-Free Survival Across the Spectrum of Chronic Heart Disease: Composite Analysis of Health Outcomes in 1226 Patients From 3 Randomized Trials. Circulation 2016; 133(19): 1867-77. STEWART S, Ball J, Horowitz JD, et al. Standard versus atrial fibrillation-specific management strategy (SAFETY) to reduce recurrent admission and prolong survival: pragmatic, multicentre, randomised controlled trial. Lancet 2015; 385(9970): 775-84.
2019-2024	SIMON STEWART NATIONAL HEALTH & MEDICAL RESEARCH COUNCIL OF AUSTRALIA SENIOR PRINCIPAL RESEARCH FELLOWSHIP (Torrens University Australia/University of Notre Dame Australia, Australia)	Strange G, Playford D, Scalia GM, et al, STEWART S. Change in ejection fraction and long-term mortality in adults referred for echocardiography. Eur J Heart Fail 2021; 23(4): 555-63.

Material not shown due to convinibility