

YALE PEABODY MUSEUM

P.O. BOX 208118 | NEW HAVEN CT 06520-8118 USA | PEABODY.YALE. EDU

JOURNAL OF MARINE RESEARCH

The *Journal of Marine Research*, one of the oldest journals in American marine science, published important peer-reviewed original research on a broad array of topics in physical, biological, and chemical oceanography vital to the academic oceanographic community in the long and rich tradition of the Sears Foundation for Marine Research at Yale University.

An archive of all issues from 1937 to 2021 (Volume 1–79) are available through EliScholar, a digital platform for scholarly publishing provided by Yale University Library at <https://elischolar.library.yale.edu/>.

Requests for permission to clear rights for use of this content should be directed to the authors, their estates, or other representatives. The *Journal of Marine Research* has no contact information beyond the affiliations listed in the published articles. We ask that you provide attribution to the *Journal of Marine Research*.

Yale University provides access to these materials for educational and research purposes only. Copyright or other proprietary rights to content contained in this document may be held by individuals or entities other than, or in addition to, Yale University. You are solely responsible for determining the ownership of the copyright, and for obtaining permission for your intended use. Yale University makes no warranty that your distribution, reproduction, or other use of these materials will not infringe the rights of third parties.



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.
<https://creativecommons.org/licenses/by-nc-sa/4.0/>



*Grazing and Control of Coral Reef Community
Structure by Diadema Antillarum
Philippi (Echinodermata: Echinoidea):
A Preliminary Study*¹

Paul W. Sammarco, Jeffrey S. Levinton, and
John C. Ogden

ABSTRACT

The grazing activities of the tropical echinoid *Diadema antillarum* Philippi can markedly influence the structure of the shallow-water epibenthic coral reef community. The elimination of an entire *Diadema* population from a patch reef in St. Croix, U.S. Virgin Islands resulted in a great increase in macro-algal biomass, alterations in species composition, shifts in dominance, an increase in species numbers, and a decrease in equitability. The increase noted in number of species may be indicative of either initial pre-competitive phases of succession or the creation of a new habitat—the micro-canopy. Experiments involving the caging of *Diadema* have shown that the organism's browsing habits may be extensively disruptive to virgin coral substrate. This evidence introduces the possibility that epifaunal community structure may be controlled through the disturbance of newly settled larvae of sessile epibenthic macro-invertebrates.

1. *Introduction.* The echinoid *Diadema* is one of the most common and conspicuous members of tropical coral reef communities on a global basis. Although a number of investigations have alluded to its ecological function in the community (Lewis, 1964; Randall, Schroeder, and Stark, 1964; Yonge, 1963), the importance of its role remains to be conclusively established.

A keystone species has been defined as the single terminal species on a high trophic level, exerting predatory control on the structure of the levels beneath it (Paine, 1966, 1969). But a wide spectrum of evidence from terrestrial, freshwater, and marine ecosystems has shown that the concept may be expanded to include even the primary-producer/primary-consumer level (see Harper, 1969; Paine and Vadas, 1969; Vadas, 1968). Predation affects prey species diversity in terms of both equitability and number of species (Tansley and

1. Received 26 June 1973; revised 31 July 1973.

Adamson, 1925; Vadas, 1968), species composition (Brooks and Dodson, 1965; Dodson, 1970; Maguire, Belk, and Wells, 1968; Vadas, 1968), longevity and reproductive success (Hope-Simpson, 1940) and succession (Paine and Vadas, 1969; Vadas, 1968). The purpose of this investigation was to preliminarily determine whether *Diadema* was acting in such a manner to maintain control of shallow-water reef community structure.

2. *Methods and Materials.* During the summer of 1972, a preliminary investigation of the effects of grazing by *Diadema antillarum* was conducted at the West Indies Laboratory of Fairleigh Dickinson University on the northern coast of St. Croix, U.S. Virgin Islands (see Sammarco, 1972). During mid-April, 1972 the *Diadema* population (>8,000 individuals) was completely removed from one of five generally homogeneous patch reefs in Tague Bay (see Ogden *et al.*, 1973); this particular patch (PR-2) was determined to be approximately 800 sq. m. in size. Between August 12 and August 28, the macro-algae² of each of the five patches were sampled by six 0.25 sq. m. quadrats laid systematically at uniform intervals along east-west transects. Biomass estimates for each species were determined by placing samples in a Fisher Iso-temp Oven usually for 18–24 hours (but sometimes up to 60 hours, depending on the time required for complete dehydration) and then obtaining dry weights on a Mettler H-10T or P-11 balance.

Another experiment was designed to help determine the degree to which *Diadema*'s grazing activities impose actual physical disturbance on the benthic substrate. Four open-bottom urchin inclusion cages measuring 1.0 × 1.0 × 0.5 m. and containing densities of 0, 2, 4, and 6 *Diadema* individuals/m² were implanted on PR-3. Large pieces of *Acropora palmata* (Coelenterata: Scleractinia) were then collected from the barrier reef, killed by immersing in freshwater for several hours, cleaned with fresh-water forced through a pipette under high pressure, and exposed to the sun to bleach-dry. The coral branches were then subdivided so that two small pieces measuring approximately 25 × 25 cm. could be introduced into each cage. Two weeks after implantation, small subsamples were taken from one of each of the coral pieces and examined for comparative qualitative differences in physical structure with the aid of a binocular dissecting microscope.

3. *Results.* The results of the macro-algae analysis are summarized in Table I. Total algal biomass of all samples for each patch in grams dry weight per square meter is presented with the number of species and Pielou's measure of equitability, J' (Pielou, 1969). A mean and standard deviation are given as a summary of the values for the control patches.

Pieces of coral substrate were examined and found to exhibit varying degrees

2. The term 'macro-algae' used in this paper is defined as those algae readily noticed with the naked eye while sampling.

of physical disturbance. The subsample from the 0/m² density cage showed, as expected, no abrasion of the calices. At a density of 2/m² evidence of grazing was apparent on various parts of the exposed surface. But at densities of 4 and 6/m² the upper portions of the calices had been abraided greatly altering the substrate's physical shape and appearance at a microscopic level.

4. *Discussion.* Although only the initial phases of succession following the perturbation caused by grazing were monitored, the ungrazed patch (PR-2) was found to differ from the controls in the following ways: First, shifts in dominance were noted. *Halimeda opuntia*, the most dominant species by dry weight (the most heavily calcified alga), was replaced by several other species on the ungrazed patch. Although the general order of magnitude of dry-weight for *Halimeda* was not altered, the vast increase in biomass of other algae caused a decline in its relative position in the dominance hierarchy. Most notably, the calcareous phaeophyte *Padina sanctae-crucis* increased in biomass by over two orders of magnitude and became the conspicuous dominant in the absence of grazing. Also found in massive abundances were *Turbinaria turbinata*, *Dictyota* sp., and *Fania adherens*. It is not yet understood whether these dominance shifts and increases in biomass were caused by a high degree of colonization, more rapid vegetative growth, or a combination of these factors (see Table I).

Changes in species diversity as measured by number of species and Pielou's measure of equitability were also striking. Table I shows that the number of species on the ungrazed patch increased by a factor of five. This increment is due to five new species: *Halimeda tuna*, *Liagora* sp., *Dictyopterus delicatula*, and two cyanophytes. None of these was located on any of the four grazed controls. The overdominance of several species, but particularly *Padina sanctae-crucis*, caused the sharp decrease in equitability.

Table I. Macro-algal biomass, species diversity, and equitability in the presence and absence of *Diadema antillarum* Philippi.

Condition	Patch Reef No.	Algal Biomass in gms. dry wt. per sq. m.	Number of Species	$J' (= H'/H'_{\max})$
Controls (Grazed)	PR-1	18.5567	17	0.6596
	PR-3	11.3915	16	0.5695
	PR-4	8.6797	14	0.6293
	PR-5	8.9165	19	0.6821
Mean		11.8861	16.5	0.6351
Standard Deviation		4.6131	2.1	0.0488
Experimental (Ungrazed)	PR-2	159.0183	23	0.4447

It is possible that the absence of *Diadema*'s extensively disruptive grazing activities resulted in the sudden availability of newly opened space and an initial colonization of new species. However, as new opportunistic dominants increase in biomass and cover they occupied the available space. Interspecific competition for space and light may eventually result in a decrease of species numbers to a point lower than that observed on the naturally grazed patches. This has been observed under similar experimental conditions in the *Agarum* kelp-bed communities of Washington (Paine and Vadas, 1969; Vadas, 1968). Increase in species numbers also may be caused by development of a micro-canopy by the leafy fronds of *Padina*, with the concurrent appearance of new understory species. These new species, necessarily sciaphilic (or 'shade-loving'; Boudouresque and Luck, 1972) would be afforded protection from predation by the *Padina* cover. *Padina* is supposedly a low preference food of the locally abundant grazing fish—the Scaridae and Acanthuridae (I. Clavijo, personal communication). In this case, it cannot be predicted that species numbers will decline, as new types of ecological space may have been created.

The results of the caging experiment have shown that *Diadema antillarum*, even at the moderate densities of 4 and 6/m², is able to alter the micro-structure of the hard substratum upon which it grazes. This introduces the possibility that *Diadema* may secondarily control epifaunal species composition, diversity, equitability, dominance, and biomass in a manner similar to its control of algae. The types of fauna most likely to be sensitive to a grazing perturbation might include many of the sessile epibenthic macro-invertebrates which compete for space: hermatypic (Lang, 1971) and ahermatypic corals, zoanthids, actinarians, gorgonians, corallimorphariids, sponges, sedentary polychaetes, etc.

Newman (1960) and Bakus (1964, 1966) suggest that marine grazers may have a strong influence on epifaunal community evolution. Dayton (1971) has experimentally demonstrated that intertidal limpets can affect the success of cirripede settlement. These grazing activities tend to disturb newly settled larvae (Connell, 1961; Hatton, 1938; Lewis, 1954; Southward, 1956; Stimson, 1970) and have been found in some cases to be differential with respect to species (Dayton, 1971). This influence on community structure may be compounded by the fact that certain invertebrate larvae are extremely sensitive to variations in the micro-structure of the benthic substrate and respond to such by altering settling behaviour (Crisp and Barnes, 1954).

Vadas (1968) has found that areas possessing different densities of *Strongylocentrotus* spp. in Washington may be associated with different levels of algal species diversity—that is, maximum diversity at intermediate urchin densities. Observations of fore-reef and back-reef lagoon areas made during April, 1973 in Discovery Bay, Jamaica, West Indies by two of us (P.W.S. and J.S.L.) tend to support Vadas's findings, but further suggest that a similar relationship may exist between *Diadema* densities and the sessile macro-invertebrate epifauna.

5. *Conclusions.* After conducting urchin exclusion experiments in Tague Bay, St. Croix, we conclude that *Diadema antillarum* Philippi may markedly influence the macro-algal community structure of coral reefs by its grazing activities which biomass, species composition, dominance, species numbers, and equitability. It is also concluded that even at moderate densities, *Diadema* can impart an ecologically significant disturbance to the substrate upon which it feeds. On this basis, it has been hypothesized that the urchin may exercise similar control over the community structure of sessile epibenthic macro-invertebrates in the shallower zones of coral reefs.

Acknowledgements. We would like to express our deeply felt gratitude to Robert L. Vadas who served as a primary source of inspiration for this study and who graciously offered personal data and helpful comments during the study's formative stages. We also wish to thank Ileana Clavijo and Rowena Kinsman for aid in collection and identification of algal specimens, and Peter M. J. Woodhead, Director, Discovery Bay Marine Laboratory for valuable advice on field localities in Discovery Bay, Jamaica. This study was supported partially by a National Science Foundation Research Grant (GA-32421) and by a research traineeship from the Division of Biological Sciences, State University of New York at Stony Brook. Contribution no. 14 of the West Indies Laboratory, Fairleigh Dickinson University, St. Croix, U.S. Virgin Islands.

REFERENCES

- BAKUS, GERALD J.
 1964. The effects of fish-grazing on invertebrate evolution in shallow-tropical waters. Allan Hancock Foundation Publication Occasional Paper No. 27, U. of So. Cal. Press, Los Angeles, Cal.
 1966. Some relationships of fishes to benthic organisms on coral reefs. *Nature* 210: 280-284.
- BOUDOURESQUE, CHARLES-FRANCOIS, and HERMANN B. LUCK
 1972. Recherches de bionomie structurale au niveau d'un peuplement benthique sciaphile. *Journal of Experimental Marine Biology and Ecology*, 8(2): 133-144.
- BROOKS, J. L. and S. I. DODSON
 1965. Predation, body size, and composition of plankton. *Science* 150: 28-35.
- CONNELL, J. H.
 1961. Effects of competition, predation by *Thais lapillus*, and other factors on natural populations of the barnacle *Balanus balanoides*. *Ecol. Monogr.*, 31: 61-104.
- CRISP, D. J. and H. BARNES
 1954. The orientation and distribution of barnacles at settlement with particular reference to surface contour. *J. Anim. Ecol.*, 23: 142-162.
- DAYTON, PAUL K.
 1971. Competition, disturbance, and community organization: The provision and subsequent utilization of space in a rocky intertidal community. *Ecol. Monogr.*, 41: 351-389.

DODSON, S. I.

1970. Complementary feeding niches sustained by size-selective predation. *Limnology and Oceanography*, 15(1): 131-137.

HARPER, J. L.

1969. The role of predation in vegetational diversity, *In: Diversity and Stability in Ecological Systems*. U.S. Brookhaven National Laboratory 22nd Symposium.

HATTON, H.

1938. Essais de biologie explicative sur quelques especes intercotidales d'algues et d'animaux. *Ann. Inst. Monaco*, 17: 241-348.

HOPE-SIMPSON, J. F.

1940. Studies of the vegetation of the English Chalk. VI. Late stages in succession leading to Chalk grassland. *J. Ecol.*, 28: 386-402.

LANG, JUDITH

1971. Interspecific aggression by scleractinian corals. 1. The rediscovery of *Scolymia cubensis* (Milne Edwards and Haime). *Bulletin of Marine Science*, 21(4): 952-959.

LEWIS, JOHN B.

1964. Feeding and digestion in the tropical sea urchin *Diadema antillarum* Philippi. *Canadian Journal of Zoology*, 42: 549-557.

LEWIS, J. R.

1954. Observations on a high-level population of limpets. *J. Anim. Ecol.*, 23: 85-100.

MAGUIRE, B. JR., D. BELK, and G. WELLS

1968. Control of community structure by mosquito larvae. *Ecology* 49: 207-210.

NEWMAN, W. A.

1960. The paucity of intertidal barnacles in the tropical western Pacific. *Veliger* 2(4): 89-94.

ODGEN, J. C., R. A. BROWN, and N. SALESKY

1973. Grazing by the echinoid *Diadema antillarum* Philippi: Formation of halos around West Indian patch reefs. *Science* 182(4113): 715-717.

PAINE, ROBERT T.

1966. Food web complexity and species diversity. *American Naturalist*, 100: 65-75.
1969. *Pisaster-Tegula* interaction: Prey patches, predator food preference, and intertidal community structure. *Ecology*, 50: 950-962.

PAINE, ROBERT T. and ROBERT L. VADAS

1969. The effects of grazing by sea urchins, *Strongylocentrotus* spp., on benthic algal populations. *Limnology and Oceanography*, 14(5): 710-719.

PIELOU, E. C.

1969. *An Introduction to Mathematical Ecology*. Wiley Interscience. John Wiley and Sons, New York.

RANDALL, J. E., R. E. SCHROEDER, and W. A. STARK

1964. Notes on the biology of the echinoid *Diadema antillarum*. *Caribb. J. Sci.*, 4(2)(3): 421-433.

SAMMARCO, PAUL W.

1972. Some aspects of the ecology of *Diadema antillarum* Philippi: Food preference and effect of grazing. Special Problems Report to the West Indies Laboratory of Fairleigh Dickinson University, Christiansted, St. Croix. 47 pp.

SOUTHWARD, A. J.

1956. The population balance between limpets and seaweeds on wave-beaten rocky shores. Rep. Mar. Biol. Sta., Pt. Erin, No. 68: 20-29.

STIMSON, J.

1970. Territorial behaviour of the owl limpet, *Lottia gigantea*. Ecology, 51: 113-118.

TANSLEY, A. G. and R. S. ADAMSON

1925. Studies of the vegetation of the English Chalk. III. The Chalk grasslands of the Hampshire-Sussex border. J. Ecol., 13: 177-223.

VADAS, ROBERT L.

1968. The ecology of Agarum and the kelp bed community. Ph. D. thesis. University of Washington, Department of Botany.

YONGE, C. M.

1963. The biology of coral reefs. Advances in Marine Biology 1: 209-260.