

ALGEBRAIC COMBINATORICS


Joseph Ben Geloun & Sanjaye Ramgoolam

Quantum mechanics of bipartite ribbon graphs: Integrality, Lattices and Kronecker coefficients

Volume 6, issue 2 (2023), p. 547-594.

<https://doi.org/10.5802/alco.254>


© The author(s), 2023.

 This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>



*Algebraic Combinatorics is published by The Combinatorics Consortium
and is a member of the Centre Mersenne for Open Scientific Publishing*
www.tccpublishing.org www.centre-mersenne.org
e-ISSN: 2589-5486





Quantum mechanics of bipartite ribbon graphs: Integrality, Lattices and Kronecker coefficients

Joseph Ben Geloun & Sanjaye Ramgoolam

ABSTRACT We define solvable quantum mechanical systems on a Hilbert space spanned by bipartite ribbon graphs with a fixed number of edges. The Hilbert space is also an associative algebra, where the product is derived from permutation group products. The existence and structure of this Hilbert space algebra has a number of consequences. The algebra product, which can be expressed in terms of integer ribbon graph reconnection coefficients, is used to define solvable Hamiltonians with eigenvalues expressed in terms of normalized characters of symmetric group elements and degeneracies given in terms of Kronecker coefficients, which are tensor product multiplicities of symmetric group representations. The square of the Kronecker coefficient for a triple of Young diagrams is shown to be equal to the dimension of a sub-lattice in the lattice of ribbon graphs. This leads to an answer to the long-standing question of a combinatorial interpretation of the Kronecker coefficients. As avenues for future research, we discuss applications of the ribbon graph quantum mechanics in algorithms for quantum computation. We also describe a quantum membrane interpretation of these quantum mechanical systems.

1. INTRODUCTION

Permutation centralizer algebras (PCAs) [73] have been found as an underlying structure which organizes the N -dependences of multi-matrix correlators in super-Yang Mills theories with $U(N)$ gauge symmetry [5, 60, 14, 11, 12, 14, 61, 83, 58, 59]. These correlators are of interest in generalizing beyond the half-BPS sector the link between BPS correlators and Young diagrams [22] in the AdS/CFT correspondence [71, 46, 94].

Permutation methods and PCAs also played a role in the enumeration of observables and the computation of correlators in Gaussian tensor models [8, 9], which have been studied in the context of applications of tensor models to random geometries and holography [1, 47, 86, 48, 95] (see reviews in [30, 63]). An important observation from [8, 9] is that 3-index tensor observables of degree n in a complex tensor model with $U(N)^{\times 3}$ symmetry can be counted using 3-tuples of permutations in S_n , subject to an equivalence relation defined by left and right multiplication by permutations in S_n . A gauge-fixed version of this formulation was described where we have pairs

Manuscript received 29th December 2020, revised 25th May 2022 and 1st July 2022, accepted 3rd July 2022.

KEYWORDS. Belyi maps, Kronecker coefficients, quantum physics, Ribbon graphs.

ACKNOWLEDGEMENTS. SR is supported by the STFC consolidated grant ST/P000754/1 “String Theory, Gauge Theory & Duality” and a Visiting Professorship at the University of the Witwatersrand, funded by a Simons Foundation grant (509116) awarded to the Mandelstam Institute for Theoretical Physics. We thank Robert de Mello Koch, Igor Frenkel, Amihay Hanany for useful discussions on the subject of this paper.

of permutations, subject to an equivalence relation defined using simultaneous conjugation of the pair by a permutation in S_n . These equivalence classes of permutation pairs are known to count bipartite ribbon graphs with n edges (a textbook reference for this subject is [66]). The permutation equivalence classes form an associative algebra, denoted $\mathcal{K}(n)$, with a symmetric non-degenerate bilinear form [9]. As a semi-simple algebra, according to the Wedderburn–Artin theorem, $\mathcal{K}(n)$ is isomorphic to a direct sum of matrix algebras [41]. The explicit isomorphism was constructed using Clebsch–Gordan coefficients of the symmetric group [73, 9]. The matrix basis for the algebra takes the form of $Q_{\tau_1, \tau_2}^{R_1 R_2 R_3}$, where R_1, R_2, R_3 are Young diagrams or partition of n and $\tau_i, i = 1, 2$, range over Clebsch–Gordan multiplicities, also known as Kronecker coefficients (the explicit formula is given in [73] and developed in detail in [9]). Further investigations of tensor models from this algebraic perspective are in [25, 4, 7, 34, 33, 31, 53, 55, 54, 2, 24]. A known connection between bipartite ribbon graphs and Belyi maps [66, 87] gives a topological version of gauge-string duality between tensor models and string theory [8], generalizing analogous correspondences between two-dimensional Yang Mills theory and topological string theory [44, 21, 50].

AdS/CFT holography gives a map between half-BPS states in $U(N)$ Yang-Mills theory at large N and the corresponding space-time geometries [69]. The study of the half-BPS sector as a toy model for questions in the black hole information loss problem [5] raised a question on how restricted sets of $U(N)$ Casimirs can distinguish Young diagrams with a fixed number n (equal to the energy of the BPS state) of boxes. This question is related, by Schur-Weyl duality, to properties of the group algebra of S_n and was studied from this perspective in [57]. A key role in this investigation was played by central elements T_k in the group algebra $\mathbb{C}(S_n)$ associated with permutations having cycle structure consisting of a single cycle of length k (for some $2 \leq k \leq n$) and remaining cycles of length 1.

In addition to these developments from theoretical physics, the investigations in this paper have been guided by the mathematical problem of determining whether there are combinatorial objects which are counted by Kronecker coefficients. While a combinatorial construction of Littlewood-Richarson coefficients, another representation theoretic multiplicity, associated with triples of Young diagrams is well known, it has been a long-standing question whether there exists a family of combinatorial objects, for each triple of Young diagrams, such that the combinatorial objects are enumerated by Kronecker coefficients. This problem was posed in [77] and placed in the context of a number of positivity problems in representation theory in [90] and is discussed in recent papers, e.g. [80, 72]. This mathematical question which may appear, at least at first sight to many physicists, to be a somewhat esoteric question, has inspired substantial recent activity and progress at the intersection of computational complexity theory, quantum information theory and representation theory. We will not attempt to give a summary of this thriving area of research, but will point the reader to some papers which give a flavour of this field [80, 72, 75, 15, 52, 79, 81].

A way to understand the problem is to compare two known computations in representation theory. The computation of characters $\chi_R(\sigma)$ of a permutation $\sigma \in S_n$ in a representation associated to Young diagram R with n boxes can be done by using the Murnaghan–Nakayama rule [76, 78]. This can be phrased in terms of the counting of a certain pattern of labellings of the boxes in R by numbers according to a rule determined by the cycle structure of σ (see for example [92][93]). In this construction, it is clear why the outcome is an integer - which is a somewhat special property of symmetric group characters, a property not shared by generic finite groups. The

Kronecker coefficient can be computed using the formula

$$(1) \quad C(R_1, R_2, R_3) = \frac{1}{n!} \sum_{\sigma \in S_n} \chi_{R_1}(\sigma) \chi_{R_2}(\sigma) \chi_{R_3}(\sigma).$$

In this formula, it is not clear why the sum over all the conjugacy classes in S_n for general n ends up giving an outcome which is a non-negative integer - although from the representation theory definition as the number of invariants in the tensor product of $R_1 \otimes R_2 \otimes R_3$, it is clear why this is the case. A combinatorial interpretation should give a new way to make it manifest that $C(R_1, R_2, R_3)$ is a non-negative integer.

The following formula which has played a role in counting tensor model invariants shows that bipartite ribbon graphs (also called ribbon graphs for short in this paper) hold some promise of progress on this problem. It is known that the total number of bipartite ribbon graphs with n edges is equal to the sum of squares of Kronecker coefficients [8, 73, 34, 9]

$$(2) \quad |\text{Rib}(n)| = \sum_{R_1, R_2, R_3 \vdash n} C(R_1, R_2, R_3)^2.$$

This formula shows that the sum of squares of Kronecker coefficients does have a combinatorial and geometric interpretation. Bipartite ribbon graphs have an elegant group theoretic characterisation in terms of pairs of permutations with an equivalence under simultaneous conjugation. A natural question is : Is it possible to refine this link to give an interpretation of a fixed $C(R_1, R_2, R_3)^2$, and a fixed $C(R_1, R_2, R_3)$, in terms of ribbon graphs? We would like an interpretation which makes the non-negative integer property of the Kronecker coefficients manifest. And are there combinatorial algorithms based on this interpretation for computing Kronecker coefficients?

The algebras $\mathcal{K}(n)$, and analogous algebras related to Littlewood-Richardson coefficients, have been studied in the theoretical physics literature primarily as a tool to understand the structure of the space of gauge invariant observables and their correlators in matrix/tensor models and in AdS/CFT (see [85] for a short review). In this paper, motivated by the mathematical question of a combinatorial interpretation of Kronecker coefficients and the connections of this question to quantum information and complexity theory, we introduce a new physical perspective on these algebras. We propose that studying solvable quantum mechanics models on algebras such as $\mathcal{K}(n)$, which are related to interesting combinatorial objects (in this case bipartite ribbon graphs) having elegant descriptions in terms of symmetric groups (in this case permutation pairs subject to an equivalence generated by conjugation with a permutation), can be a fruitful avenue to explore interesting interfaces between physics, mathematics and computational complexity theory.

Section 2 develops the quantum mechanics on $\mathcal{K}(n)$. $\mathcal{K}(n)$ is a subspace of $\mathbb{C}(S_n) \otimes \mathbb{C}(S_n)$ which is invariant under conjugation by $\gamma \otimes \gamma$ for $\gamma \in S_n$. As a vector space, it has two interesting bases. There is a basis E_r of elements labelled by an index r ranging over equivalence classes of pairs $(\sigma_1, \sigma_2) \in S_n \times S_n$, with the equivalence relation

$$(3) \quad (\sigma_1, \sigma_2) \sim (\gamma \sigma_1 \gamma^{-1}, \gamma \sigma_2 \gamma^{-1}),$$

defined using $\gamma \in S_n$. We refer to this basis as the geometric ribbon graph basis. There is another basis labelled by triples of Young diagrams (R_1, R_2, R_3) , where each Young diagram has n boxes, such that the Kronecker coefficient $C(R_1, R_2, R_3)$ is non-zero. We refer to this as the Fourier basis for $\mathcal{K}(n)$. In Section 2.1, we review (from [8, 73, 9]) the formula (14) for the Fourier basis elements in terms of matrix elements and Clebsch-Gordan coefficients of S_n . The Fourier basis also makes the Wedderburn-Artin decomposition of $\mathcal{K}(n)$ into matrix algebras manifest. We define

a natural inner product on $\mathcal{K}(n)$ inherited from $\mathbb{C}(S_n) \otimes \mathbb{C}(S_n)$ and prove that $\mathcal{K}(n)$ is a Hilbert space (Proposition 2.1). We prove that the product structure on $\mathcal{K}(n)$ in the geometric ribbon graph basis is given by integers (Section 2.3).

The fact that $\mathcal{K}(n)$ is a vector space as well as an algebra (i.e. vector space equipped with an associative product) with a known Wedderburn–Artin decomposition can be exploited to write down interesting solvable Hamiltonians for quantum mechanical systems having $\mathcal{K}(n)$ as a Hilbert space. We introduce a set of Hermitian operators $T_k^{(i)}$ on $\mathcal{K}(n)$ which are central elements of $\mathcal{K}(n)$ and act on $\mathcal{K}(n)$ using the product operation in the algebra. The indices take values $i \in \{1, 2, 3\}$ and $k \in \{2, 3, \dots, \tilde{k}_*\}$. The number \tilde{k}_* is chosen to obey $\tilde{k}_* \geq k_*(n)$, where $k_*(n)$ is an integer between 2 to n . $k_*(n)$ is defined [57] as the minimum integer such that the central elements T_k in $\mathbb{C}(S_n)$ with k ranging in $\{2, 3, \dots, k_*(n)\}$ generate the centre. The precise definition of the operators $T_k^{(i)}$, which we call reconnection operators, is given in Section 2.4. It is shown (Proposition 2.2) that the matrix elements of these operators in the geometric ribbon graph basis are non-negative integers.

In Section 3, we introduce the notion of the Fourier subspace of $\mathcal{K}(n)$ associated with a triple of Young diagrams (R_1, R_2, R_3) . This subspace has dimension $C(R_1, R_2, R_3)^2$. Proposition 3.1 shows that the Fourier basis elements are eigenvectors of the reconnection operators, with eigenvalues given by normalized characters of symmetric groups. Proposition 3.2 shows that the eigenvalue sets of reconnection operators chosen with $k \in \{2, \dots, \tilde{k}_*\}$ can be used to distinguish Fourier subspaces associated with distinct triples of Young diagrams. These results are used (Section 3.2) to construct for each n , and each triple (R_1, R_2, R_3) , a rectangular matrix of integers having a null space which spans the Fourier subspace of the specified triple. Section 3.3 constructs Hamiltonians as linear combinations of the reconnection matrices, which are square (non-negative) integer matrices in the geometric basis and distinguish Fourier subspaces with distinct Young diagram triples. Using Proposition 3.1, the eigenvalues of these Hamiltonians are expressed as linear combinations of normalized symmetric group characters. The eigenspaces for distinct eigenvalues are the Fourier subspaces for distinct Young diagram triples.

The realisation of Fourier subspaces in $\mathcal{K}(n)$ labelled by Young diagram triples (R_1, R_2, R_3) as eigenspaces of integer reconnection matrices is thus one of two important inputs in our discussion. It means that while the formula (14) for Fourier basis elements uses detailed representation theoretic data such as matrix elements of permutations in some chosen basis for symmetric group representations along with Clebsch–Gordan coefficients, there is a new approach to the Fourier subspace of a triple of Young diagrams based on integer reconnection matrices. Now generic integer matrices do not necessarily have integer or rational eigenvalues (see for example [36]). For the reconnection matrices at hand however we know, using symmetric group representation theory (Proposition 3.1 along with Lemma 3.3), that the eigenvalues are integers. These eigenvalues are known to be calculable using combinatorial algorithms, notably the Murnaghan–Nakayama rule. Thus, we are able to replace the more obvious (but computationally expensive) computation of the Fourier subspace using direct implementation of the formula (14) with the calculation of null spaces of integer matrices which takes two combinatorial inputs: the combinatorics of reconnection matrices and the Murnaghan–Nakayama algorithm. This allows us to express the problem of finding the Fourier subspaces of Young diagram triples as a question about null spaces of integer matrices. This in turn allows us to access results from the subject of integer matrices and lattice algorithms.

Section 4 recalls a key result from the integer matrices and lattice algorithms. Any integer matrix, square or rectangular, has a unique Hermite normal form (HNF). There are standard algorithms in computational number theory for finding the HNF (see e.g. [20, 88, 74]) and such algorithms are also accessible in group theoretic software such as SAGE or GAP [39]. A consequence is that, for the Fourier subspaces associated with Young diagram triples defined in Section 3, there are bases which are integer linear combinations of the geometric ribbon graph basis vectors. For each triple (R_1, R_2, R_3) , given a choice of the rectangular matrix (which can be specified using a choice of k_* as in Section 3.2) or square matrix (specified using a Hamiltonian as in Section 3.3), any HNF algorithm leads to a list of linearly independent integer null vectors, which are $C(R_1, R_2, R_3)^2$ in number. This list of integer null vectors specifies a sub-lattice in the lattice $\mathbb{Z}^{|\text{Rib}(n)|}$ in $\mathcal{K}(n)$ generated by all integer linear combinations of the geometric ribbon graph vectors. This provides (Theorem 4.1 and Corollary 4.2) a positive answer to the questions of a combinatorial interpretation and construction for the square of the Kronecker coefficient.

It is natural to ask if a construction of $C(R_1, R_2, R_3)$ rather than its square can be given along these lines. To this end, we consider an operation on bipartite ribbon graphs, which has previously been studied in the context of Belyi maps [56]. In the permutation pair description of ribbon graphs, this operation amounts to inverting both permutations. In Section 4.3 we study a linear involution S (also called conjugation) on $\mathcal{K}(n)$ defined using this inversion. Comparing the action of the involution on the ribbon graph basis with its action on the Fourier basis elements (14) leads to the result that the sum of Kronecker coefficients is equal to the number of self-conjugate ribbon graphs. Considering linear operators acting on $\mathcal{K}(n)$ constructed from the reconnection operators $T_k^{(i)}$ as well as the conjugation operator S leads to sub-lattices of dimension $C(R_1, R_2, R_3)(C(R_1, R_2, R_3) + 1)/2$, $C(R_1, R_2, R_3)(C(R_1, R_2, R_3) - 1)/2$, both of which come equipped with a list of linearly independent integer basis vectors from an HNF construction. Choosing an injection from the set of basis vectors of the smaller sub-lattice into the set of basis vectors of the larger sub-lattice yields a subset of basis vectors of the larger sub-lattice, which equal $C(R_1, R_2, R_3)$ in number. This realises $C(R_1, R_2, R_3)$ as the dimension of a sub-lattice in $\mathbb{Z}^{|\text{Rib}(n)|}$.

In the concluding section we give a summary of our results. While the content of this paper is primarily mathematical, its motivations come from the physics of strings and tensor models. The concluding section thus includes a description of future research directions based on the links to physics. There is a more extended discussion setting up the first steps for these future directions in the arxiv version of this paper [10]. The appendices give some detailed steps in the proofs and examples of results from the computation of Fourier basis vectors using reconnection operators. The last appendix gives key parts of the GAP code used.

2. QUANTUM MECHANICS OF RIBBON GRAPHS: COMMUTING HAMILTONIANS FROM CENTRES OF ALGEBRAS $\mathcal{K}(n)$

In this section we set up the quantum mechanics of bipartite ribbon graphs using their description in terms of permutation groups. We introduce the space of states, two bases for the space (a geometric basis and a Fourier basis), an inner product and Hermitian operators on the state space which have eigenvalues expressible in terms of normalized symmetric group characters.

2.1. REVIEW OF PREVIOUS RESULTS ON THE ALGEBRA $\mathcal{K}(n)$ OF BIPARTITE RIBBON GRAPHS. We give an overview of the description of bipartite ribbon graphs in terms of symmetric groups. A useful textbook reference is [66] which gives references to the

original mathematical literature. We will also be making extensive use of formulae from the representation theory of symmetric groups. A mathematical physics reference is [49]. The key formulae are summarised the appendices of [9].

2.1.1. *Counting bipartite ribbon graphs.* A bipartite ribbon graph, also called a hypermap, is a graph embedded on a two-dimensional surface with black and white vertices, such that edges connect black to white vertices and cutting the surface along the edges leaves a disjoint union of regions homeomorphic to open discs. Bipartite ribbon graphs, denoted ribbon graphs for short in this paper, with n edges can be described using permutations of $\{1, 2, \dots, n\}$ forming the symmetric group S_n . Label the edges with integers $\{1, 2, \dots, n\}$. Reading the edges around the black vertices following a chosen orientation on the surface gives the cycles of a permutation τ_1 , while the white vertices similarly give a permutation τ_2 . Relabelling the edges, $i \rightarrow \mu(i)$ using $\mu \in S_n$, amounts to conjugating the pair $(\tau_1, \tau_2) \rightarrow (\mu\tau_1\mu^{-1}, \mu\tau_2\mu^{-1})$. Distinct ribbon graphs are thus equivalence classes of pairs $(\tau_1, \tau_2) \in S_n \times S_n$ under the equivalence relation

$$(4) \quad (\tau_1, \tau_2) \sim (\tau'_1, \tau'_2) \text{ iff } \exists \mu \in S_n, (\tau'_1, \tau'_2) = (\mu\tau_1\mu^{-1}, \mu\tau_2\mu^{-1})$$

The set of permutation pairs within a fixed equivalence class forms an orbit for the action of S_n on $S_n \times S_n$ given in (4). We define $\text{Rib}(n)$ to be the set of equivalence classes, or the set of orbits. There are commands in group theoretic software GAP [39] that directly generate these orbits for any n , see `RibbSetFunction(n)` appendix C. As an example, consider the case $n = 3$. These are the 11 ribbon graphs shown in Figure 1. The label appearing below each ribbon graph is an index running from 1 to 11. The sole non-planar (genus one) ribbon graph is the equivalence class containing the pair $[(123), (123)]$.




FIGURE 1. Bipartite ribbon graphs with $n = 3$ edges

The counting of ribbon graphs is also related to the counting of bipartite graphs with n trivalent vertices with three incoming colored edges and n trivalent vertices with three outgoing colored edges [8, 9]. This counting problem corresponds to equivalence classes of triples $(\sigma_1, \sigma_2, \sigma_3) \sim (\mu_1\sigma_1\mu_2, \mu_1\sigma_2\mu_2, \mu_1\sigma_3\mu_3)$ for $\sigma_i \in S_n$ and $\mu_1, \mu_2 \in S_n$. In turn this counting also gives the number of linearly independent degree n polynomial functions of tensor variables Φ_{i_1, i_2, i_3} and $\bar{\Phi}^{\bar{i}_1, \bar{i}_2, \bar{i}_3}$ invariant the action of $U(N)^{\times 3}$, for $n \leq N$ and with (i_1, i_2, i_3) transforming as the fundamental of the unitary group and $(\bar{i}_1, \bar{i}_2, \bar{i}_3)$, transforming in the anti-fundamental. Our focus in this paper will be on ribbon graphs, and we will discuss tensor model observables further in the outlook Section 5.

2.1.2. *The permutation centralizer algebra (PCA) $\mathcal{K}(n)$ and its geometric basis.* Introducing the group algebra $\mathbb{C}(S_n)$, consider the elements of $\mathbb{C}(S_n) \otimes_{\mathbb{C}} \mathbb{C}(S_n)$, written more simply $\mathbb{C}(S_n) \otimes \mathbb{C}(S_n)$, obtained by starting with a tensor product $\sigma_1 \otimes \sigma_2$ and summing all their diagonal conjugates as

$$(5) \quad \sigma_1 \otimes \sigma_2 \rightarrow \sum_{\gamma \in S_n} \gamma \sigma_1 \gamma^{-1} \otimes \gamma \sigma_2 \gamma^{-1}.$$

Two pairs (σ_1, σ_2) and (σ'_1, σ'_2) related by the equivalence (4) produce the same sum. Now, consider the \mathbb{C} -vector subspace $\mathcal{K}(n) \subset \mathbb{C}(S_n) \otimes \mathbb{C}(S_n)$ spanned by all $\sum_{\gamma \in S_n} \gamma \sigma_1 \gamma^{-1} \otimes \gamma \sigma_2 \gamma^{-1}$, σ_1 and $\sigma_2 \in S_n$:

$$(6) \quad \mathcal{K}(n) = \text{Span}_{\mathbb{C}} \left\{ \sum_{\gamma \in S_n} \gamma \sigma_1 \gamma^{-1} \otimes \gamma \sigma_2 \gamma^{-1}, \sigma_1, \sigma_2 \in S_n \right\}.$$

The dimension of $\mathcal{K}(n)$ is equal to the number of ribbon graphs with n edges, i.e. $|\text{Rib}(n)|$. In [9], it is shown that $\mathcal{K}(n)$ is an associative algebra, with the product being inherited from $\mathbb{C}(S_n) \otimes \mathbb{C}(S_n)$. $\mathcal{K}(n)$ is a permutation centralizer algebra (PCA) - a subspace of an algebra with basis given by permutations forming a group (here permutation pairs (σ_1, σ_2) forming the group $S_n \times S_n$), which commutes with a subgroup of the permutations, here (γ, γ) forming the diagonal subgroup $S_n \subset S_n \times S_n$. $\mathcal{K}(n)$ is also semi-simple: it has a non-degenerate symmetric bilinear pairing given by

$$(7) \quad \delta_2 : \mathbb{C}(S_n)^{\otimes 2} \times \mathbb{C}(S_n)^{\otimes 2} \rightarrow \mathbb{C}$$

where

$$(8) \quad \delta_2(\otimes_{i=1}^2 \sigma_i; \otimes_{i=1}^2 \sigma'_i) = \prod_{i=1}^2 \delta(\sigma_i \sigma'_i{}^{-1})$$

which extends to linear combinations with complex coefficients. Semi-simplicity implies that, by the Wedderburn–Artin theorem [41, 84], $\mathcal{K}(n)$ admits a decomposition in simple matrix algebras. This decomposition is made manifest using what we denote as the Fourier basis, which we discuss shortly in Section 2.1.3.

Start with a ribbon graph with label $r \in \{1, \dots, |\text{Rib}(n)|\}$. As discussed in Section 2.1.1, the set of ribbon graphs is in 1-1 correspondence with orbits of the action of S_n on $S_n \times S_n$, by conjugation as given in (4). Pick a pair of permutations $(\tau_1^{(r)}, \tau_2^{(r)})$ among the permutation pairs representing the ribbon graph r . The orbit $\text{Orb}(r)$ is the set of elements in $S_n \times S_n$ which can be written as $(\mu\tau_1^{(r)}\mu^{-1}, \mu\tau_2^{(r)}\mu^{-1})$ for some $\mu \in S_n$. Hence, consider the basis element in $\mathcal{K}(n)$ associated with r as:

$$(9) \quad E_r = \frac{1}{n!} \sum_{\mu \in S_n} \mu\tau_1^{(r)}\mu^{-1} \otimes \mu\tau_2^{(r)}\mu^{-1}.$$

Let $\text{Aut}(\tau_1^{(r)}, \tau_2^{(r)})$ be the subgroup of S_n which leaves fixed the pair $(\tau_1^{(r)}, \tau_2^{(r)})$. The order of this group is $|\text{Aut}(\tau_1^{(r)}, \tau_2^{(r)})|$ and is independent of the choice of representative, so we can write this as $|\text{Aut}(r)|$. The orbit-stabilizer theorem (see for example [16]) gives an isomorphism between $\text{Orb}(r)$ and the coset $S_n/\text{Aut}(\tau_1^{(r)}, \tau_2^{(r)})$. Let a be a label for the distinct permutation pairs in $\text{Orb}(r)$:

$$\begin{aligned}
 E_r &= \frac{1}{n!} \sum_{\mu \in S_n} \mu \tau_1^{(r)} \mu^{-1} \otimes \mu \tau_2^{(r)} \mu^{-1} \\
 &= \frac{|\text{Aut}(r)|}{n!} \sum_{a \in \text{Orb}(r)} \tau_1^{(r)}(a) \otimes \tau_2^{(r)}(a) \\
 (10) \quad &= \frac{1}{|\text{Orb}(r)|} \sum_{a \in \text{Orb}(r)} \tau_1^{(r)}(a) \otimes \tau_2^{(r)}(a).
 \end{aligned}$$

Both these expressions for E_r will be useful. We will refer to the E_r as the geometric or ribbon graph basis vectors for $\mathcal{K}(n)$.

The pairing (7) evaluated on this basis is

$$\begin{aligned}
 \delta_2(E_r, E_s) &= \frac{1}{n!} \sum_{\gamma} \delta(\sigma_1^{(r)} \gamma (\sigma_1^{(s)})^{-1} \gamma^{-1}) \delta(\sigma_2^{(r)} \gamma (\sigma_2^{(s)})^{-1} \gamma^{-1}) \\
 (11) \quad &= \frac{1}{n!} |\text{Aut}(r)| \delta_{sr} = \frac{1}{|\text{Orb}(r)|} \delta_{rs}.
 \end{aligned}$$

The basis vectors associated with distinct orbits $r \neq s$ are orthogonal.

2.1.3. *A Fourier basis for $\mathcal{K}(n)$.* The number of bipartite ribbon graphs with n edges, which is the dimension of $\mathcal{K}(n)$, can be given as a sum of partitions of n [8, 9] or as a sum over triples R_1, R_2, R_3 of irreducible representations (irreps) of S_n :

$$(12) \quad |\text{Rib}(n)| = \text{Dim}(\mathcal{K}(n)) = \sum_{R_1, R_2, R_3 \vdash n} C(R_1, R_2, R_3)^2 = \sum_{p \vdash n} \text{Sym}(p).$$

R_1, R_2, R_3 are partitions of n (denoted by $R_i \vdash n$) which correspond to Young diagrams with n boxes. We will denote their dimension as $d(R_i)$.

Describing p in terms of a set of numbers p_i giving the multiplicity of parts i in the partition,

$$(13) \quad \text{Sym}(p) = \prod_i i^{p_i} p_i!.$$

The form of this sum of squares is explained by the Wedderburn–Artin decomposition of $\mathcal{K}(n)$: an explicit basis, which we refer to as the Fourier basis and which exhibits the decomposition, can be constructed using Clebsch–Gordan coefficients and matrix elements of permutation groups [73, 9]. This basis takes the form

$$(14) \quad Q_{\tau_1, \tau_2}^{R_1, R_2, R_3} = \kappa_{R_1, R_2} \sum_{\sigma_1, \sigma_2 \in S_n} \sum_{i_1, i_2, i_3, j_1, j_2} C_{i_1, i_2; i_3}^{R_1, R_2; R_3, \tau_1} C_{j_1, j_2; i_3}^{R_1, R_2; R_3, \tau_2} D_{i_1 j_1}^{R_1}(\sigma_1) D_{i_2 j_2}^{R_2}(\sigma_2) \sigma_1 \otimes \sigma_2.$$

$D_{ij}^R(\sigma)$ are the matrix elements of the linear operator $D^R(\sigma)$ in an orthonormal basis for the irrep R . The indices τ_1, τ_2 run over an orthonormal basis for the multiplicity space of R_3 appearing in the tensor decomposition of $R_1 \otimes R_2$. This multiplicity is equal to the Kronecker coefficient $C(R_1, R_2, R_3)$ which is also the multiplicity of the trivial representation in the tensor product decomposition of $R_1 \otimes R_2 \otimes R_3$. $\kappa_{R_1, R_2} = \frac{d(R_1)d(R_2)}{(n!)^2}$ is a normalization factor, where $d(R_i)$ is the dimension of the irrep R_i . $C_{i_1, i_2; i_3}^{R_1, R_2; R_3, \tau_1}$ are Clebsch–Gordan coefficients of the representations of S_n

(see the appendices of [9] for the properties needed to prove that this expression gives a Wedderburn–Artin basis for $\mathcal{K}(n)$).

These elements $Q_{\tau_1, \tau_2}^{R_1, R_2, R_3} \in \mathbb{C}(S_n) \otimes \mathbb{C}(S_n)$ are invariant under diagonal conjugation

$$(15) \quad (\gamma \otimes \gamma) \cdot Q_{\tau_1, \tau_2}^{R_1, R_2, R_3} \cdot (\gamma^{-1} \otimes \gamma^{-1}) = Q_{\tau_1, \tau_2}^{R_1, R_2, R_3},$$

and therefore belong to $\mathcal{K}(n)$. It was verified [9] that they define the Wedderburn–Artin matrix bases of $\mathcal{K}(n)$:

$$(16) \quad Q_{\tau_1, \tau_2}^{R_1, R_2, R_3} Q_{\tau'_2, \tau'_3}^{R'_1, R'_2, R'_3} = \delta_{R_1 R'_1} \delta_{R_2 R'_2} \delta_{R_3 R'_3} \delta_{\tau_2 \tau'_2} Q_{\tau_1, \tau_3}^{R_1, R_2, R_3}.$$

The normalization κ_{R_1, R_2} is chosen to ensure that the RHS has the standard form for multiplication of elementary matrices, for each block labelled by triples (R_1, R_2, R_3) with non-vanishing Kronecker coefficient $C(R_1, R_2, R_3)$. Noting that $C(R_1, R_2, R_3)$ is at most 1 for $n \leq 4$, then the matrices $Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}$ are 1×1 hence are commuting for $n \leq 4$. The set $\{Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}\} \in \mathcal{K}(n)$ are orthogonal with respect to the bilinear pairing δ_2

$$(17) \quad \delta_2(Q_{\tau_1, \tau'_1}^{R_1, R_2, R_3}; Q_{\tau_2, \tau'_2}^{R'_1, R'_2, R'_3}) = \kappa_{R_1, R_2} d(R_3) \delta_{R_1 R'_1} \delta_{R_2 R'_2} \delta_{R_3 R'_3} \delta_{\tau_1 \tau_2} \delta_{\tau'_1 \tau'_2}.$$

The cardinality of this set of orthogonal elements in $\mathcal{K}(n)$ is

$$(18) \quad \sum_{R_1, R_2, R_3} \sum_{\tau_1, \tau_2} 1 = \sum_{R_1, R_2, R_3} C(R_1, R_2, R_3)^2 = |\text{Rib}(n)|$$

which allows us to confirm that these elements $\{Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}\}$ form an orthogonal basis of $\mathcal{K}(n)$.

The sets $\{Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}\}$ and $\{E_r\}$ define orthogonal bases of $\mathcal{K}(n)$. We refer to the set $\{E_r\}$ as the geometric or ribbon graph basis and to the set $\{Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}\}$ as the representation theoretic or Fourier basis of $\mathcal{K}(n)$. The change of basis from the Fourier basis to the geometric basis is made explicit in the appendix B. The existence of these two bases and their interplay is an important resource exploited in this paper.

2.2. INNER PRODUCT AND HILBERT SPACE.

2.2.1. $\mathcal{K}(n)$ as a Hilbert space.

PROPOSITION 2.1. *The algebra $\mathcal{K}(n)$ is a Hilbert space with the ribbon graph vectors E_r forming an orthogonal basis; the vectors $\sqrt{|\text{Orb}(r)|} E_r \equiv e_r$ form an orthonormal basis.*

Proof. Define the inner product g on $\mathbb{C}(S_n) \otimes \mathbb{C}(S_n)$, using the basis of permutation pairs and extend it by linearity. For two pairs $\alpha = (\alpha_1, \alpha_2), \beta = (\beta_1, \beta_2)$ in $S_n \times S_n$ we define

$$(19) \quad g(\alpha, \beta) = g(\alpha_1 \otimes \alpha_2, \beta_1 \otimes \beta_2) = \delta(\alpha_1^{-1} \beta_1) \delta(\alpha_2^{-1} \beta_2)$$

where δ is the delta function on S_n ($\delta(\sigma) = 1$ if and only if $\sigma = \text{id}$, otherwise $\delta(\sigma) = 0$). This extends by linearity to a sesquilinear form on $\mathbb{C}(S_n) \otimes \mathbb{C}(S_n)$ as

$$(20) \quad g\left(\sum_i a_i \alpha_{1i} \otimes \alpha_{2i}, \sum_j b_j \beta_{1j} \otimes \beta_{2j}\right) = \sum_{i,j} \bar{a}_i b_j \delta(\alpha_{1i}^{-1} \beta_{1j}) \delta(\alpha_{2i}^{-1} \beta_{2j})$$

where $a_i, b_i \in \mathbb{C}$ and where the bar means complex conjugation.

We can show that g satisfies conjugation property $g(\alpha, \beta) = \overline{g(\beta, \alpha)}$ and is positive definite. It therefore gives an inner product on $\mathbb{C}(S_n) \otimes \mathbb{C}(S_n)$.

We compute the inner product of two ribbon graph basis vectors

$$\begin{aligned}
 g(E_r, E_s) &= \frac{1}{|\text{Orb}(r)|} \frac{1}{|\text{Orb}(s)|} \sum_{a \in \text{Orb}(r)} \sum_{b \in \text{Orb}(s)} g(\sigma_1^{(r)}(a) \otimes \sigma_2^{(r)}(a), \sigma_1^{(s)}(b) \otimes \sigma_2^{(s)}(b)) \\
 (21) \quad &= \frac{1}{|\text{Orb}(r)|} \frac{1}{|\text{Orb}(s)|} \sum_{a \in \text{Orb}(r)} \sum_{b \in \text{Orb}(s)} \delta_{a,b} \delta_{r,s}
 \end{aligned}$$

where the only way that $\delta((\sigma_1^{(r)}(a))^{-1} \sigma_1^{(s)}(b)) \delta((\sigma_2^{(r)}(a))^{-1} \sigma_2^{(s)}(b)) = 1$ for two orbit elements a and b is when $a = b$. As a couple $(\sigma_1^{(r)}(a), \sigma_2^{(r)}(a))$ can only appear in a unique orbit, we therefore have

$$(22) \quad g(E_r, E_s) = \frac{1}{|\text{Orb}(s)|} \delta_{r,s}.$$

Then the set of $\{E_r\}$ for $r \in \{1, \dots, |\text{Rib}(n)|\}$ defines an orthogonal basis of $\mathcal{K}(n)$ which becomes a Hilbert space of states with inner product g . We can define orthonormal bases for $\mathcal{K}(n)$ as states of the form:

$$(23) \quad e_r = \sqrt{|\text{Orb}(r)|} E_r.$$

This completes the proof of the proposition. □

2.2.2. *Involution on $\mathcal{K}(n)$ from permutation inversion.* We also define the linear conjugation operator $S : \mathbb{C}(S_n) \rightarrow \mathbb{C}(S_n)$ that maps a linear combination $A = \sum_i c_i \sigma_i \in \mathbb{C}(S_n)$ to

$$(24) \quad S(A) := \sum_i c_i \sigma_i^{-1}.$$

Extend this operation to $\mathbb{C}(S_n) \otimes \mathbb{C}(S_n)$ by inverting the permutation in each tensor factor: $S(\sigma_1 \otimes \sigma_2) = \sigma_1^{-1} \otimes \sigma_2^{-1}$ and using linearity

$$(25) \quad S\left(\sum_i a_i \sigma_{1i} \otimes \sigma_{2i}\right) = \sum_i a_i \sigma_{1i}^{-1} \otimes \sigma_{2i}^{-1}.$$

S is an involution: $S^2 = \text{id}$, and obeys $S(AB) = S(B)S(A)$.

The conjugation S gives a well defined involution from the set of equivalence classes forming $\mathcal{K}(n)$ to itself. To see this note that if $(\sigma_1, \sigma_2) \in \text{Orb}(r)$ maps under inversion to a pair $(\sigma_1^{-1}, \sigma_2^{-1}) \in \text{Orb}(s)$ (where s may or may not be equal to r), then for any $\mu \in S_n$

$$(26) \quad S(\mu \sigma_1 \mu^{-1}, \mu \sigma_2 \mu^{-1}) = (\mu \sigma_1^{-1} \mu^{-1}, \mu \sigma_2^{-1} \mu^{-1}) \in \text{Orb}(s).$$

If for a given r , S maps the pairs $\text{Orb}(r)$ back to $\text{Orb}(r)$ we have

$$(27) \quad S(E_r) = E_r$$

Such ribbon graphs will be called self-conjugate. If $\text{Orb}(r)$ is mapped to $\text{Orb}(s)$ with $s \neq r$, then

$$(28) \quad \begin{aligned} S(E_r) &= E_s \\ S(E_s) &= E_r \end{aligned}$$

and we call (E_r, E_s) a conjugate pair. In Section 4.3 we compute the action of S on the Fourier basis. The interplay between these two actions is used to show that the total number of self-conjugate ribbon graphs with n edges is equal to the sum of Kronecker coefficients $C(R_1, R_2, R_3)$ for $R_1, R_2, R_3 \vdash n$. The operator S is also useful in proving the hermiticity of the reconnection operators $T_k^{(i)}$ (Proposition 2.3) which are used to construct Hamiltonians on $\mathcal{K}(n)$ in Section 3.3.

2.3. THE INTEGRALITY STRUCTURE OF THE PRODUCT ON $\mathcal{K}(n)$. The product in the algebra $\mathcal{K}(n)$ gives an expansion of the product $E_r E_s$ of two geometric basis vectors

$$(29) \quad E_r E_s = \sum_{t=1}^{|\text{Rib}(n)|} C_{rs}^t E_t.$$

We will express the structure constants

$$(30) \quad C_{rs}^t = \text{Coeff}(E_t, E_r E_s)$$

in terms of non-negative integers. Recall from Section 2.1.2 the two expressions for E_r

$$(31) \quad \begin{aligned} E_r &= \frac{1}{n!} \sum_{\mu \in S_n} \mu \sigma_1^{(r)} \mu^{-1} \otimes \mu \sigma_2^{(r)} \mu^{-1} \\ &= \frac{1}{|\text{Orb}(r)|} \sum_{a \in \text{Orb}(r)} \sigma_1^{(r)}(a) \otimes \sigma_2^{(r)}(a). \end{aligned}$$

For notational convenience, we define

$$(32) \quad \sigma^{(r)} = \sigma_1^{(r)} \otimes \sigma_2^{(r)}, \quad \mu \sigma^{(r)} \mu^{-1} = \mu \sigma_1^{(r)} \mu^{-1} \otimes \mu \sigma_2^{(r)} \mu^{-1}.$$

We express (10) in a simpler form as

$$(33) \quad E_r = \frac{1}{n!} \sum_{\mu \in S_n} \mu \sigma^{(r)} \mu^{-1} = \frac{1}{|\text{Orb}(r)|} \sum_{a \in \text{Orb}(r)} \sigma^{(r)}(a).$$

Consider the product of two elements of $\mathcal{K}(n)$ associated with orbits r and s :

$$(34) \quad \begin{aligned} E_r E_s &= \frac{1}{|\text{Orb}(r)|} \frac{1}{|\text{Orb}(s)|} \sum_{a \in \text{Orb}(r)} \sum_{b \in \text{Orb}(s)} \sigma^{(r)}(a) \sigma^{(s)}(b) \\ &= \frac{1}{(n!)^2} \sum_{\mu_1, \mu_2 \in S_n} \mu_1 \sigma^{(r)} \mu_1^{-1} \mu_2 \sigma^{(s)} \mu_2^{-1}. \end{aligned}$$

We can write $\nu = \mu_1^{-1} \mu_2$ and then solve for $\mu_1 = \mu_2 \nu^{-1}$ to write

$$(35) \quad \begin{aligned} E_r E_s &= \frac{1}{n!} \frac{1}{|\text{Orb}(r)|} \sum_{\mu_2 \in S_n} \sum_{a \in \text{Orb}(r)} \mu_2 \sigma^{(r)}(a) \sigma^{(s)} \mu_2^{-1} \\ &= \frac{1}{|\text{Orb}(r)|} \sum_{a \in \text{Orb}(r)} \frac{1}{|\text{Orb}(\sigma^{(r)}(a) \sigma^{(s)})|} \sum_{b \in \text{Orb}(\sigma^{(r)}(a) \sigma^{(s)})} \sigma(b) \\ &= \sum_t \frac{1}{|\text{Orb}(r)|} \sum_{b \in \text{Orb}(t)} \frac{\sigma^{(t)}(b)}{|\text{Orb}(t)|} \sum_{a \in \text{Orb}(r)} \delta(\text{Orb}(t), \text{Orb}(\sigma^{(r)}(a) \sigma^{(s)})) \\ &= \sum_t \frac{1}{|\text{Orb}(r)|} E_t \sum_{a \in \text{Orb}(r)} \delta(\text{Orb}(t), \text{Orb}(\sigma^{(r)}(a) \sigma^{(s)})), \end{aligned}$$

where $\delta(\text{Orb}(s), \text{Orb}(t))$ is the Kronecker symbol δ_{st} for the labels s and t . We have thus expressed the product of $E_r E_s$ in terms of the non-negative integer

$$(36) \quad \begin{aligned} &\sum_{a \in \text{Orb}(r)} \delta(\text{Orb}(t), \text{Orb}(\sigma^{(r)}(a) \sigma^{(s)})) \\ &= \text{Number of times the multiplication of elements from orbit } r \\ &\quad \text{with a fixed element in orbit } s \text{ to the right produces an element in orbit } t. \end{aligned}$$

If we solve for μ_2 instead as $\mu_2 = \mu_1\nu$, then we get

$$\begin{aligned}
 E_r E_s &= \sum_t \frac{1}{|\text{Orb}(s)|} \sum_{b \in \text{Orb}(t)} \frac{\sigma^{(t)}(b)}{|\text{Orb}(t)|} \sum_{a \in \text{Orb}(s)} \delta(\text{Orb}(t), \text{Orb}(\sigma^{(r)}\sigma^{(s)}(a))) \\
 (37) \quad &= \sum_t \frac{1}{|\text{Orb}(s)|} E_t \sum_{a \in \text{Orb}(s)} \delta(\text{Orb}(t), \text{Orb}(\sigma^{(r)}\sigma^{(s)}(a))).
 \end{aligned}$$

Here we have expressed the same product in terms of the non-negative integers

$$\begin{aligned}
 &\sum_{a \in \text{Orb}(s)} \delta(\text{Orb}(t), \text{Orb}(\sigma^{(r)}\sigma^{(s)}(a))) \\
 &= \text{Number of times the multiplication of elements from orbit } s \\
 &\quad \text{with a fixed element in orbit } r \text{ from the left produces an element in orbit } t. \\
 (38)
 \end{aligned}$$

Equivalently, we can express these by saying that the coefficient $\text{Coeff}(E_t, E_r E_s)$ of E_t in the expansion of $E_r E_s$ is given by

$$\begin{aligned}
 \text{Coeff}(E_t, E_r E_s) &= \frac{1}{|\text{Orb}(s)|} \sum_{a \in \text{Orb}(s)} \delta(\text{Orb}(t), \text{Orb}(\sigma^{(r)}\sigma^{(s)}(a))) \\
 (39) \quad &= \frac{1}{|\text{Orb}(r)|} \sum_{a \in \text{Orb}(r)} \delta(\text{Orb}(t), \text{Orb}(\sigma^{(r)}(a)\sigma^{(s)})).
 \end{aligned}$$

If we just keep μ_1, μ_2 in (34) we can write (treating r and s more symmetrically)

$$\text{Coeff}(E_t, E_r E_s) = \frac{1}{|\text{Orb}(s)| |\text{Orb}(r)|} \sum_{a \in \text{Orb}(r)} \sum_{b \in \text{Orb}(s)} \delta(\text{Orb}(t), \text{Orb}(\sigma^{(r)}(a)\sigma^{(s)}(b))). \tag{40}$$

Recalling that (22) holds, we therefore have

$$\text{Coeff}(E_t, E_r E_s) = g(E_r E_s, E_t) |\text{Orb}(t)|. \tag{41}$$

Hence

$$\begin{aligned}
 (42) \quad \delta_2(E_r E_s S(E_t)) &= g(E_r E_s, E_t) \\
 &= \frac{1}{|\text{Orb}(s)| |\text{Orb}(r)| |\text{Orb}(t)|} \sum_{a \in \text{Orb}(r)} \sum_{b \in \text{Orb}(s)} \delta(\text{Orb}(t), \text{Orb}(\sigma^{(r)}(a)\sigma^{(s)}(b))).
 \end{aligned}$$

These formulae show that the structure constants of the algebra $\mathcal{K}(n)$ are expressed in terms of non-negative integers obtained from the combinatorial multiplications of elements in the orbits, which provide the geometrical ribbon graph basis vectors E_r of $\mathcal{K}(n)$. In general, the product is not commutative $E_r E_s \neq E_s E_r$. In the next section, we will exploit this integral structure, for the particular cases where E_r are chosen to be central elements in $\mathcal{K}(n)$, associated with permutations having a cycle of length k and remaining cycles of length 1. These central elements will be used to construct Hamiltonians and the eigenproblems of these Hamiltonians will become questions about non-negative integer matrices.

2.4. THE CENTRE OF $\mathcal{K}(n)$ AND RECONNECTION OPERATORS $T_k^{(i)}$. In this section, we will review some properties of the centre of $\mathcal{K}(n)$ and introduce central elements $T_k^{(i)} \in \mathcal{K}(n)$ labelled by $k \in \{2, 3, \dots, n\}$ and $i \in \{1, 2, 3\}$. These central elements act on $\mathcal{K}(n)$ by multiplication. Since $\mathcal{K}(n)$ (when equipped with the inner product specified) is also the Hilbert space of our quantum mechanical systems, the elements $T_k^{(i)}$ also define linear operators when they act on $\mathcal{K}(n)$ by multiplication. We are taking advantage of a state-operator correspondence which is possible when the Hilbert space of a quantum mechanical space is also an algebra. We prove that these linear operators

are Hermitian with respect to the inner product defined in Section 2.2. We refer to the $T_k^{(i)}$ as reconnection operators, since they act, as we will see shortly, on the permutations defining elements of $\mathcal{K}(n)$ by multiplication of permutations. In the diagrammatic description of tensor model observables associated with $\mathcal{K}(n)$ [8] this operation involves reconnecting the index lines of the tensor variables $\Phi^{\otimes n}$ with those of $\bar{\Phi}^{\otimes n}$. In terms of ribbon graphs, the action of $T_k^{(i)}$ amounts to splitting and joining of vertices: such operators have also been discussed in [53, 55].

Let us first recall some properties of the centre $\mathcal{Z}(\mathbb{C}(S_n))$ of $\mathbb{C}(S_n)$. The centre is defined as the sub-algebra of elements which commute with all $\mathbb{C}(S_n)$. $\mathcal{Z}(\mathbb{C}(S_n))$ is a commutative algebra of dimension $p(n)$, the number of partitions of n . The conjugacy classes of S_n are specified by cycle structures of permutations which define partitions of n . The sum of elements in a conjugacy class is a central element in the group algebra. A linear basis for the centre is given by these class sums. For any integer k , such that $2 \leq k \leq n$, let \mathcal{C}_k to be the conjugacy class of permutations $\sigma \in S_n$ made of a single cycle of length k and remaining cycles of length 1. As an example, for $n = 3$, $k = 2$, the conjugacy class \mathcal{C}_2 is the set of permutations $\{(1, 2)(3), (2, 3)(1), (1, 3)(2)\}$. Define T_k as the sum

$$(43) \quad T_k = \sum_{\sigma \in \mathcal{C}_k} \sigma.$$

$|T_k|$ will refer to as the number of terms in that sum, equivalently the number of terms in \mathcal{C}_k , which is $\frac{n!}{k(n-k)!}$. For any n , the set $\{T_2, T_3, \dots, T_n\}$ generates the centre [57], i.e. linear combinations of products of these T_k span $\mathcal{Z}(\mathbb{C}(S_n))$. In fact there is no need to consider the entire set to generate $\mathcal{Z}(\mathbb{C}(S_n))$. Indeed, there exists $k_*(n) \leq n$, such that the subset $\{T_2, \dots, T_{k_*(n)}\}$ spans the center [57]. This is related to the fact that the ordered list of normalized characters $(\hat{\chi}_R(T_2), \hat{\chi}_R(T_3), \dots, \hat{\chi}_R(T_{k_*(n)}))$ uniquely identifies the Young diagram R . These normalized characters are defined as

$$(44) \quad \hat{\chi}_R(T_k) = \frac{\chi_R(T_k)}{d(R)}$$

where $d(R)$ is the dimension of the irrep R . The sequence $k_*(n)$ was explicitly computed [57], with the help of character formulae in [67, 23], to be

$$(45) \quad \begin{aligned} k_*(n) &= 2 \quad \text{for } n \in \{2, 3, 4, 5, 7\} \\ k_*(n) &= 3 \quad \text{for } n \in \{6, 8, 9, \dots, 14\} \\ k_*(n) &= 4 \quad \text{for } n \in \{15, 16, \dots, 23, 25, 26\} \\ k_*(n) &= 5 \quad \text{for } n \in \{24, 27, \dots, 41\} \\ k_*(n) &= 6 \quad \text{for } n \in \{42, \dots, 78, 79, 81\}. \end{aligned}$$

At any $n \geq 2$, we will define elements in $\mathbb{C}(S_n) \otimes \mathbb{C}(S_n)$

$$(46) \quad \begin{aligned} T_k^{(1)} &= T_k \otimes 1 = \sum_{\sigma \in \mathcal{C}_k} \sigma \otimes 1, \\ T_k^{(2)} &= 1 \otimes T_k = \sum_{\sigma \in \mathcal{C}_k} 1 \otimes \sigma, \\ T_k^{(3)} &= \sum_{\sigma \in \mathcal{C}_k} \sigma \otimes \sigma. \end{aligned}$$

These commute with permutations $\gamma \otimes \gamma$ and are thus in the sub-algebra $\mathcal{K}(n) \subset \mathbb{C}(S_n) \otimes \mathbb{C}(S_n)$. Using the correspondence between permutation pairs and ribbon graphs described in Section 2.1.1, it is straightforward to describe the ribbon graphs corresponding to $T_k^{(i)}$. In the case $n = 3$, $T_2^{(1)}, T_2^{(2)}, T_2^{(3)}$ correspond to the ribbon graphs labelled 4, 2, 5 respectively in Figure 1. The elements $T_3^{(1)}, T_3^{(2)}, T_3^{(3)}$ correspond

to the graphs labelled 8, 3, 10 respectively. For general n , $T_k^{(1)}$ corresponds to a ribbon graph with genus zero, having one black vertex of valency k , $n - k$ black vertices of valency one, and n white vertices of valency one. For $T_k^{(2)}$, we have a ribbon graph with one white vertex of valency k , $n - k$ vertices of valency one, and n black vertices of valency one. For $T_k^{(3)}$ we have a graph with genus $\lfloor \frac{k-1}{2} \rfloor$ with a black k -valent vertex connected to a white k -valent vertex, along with $n - k$ one-valent black and white vertices. These graphs are disconnected for $n > k$.

The $T_k^{(i)}$'s act as linear operators on $\mathbb{C}(S_n) \otimes \mathbb{C}(S_n)$ by left multiplication. Right multiplication gives the same operators because these are central operators in $\mathbb{C}(S_n) \otimes \mathbb{C}(S_n)$. They are also central in $\mathcal{K}(n)$, since this is a sub-algebra of $\mathbb{C}(S_n) \otimes \mathbb{C}(S_n)$. Let $(\mathcal{M}_k^{(i)})_r^s$ be the matrix elements of $T_k^{(i)}$

$$(47) \quad T_k^{(i)} E_s = \sum_s (\mathcal{M}_k^{(i)})_s^t E_t$$

in the geometric basis.

PROPOSITION 2.2. *The matrix elements $(\mathcal{M}_k^{(i)})_r^s$ are non-negative integers.*

Proof. The $T_k^{(i)}$ are proportional to instances of the geometric basis vectors E_r (31) obtained by summing over diagonal conjugations of permutations of the form $\sigma \otimes 1, 1 \otimes \sigma, \sigma \otimes \sigma$, where σ is a cyclic permutation of a subset of k numbers from $\{1, 2, \dots, n\}$. Using the correspondence (Section 2.1.1) between ribbon graphs and permutations, they each correspond to a ribbon graph. Each $T_k^{(i)}$ corresponds to a ribbon graph, with some label r which we will call $r(k, i)$. The proportionality constant is given as

$$(48) \quad T_k^{(i)} = |\text{Orb}(r(k, i))| E_{r(k, i)}$$

since the $T_k^{(i)}$ are equal to a sum of elements in an orbit generated by the diagonal conjugations, while E_r are defined to be such sums normalized by the orbit size. The formula (39) for the algebra product in the geometric basis then implies that

$$(49) \quad (\mathcal{M}_k^{(i)})_s^t = \text{Number of times the multiplication of elements in the sum } T_k^{(i)} \text{ with a fixed element in orbit } s \text{ to the right produces an element in orbit } t.$$

□

PROPOSITION 2.3. *$T_k^{(i)}$ are Hermitian operators on $\mathcal{K}(n)$ in the inner product defined by (20) :*

$$(50) \quad g(E_s, T_k^{(i)} E_r) = g(T_k^{(i)} E_s, E_r).$$

Proof. Using (22) and (47) we evaluate

$$(51) \quad g(E_s, T_k^{(i)} E_r) = (\mathcal{M}_k^{(i)})_r^s \frac{1}{|\text{Orb}(s)|}.$$

By renaming r, s , we have

$$(52) \quad g(E_r, T_k^{(i)} E_s) = (\mathcal{M}_k^{(i)})_s^r \frac{1}{|\text{Orb}(r)|}.$$

Using definition of g in (20) and of δ_2 in (8), the following is true

$$(53) \quad g(\alpha, \beta) = \delta_2(\bar{\alpha}, \beta).$$

Using this relation between the inner product and the delta function, we have

$$(54) \quad g(E_r, T_k^{(i)} E_s) = \delta_2(\bar{E}_r, (T_k^{(i)} E_s)).$$

From the definition of S (24), note that $\delta_2(S(a), id) = \delta_2(a, id)$, and

$$\delta_2(a, b) = \delta_2(aS(b), id) = \delta_2(S(b)a, id).$$

Moreover, $S(T_k) = T_k$ and $S(AB) = S(B)S(A)$ imply

$$\begin{aligned} (55) \quad \delta_2(\overline{E_r}, T_k^{(i)} E_s) &= \delta_2(S(T_k^{(i)} E_s) \overline{E_r}, id) = \delta_2(S(E_s) T_k^{(i)} \overline{E_r}, id) \\ &= \delta_2(T_k^{(i)} \overline{E_r}, E_s) = g(\overline{T_k^{(i)} \overline{E_r}}, E_s). \end{aligned}$$

Next observe that $\alpha = T_k^{(1)} E_r$ and $\beta = E_s$ have all real coefficients in the geometric ribbon graph basis,

$$(56) \quad g(\overline{T_k^{(i)} \overline{E_r}}, E_s) = g(T_k^{(i)} E_r, E_s).$$

This sequence of steps starting from (54) shows that

$$(57) \quad g(E_r, T_k^{(i)} E_s) = g(T_k^{(i)} E_r, E_s).$$

This proves the proposition. □

Remark

The matrix elements of $T_k^{(i)}$ in the non-orthogonal basis E_r are not symmetric. Indeed from (51), (52), and (50) we have

$$(58) \quad (\mathcal{M}_k^{(i)})_r^s = |\text{Orb}(s)| (\mathcal{M}_k^{(i)})_s^r \frac{1}{|\text{Orb}(r)|}.$$

If we consider instead the matrix elements of $T_k^{(i)}$ on the orthonormal basis vectors $e_r = \sqrt{|\text{Orb}(r)|} E_r$,

$$(59) \quad T_k^{(i)} e_r = \sqrt{|\text{Orb}(r)|} (\mathcal{M}_k^{(i)})_r^s \frac{1}{\sqrt{|\text{Orb}(s)|}} e_s.$$

These matrix elements are symmetric under exchange of r and s .

Remark

The operators $T_k^{(i)}$, as i ranges over $\{1, 2, 3\}$ and k ranges over some subset of $\{2, 3, \dots, n\}$ form a set of commuting Hermitian operators on $\mathcal{K}(n)$. The commutativity follows from the fact that they are central elements of $\mathcal{K}(n)$, the hermiticity from Proposition 2.3. Considering such sets of operators as Hamiltonians defining a time evolution of states in $\mathcal{K}(n)$ we have time-dependent ribbon graph states of the form

$$(60) \quad E_r(t) = e^{-itT_k^{(i)}} E_r.$$

In Section 3.3 we will construct Hamiltonians which are particular linear combinations of these operators, and have the property that their eigenvalue degeneracies are Kronecker coefficients. In order to build up to this, we will now consider the action of the $T_k^{(i)}$ operators on the Fourier basis of $\mathcal{K}(n)$.

3. INTEGER MATRICES AND KRONECKER COEFFICIENTS

In this section we will consider the action of the reconnection operators $T_k^{(i)}$ introduced in Section 2.4 on the elements $Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}$ of the Fourier basis set for $\mathcal{K}(n)$ described in Section 2.1.3. The subspace of $\mathcal{K}(n)$ spanned by the Fourier basis elements for a fixed ordered Young diagram triple (R_1, R_2, R_3) has dimension equal to $C(R_1, R_2, R_3)^2$, the square of the Kronecker coefficient for the triple. We will refer to such a subspace as the Fourier subspace of $\mathcal{K}(n)$ associated with the triple (R_1, R_2, R_3) . We will show (Section 3.2) that the Fourier subspace for a triple form an eigenspace of the reconnection operators. The eigenvalues are normalized characters of the symmetric group, with rational values known from symmetric group representation theory.

We work with a set of reconnection operators chosen such that their eigenvalues uniquely specify the Young diagram triple. This means that, although the Fourier basis was initially defined using matrix elements and Clebsch–Gordan coefficients for symmetric groups (equation (14)), we can use the reconnection operators and the eigenvalues as input, and compute the Fourier subspace for a fixed triple of Young diagrams directly as the null space of an integer matrix built from the reconnection operators and the eigenvalues specifying a Young diagram triple. This gives a computational approach for the Fourier subspace associated with a Young diagram triple without using the detailed representation theory input of matrix elements and Clebsch–Gordan coefficients: we are only using the coarser input of character formulae (having known combinatorial algorithms for their computation) alongside the combinatorially defined reconnection operators. The first algorithm based on this approach (Section 3.2) amounts to finding the null vectors of a rectangular integer matrix. In Section 3.3 we give the construction of quantum mechanical Hamiltonians which are integer linear combinations of the reconnection operators $T_k^{(i)}$ and have eigenvalues that uniquely specify a Young diagram triple. This leads to an algorithm which obtains the Fourier subspace for a specified triple as the null space of a square integer matrix.

3.1. FOURIER SUBSPACE OF A YOUNG DIAGRAM TRIPLE AS EIGENSPACE OF RECONNECTION OPERATORS.

PROPOSITION 3.1. *For all $k \in \{2, 3, \dots, n\}$, $\{R_i \vdash n : i \in \{1, 2, 3\}\}$, $\tau_1, \tau_2 \in \llbracket 1, C(R_1, R_2, R_3) \rrbracket$, the Fourier basis elements $Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}$ are eigenvectors of $T_k^{(i)}$:*

$$(61) \quad T_k^{(1)} Q_{\tau_1, \tau_2}^{R_1, R_2, R_3} = \left(\sum_{\sigma \in C_k} \sigma \otimes 1 \right) Q_{\tau_1, \tau_2}^{R_1, R_2, R_3} = \frac{\chi_{R_1}(T_k)}{d(R_1)} Q_{\tau_1, \tau_2}^{R_1, R_2, R_3},$$

$$(62) \quad T_k^{(2)} Q_{\tau_1, \tau_2}^{R_1, R_2, R_3} = \left(\sum_{\sigma \in C_k} 1 \otimes \sigma \right) Q_{\tau_1, \tau_2}^{R_1, R_2, R_3} = \frac{\chi_{R_2}(T_k)}{d(R_2)} Q_{\tau_1, \tau_2}^{R_1, R_2, R_3},$$

$$(63) \quad T_k^{(3)} Q_{\tau_1, \tau_2}^{R_1, R_2, R_3} = \left(\sum_{\sigma \in C_k} \sigma \otimes \sigma \right) Q_{\tau_1, \tau_2}^{R_1, R_2, R_3} = \frac{\chi_{R_3}(T_k)}{d(R_3)} Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}.$$

The proof is given in appendix B. Note that the eigenvalues do not depend on the multiplicity indices τ_1 and τ_2 , but only on the Young diagram labels (R_1, R_2, R_3) . The proof of Proposition 3.1 relies on representation theoretic arguments.

PROPOSITION 3.2. *For any $\tilde{k}_* \in \{k_*(n), k_*(n) + 1, \dots, n\}$ the list of eigenvalues of the reconnection operators $\{T_2^{(1)}, T_3^{(1)}, \dots, T_{k_*}^{(1)}; T_2^{(2)}, T_3^{(2)}, \dots, T_{k_*}^{(2)}; T_2^{(3)}, T_3^{(3)}, \dots, T_{k_*}^{(3)}\}$ uniquely determines the Young diagram triples (R_1, R_2, R_3) .*

Proof. It was shown in [57] that the normalized characters $\{\frac{\chi_R(T_2)}{d(R)}, \frac{\chi_R(T_3)}{d(R)}, \dots, \frac{\chi_R(T_n)}{d(R)}\}$ form ordered lists of numbers which distinguish Young diagrams R with n boxes. For all $n > 2$ it was shown that the shorter list $\{\frac{\chi_R(T_2)}{d(R)}, \frac{\chi_R(T_3)}{d(R)}, \dots, \frac{\chi_R(T_{n-1})}{d(R)}\}$ distinguishes Young diagrams. This result follows from the fact that the central elements $\{T_2, \dots, T_{n-1}\}$ generate the centre of $C(S_n)$ for $n > 2$. It was found that there generically exist $k_*(n) < n - 1$ such that the shorter lists $\{\frac{\chi_R(T_2)}{d(R)}, \frac{\chi_R(T_3)}{d(R)}, \dots, \frac{\chi_R(T_{k_*(n)})}{d(R)}\}$ distinguish Young diagrams. The values of $k_*(n)$ computed for all n up to 79 are given in (45). While the general $k_*(n)$ are not currently known, any $n \geq \tilde{k}_* \geq k_*(n)$ gives a longer list which distinguishes Young diagrams. A triple of lists of length \tilde{k}_* distinguishes a triple of Young diagrams. Using Proposition 3.1, these are the sets

of eigenvalues of the operators $\{T_2^{(1)}, T_3^{(1)}, \dots, T_{k_*}^{(1)}, T_2^{(2)}, T_3^{(2)}, \dots, T_{k_*}^{(2)}, T_2^{(3)}, T_3^{(3)}, \dots, T_{k_*}^{(3)}\}$. □

LEMMA 3.3. *The sum of all permutations σ in the conjugacy class C_p in S_n for partition p are central elements in $\mathcal{Z}(\mathbb{C}(S_n))$. The irreducible normalized characters of these central elements are integers :*

$$(64) \quad \frac{\chi^R(T_p)}{d(R)} \in \mathbb{Z}.$$

The proof combines a known number theoretic fact about the normalized characters of a finite group being algebraic integers [89], along with the rationality of characters of irreducible representations of S_n which follows from the Murnaghan–Nakayama Lemma.

Proof. The elements T_p as p runs over the classes form a basis for $\mathcal{Z}(\mathbb{C}(S_n))$. The structure constants of the multiplication are defined by

$$(65) \quad T_p T_q = \sum_r C_{pq}^r T_r.$$

These structure constants C_{pq}^r are integers. The normalized characters $\frac{\chi^R(T_p)}{d(R)}$ are eigenvalues of the matrix defined by C_{pq}^r for fixed p . The eigenvalues of an integer matrix are algebraic integers (see e.g. Proposition III.4.3 [89]). In the case of symmetric groups, we know that $\chi^R(\sigma)$ for $\sigma \in C_p$ is an integer by using the Murnaghan–Nakayama Lemma. It follows that the normalized characters $\frac{\chi^R(T_p)}{d(R)}$ are rational. A rational number which is also an algebraic integer is necessarily an integer. This means that the normalized characters are integers, for any conjugacy class p . □

In particular for the partitions of the form $[k, 1^{n-k}]$ the normalized characters

$$(66) \quad \frac{\chi^R(T_k)}{d(R)} \in \mathbb{Z}.$$

Using the formulae for the normalized characters in [67, 23] the normalized characters for T_2 (as well as T_3, T_4, T_5, T_6) are evidently integers for any Young diagram. For higher higher T_k , the integrality is not evident from the formulae, but hold from the above argument.

3.2. FOURIER SUBSPACE OF TRIPLE AS NULL-SPACE OF RECTANGULAR INTEGER MATRICES. It is useful to recall from equation (49) and Proposition 2.2 reproduced here for convenience (with a slight change in index labels) :

$$(67) \quad T_k^{(i)} E_r = \sum_s (\mathcal{M}_k^{(i)})_r^s E_s$$

with

$$(68) \quad (\mathcal{M}_k^{(i)})_r^s = \text{Number of times the multiplication of elements in the sum } T_k^{(i)} \text{ with a fixed element in orbit } r \text{ to the right produces an element in orbit } s.$$

Using the definition of $T_k^{(i)}$, this means that

$$\begin{aligned} (\mathcal{M}_k^{(1)})_r^s &= \sum_{\gamma \in \mathcal{C}_k} \delta(\text{Orb}(s), \text{Orb}(\gamma \tau_1^{(r)} \otimes \tau_2^{(r)})), \\ (\mathcal{M}_k^{(2)})_r^s &= \sum_{\gamma \in \mathcal{C}_k} \delta(\text{Orb}(s), \text{Orb}(\tau_1^{(r)} \otimes \gamma \tau_2^{(r)})), \end{aligned}$$

$$(69) \quad (\mathcal{M}_k^{(3)})_r^s = \sum_{\gamma \in \mathcal{C}_k} \delta(\text{Orb}(s), \text{Orb}(\gamma\tau_1^{(r)} \otimes \gamma\tau_2^{(r)})).$$

The integer matrices $\mathcal{M}_k^{(i)}$ of $T_k^{(i)}$ are constructed in Code2 of the appendix C, see the function `ArrayTi(n, kmax)`. Note also that we have the following relations

$$(70) \quad \sum_s (\mathcal{M}_k^{(i)})_r^s = |T_k| = \frac{n!}{k(n-k)!},$$

where $|T_k|$ is the number of terms in the sum T_k , see the discussion after (43). Thus each column of $(\mathcal{M}_k^{(i)})_r^s$ is a list of non-negative integers adding up to $|T_k|$.

3.2.1. *Stacking $T_k^{(i)}$ matrices and common eigenspace.* Using Proposition 3.2, the Fourier subspace for a given triple (R_1, R_2, R_3) is uniquely specified as common eigenspace of the operators $T_k^{(i)}$, for $k \in \{2, \dots, \tilde{k}_*(n)\}$ and $i \in \{1, 2, 3\}$; with $k_*(n) \leq \tilde{k}_* \leq n$, with specified eigenvalues for these reconnection operators, which are known from symmetric group representation theory. These eigenvalues are normalized characters which can be combinatorially computed in at least two known ways. The numerator $\chi_R(T_k)$ is given by $\chi_R(T_k) = |T_k| \chi_R(\sigma)$ for $\sigma \in \mathcal{C}_k$. The character $\chi_R(\sigma)$ can be computed with the combinatorial Murnaghan–Nakayama rule [76] [78]. The dimension $d(R)$ is obtained from the hook formula for dimensions. Another combinatorial formula gives $\frac{k \chi_R(T_k)}{d(R)}$ [91][37].

The vectors in the Fourier subspace for a triple (R_1, R_2, R_3) solve the following matrix equation

$$(71) \quad \begin{bmatrix} \mathcal{M}_2^{(1)} - \frac{\chi_{R_1}(T_2)}{d(R_1)} \\ \vdots \\ \mathcal{M}_{\tilde{k}_*}^{(1)} - \frac{\chi_{R_1}(T_{\tilde{k}_*})}{d(R_1)} \\ \mathcal{M}_2^{(2)} - \frac{\chi_{R_2}(T_2)}{d(R_2)} \\ \vdots \\ \mathcal{M}_{\tilde{k}_*}^{(2)} - \frac{\chi_{R_2}(T_{\tilde{k}_*})}{d(R_2)} \\ \mathcal{M}_2^{(3)} - \frac{\chi_{R_3}(T_2)}{d(R_3)} \\ \vdots \\ \mathcal{M}_{\tilde{k}_*}^{(3)} - \frac{\chi_{R_3}(T_{\tilde{k}_*})}{d(R_3)} \end{bmatrix} \cdot v = \mathbf{0}.$$

This rectangular array gives the matrix elements of a linear operator mapping $\mathcal{K}(n)$ to $3(\tilde{k}_* - 1)$ copies of $\mathcal{K}(n)$, using the geometric basis of ribbon graph vectors for $\mathcal{K}(n)$. From Lemma 3.3, the normalized characters are integers. Renaming as $\mathcal{L}_{R_1, R_2, R_3}$ the integer matrix in (71) we have

$$(72) \quad \mathcal{L}_{R_1, R_2, R_3} \cdot v = 0.$$

We then have, for each triple of Young diagrams, the problem of finding the null space of an integer matrix. Null spaces of integer matrices have integer null vector bases. These can be interpreted in terms of lattices and can be constructed using integral algorithms. We will discuss the integrality properties of the null vectors and the associated interpretation in terms of lattices further in Section 4.

3.2.2. *Computational implementation and examples.* The construction of this rectangular matrix using reconnections on ribbon graph equivalence classes along with normalized symmetric group characters is implemented using the software [39]. This is described in appendix C. For the case $n = 3$, $k_*(n) = 2$ and we choose $\tilde{k}_* = k_*(3) = 2$. The three reconnection operators $T_2^{(1)}, T_2^{(2)}, T_2^{(3)}$ suffice to distinguish the Young diagram triples. The matrices for each operator are given in appendix A.2. In this case the Kronecker coefficients $C(R_1, R_2, R_3)$ are either 0 or 1. An integer null basis vector for the Fourier subspace associated with each triple having non-vanishing Kronecker coefficient is given in appendix A.2. The vectors shown give the coefficients of the vectors E_r for the index $r \in \{1, 2, \dots, 11\}$ and the graph associated with each index is shown in Figure 1.

3.3. FOURIER SUBSPACE OF TRIPLE AS NULL SPACE OF SQUARE INTEGER MATRICES. We now show that Fourier subspace for a triple of Young diagrams can be obtained as the null space of a square matrix. Rather than stacking the matrices for reconnection operators $T_k^{(i)}$ in a rectangular matrix, we will take linear combinations of these linear operators with integer coefficients. These linear combinations define Hamiltonians in the quantum mechanics of ribbon graphs. The coefficients are chosen with some care, using a procedure we explain, to ensure that the eigenvalues of the Hamiltonian for distinct Young diagram triples are distinct.

Distinguishing Young diagram triples with quantum mechanical Hamiltonians. We consider Hermitian Hamiltonians of the form

$$(73) \quad \mathcal{H} = \sum_{i=1}^3 \sum_{k=2}^{k_*(n)} a_{i,k} T_k^{(i)}$$

with coefficients $a_{i,k}$ which we will discuss shortly. For simplicity we have taken $\tilde{k}_* = k_*(n)$ which is the minimum needed for the list of normalized characters to distinguish Young diagrams with n boxes. Using Proposition 3.1 we have

$$(74) \quad \mathcal{H} Q_{\tau_1, \tau_2}^{R_1, R_2, R_3} = \left(\sum_{i,k} a_{i,k} \frac{\chi_{R_i}(T_k)}{d(R_i)} \right) Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}.$$

The Fourier subspace of $\mathcal{K}(n)$ for the ordered triple (R_1, R_2, R_3) , which has dimension $C(R_1, R_2, R_3)^2$, is an eigenspace of these Hamiltonians. We will show that the coefficients $a_{i,k}$ can be chosen as integers which ensure that the eigenvalues, which we denote as ω_{R_1, R_2, R_3}

$$(75) \quad \omega_{R_1, R_2, R_3} = \sum_{i,k} a_{i,k} \frac{\chi_{R_i}(T_k)}{d(R_i)}$$

distinguish the ordered triples (R_1, R_2, R_3) . In other words, we can choose integers $a_{i,k}$ to have

triple-distinguishing Hamiltonians with the property :

$$(76) \quad \text{If } (R_1, R_2, R_3) \neq (R'_1, R'_2, R'_3) \text{ then } \omega_{R_1, R_2, R_3} \neq \omega_{R'_1, R'_2, R'_3}.$$

Characterising the general choice of integers $a_{i,k}$ which defines a Hamiltonian (73) with the property (76) is an interesting problem: here we will only show that these Hamiltonians exist, using a particular construction. As an operator on $\mathcal{K}(n)$, using the geometric ribbon graph basis, the matrix elements of \mathcal{H} are integers

$$(77) \quad \text{Coeff}(E_s, \mathcal{H}E_r) = \sum_{i=1}^3 \sum_{k=2}^{k_*(n)} a_{i,k} (\mathcal{M}_k^{(i)})_r^s$$

using Proposition 2.2 and (49). For convenience, we will use the notation $\widehat{\chi}_R(T_k)$ for the normalized characters $\frac{\chi_R(T_k)}{d(R)}$ which are integers by Lemma 3.3. The eigenvalues of the Hamiltonians are

$$(78) \quad \omega_{R_1, R_2, R_3} = \sum_{i=1}^3 \sum_{k=2}^{k_*} a_{i,k} \widehat{\chi}_{R_i}(T_k) = \sum_{k=2}^{k_*(n)} a_{1,k} \widehat{\chi}_{R_1}(T_k) + a_{2,k} \widehat{\chi}_{R_2}(T_k) + a_{3,k} \widehat{\chi}_{R_3}(T_k).$$

For each triple of Young diagrams (R_1, R_2, R_3) the operator

$$(79) \quad (\mathcal{H} - \omega_{R_1, R_2, R_3}) \equiv \mathcal{H}_{R_1, R_2, R_3}$$

is, in the geometric ribbon graph basis, a square integer matrix. The Fourier subspace of $\mathcal{K}(n)$ associated with the triple (R_1, R_2, R_3) is the space spanned by the null vectors v of this operator

$$(80) \quad \mathcal{H}_{R_1, R_2, R_3} \cdot v = 0.$$

We now turn to the demonstration that such triple-distinguishing Hamiltonians constructed from integer $a_{i,k}$ indeed exist in general.

Triple distinguishing property for low n

Consider the problem of establishing the property in (76) for the cases $n = 2, 3, 4, 5, 7$ where $k_*(n) = 2$. In this case, simplify the notation

$$(81) \quad a_{1,2} = a_1 ; a_{2,2} = a_2 ; a_{3,2} = a_3$$

Introduce a label q which indexes partitions of n , so R_q is a Young diagram with n boxes. With the simplified notation

$$(82) \quad X_q = \widehat{\chi}_{R_q}(T_2)$$

the eigenvalues are

$$(83) \quad \omega_{R_{q_1}, R_{q_2}, R_{q_3}} = (a_1 X_{q_1} + a_2 X_{q_2} + a_3 X_{q_3}).$$

These X_q are integers using Lemma 3.3.

Conditions on a_i

The problem of finding a_i to have Hamiltonians with the triple-distinguishing property is now the problem of finding a_1, a_2, a_3 such that for any distinct triples $(R_{q_1}, R_{q_2}, R_{q_3}) \neq (R_{q'_1}, R_{q'_2}, R_{q'_3})$ the eigenvalues in (83) $\omega_{R_{q_1}, R_{q_2}, R_{q_3}} \neq \omega_{R_{q'_1}, R_{q'_2}, R_{q'_3}}$. In other words, the problem is to

Find integers a_1, a_2, a_3 such that

$$(84) \quad \begin{aligned} & a_1(X_{q_1} - X_{q'_1}) + a_2(X_{q_2} - X_{q'_2}) + a_3(X_{q_3} - X_{q'_3}) = 0 \\ & \text{is only satisfied when } X_{q_1} = X_{q'_1}, X_{q_2} = X_{q'_2}, X_{q_3} = X_{q'_3}. \end{aligned}$$

Equivalently, the problem is to find integers (a_1, a_2, a_3) such that the sum

$$(85) \quad a_1(X_{q_1} - X_{q'_1}) + a_2(X_{q_2} - X_{q'_2}) + a_3(X_{q_3} - X_{q'_3})$$

is never zero for any choice of distinct triples (q_1, q_2, q_3) and (q'_1, q'_2, q'_3) . Note that two triples are considered distinct if they differ in any of the 3 slots. E.g. if $q_1 \neq q'_1$ then (q_1, q_2, q_3) and (q'_1, q_2, q_3) are distinct triples.

Given the condition on the a_i , none of the a_i can be zero. Suppose the contrary, e.g. $a_1 = 0$. Then we can take $X_{q_1} \neq X_{q'_1}$ but $X_{q_2} = X_{q'_2}, X_{q_3} = X_{q'_3}$, and get a solution to (84). Let us look for a solution where $a_1 = 1$.

The possible differences $X_q - X_{q'}$ at fixed n form a finite set of values. For example at $n = 3$, they can be $(0, 3, -3, 6, -6)$. The list of non-zero $X_q - X_{q'}$ has prime factors 2, 3. Take a prime p_1 which is not one of these prime factors. E.g. in this $n = 3$ case,

take $p_1 = 5$. Let $a_2 = p_1$. This ensures that, when $X_{q_3} = X_{q'_3}$, we cannot solve (84) : the second term has a prime factor p_1 while the first does not, so they cannot add up to zero. For a_3 we pick another prime p_2 which is not p_1 and does not appear among the prime factors of $X_q - X_{q'}$ for any q, q' . This ensures that the condition (84) on the a_i is satisfied for all triples where $X_{q_2} = X_{q'_2}$.

Now we consider the generic case where

$$(86) \quad X_{q_1} \neq X_{q'_1}, X_{q_2} \neq X_{q'_2}, X_{q_3} \neq X_{q'_3}.$$

To continue satisfying (84) we can choose a_3 large enough that the last term cannot be cancelled by the sum of first two terms. So pick p_2 such that

$$(87) \quad p_2 \text{Min}_{q,q'} |X_q - X_{q'}| > \text{Max}(|p_1(X_{q_2} - X_{q'_2}) + (X_{q_1} - X_{q'_1})|).$$

Using the inequality,

$$(88) \quad p_1 \text{Max}_{q_2,q'_2} |X_{q_2} - X_{q'_2}| + \text{Max}_{q_1,q'_1} |X_{q_1} - X_{q'_1}| \geq \text{Max}(|p_1(X_{q_2} - X_{q'_2}) + (X_{q_1} - X_{q'_1})|).$$

we can write a computationally simpler condition

$$(89) \quad p_2 \text{Min}_{q,q'} |X_q - X_{q'}| > (p_1 + 1) \text{Max}_{q,q'} |X_q - X_{q'}|.$$

Since we are in the case (86), $\text{Min}_{q,q'} |X_q - X_{q'}| > 0$.

In the example of $n = 3$, choosing $p_1 = 5$ as explained above, pick $p_2 = 13$, which satisfies (89) because $13 * 3 > 5 * 6 + 6$. We conclude that the choice $(a_1, a_2, a_3) = (1, 5, 13)$ at $n = 3$, satisfies the condition (84) and the eigenvalues of $\mathcal{H} = T_2^{(1)} + 5T_2^{(2)} + 13T_2^{(3)}$ distinguish the triples (R_1, R_2, R_3) which label the Q -basis, and the degeneracies of the eigenspaces are precisely the squares of Kronecker coefficients.

Triple distinguishing property for general n

We now have eigenvalues of \mathcal{H} equal to

$$(90) \quad \omega_{R_{q_1}, R_{q_2}, R_{q_3}} = \sum_{k=2}^{k_*} (a_{1,k} X_{q_1,k} + a_{2,k} X_{q_2,k} + a_{3,k} X_{q_3,k})$$

with

$$(91) \quad X_{q_i,k} = \widehat{\chi}_{R_{q_i}}(T_k).$$

We choose $a_{i,k}$ to have Hamiltonians with the triple-distinguishing property, i.e. $\omega_{R_{q_1}, R_{q_2}, R_{q_3}} \neq \omega_{R_{q'_1}, R_{q'_2}, R_{q'_3}}$ for $(R_{q_1}, R_{q_2}, R_{q_3}) \neq (R_{q'_1}, R_{q'_2}, R_{q'_3})$. This the problem

Find integers $a_{1,k}, a_{2,k}, a_{3,k}$ such that

$$(92) \quad \sum_{k=2}^{k_*} a_{1,k} (X_{q_1,k} - X_{q'_1,k}) + a_{2,k} (X_{q_2,k} - X_{q'_2,k}) + a_{3,k} (X_{q_3,k} - X_{q'_3,k}) = 0$$

is only satisfied when $X_{q_1} = X_{q'_1}, X_{q_2} = X_{q'_2}, X_{q_3} = X_{q'_3}$.

The previous strategy for low n extends here. Suppose $q_1 \neq q'_1$ but $q_2 = q'_2, q_3 = q'_3$. In this case, we need to make sure that the $a_{1,k}$ are chosen such that for any pair q_1, q'_1

$$(93) \quad \sum_{k=2}^{k_*} a_{1,k} (X_{q_1,k} - X_{q'_1,k}) \neq 0.$$

One scheme for producing such a collection of $a_{1,k}$ is to use prime decompositions again. Consider the differences $X_{q_1,k} - X_{q'_1,k}$ as q_1, q'_1 range over distinct pairs. Consider the set of prime factors, denoted $\text{PrimesDiffs}(n, k)$ appearing in the integer differences $X_{q_1,k} - X_{q'_1,k}$ as q_1, q'_1 range over distinct pairs. Choose $a_{1,2} = 1$ and

$a_{1,3} = p_1$, with $p_1 \notin \text{PrimesDiffs}(n, 2)$. Then $a_{1,4} = p_2$ is a bigger prime chosen such that $p_2 \notin \{p_1\} \cup \text{PrimesDiffs}(n, 2) \cup \text{PrimesDiffs}(n, 3)$ and

$$(94) \quad \text{Max}_{q,q'} |(X_{q,2} - X_{q',2})| + p_1 \text{Max}_{q,q'} |(X_{q,3} - X_{q',3})| < p_2 \text{Min}_{q,q'} |(X_{q,4} - X_{q',4})|.$$

By iterating this procedure, we select increasing primes $p_1, p_2, \dots, p_{k_*-2}$ to ensure (93).

Back to considering (92): the case $q_1 = q'_1, q_2 \neq q'_2, q_3 = q'_3$ requires

$$(95) \quad \sum_{k=2}^{k_*} a_{2,k}(X_{q_2,k} - X_{q'_2,k}) \neq 0.$$

The case $q_1 = q'_1, q_2 = q'_2, q_3 \neq q'_3$ requires

$$(96) \quad \sum_{k=2}^{k_*} a_{3,k}(X_{q_3,k} - X_{q'_3,k}) \neq 0.$$

We also need to ensure that the condition (92) holds when two of the q 's are distinct and when all three are distinct. We can pick

$$(97) \quad \begin{aligned} (a_{1,2}, a_{1,3}, \dots, a_{1,k_*}) &= (1, p_1, p_2, \dots, p_{k_*-2}) \\ (a_{2,1}, a_{2,2}, \dots, a_{2,k_*-2}) &= p_{k_*-1}(1, p_1, \dots, p_{k_*-2}) \\ (a_{3,1}, a_{3,2}, \dots, a_{3,k_*-2}) &= p_{k_*}(1, p_1, \dots, p_{k_*-2}). \end{aligned}$$

The primes are chosen such that $p_1 < p_2 < \dots < p_{k_*-2} < p_{k_*-1} < p_{k_*}$, with p_{k_*-1} such that

$$(98) \quad \text{Max}_{q,q'} \left(\left| \sum_{k=2}^{k_*} p_{k-2}(X_{q,k} - X_{q',k}) \right| \right) \leq \text{Max}_{q,q'} \left(\sum_{k=2}^{k_*} p_{k-2} |X_{q,k} - X_{q',k}| \right) < p_{k_*-1} \text{Min}_{q,q'} \left(\left| \sum_{k=2}^{k_*} p_{k-2}(X_{q,k} - X_{q',k}) \right| \right)$$

where we extend the sequence p_l to $p_0 = 1$, and also such that p_{k_*} obeys (99)

$$(1 + p_{k_*-1}) \text{Max}_{q,q'} \left(\sum_{k=2}^{k_*} p_{k-2} |X_{q,k} - X_{q',k}| \right) < p_{k_*} \text{Min}_{q,q'} \left(\left| \sum_{k=2}^{k_*} p_{k-2}(X_{q,k} - X_{q',k}) \right| \right).$$

The Min on the RHS is non-zero since this condition is coming from the vase $q_1 \neq q'_1, q_2 \neq q'_2, q_3 \neq q'_3$. With these integer choices of $a_{1,k}, a_{2,k}, a_{3,k}$ we can ensure that \mathcal{H} has eigenvalues which distinguish the triples (R_1, R_2, R_3) in the Fourier basis elements $Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}$. The dimensions of the distinct eigenspaces are $C(R_1, R_2, R_3)^2$.

Examples. In fact, for $k = 2$, $\text{Max}_{q,q'} |X_{q,2} - X_{q',2}|$ is known and equals $2 \cdot |\widehat{\chi}_{[1^n]}(T_2)| = n(n-1)$ and $\text{Min}_{q,q'} |X_{q,2} - X_{q',2}|$ cannot be lower than 1 since we know that $X_{q,2}, X_{q',2}$ are integers. Using this lower bound

$$(100) \quad p_2 \text{Min}_{q,q'} |X_{q,2} - X_{q',2}| > p_2 > (p_1 + 1)n(n-1).$$

Thus picking the minimal prime p_2 larger than $(p_1 + 1)n(n-1)$ would solve the inequality in (89). When $\text{Min}_{q,q'} |X_{q,2} - X_{q',2}| > 1$ then the above is a still sufficient condition but does not lead to the smallest p_2 . After some illustrations, we will discuss sufficient conditions that leads to other solutions of the problem.

Case $n = 5$. Here $k_* = 2$ and we have

$$(101) \quad X_{q,2} - X_{q',2} \in \{-20, -15, -12, -10, -8, -7, -5, -4, -3, -2, 0, 2, 3, 4, 5, 7, 8, 10, 12, 15, 20\}$$

with $\widehat{\chi}_{[1^n]}(T_2) = -10$ so that $\text{Max}_{q,q'}|X_{q,2} - X_{q',2}| = 20$, and $\text{Min}_{q,q'}|X_{q,2} - X_{q',2}| = 2 > 0$. The set of prime divisors of the above set is $\{2, 3, 5, 7\}$. We choose $p_1 = 11$ and therefore the inequality (89) becomes

$$(102) \quad 2p_2 > 12 \times 20 = 240.$$

Hence we choose $p_2 = 127$, and the triple $(p_0, p_1, p_2) = (1, 11, 127)$ solves (84). The Hamiltonian $\mathcal{H} = T_2^{(1)} + 11T_2^{(2)} + 127T_2^{(3)}$ is an integer matrix in the geometric ribbon graph basis, with the property that distinct Young diagram triples are associated with distinct eigenvalues, and the eigenvalue degeneracies are given by $C(R_1, R_2, R_3)^2$.

Case $n = 7$. Again $k_* = 2$ should be the max of k . We have

$$(103) \quad \begin{aligned} X_{q,2} - X_{q',2} \in \{ & -42, -35, -30, -28, -27, -24, -23, -22, -21, -20, -18, -17, -16, \\ & -15, -14, -13, -12, -11, -10, -9, -8, -7, -6, -5, -4, -3, -2, -1, 0, 1, 2, 3, \\ & 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 27, 28, 30, 35, 42 \}, \end{aligned}$$

with $\widehat{\chi}_{[1^n]}(T_2) = -21$, $\text{Max}_{q,q'}|X_{q,2} - X_{q',2}| = 42$, and $\text{Min}_{q,q'}|X_{q,2} - X_{q',2}| = 1$. The set of prime divisors is $\{2, 3, 5, 7, 23, 17, 13, 11\}$. Choose $p_1 = 19$, and then we seek

$$(104) \quad p_2 > 20 * 42 = 840.$$

We fix $p_2 = 853$ and $(a_1, a_2, a_3) = (p_0, p_1, p_2) = (1, 19, 853)$ is one correct triple solves (84). Thus $\mathcal{H} = T_2^{(1)} + 19T_2^{(3)} + 853T_2^{(3)}$ is an integer matrix in the geometric ribbon graph basis, with the property that distinct Young diagram triples are associated with distinct eigenvalues, and the eigenvalue degeneracies are given by $C(R_1, R_2, R_3)^2$.

Case $n = 6$. In this case $k_* = 3$ and

$$X_{q,2} - X_{q',2} \in \{ -30, -24, -20, -18, -15, -14, -12, -10, -9, -8, -6, -5, -4, -3, -2, 0, 2, 3, 4, 5, 6, 8, 9, 10, 12, 14, 15, 18, 20, 24, 30 \}$$

$$\text{PrimesDiffs}(6, 2) = \{2, 3, 5, 7\}, \quad \text{Min}_{q,q'}|X_{q,2} - X_{q',2}| = 2;$$

$$X_{q,3} - X_{q',3} \in \{ -48, -45, -40, -36, -24, -21, -16, -12, -9, -8, -5, -4, -3, 0, 3, 4, 5, 8, 9, 12, 16, 21, 24, 36, 40, 45, 48 \}$$

$$(105) \quad \text{PrimesDiffs}(6, 3) = \{2, 3, 5, 7\}, \quad \text{Min}_{q,q'}|X_{q,3} - X_{q',3}| = 3.$$

We follow the procedure and require $p_1 \notin \text{PrimesDiffs}(6, 2)$, hence, for instance $p_1 = 11$. Then we seek p_2 that obeys

$$(106) \quad \begin{aligned} & \text{Max}_{q,q'} \left(|X_{q,2} - X_{q',2}| + p_1 |X_{q,3} - X_{q',3}| \right) \\ & < p_2 \text{Min}_{q,q'} \left| (X_{q,2} - X_{q',2}) + p_1 (X_{q,3} - X_{q',3}) \right| \\ & \text{Max}_{q,q'} \left(|X_{q,2} - X_{q',2}| + p_1 |X_{q,3} - X_{q',3}| \right) = 30 + p_1 * 48 = 30 + 11 * 48 = 558, \\ & \text{Min}_{q,q'} \left| (X_{q,2} - X_{q',2}) + p_1 (X_{q,3} - X_{q',3}) \right| = 2. \end{aligned}$$

Thus we seek a prime p_2 such that

$$(107) \quad 558 < 2p_2, \quad p_2 > 279.$$

We then use $p_2 = 289$. It remains to determine p_3 satisfying

$$(1 + p_2) \times 558 < 2p_3$$

$$(108) \quad (1 + 289) \times 558 = 161820 < 2p_3, \quad p_3 > 80910.$$

that gives $p_3 = 80917$. Thus the quadruple is $(1, p_1, p_2, p_3) = (1, 11, 289, 80917)$ with $a_{1,2} = 1, a_{1,3} = 11, a_{2,2} = 289, a_{2,3} = 289 \times 11 = 3179, a_{3,2} = 80917, a_{3,3} = 80917 \times 11 = 890087$ solves the condition (92). As a result the Hamiltonian $\mathcal{H} = \sum_{i,k} a_{i,k} T_k^{(i)}$ with these coefficients has integer matrix elements in the ribbon graph basis, has distinct eigenvalues for distinct triples of Young diagrams, and eigenvalue degeneracies given by $C(R_1, R_2, R_3)^2$.

Sufficient conditions. At smallest order of $k_* = 2, 3$, there are quick sufficient conditions that solve the problem, for all $n = 2, 3, 4, \dots, 14$. We are confident that similar identities holds for higher order in k_* . Note that the solutions p_k 's provided below need not be the smallest possible but we arrive at easy programming equalities.

Consider first $k_* = 2$, pick p_1 as the first prime number above $n(n - 1)$ (as $\text{Max}_{q,q'} |X_{q,2} - X_{q',2}| = n(n - 1)$). This already guarantees that it does not belong to the set of prime divisors of the set $\{X_q - X_{q'}\}$. Then choose the prime $p_2 > (p_1 + 1) * n(n - 1)$ then $(a_1, a_2, a_3) = (1, p_1, p_2)$ solves the condition (84).

Addressing $k_* = 3$, we can replace $\text{Min}_{q,q'} |\cdot|$ by the lower bound 1 and the $\text{Max}_{q,q'} |\cdot| = 2\hat{\chi}_{[1^n]}(T_k)$. Let us illustrate this idea at $n = 6$. Already, p_1 has been fixed to be the smallest prime $p_1 > n(n - 1)$. We consider $\text{Max}_q X_{q,3} = \hat{\chi}_{[1^n]}(T_3) = \frac{n(n-1)(n-2)}{3}$, and thus $\text{Max}_{q,q'} |X_{q,3} - X_{q',3}| \leq 2\frac{n(n-1)(n-2)}{3}$. We choose

$$(109) \quad \begin{aligned} p_2 &> \text{Max}_{q,q'} (|X_{q,2} - X_{q',2}|) + p_1 \text{Max}(|X_{q,3} - X_{q',3}|) \\ &> n(n - 1) + p_1 \frac{2}{3} n(n - 1)(n - 2). \end{aligned}$$

Thus we choose p_2 to be the next prime after $\frac{n}{3}(n - 1)(3 + 2p_1(n - 2))$. Last p_3 should obey the bound

$$(110) \quad \begin{aligned} p_3 &> (1 + p_2) \text{Max}_{q,q'} (|X_{q,2} - X_{q',2}| + p_1 |X_{q,3} - X_{q',3}|) \\ &> (1 + p_2)(n(n - 1) + p_1 \frac{2}{3} n(n - 1)(n - 2)). \end{aligned}$$

Therefore picking p_3 as the next prime larger than $\frac{n}{3}(n - 1)(1 + p_2)(3 + 2p_1(n - 2))$ will solve the issue.

At $n = 6$, $\text{Max}_{q,q'} |X_{q,2} - X_{q',2}| = 30 = 2\hat{\chi}_{[1^n]}(T_2)$, We can pick $p_1 = 31$. Then $\text{Max}_{q,q'} |X_{q,3} - X_{q',3}| = 48 < 80 = 2\hat{\chi}_{[1^n]}(T_3)$. This will fixe p_2 and p_3 . Then an alternative quadruple that solves the problem is given by $(1, 31, 2521, 6330223)$ (to be compared with the previous quadruple $(1, 11, 289, 80917)$ in equation (108)).

4. KRONECKER COEFFICIENTS AND RIBBON GRAPH SUB-LATTICES

In Section 3, we constructions of an integer matrix for each ordered triple of Young diagrams (R_1, R_2, R_3) with n boxes, with the property that their null space gives a basis for the Fourier subspace of $\mathcal{K}(n)$ associated with that triple. This subspace has dimension equal to the square of the Kronecker coefficient : $C(R_1, R_2, R_3)^2$. These matrices are constructed from central elements $T_k^{(i)}$ (introduced in Section 2.4) of the algebra $\mathcal{K}(n)$ of bipartite ribbon graphs with n edges, where $i \in \{1, 2, 3\}$ and $k \in \{2, 3, \dots, \tilde{k}_*\}$. The parameter $\tilde{k}_* \in \{k_*(n), k_*(n) + 1, \dots, n\}$, where $k_*(n)$ is the minimal integer such that the central elements $\{T_2, \dots, T_{k_*(n)}\}$ generate the centre of $\mathbb{C}(S_n)$ and it has been computed for n up to 79 [57]. We have two constructions for each Young diagram triple, one producing a rectangular matrix $\mathcal{L}_{R_1, R_2, R_3}$ (72) and another producing a square matrix $\mathcal{H}_{R_1, R_2, R_3}$ (80). In each case, we are solving the

linear equation

$$(111) \quad X \cdot v = 0$$

where $X = \mathcal{L}_{R_1, R_2, R_3}$ or $X = \mathcal{H}_{R_1, R_2, R_3}$.

The null spaces of integer matrices have bases given as integer vectors. This follows from the theory of Hermite normal forms and has an interpretation in terms of sub-lattices. In the present application we have a lattice

$$(112) \quad \mathbb{Z}^{|\text{Rib}(n)|} \subset \mathbb{R}^{|\text{Rib}(n)|}$$

which is interpreted as the space of integer linear combinations of the geometric ribbon graph basis vectors E_r of the ribbon graph algebra $\mathcal{K}(n)$. We will refer to $\mathbb{Z}^{|\text{Rib}(n)|}$ as the lattice of ribbon graphs. In this section, we will explain the key facts about integer matrices and sub-lattices that we will need and state the first main result of this paper, Theorem 4.1. This is the construction of $C(R_1, R_2, R_3)^2$ as the dimension of a sub-lattices of the lattice of ribbon graphs.

A classic problem asks for a combinatorial construction of the Kronecker coefficient associated with every triple of Young diagrams [77, 90]. Recent progress on this problem from a number of directions and its connections to computational complexity is summarised in [80]. Theorem 4.1 gives a *combinatorial interpretation* of the square of the Kronecker coefficients. The theory of Hermite normal forms for integer matrices also offers *combinatorial algorithms* for finding the null spaces (Corollary 4.2). It is also interesting to ask if there is a purely combinatorial proof - without using representation theory - of the formula for Kronecker coefficients in terms of characters, which can be viewed as combinatorial objects, for example, by the Murnaghan–Nakayama algorithm. Our proof of Theorem 4.1 relies in an important way on representation theory, e.g. in the derivation of Proposition 3.1 which enters the proof, and is therefore not purely combinatorial. In Section 4.2 we discuss how the question of a purely combinatorial proof of the Theorem 4.1 raises interesting questions on integer matrices.

In Section 4.3, we consider the $S = \pm 1$ eigenspaces of the conjugation operator defined in 2.2.2. This leads to the definition of sub-lattices of the lattice of ribbon graphs with dimensions $C(R_1, R_2, R_3)(C(R_1, R_2, R_3) + 1)/2$ and $C(R_1, R_2, R_3)(C(R_1, R_2, R_3) - 1)/2$, constructed as null spaces of integer matrices. The difference of these dimensions is $C(R_1, R_2, R_3)$ which can therefore be constructed by choosing a map from a basis set for the null vectors, determined for example by a Hermite normal form algorithm, for the smaller sub-lattice to a basis set for the larger sub-lattice (Theorem 4.4).

4.1. NULL-VECTORS OF INTEGER MATRICES AND LATTICES. The null space of the integer matrix X defined by (111) is the span of a set of null vectors which can be chosen to be integer vectors, i.e. integral linear combinations of the E_r . A key result from the theory of integer matrices and lattices is that any integer matrix A (square or rectangular) has a unique Hermite normal form (HNF) (as explained in textbooks such as [20][88] or online notes such as [74]). These can be computed using mathematical software such as GAP, SAGE or Mathematica. Thus A has a decomposition $A = Uh$: U is a unimodular matrix, i.e. an integer matrix of determinant ± 1 . In the following discussion we will use $A = X^T$. h is an integer matrix with the following properties :

- h is upper triangular (that is, $h_{ij} = 0$ for $i > j$), and any rows of zeros are located below any other row.
- The leading coefficient (the first non-zero entry from the left, also called the pivot) of a non-zero row is always strictly to the right of the leading coefficient of the row above it; moreover, it is positive.

- The elements below pivots are zero and elements above pivots are non-negative and strictly smaller than the pivot.

The construction of h proceeds by applying a sequence of operations involving one of the following in each step.

- Swop two rows.
- Multiply a row by -1 .
- Add an integer multiple of a row to another row of A .

Each of these operations corresponds to left multiplication of A by a unimodular matrix U : $A \rightarrow UA$. Every integer matrix A can be brought into HNF by a sequence of these elementary integer row operations. Suppose we want to find the vectors v which obey $Xv = 0$. Equivalently $v^T X^T = 0$. We apply the elementary integer row operations to bring X^T to HNF. This means $X^T = Uh$. The number N of lower rows of zeroes in h is the dimension of the null space of X . The integer null vectors can be read off from the lower N rows of U . The non-zero rows of h give an integer basis for the image of X .

As a simple example to illustrate these properties, take

$$(113) \quad X = \begin{pmatrix} 1 & 1 \\ 2 & 2 \end{pmatrix}, \quad X^T = \begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix},$$

with X^T being the transpose of X . By applying the row operation of replacing the second row R_2 by $R_2 - R_1$ we get the HNF

$$(114) \quad X^T \rightarrow h = \begin{pmatrix} 1 & 2 \\ 0 & 0 \end{pmatrix}.$$

The unimodular matrix which implements this row operation is

$$(115) \quad U = \begin{pmatrix} 1 & 0 \\ 1 & -1 \end{pmatrix}$$

i.e

$$(116) \quad UX^T = \begin{pmatrix} 1 & 2 \\ 0 & 0 \end{pmatrix} = h.$$

The lower row of U , when transposed, gives the null vector for the action of $X : v \rightarrow Xv$

$$(117) \quad X \begin{pmatrix} 1 \\ -1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

The non-vanishing row of h , transposes to the column vector which gives the image of X since

$$(118) \quad X \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = (x_1 + x_2) \begin{pmatrix} 1 \\ 2 \end{pmatrix}.$$

To see that the connection between the lower rows of the unimodular matrix U in the decomposition $UX^T = h$ corresponds to null vectors, observe that

$$(119) \quad UX^T = h$$

can be written as

$$(120) \quad \sum_k U_{ik} X_{kj}^T = h_{ij}.$$

The vanishing rows of h correspond to values of i such that $h_{ij} = 0$ for all j . Fixing one of these i we have vectors U_{ik} as k varies, with the property:

$$(121) \quad \sum_k X_{jk} U_{ik} = 0.$$

Note that we could have equivalently worked with elementary column operations on X rather than elementary row operations on X^T .

By definition the matrix U has integer entries, so this construction gives an integer basis for the null space of X^T . The null vectors of X , found as integer linear combinations of E_r , define a sub-lattice of the lattice $\mathbb{Z}^{|\text{Rib}(n)|}$. The dimension of the sub-lattice is the square $(C(R_1, R_2, R_3))^2$ of the Kronecker coefficient. The square of the Kronecker coefficient thus has the combinatorial interpretation as the dimension of a sub-lattice of the lattice of ribbon graphs. We have thus arrived at the first main theorem of this paper.

THEOREM 4.1. *For every triple of Young diagrams (R_1, R_2, R_3) with n boxes, the lattice*

$$(122) \quad \mathbb{Z}^{|\text{Rib}(n)|}$$

of integer linear combinations of the geometric basis vectors E_r of $\mathcal{K}(n)$ contains a sub-lattice of dimension $(C(R_1, R_2, R_3))^2$ spanned by a basis of integer null vectors of the operator X , which is $\mathcal{L}_{R_1, R_2, R_3}$ from (72) in the rectangular matrix construction or $\mathcal{H}_{R_1, R_2, R_3}$ from (80) in the square matrix construction.

Solving for the null vectors of X using the HNF shows that there is sub-lattice in the lattice of ribbon graphs whose dimension is $(C(R_1, R_2, R_3))^2$. This gives a combinatorial interpretation for the square of the Kronecker coefficient.

But the theory of lattices is even more powerful. The columns of X (equivalently the rows of X^T) are a set of vectors in the lattice of ribbon graphs. The space spanned by the integer linear combinations of these vectors is a sub-lattice of dimension $|\text{Rib}(n)| - C(R_1, R_2, R_3)^2$ (that is the column rank of X or the row rank X^T). The process of arriving at the HNF through row operations on X^T amounts to simplifying the description of this sub-lattice until it is given as the integer linear combinations of a linearly independent set of integer vectors which sit in the rows of h . This process also defines a unimodular matrix which encodes the integer null vectors of X . Each step in the process of row operations acts on the set of lattice vectors in X , and can thus be viewed as constructive combinatorial steps.

COROLLARY 4.2. *There is a constructive procedure for the sub-lattice in Theorem 4.1 consisting of integer row operations on the list of integer rows of X^T , which produce the HNF of X^T .*

Proof. The treatment of rows of X^T to put it in a HNF form $X^T = Uh$ is a combinatorial construction consisting of a discrete sequence of integer row operations (swop, multiplying by -1 and integer linear combinations of rows). The outcome h of the HNF construction gives a basis for the sub-lattice of dimension $|\text{Rib}(n)| - C(R_1, R_2, R_3)^2$ spanned by integer linear combinations of the rows of X^T . The outcome U is built in successive steps by matrices implementing the integer elementary row operations on X^T . At the start of an algorithm for the HNF of X^T , the rows give a generically over-complete basis for the lattice generated by these rows. X^T is modified step by step until the last step produces h . At each step the intermediate matrix has a list of lattice vectors. At the end of an algorithm for the HNF, there is a sequence of rows of zeros in h and the corresponding rows of U record the integer null vectors, the number of which is $C(R_1, R_2, R_3)^2$. The construction of U associated with a given X is thus a sequence of combinatorial operations on lattice vectors in $\mathbb{Z}^{|\text{Rib}(n)|}$. □

The key fact from the theory of integer matrices and lattices we have used is the existence and uniqueness of the HNF. In the above we have focused on the fact that

integral algorithms exist which produce from X , the null vectors and the HNF. We have not focused on the computational complexity of the problem. We make some initial remarks in this direction. The LLL algorithm [68] is known to calculate HNF's in a time that is polynomial in the size of the matrix. Our matrices are very large - grow as the number of ribbon graphs. We know from [8] for a partition $\sum_i i p_i = n$

$$(123) \quad |\text{Rib}(n)| = \sum_{p \vdash n} \prod_{i=1}^n i^{p_i} (p_i!).$$

The leading orders of the asymptotics of this number are known (see [64]): $|\text{Rib}(n)| \sim n! * (1 + 2/n^2 + 5/n^3 + 23/n^4 + 106/n^5 + 537/n^6 + 3143/n^7 + 20485/n^8 + 143747/n^9 + 1078660/n^{10})$. Thus the data size of our problem already grows like $\mathcal{O}(n!)$ (assuming that $k_*(n) \ll n$ as $n \rightarrow \infty$). It seems that combined with a problem of time complexity, our problem entails a super-exponential complexity in (memory) space. Hence, although the time complexity of HNF could be polynomial in the data size, it would remain $\mathcal{O}(n!)$. A more thorough discussion of the complexity of the algorithm for construction the sub-lattice in Theorem 4.1 taking into account the group theoretic characteristics of the integer matrix X is left for the future.

4.2. COMBINATORIAL INTERPRETATIONS, ALGORITHMS AND PROOFS. An interesting question for a combinatorial construction of Kronecker coefficients posed in [90] is whether it gives a new proof of the fact that these coefficients are non-negative integers. It is of course obvious from representation theory that $C(R_1, R_2, R_3)$ is non-negative-integer - it is the number of times R_3 appears in the tensor product decomposition $R_1 \otimes R_2$ when viewed as a representation using the diagonal action of permutations. The character formula

$$(124) \quad C(R_1, R_2, R_3) = \frac{1}{n!} \sum_{\sigma \in S_n} \chi_{R_1}(\sigma) \chi_{R_2}(\sigma) \chi_{R_3}(\sigma)$$

- although it can be derived from representation theory - can also be viewed as a purely combinatorial formula, where the characters are given for example by the Murnaghan–Nakayama combinatorial rule. From the purely combinatorial point of view, the non-negative integer property is not manifest.

The sub-lattice interpretation of Kronecker coefficients (Theorem 4.1) makes it manifest that they are non-negative integers. Algorithms for computing the HNFs are combinatorial operations on lists of lattice vectors. Our proof of the interpretation and of the validity of the algorithms has relied on an important input from representation theory (Proposition 3.1). An interesting question is whether lattices of ribbon graphs offer an avenue to provide a purely combinatorial understanding, with no representation theory input, for the non-negativity of $C(R_1, R_2, R_3)$, defined by the formula (124) in terms of characters computable by the combinatorial Murnaghan–Nakayama rule. To give some context to this question, consider the equality of $n!$ with the sum of squares of the dimensions of irreducible representations of S_n , which can be derived using representation theory. This can also be derived purely combinatorially using the Robinson-Schensted correspondence which gives a bijection between permutations in S_n and pairs of standard Young tableaux having the same shape and n boxes (see for example a textbook reference [38]).

This raises some questions on the non-negative integer matrices $T_k^{(i)}$, the rectangular integer matrices in Section 3.2 and the square matrices of Hamiltonian matrix elements in Section 3.3. The first step would be to derive formulae for the eigenvalues $T_k^{(i)}$ recovering the Murnaghan–Nakayama combinatorics of these eigenvalues directly from these matrices built using the reconnection matrices $T_k^{(i)}$. The second

step would be to show that the eigenvalue degeneracies are given by (124), viewed as an expression for the degeneracies in terms of the eigenvalues. Any integer matrix is known to have a Smith normal form (SNF) which can be calculated by algorithms generalizing to those for HNFs [20]. In the SNF for X , we have $X = UDV$, where the matrix D is a diagonal matrix of singular values. The relation between these singular values and the eigenvalues of an integer matrix X has been studied in [70]. Singular values in the SNF are accessible to integer-matrix algorithms while eigenvalues enter the link between the integer matrices at hand and the Kronecker coefficients. Better understanding this link could potentially help towards a purely combinatorial proof of the interpretation and algorithms for Kronecker coefficients based on Theorem 4.1.

4.3. CONJUGATION ACTION AND ADDITIONAL SUB-LATTICES. In section (4.1), the HNF of integer matrices to a $C(R_1, R_2, R_3)^2$ dimensional sublattice of ribbon graphs and has provided a refinement of the counting of all ribbon graphs. We now describe integer matrices which will lead us to a sublattice interpretation of $C(R_1, R_2, R_3)$.

In Section 2.2.2 we defined a conjugation operator S (24) which satisfies $S^2 = \text{id}$. The conjugation either maps a ribbon graph to itself $S(E_r) = E_r$, or distinct pairs $E_s \neq E_t$ are related by $S(E_s) = E_t; S(E_t) = E_s$. We refer to the former as self-conjugate ribbon graphs and the the latter as conjugate pairs.

In order to illustrate the action of S consider $n = 3$. Inversion of the permutation pairs representing a ribbon graph leaves the pair unchanged unless one of the permutations has a cycle of length 3. For this $n = 3$ case, all ribbon graph vectors E_r are self-conjugate : inversion maps any representative pair or permutations to another pair within the same orbit. We list the orbits at $n = 3$ which involve a cycle of length 3, to illustrate this property

$$\begin{aligned}
 3 : & \quad [((1, 2, 3)), ((1, 3, 2))] \\
 7 : & \quad [((2, 3), (1, 2, 3)), ((2, 3), (1, 3, 2)), ((1, 2), (1, 2, 3)), \\
 & \quad ((1, 2), (1, 3, 2)), ((1, 3), (1, 2, 3)), ((1, 3), (1, 3, 2))] \\
 8 : & \quad [((1, 2, 3), ()), ((1, 3, 2), ())] \\
 9 : & \quad [((1, 2, 3), (2, 3)), ((1, 2, 3), (1, 2)), ((1, 2, 3), (1, 3)), \\
 & \quad ((1, 3, 2), (2, 3)), ((1, 3, 2), (1, 2)), ((1, 3, 2), (1, 3))] \\
 10 : & \quad [((1, 2, 3), (1, 2, 3)), ((1, 3, 2), (1, 3, 2))] \\
 11 : & \quad [((1, 2, 3), (1, 3, 2)), ((1, 3, 2), (1, 2, 3))]
 \end{aligned}
 \tag{125}$$

where the first column contains the labels (i.e 3,7,8, etc.) of ribbon graphs of Figure 1. As we will see shortly, this self-conjugation property can be understood using the action of S on the Fourier basis of $\mathcal{K}(n)$.

The following statement holds:

PROPOSITION 4.3. Under the conjugation action, we have

$$S(Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}) = Q_{\tau_2, \tau_1}^{R_1, R_2, R_3}.
 \tag{126}$$

Proof. Consider $Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}$ given by (14), then

$$\begin{aligned}
 S(Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}) &= \\
 \kappa_{R_1, R_2} \sum_{\sigma_1, \sigma_2 \in S_n} \sum_{i_1, j_1} C_{i_1, i_2; i_3}^{R_1, R_2; R_3, \tau_1} C_{j_1, j_2; i_3}^{R_1, R_2; R_3, \tau_2} D_{i_1 j_1}^{R_1}(\sigma_1) D_{i_2 j_2}^{R_2}(\sigma_2) \sigma_1^{-1} \otimes \sigma_2^{-1} \\
 &= \kappa_{R_1, R_2} \sum_{\sigma_1, \sigma_2 \in S_n} \sum_{i_1, j_1} C_{i_1, i_2; i_3}^{R_1, R_2; R_3, \tau_1} C_{j_1, j_2; i_3}^{R_1, R_2; R_3, \tau_2} D_{i_1 j_1}^{R_1}(\sigma_1^{-1}) D_{i_2 j_2}^{R_2}(\sigma_2^{-1}) \sigma_1 \otimes \sigma_2 \\
 &= \kappa_{R_1, R_2} \sum_{\sigma_1, \sigma_2 \in S_n} \sum_{i_1, j_1} C_{i_1, i_2; i_3}^{R_1, R_2; R_3, \tau_1} C_{j_1, j_2; i_3}^{R_1, R_2; R_3, \tau_2} D_{j_1 i_1}^{R_1}(\sigma_1) D_{j_2 i_2}^{R_2}(\sigma_2) \sigma_1 \otimes \sigma_2 \\
 &= \kappa_{R_1, R_2} \sum_{\sigma_1, \sigma_2 \in S_n} \sum_{i_1, j_1} C_{j_1, j_2; i_3}^{R_1, R_2; R_3, \tau_1} C_{i_1, i_2; i_3}^{R_1, R_2; R_3, \tau_2} D_{i_1 j_1}^{R_1}(\sigma_1) D_{i_2 j_2}^{R_2}(\sigma_2) \sigma_1 \otimes \sigma_2
 \end{aligned}$$

$$(127) \quad = Q_{\tau_2, \tau_1}^{R_1, R_2, R_3}.$$

We used the fact that $D_{ij}^R(\sigma^{-1}) = D_{ji}^R(\sigma)$ and a relabelling of indices $i_1, i_2 \leftrightarrow j_1, j_2$. \square

Remark Proposition 4.3 implies that at $n = 3$, where $C(R_1, R_2, R_3)$ is either 1 or 0, the only possible eigenvalue of S is 1. Considering then the action of S on the geometric basis, we deduce that all the ribbon graphs must be self-conjugate. This is indeed confirmed by (125).

On the geometrical ribbon graph basis for $\mathcal{K}(n)$, the action of S can leave a ribbon basis element E_r invariant, or it can pair the ribbon with another ribbon. Let us denote by $E_r^{(s)}$ the self-conjugate ribbons, which stay invariant under conjugation. The non-self conjugate pairs are $(E_r^{(n)}, E_r^{(\bar{n})})$. The $+1$ eigenspace of S in $\mathcal{K}(n)$ is spanned by $\{E_r^{(s)}, (E_r^{(n)} + E_r^{(\bar{n})})\}$. The -1 eigenspace of S is spanned by $\{(E_r^{(n)} - E_r^{(\bar{n})})\}$. Let us denote the vector space of ribbon graphs, which is the underlying vector space of the algebra $\mathcal{K}(n)$ by $V^{\text{Rib}(n)}$. $V^{\text{Rib}(n)}$ has a decomposition according to the eigenvalues of S

$$(128) \quad V^{\text{Rib}(n)} = V_{S=1}^{\text{Rib}(n)} \oplus V_{S=-1}^{\text{Rib}(n)}.$$

The $S = 1$ -eigenspace is the direct sum

$$(129) \quad V_{S=1}^{\text{Rib}(n)} = V_{\text{pairs}^+}^{\text{Rib}(n)} \oplus V_{\text{singlets}}$$

where $V_{\text{pairs}^+}^{\text{Rib}(n)}$ is spanned by $\{(E_r^{(n)} + E_r^{(\bar{n})})\}$ and V_{singlets} by $\{E_r^{(s)}\}$, whereas the $S = (-1)$ -eigenspace is

$$(130) \quad V_{S=-1}^{\text{Rib}(n)} = V_{\text{pairs}^-}^{\text{Rib}(n)}.$$

Using the Wedderburn–Artin decomposition of $\mathcal{K}(n)$ we also have

$$(131) \quad V^{\text{Rib}(n)} = \bigoplus_{R_1, R_2, R_3} V^{\text{Rib}(n): R_1, R_2, R_3}$$

where $V^{\text{Rib}(n): R_1, R_2, R_3}$ has dimension $C(R_1, R_2, R_3)^2$ and is spanned by the $Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}$ for all τ_1 and τ_2 . The projection of $V^{\text{Rib}(n)}$ to a fixed R_1, R_2, R_3 commutes with the operator S . This is evident from Proposition 4.3. Using this proposition, it is also obvious that the $S = 1$ subspace of $V^{\text{Rib}(n): R_1, R_2, R_3}$ is given by

$$(132) \quad \begin{aligned} V_{S=1}^{\text{Rib}(n): R_1, R_2, R_3} = & \text{Span}\{Q_{\tau, \tau}^{R_1, R_2, R_3} : 1 \leq \tau \leq C(R_1, R_2, R_3)\} \\ & \oplus \text{Span}\{Q_{\tau_1, \tau_2}^{R_1, R_2, R_3} + Q_{\tau_2, \tau_1}^{R_1, R_2, R_3} : 1 \leq \tau_1 < \tau_2 \leq C(R_1, R_2, R_3)\} \end{aligned}$$

and its $S = -1$ subspace is

$$(133) \quad V_{S=-1}^{\text{Rib}(n): R_1, R_2, R_3} = \text{Span}\{Q_{\tau_1, \tau_2}^{R_1, R_2, R_3} - Q_{\tau_2, \tau_1}^{R_1, R_2, R_3} : 1 \leq \tau_1 < \tau_2 \leq C(R_1, R_2, R_3)\}.$$

Then $V^{\text{Rib}(n): R_1, R_2, R_3} = V_{S=1}^{\text{Rib}(n): R_1, R_2, R_3} \oplus V_{S=-1}^{\text{Rib}(n): R_1, R_2, R_3}$. Combining this with (131) we then have

$$(134) \quad V^{\text{Rib}(n)} = \bigoplus_{R_1, R_2, R_3} \left(V_{S=1}^{\text{Rib}(n): R_1, R_2, R_3} \oplus V_{S=-1}^{\text{Rib}(n): R_1, R_2, R_3} \right).$$

From (133) we deduce that

$$(135) \quad \begin{aligned} \text{Dim} \left(V_{S=-1}^{\text{Rib}(n): R_1, R_2, R_3} \right) &= \frac{C(R_1, R_2, R_3)(C(R_1, R_2, R_3) - 1)}{2} \\ &= \text{Dim} \left(P^{R_1, R_2, R_3} V_{\text{pairs}^-}^{\text{Rib}(n)} \right) \end{aligned}$$

with P^{R_1, R_2, R_3} the projector onto $V^{\text{Rib}(n):R_1, R_2, R_3}$. Similarly from (132) we have

$$\begin{aligned} \text{Dim} \left(V_{S=+1}^{\text{Rib}(n):R_1, R_2, R_3} \right) &= \frac{C(R_1, R_2, R_3)(C(R_1, R_2, R_3) + 1)}{2} \\ &= \text{Dim} \left(P^{R_1, R_2, R_3} V_{\text{pairs}^+}^{\text{Rib}(n)} \right) + \text{Dim} \left(P^{R_1, R_2, R_3} V_{\text{singlets}}^{\text{Rib}(n)} \right). \end{aligned} \tag{136}$$

Note that we do not have separate expressions for the two terms in the sum above in terms of Kronecker coefficients, since we do not expect the P^{R_1, R_2, R_3} to commute with the projection of $V_{S=1}^{\text{Rib}(n)}$ into the separate summands $V_{\text{singlets}}^{\text{Rib}(n)}$ and $V_{\text{pairs}^+}^{\text{Rib}(n)}$.

If we do the sum over R_1, R_2, R_3 , we have

$$\begin{aligned} \text{Dim} \left(V_{S=+1}^{\text{Rib}(n)} \right) &= \sum_{R_1, R_2, R_3} \frac{C(R_1, R_2, R_3)(C(R_1, R_2, R_3) + 1)}{2} \\ &= \text{Dim} \left(V_{\text{pairs}^+}^{\text{Rib}(n)} \right) + \text{Dim} \left(V_{\text{singlets}} \right) \end{aligned} \tag{137}$$

and
(138)

$$\text{Dim} \left(V_{S=-1}^{\text{Rib}(n)} \right) = \sum_{R_1, R_2, R_3} \frac{C(R_1, R_2, R_3)(C(R_1, R_2, R_3) - 1)}{2} = \text{Dim} \left(V_{\text{pairs}^-}^{\text{Rib}(n)} \right).$$

Since

$$\text{Dim} \left(V_{\text{pairs}^+}^{\text{Rib}(n)} \right) = \text{Dim} \left(V_{\text{pairs}^-}^{\text{Rib}(n)} \right) \tag{139}$$

we have

$$\text{Dim} \left(V_{\text{pairs}^+}^{\text{Rib}(n)} \right) = \sum_{R_1, R_2, R_3} \frac{C(R_1, R_2, R_3)(C(R_1, R_2, R_3) - 1)}{2}, \tag{140}$$

$$\text{Dim} \left(V_{\text{singlets}}^{\text{Rib}(n)} \right) = \sum_{R_1, R_2, R_3} C(R_1, R_2, R_3). \tag{141}$$

While the sum over triples of Young diagrams with n boxes of the square of Kronecker coefficients gives the number of ribbon graphs with n edges, the sum of the Kronecker coefficients gives the number of singlet ribbon graphs.

The sequence of sums of the Kroneckers for $n = 1, \dots, 10$ is

$$1, 4, 11, 43, 149, 621, 2507, 11174, 49972, 237630 \tag{142}$$

that coincide with the number of self-conjugate ribbon graphs for $n = 1, \dots, 6$

$$1, 4, 11, 43, 149, 621. \tag{143}$$

For $n \geq 7$ our current GAP program for enumerating the self-conjugate ribbons is no longer very efficient, but by the derivation we have given of (141) these two sequences will agree.

The projection from $\mathcal{K}(n)$ to $V_{S=1}^{\text{Rib}(n):R_1, R_2, R_3}$ can be done by using the $T_k^{(i)}$ for $i \in \{1, 2, 3\}; k \in \{2, 3, \dots, \tilde{k}_*\}$ to build a rectangular matrix as in (72) which projects to R_1, R_2, R_3 and further stacking the matrix $S - 1$. This gives an integer matrix of size $(3(\tilde{k}_* - 1) + 1)|\text{Rib}(n)| \times |\text{Rib}(n)|$ with null space spanning $V_{S=1}^{\text{Rib}(n):R_1, R_2, R_3}$. We can also use the Hamiltonian square matrix construction (80) along with the $S - 1$ matrix to build an integer matrix of size $2|\text{Rib}(n)| \times |\text{Rib}(n)|$ which projects to $V_{S=1}^{\text{Rib}(n):R_1, R_2, R_3}$. By replacing $(S - 1)$ with $(S + 1)$ in these constructions we can obtain the subspace $V_{S=-1}^{\text{Rib}(n):R_1, R_2, R_3}$ of $\mathcal{K}(n)$ as null spaces of integer rectangular or square matrices.

As in Section 4.1 the HNF construction of $V_{S=\pm 1}^{\text{Rib}(n):R_1,R_2,R_3}$ determines sub-lattices of $\mathbb{Z}^{|\text{Rib}(n)|}$. Thus, on one hand, we have lattice constructions for

$$(144) \quad \frac{C(R_1, R_2, R_3)(C(R_1, R_2, R_3) - 1)}{2}$$

as the dimension of $V_{\text{pairs}^-}^{\text{Rib}(n):R_1,R_2,R_3}$ and, on the other, we also have a construction of

$$(145) \quad \frac{C(R_1, R_2, R_3)(C(R_1, R_2, R_3) + 1)}{2}$$

as the dimension of $V_{S=1}^{\text{Rib}(n):R_1,R_2,R_3}$. The difference of these is the number $C(R_1, R_2, R_3)$. By choosing an injection between the smaller sub-lattice and the larger sub-lattice, we can get a constructive interpretation of $C(R_1, R_2, R_3)$. It will be interesting to investigate if there is a canonical choice of such an injection.

We summarise the outcome of the above discussion as a theorem

THEOREM 4.4. *For every triple of Young diagrams (R_1, R_2, R_3) with n boxes, there are three constructible sub-lattices of $\mathbb{Z}^{|\text{Rib}(n)|}$ of respective dimensions $C(R_1, R_2, R_3)(C(R_1, R_2, R_3) + 1)/2$, $C(R_1, R_2, R_3)(C(R_1, R_2, R_3) - 1)/2$, and $C(R_1, R_2, R_3)$.*

As an illustration, there are two interesting cases at $n = 5$ with $C(R_1, R_2, R_3) = 2$, $(R_1, R_2, R_3) = ([3, 2], [3, 1, 1], [3, 1, 1])$ and $(R_1, R_2, R_3) = ([3, 1, 1], [3, 1, 1], [2, 2, 1])$, their permutations. We have $([3, 2], [3, 1, 1], [3, 1, 1])$:

$$(146) \quad \begin{aligned} \text{Dim} \left(V^{\text{Rib}(n):R_1,R_2,R_3} \right) &= 4 \\ \text{Dim} \left(V_{S=+1}^{\text{Rib}(n):R_1,R_2,R_3} \right) &= 3 \\ \text{Dim} \left(V_{S=-1}^{\text{Rib}(n):R_1,R_2,R_3} \right) &= 1. \end{aligned}$$

The same equations hold for $([3, 1, 1], [3, 1, 1], [2, 2, 1])$.

5. CONCLUSIONS

We give a summary of our main results and outline important directions for future research. Section 5.2 uses the link between bi-partite ribbon graphs and Belyi maps. Section 5.3 outlines quantum algorithms motivated by links between the algebra $\mathcal{K}(n)$ and Kronecker coefficients. Section 5.4 describes physically motivated generalizations of the present work based on algebras related to $\mathcal{K}(n)$ which also have interesting geometric interpretations.

5.1. SUMMARY. In this paper we have developed quantum mechanics on a class of state spaces which are also algebras (in the present case the algebras $\mathcal{K}(n)$), and have a distinguished geometrical/combinatorial basis associated with combinatorial objects (in the case at hand bipartite ribbon graphs). The combinatorial objects have a description in terms of equivalence classes defined using permutations and the algebra can be realised as a subspace of a tensor product of group algebras (in this case $\mathbb{C}(S_n) \otimes \mathbb{C}(S_n)$ - and there is a gauge equivalent formulation in terms of $\mathbb{C}(S_n)^{\otimes 3}$ as explained in [9]). By exploiting the algebra structure, we are able to relate the eigenvalues of Hermitian Hamiltonians on these state spaces to characters of symmetric groups, and the multiplicities to group theoretic multiplicities (in this case Kronecker coefficients). The integrality structure of the algebra when expressed in terms of the geometrical basis means that the solving the Hamiltonians is a problem that can draw on techniques from the mathematics of integer matrices and lattices. It follows that

the square of Kronecker coefficients (C^2) can be realised as the dimension of a sub-lattice in the lattice generated by ribbon graphs. The algebra has an involution which is inherited from the inversion of permutations and commutes with the Hamiltonians considered here. The involution is used to define sub-lattices of dimensions $C(C+1)/2$ and $C(C-1)/2$. Choosing an injection of the set of sub-lattice basis vectors of the smaller sub-lattice, selected by the HNF construction of integer matrices we built, into the set of sub-lattice basis vectors of the larger sub-lattice also fixed by the HNF construction, leads to a sub-lattice of dimension C .

5.2. BELYI MAPS AND QUANTUM MEMBRANE INTERPRETATION OF QUANTUM MECHANICS ON $\mathcal{K}(n)$. Bipartite ribbon graphs have a rich geometrical structure related to Belyi maps and number theory. In this section, we use this connection to Belyi maps to give an interpretation of quantum mechanical evolution in the ribbon graph quantum mechanics in terms of membranes : the covering surfaces arising in Belyi maps appear at fixed time and can be interpreted as string worldsheets in topological strings - the quantum mechanical time is an additional coordinate which can be viewed as part of a membrane worldvolume.

Bipartite ribbon graphs with n edges are in 1-1 correspondence with holomorphic maps f (branched covers) from a Riemann surface Σ_g to two-dimensional Riemann sphere with degree n and 3 branch points :

$$(147) \quad f : \Sigma_g \rightarrow \mathbb{CP}^1.$$

These branch points can be taken to be $\{0, 1, \infty\}$ (see Chapter 2 of [66]). If we label the inverse images of a generic point on the sphere as $\{1, 2, \dots, n\}$, then the branching at the three branch points are described by the three permutations $\sigma_1, \sigma_2, \sigma_3 = (\sigma_1\sigma_2)^{-1}$. The genus g of the covering surface is given by the Riemann-Hurwitz formula

$$(148) \quad (2g - 2) = n - C_{\sigma_1} - C_{\sigma_2} - C_{\sigma_3}$$

where C_σ is the number of cycles of the permutation σ . Branched covers with exactly three branch points, also called Belyi maps, have the property that the covering surface Σ_g as well as the covering map can be defined in terms of equations with coefficients which are algebraic numbers, complex numbers which are solutions of polynomials with integer coefficients [6]. Conversely any such algebraic surface can be realised as a branched cover of the sphere with 3 branch points. The inverse image of the interval $[0, 1]$ on the Riemann sphere defines a graph embedded in the surface Σ_g , also called a map. These maps were called Dessins d'Enfants by Grothendieck who proposed their combinatorial study as a tool to understand representations of the absolute Galois Group, an object of fundamental importance in number theory [45]. A survey of mathematical work in this area is in [87]. It is interesting that the conjugation operation S which has played an important role in this paper (Section 4.3) has previously appeared in the study of "operations on maps". The self-conjugate graphs correspond to reflexible Belyi maps in the terminology of [56]. The number of distinct terms in E_r , when expanded as a sum in $\mathbb{C}(S_n) \otimes \mathbb{C}(S_n)$ is $n!$ divided by the order of the automorphism group of the Belyi map, i.e. the number of holomorphic invertible maps $\phi : \Sigma_g \rightarrow \Sigma_g$ obeying $f \circ \phi = f$.

Each ribbon graph defines an element $E_r \in \mathcal{K}(n) = \mathbb{C}(S_n) \otimes \mathbb{C}(S_n)$. The quantum mechanical evolution of such a state produces

$$(149) \quad E_r(t) = e^{-i\mathcal{H}t} E_r.$$

At generic t , this is a superposition of different basis elements $E_s \in \mathcal{K}(n)$. Such a superposition determines a linear combination of Belyi curves and Belyi maps. The evolution over all $t \geq 0$ determines a quantum membrane worldvolume mapping to $S^2 \times \mathbb{R}^+$ which restricts to a single Belyi map at $t = 0$ and subsequent periodic

intervals, but is generically a superposition of covering surfaces mapping to S^2 . Belyi maps have been linked to matrix models and topological strings [26, 42]. The discrete spin states of a particle such as an electron (two-dimensional state space) which are the subject of the quantum mechanics of spin are being generalized in the quantum mechanics of ribbon graphs to discrete states of a two-dimensional surface. These discrete states have the rich structure of an algebra, at each n the algebra $\mathcal{K}(n)$, and they have the rich geometrical structure in terms of an algebraic number realization as Belyi curves. The additional time direction of the quantum mechanics forms the $2 + 1$ dimensional worldvolume of a membrane generalizing the $0 + 1$ dimensional worldline of a particle. It would be interesting to develop the realisations of such quantum mechanical evolutions using worldvolume membrane actions (see for example [35, 29, 40, 13, 51]) in a topological and non-relativistic limit.

The link to tensor models also leads to other quantum mechanical systems which can be viewed as generalizations of the systems discussed here. Our quantum systems have been discussed in terms of state spaces $\mathcal{K}(n)$ for a fixed n . We can generalize, somewhat trivially, to the infinite dimensional state space

$$(150) \quad \mathcal{K}(\infty) = \bigoplus_{n=0} \mathcal{K}(n)$$

where $\mathcal{K}(0)$ is defined to be the one-dimensional vector space \mathbb{C} . We can use a Hamiltonian of the form $\mathcal{H} = \sum_{n=0}^{\infty} \mathcal{H}^{(n)}$, where $\mathcal{H}^{(n)}$ is a Hamiltonian of the form we discussed at fixed n . This Hamiltonian generates time evolutions which mix ribbon graphs with a given number of edges, or Belyi maps with a given degree. Generic Hamiltonians for tensor models, when expressed in terms of $\mathcal{K}(\infty)$ would be expected to generate interactions which mix different values of n . Some recent literature solving quantum mechanical Hamiltonians for tensor models is in [65, 62].

5.3. QUANTUM COMPUTING AND KRONECKER COEFFICIENTS. In this section we describe how quantum mechanical systems on ribbon graphs, hypothetically engineered in the laboratory, can be used to detect the non-vanishing of Kronecker coefficients. It has been shown [52] that the question of deciding the positivity of the Kronecker coefficient for a triple of Young diagrams is NP-hard. The question of whether a quantum computer can outperform classical computers on some chosen task of interest is the problem of quantum supremacy. For recent progress, on specific tasks of generating random number sequences, see [3]. A quantum mechanical system of ribbon graphs can conceivably be engineered directly by identifying physical objects with the properties of ribbon graphs, or perhaps more realistically for the near future, we may consider the idea of quantum simulation where an experimentally controllable quantum system, such as superconducting circuits, is used to simulate another quantum system of interest. A recent review on quantum simulators is [82]. A physical or simulated quantum mechanical system of ribbon graphs would allow, using the connections we have developed here between ribbon graphs and Kronecker coefficients for any specified triple of Young diagrams, the possibility of detecting non-vanishing Kronecker coefficients by observing the time evolution of ribbon graphs. We refer to [10] for further details and scenarios inspecting this question.

An important role is played in our construction of integer matrices with integer null spaces of ribbon graph vectors by the number $k_*(n)$ defined in Section 2.4. To get precise estimates of the computational complexity of our algorithms viewed as a way to calculate Kronecker coefficients at large n , it is desirable to find estimates of the growth of this number with n as n tends to infinity. As discussed in [57, 5] this asymptotic behaviour is also relevant to understanding information loss in toy models of black holes arising in AdS/CFT.

5.4. GENERALIZATIONS. Permutation equivalence classes provide a general approach to the counting of a variety of combinatorial objects of interest in theoretical physics, e.g. Feynman diagrams [28, 27, 43, 19], gauge invariants in matrix and tensor models (see a review in [85]), and frequently these equivalence classes have an algebra structure. The development of quantum mechanical systems where these combinatorial objects become quantum states, their associated permutation algebras are used to express quantum mechanical problems in terms of representation theoretic objects is a promising avenue for further fruitful investigations. We expect the integrality structure of the algebras, when expressed in terms of the geometric basis, to be generic. This will allow a realisation of the representation theoretic quantities in terms of sub-lattices of the lattice generated by the combinatorial objects. An example of such combinatorial algebra studied in detail in [73] is associated with colored necklaces having m beads of one colour and n beads of another color.

Another interesting direction for research is the generalization of the present study to real tensor invariants in particular the $O(N)^3$ invariants [18, 17]. In this case, the ribbon graphs are not bipartite and their counting gives the sum of Kronecker coefficients with Young diagram restricted to even partitions [4, 7].

Although our investigations of $\mathcal{K}(n)$ were motivated by the study of correlators of tensor models in [8, 9] the 3-index tensor variables have not played a direct role in this paper. The space of all gauge invariants constructed from complex tensors $\Phi_{ijk}, \bar{\Phi}^{ijk}$ is isomorphic as a vector space to $\mathcal{K}(\infty)$ defined in 150. On this space the operators $T_k^{(i)}$ we used in this paper should be expressible in terms of differential operators. The map between permutation algebra elements analogous to $T_k^{(i)}$ and differential operators was given in the context of multi-matrix invariants in [61]. The operators $T_k^{(i)}$ are also related to the cut-and-join operators considered in tensor model context in [32, 53]. An interesting problem is to use the connection between differential operators and the reconnection operators $T_k^{(i)}$ to develop Hamiltonian and Lagrangian formulations of the quantum mechanical systems discussed here. The link between such Lagrangian formulations in terms of tensor variables and possible membrane world-volume Lagrangians connecting with the membrane interpretation based on Belyi maps (discussed in Section 5.2) would be illuminating.

APPENDIX

APPENDIX A. RECONNECTION OPERATORS $T_2^{(i)}$ AS MATRICES AT $n = 3$

In this appendix, we work at $n = 3$ and give the construction of the matrices for reconnection operators $T_2^{(i)}$, $i = 1, 2, 3$. In this case there are 11 bipartite ribbon graphs. The matrix elements $(\mathcal{M}_2^{(i)})_r^s$ of the reconnection operators in the geometric ribbon graph basis are 11×11 integer matrices and can be used to determine the Young diagram triples with non-vanishing Kronecker coefficient. Each of these non-vanishing Kronecker coefficients is 1 and we construct the corresponding integer vector in the ribbon graph lattice which generates a one-dimensional sub-lattice.

A.1. $T_2^{(i)}$ MATRICES. Note that we have provided a code to produce all entries $(\mathcal{M}_k^{(i)})_r^s$, see Code2 in appendix C. For $T_2^{(1)}, T_2^{(2)}$ and $T_2^{(3)}$, we have the following

non-negative integer matrices, respectively,

$$(151) \quad \mathcal{M}_2^{(1)} = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 3 & 0 & 0 & 0 & 0 & 0 & 0 & 3 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 3 & 0 & 0 & 0 & 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \end{pmatrix}$$

$$(152) \quad \mathcal{M}_2^{(2)} = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 3 & 0 & 3 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 & 0 & 2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 3 & 0 & 3 & 3 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \end{pmatrix}$$

$$(153) \quad \mathcal{M}_2^{(3)} = \begin{pmatrix} 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 3 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 3 & 0 \\ 0 & 0 & 3 & 0 & 0 & 0 & 0 & 3 & 0 & 0 & 3 \\ 0 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 & 0 & 2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

A.2. NULLSPACE AT $n = 3$. In this section we give, for $n = 3$, the common nullspace of all operators $(T_2^{(i)} - \chi_{R_i}(T_2)/d(R_i))$, for all $i = 1, 2, 3$ at fixed $R_i \vdash n$. This is the rectangular construction of Section 3.2.

The vector v generating the null space for each triple of Young diagram (R_1, R_2, R_3) with non-vanishing Kronecker coefficient is :

	(R_1, R_2, R_3)	v
1	$([1,1,1], [1,1,1], [3])$	$(1, -3, 2, -3, 3, 6, -6, 2, -6, 2, 2)$
2	$([1,1,1], [2,1], [2,1])$	$(-2, 0, 2, 6, 0, 0, -6, -4, 0, 2, 2)$
3	$([1,1,1], [3], [1,1,1])$	$(1, 3, 2, -3, -3, -6, -6, 2, 6, 2, 2)$
4	$([2,1], [1,1,1], [2,1])$	$(-2, 6, -4, 0, 0, 0, 0, 2, -6, 2, 2)$
5	$([2,1], [2,1], [1,1,1])$	$(-2, 0, 2, 0, 6, -6, 0, 2, 0, -4, 2)$
6	$([2,1], [2,1], [2,1])$	$(1, 0, -1, 0, 0, 0, 0, -1, 0, -1, 2)$
7	$([2,1], [2,1], [3])$	$(-2, 0, 2, 0, -6, 6, 0, 2, 0, -4, 2)$
8	$([2,1], [3], [2,1])$	$(-2, -6, -4, 0, 0, 0, 0, 2, 6, 2, 2)$
9	$([3], [1,1,1], [1,1,1])$	$(1, -3, 2, 3, -3, -6, 6, 2, -6, 2, 2)$
10	$([3], [2,1], [2,1])$	$(-2, 0, 2, -6, 0, 0, 6, -4, 0, 2, 2)$
11	$([3], [3], [3])$	$(1, 3, 2, 3, 3, 6, 6, 2, 6, 2, 2)$

One recognizes that the last vector $(1, 3, 2, 3, 3, 6, 6, 2, 6, 2, 2)$ is $\sum_r |\text{Orb}(r)| E_r$, where $|\text{Orb}(r)|$ is the orbit size of the r 'th ribbon graph equivalence class. All $C(R_1, R_2, R_3) = 1$ for the triples (R_1, R_2, R_3) given above. For a each triple, we actually see that the null space is one dimensional at $n = 3$. For $n > 4$, there are $C(R_1, R_2, R_3) > 1$ and therefore the nullspace become of dimension higher than 1 as expected.

APPENDIX B. GEOMETRIC AND FOURIER BASIS

This appendix elaborates on the transformation between the Fourier basis and the geometric basis of $\mathcal{K}(n)$. We check that the Fourier base $\{Q_{\tau, \tau'}^{R_1, R_2, R_3}\}$ expands in terms of ribbon graph base $\{E_r\}$ and vice-versa. We also give a proof of Proposition 3.1.

B.1. CHANGE OF BASIS. Ribbon graph expansion of $Q_{\tau, \tau'}^{R_1, R_2, R_3}$. Start with the base $Q_{\tau, \tau'}^{R_1, R_2, R_3}$ (14) that we re-expand using the orbit decomposition in the same way as in (10) as:

$$\begin{aligned}
 Q_{\tau_1, \tau_2}^{R_1, R_2, R_3} &= \frac{1}{n!} \kappa_{R, S} \sum_r \sum_{a \in \text{Orb}(r)} \sum_{i_1, i_2, i_3, j_1, j_2} C_{i_1, i_2; i_3}^{R_1, R_2; R_3, \tau_1} C_{j_1, j_2; i_3}^{R_1, R_2; R_3, \tau_2} \\
 &\times D_{i_1 j_1}^{R_1}(\sigma_1^{(r)}(a)) D_{i_2 j_2}^{R_2}(\sigma_2^{(r)}(a)) \sigma_1^{(r)}(a) \otimes \sigma_2^{(r)}(a)
 \end{aligned}
 \tag{154}$$

where $(\sigma_1^{(r)}(a), \sigma_2^{(r)}(a))$ is a representative pair in the orbit $\text{Orb}(r)$ that defines the ribbon graph r .

Therefore, we can reorganize the sum and collect for each ribbon graph base element, its coefficient in the above expansion

$$\begin{aligned}
 Q_{\tau_1, \tau_2}^{R_1, R_2, R_3} &= \kappa_{R, S} \sum_r \left[\sum_{i_1, i_2, i_3, j_1, j_2} C_{i_1, i_2; i_3}^{R_1, R_2; R_3, \tau_1} C_{j_1, j_2; i_3}^{R_1, R_2; R_3, \tau_2} D_{i_1 j_1}^{R_1}(\sigma_1^{(r)}) D_{i_2 j_2}^{R_2}(\sigma_2^{(r)}) \right] |\text{Orb}(r)| E_r
 \end{aligned}
 \tag{155}$$

where $D_{i_1 j_1}^{R_1}(\sigma_1^{(r)}(a)) D_{i_2 j_2}^{R_2}(\sigma_2^{(r)}(a))$ has been replaced with $D_{i_1 j_1}^{R_1}(\sigma_1^{(r)}) D_{i_2 j_2}^{R_2}(\sigma_2^{(r)})$ where $(\sigma_1^{(r)}, \sigma_2^{(r)})$ is any representative pair in $\text{Orb}(r)$. This can be done because the coefficient in square brackets is invariant under simultaneous conjugation of σ_1, σ_2 by a permutation γ .

Fourier expansion of E_r . Consider the following expansion of some E_r (9) in terms of the basis $Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}$:

$$(156) \quad E_r = \sum_{R_i, \tau_i} \mathbf{C}_{R_1, R_2, R_3}(\sigma_1^{(r)}, \sigma_2^{(r)}) Q_{\tau_1, \tau_2}^{R_1, R_2, R_3},$$

where $(\sigma_1^{(r)}, \sigma_2^{(r)})$ form a permutation pair in the orbit r , and the coefficient $\mathbf{C}_{R_1, R_2, R_3}(\sigma_1^{(r)}, \sigma_2^{(r)})$ is to be determined. Use the orthogonality relation (17) and evaluate $\mathbf{C}_{R_1, R_2, R_3}(\sigma_1^{(r)}, \sigma_2^{(r)})$:

$$(157) \quad \begin{aligned} \delta_2(E_r, Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}) &= \sum_{R'_i, \tau'_i} \mathbf{C}_{R'_1, R'_2, R'_3}(\sigma_1^{(r)}, \sigma_2^{(r)}) \delta_2(Q_{\tau'_1, \tau'_2}^{R'_1, R'_2, R'_3}, Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}) \\ &= \mathbf{C}_{R_1, R_2, R_3}(\sigma_1^{(r)}, \sigma_2^{(r)}) \kappa_{R_1, R_2} d(R_3). \end{aligned}$$

On the other hand, using (14), we also compute

$$(158) \quad \begin{aligned} \delta_2(E_r, Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}) &= \kappa_{R_1, R_2} \sum_{\gamma_1, \gamma_2 \in S_n} \sum_{i_1, i_2, i_3, j_1, j_2} C_{i_1, i_2; i_3}^{R_1, R_2; R_3, \tau_1} C_{j_1, j_2; i_3}^{R_1, R_2; R_3, \tau_2} \\ &\times D_{i_1 j_1}^{R_1}(\gamma_1) D_{i_2 j_2}^{R_2}(\gamma_2) \delta_2(\sigma_1^{(r)} \otimes \sigma_2^{(r)}, \gamma_1 \otimes \gamma_2) \\ &= \kappa_{R_1, R_2} \sum_{i_1, i_2, i_3, j_1, j_2} C_{i_1, i_2; i_3}^{R_1, R_2; R_3, \tau_1} C_{j_1, j_2; i_3}^{R_1, R_2; R_3, \tau_2} D_{i_1 j_1}^{R_1}(\sigma_1^{(r)}) D_{i_2 j_2}^{R_2}(\sigma_2^{(r)}) \end{aligned}$$

from which we conclude

$$(159) \quad \mathbf{C}_{R_1, R_2, R_3}(\sigma_1^{(r)}, \sigma_2^{(r)}) = \frac{1}{d(R_3)} \sum_{i_l, j_l} D_{i_1 j_1}^{R_1}(\sigma_1^{(r)}) D_{i_2 j_2}^{R_2}(\sigma_2^{(r)}) C_{i_1, i_2; i_3}^{R_1, R_2; R_3, \tau_1} C_{j_1, j_2; i_3}^{R_1, R_2; R_3, \tau_2}.$$

One checks that $\mathbf{C}_{R_1, R_2, R_3}(\sigma_1, \sigma_2)$ is invariant under diagonal conjugation

$$\mathbf{C}_{R_1, R_2, R_3}(\gamma \sigma_1 \gamma^{-1}, \gamma \sigma_2 \gamma^{-1}) = \mathbf{C}_{R_1, R_2, R_3}(\sigma_1, \sigma_2)$$

(this can be shown using the so-called $DDC = CD$ relation and the orthogonality of representation matrices, see appendix A.1 and A.2 of [9]).

An immediate consequence of these formulae is that

$$(160) \quad Q_{\tau_1, \tau_2}^{R_1, R_2, R_3} = \kappa_{R, S} d(R_3) \sum_{\tau} \mathbf{C}_{R_1, R_2, R_3}(\sigma_1^{(r)}, \sigma_2^{(r)}) |\text{Orb}(r)| E_r.$$

Thus the orthogonal Fourier basis elements $Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}$ and expressible in terms of E_r and vice-versa.

B.2. FOURIER BASIS AS EIGENVECTORS OF RECONNECTION OPERATORS $T_k^{(i)}$. To prove Proposition 3.1, we start with some preliminary observations about the group algebra of the symmetric group.

Let $\mathbb{C}(S_n)$ the group algebra of S_n . An inner product on the group algebra is defined by specifying on basis elements $\sigma, \tau \in S_n$, $\delta(\sigma; \tau) = \delta(\sigma \tau^{-1})$. Consider the Fourier basis set $Q_{ij}^R \in \mathbb{C}(S_n)$

$$(161) \quad Q_{ij}^R = \frac{\kappa_R}{n!} \sum_{\sigma \in S_n} D_{ij}^R(\sigma) \sigma, \quad \kappa_R^2 = n! d(R)$$

where $D_{ij}^R(\sigma)$ are matrix elements of σ in an orthonormal basis for the irreducible representation R . κ_R is fixed such that $\delta(Q_{ij}^R; Q_{i'j'}^R) = \delta_{RR'} \delta_{ii'} \delta_{jj'}$ making $\{Q_{ij}^R\}$ an orthonormal basis of $\mathbb{C}(S_n)$. Furthermore, these elements also obey

$$(162) \quad \tau Q_{ij}^R = \sum_l D_{li}^R(\tau) Q_{lj}^R, \quad Q_{ij}^R \tau = \sum_l Q_{il}^R D_{jl}^R(\tau).$$

The following statement holds:

LEMMA B.1.

$$(163) \quad T_k Q_{ij}^R = \frac{\chi_R(T_k)}{d(R)} Q_{ij}^R.$$

Proof. We let act T_k on Q_{ij}^R and using (162) we write:

$$(164) \quad T_k Q_{ij}^R = \sum_{\sigma \in \mathcal{C}_k} \sigma Q_{ij}^R = \sum_l \left(\sum_{\sigma \in \mathcal{C}_k} D_{li}^R(\sigma) \right) Q_{ij}^R.$$

The sum $\sum_{\sigma \in \mathcal{C}_k} D_{li}^R(\sigma)$ may be also written $D_{li}^R(T_k)$. As T_k is central and commute with all elements, $D_{li}^R(T_k) = \alpha \delta_{li}$ a constant diagonal matrix by Schur lemma. We also have $\sum_i D_{ii}^R(T_k) = \chi_R(T_k) = \alpha d(R)$, that yields $\alpha = \chi_R(T_k)/d(R)$. Then back to our previous expression (164)

$$(165) \quad T_k Q_{ij}^R = \sum_l \left(\frac{\chi_R(T_k)}{d(R)} \delta_{li} \right) Q_{ij}^R = \frac{\chi_R(T_k)}{d(R)} Q_{ij}^R$$

which proves the lemma. □

Thus Q_{ij}^R is an eigenvector of T_k with eigenvalue $\chi_R(T_k)/d(R)$. We get back to our main concern, namely the action of $T_k^{(i)}$ on $Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}$.

Proof of Proposition 3.1. We want to prove that $Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}$ define eigenvectors of $T_k^{(i)}$. $Q_{\tau_1, \tau_2}^{R_1, R_2, R_3}$ (14) can be written in terms of the Fourier basis set for $\mathbb{C}(S_n)$ as

$$(166) \quad Q_{\tau_1, \tau_2}^{R_1, R_2, R_3} = \kappa'_{R_1, R_2} \sum_{i_l, j_l} C_{i_1, i_2; i_3}^{R_1, R_2; R_3, \tau_1} C_{j_1, j_2; i_3}^{R_1, R_2; R_3, \tau_2} Q_{i_1 j_1}^{R_1} \otimes Q_{i_2 j_2}^{R_2},$$

where $\kappa'_{R, S}$ is a normalization factor

$$(167) \quad \kappa'_{R, S} = \sqrt{\frac{d(R)d(S)}{n!}}.$$

Then, it becomes obvious that, by Lemma B.1, (61) and (62) hold as $T_k^{(1)}$ and $T_k^{(2)}$ are defined by the actions on T_k of the left or right factors of $\mathbb{C}(S_n) \otimes \mathbb{C}(S_n)$. The third relation requires a bit more work. Use (162) to rewrite:

$$(168) \quad \begin{aligned} T_k^{(3)} Q_{\tau_1, \tau_2}^{R_1, R_2, R_3} &= \kappa'_{R_1, R_2} \sum_{\sigma \in \mathcal{C}_k} \sum_{i_l, j_l} C_{i_1, i_2; i_3}^{R_1, R_2; R_3, \tau_1} C_{j_1, j_2; i_3}^{R_1, R_2; R_3, \tau_2} \sigma Q_{i_1 j_1}^{R_1} \otimes \sigma Q_{i_2 j_2}^{R_2} \\ &= \kappa'_{R_1, R_2} \sum_{\sigma \in \mathcal{C}_k} \sum_{\sigma_1, \sigma_2 \in S_n} \sum_{i_l, j_l} C_{i_1, i_2; i_3}^{R_1, R_2; R_3, \tau_1} C_{j_1, j_2; i_3}^{R_1, R_2; R_3, \tau_2} \\ &\quad \times \sum_{m_1, m_2} D_{m_1 i_1}^{R_1}(\sigma) Q_{m_1 j_1}^{R_1} \otimes D_{m_2 i_2}^{R_2}(\sigma) Q_{m_2 j_2}^{R_2}. \end{aligned}$$

We use the identity

$$(169) \quad \sum_{j_1, j_2} D_{i_1 j_1}^{R_1}(\gamma) D_{i_2 j_2}^{R_2}(\gamma) C_{j_1, j_2; j_3}^{R_1, R_2; R_3, \tau} = \sum_{i_3} C_{i_1, i_2; i_3}^{R_1, R_2; R_3, \tau} D_{i_3 j_3}^{R_3}(\gamma)$$

which holds for any $\gamma \in S_n$ and expresses the fact that the Clebsch–Gordan coefficients intertwine the action of γ in $R_1 \otimes R_2$ with the action in R_3 . We re-express the above as

$$\begin{aligned} T_k^{(3)} Q_{\tau_1, \tau_2}^{R_1, R_2, R_3} &= \kappa'_{R_1, R_2} \sum_{\sigma \in \mathcal{C}_k} \sum_{\sigma_1, \sigma_2 \in S_n} \sum_{m_1, i_3, j_1} \left(\sum_{i_1, i_2} D_{m_1 i_1}^{R_1}(\sigma) D_{m_2 i_2}^{R_2}(\sigma) C_{i_1, i_2; i_3}^{R_1, R_2; R_3, \tau_1} \right) \\ &\quad \times C_{j_1, j_2; i_3}^{R_1, R_2; R_3, \tau_2} Q_{m_1 j_1}^{R_1} \otimes Q_{m_2 j_2}^{R_2} \end{aligned}$$

$$\begin{aligned}
 &= \kappa'_{R_1, R_2} \sum_{\sigma \in \mathcal{C}_k} \sum_{\sigma_1, \sigma_2 \in S_n} \sum_{m_1, i_3, j_1} \left(\sum_{m_3} C_{m_1, m_2; m_3}^{R_1, R_2; R_3, \tau_1} D_{m_3 i_3}^{R_3}(\sigma) \right) \\
 &\times C_{j_1, j_2; i_3}^{R_1, R_2; R_3, \tau_2} Q_{m_1 j_1}^{R_1} \otimes Q_{m_2 j_2}^{R_2} \\
 &= \kappa'_{R_1, R_2} \sum_{m_1, i_3, j_1} \left(\sum_{m_3} C_{m_1, m_2; m_3}^{R_1, R_2; R_3, \tau_1} D_{m_3 i_3}^{R_3}(T_k) \right) \\
 &\times C_{j_1, j_2; i_3}^{R_1, R_2; R_3, \tau_2} Q_{m_1 j_1}^{R_1} \otimes Q_{m_2 j_2}^{R_2} \\
 &= \kappa'_{R_1, R_2} \sum_{\sigma_1, \sigma_2 \in S_n} \sum_{m_1, i_3, j_1} \left(\sum_{m_3} C_{m_1, m_2; m_3}^{R_1, R_2; R_3, \tau_1} \frac{\chi_{R_2}(T_k)}{d(R_3)} \delta_{m_3 i_3} \right) \\
 &\times C_{j_1, j_2; i_3}^{R_1, R_2; R_3, \tau_2} Q_{m_1 j_1}^{R_1} \otimes Q_{m_2 j_2}^{R_2} \\
 (170) \quad &= \frac{\chi_{R_3}(T_k)}{d(R_3)} \left[\kappa'_{R_1, R_2} \sum_{m_1, i_3, j_1} C_{m_1, m_2; i_3}^{R_1, R_2; R_3, \tau_1} C_{j_1, j_2; i_3}^{R_1, R_2; R_3, \tau_2} Q_{m_1 j_1}^{R_1} \otimes Q_{m_2 j_2}^{R_2} \right]
 \end{aligned}$$

where, at an intermediate step, we use again $D_{m_3 i_3}^{R_3}(T_k) = \alpha \delta_{m_3 i_3}$ (with α worked out in Lemma 164). This ends the proof the proposition. □

APPENDIX C. GAP CODES

In this appendix, we give several GAP functions that lead to the calculation of common nullspace of the operators T_2 and T_3 at rank 3 and for arbitrary n . This code is computed with Sage calling the GAP package. Hence the `%gap` appearing at the beginning of each function. Such command could be replaced by a single `%gap` depending on the environment. The comments, or lines starting by `#`, inside the code are self-explanatory and help to understand of each part of the current function.

For $n \leq 14$, $kmax \leq 3$, here is the overall strategy of the calculation:

- (1) generate the set of ribbon graphs (denoted as $\text{Rib}(n)$ in the bulk of the paper) as list with `RibbSetFunction(n)`; their number is ℓ (this is $|\text{Rib}(n)|$ in the bulk of paper);
- (2) construct T_2 and T_3 , and their different action $T_2^{(i)}$ and $T_3^{(i)}$, $i = 1, 2, 3$ acting on different slots of pairs of permutation representing ribbon graphs ;
- (3) calculate the number of time that a ribbon graph b appears in the expansion of $T_2^{(i)} a$ or $T_3^{(i)} a$, for all ribbon graph a ;
- (4) generate the $\ell \times \ell$ -matrix $(L_l^{(i)})_{ab}$, $l = 2, 3$, $i = 1, 2, 3$, of all $T_2^{(i)}$ and $T_3^{(i)}$;
- (5) introduce the list of normalized characters $\widehat{\chi}_R(T_p) := \chi_R(T_p)/d(R)$ using the formulae from [67] for $p := 2, 3$;
- (6) then solve the nullspace of the transpose of the stack of matrices $M_\ell^{(i)}(R) = L_\ell^{(i)} - \widehat{\chi}_R(T_p) * Id_\ell$, Id_ℓ being the $\ell \times \ell$ -identity matrix. The dimension of this space is $C(R, S, T)^2$.
- (7) Alternatively, we generate the sequence of prime numbers $a_{i,k}$ (97) used in the construction of the Hamiltonian $\mathcal{H} := \sum_{k=2}^{kmax} \sum_{i=1}^3 a_{i,k} T_k^{(i)}$ as a matrix sum; and solve the nullspace of the sum of matrices

$$(171) \quad M_2^{(1)}(R) + a_1 M_3^{(1)}(R) + a_2 (M_2^{(2)}(S) + a_1 M_3^{(2)}(S)) + a_3 (M_2^{(3)}(T) + a_1 M_3^{(3)}(T)).$$

We compute the stack of matrices $M_\ell^{(i)}(R)$ in Gap and solve for its null space. There is a corresponding equation for \mathcal{H} that we also put in comments.

Code1: Generating all ribbons with n -edges. The function `RibbSetFunction(n)` returns the list of bipartite ribbon graphs made n ribbon edges and at most n black vertices, and at most n white edges. We use the PCA formulation where each ribbon

graph is represented by its equivalent class, namely an orbit under diagonal S_n group action on a pair of S_n permutations (σ_1, σ_2) .

```

%%gap
#-----
# Function returning an ordered list of ribbon graphs - each ribbon
# graph represented as a set of permutations within
# an  $S_n$  orbit by diagonal conjugation of pairs of permutations

RibbSetFunction := function( n )
  local G, Pairs, Ribb, RibbSets, a, b;
  G := SymmetricGroup(n);
  Pairs := [];

  for a in G do
    for b in G do
      Add (Pairs, [a, b]);
    od;
  od;

  ## In Gap, group G action on list and groups is always by conjugation
  # Ribb list of ribbons as G orbits on pairs (tau_1, tau_2)
  # OnPairs option of function Orbit means G acts diagonally on pairs
  # (g tau_1 g^(-1), g tau_2 g^(-1))

  Ribb := Orbits (G, Pairs, OnPairs);

  # RibbSets is now list of sets of pairs within an orbit, the order within
  # the set does not matter
  RibbSets := [] ;
  for a in [ 1 .. Length ( Ribb ) ] do
    Add ( RibbSets , Set ( Ribb[a] ) ) ;
  od ;
  return RibbSets;
end;

```

Code2: Constructing T -operators. We construct the operators $T_l^{(i)}$.

```

%%gap
#-----
# Given a number n, and kmax
# returns the (kmax -1)x3 arrays  $T(i)_{\{a, b\}}$  - of size
# |RibSetLoc| * |RibSetLoc|.
# at fixed i and p fixed this is the matrix of the action of  $T^{(p)}(i)$  ;
# |RibSetLoc| is the size of the set of ribbons generated
# by RibbSetFunction(n);
# for any a, a ribbon graph orbit RibbSetLoc[a] is represented
# by RibSetLoc[a][1].
# Depending on i, we operate on the first permutation, the second or both
# in RibSetLoc[a][1] and return b, namely the position in RibSetLoc
# of the outcome; RibSetLoc[a][1] is the representative perm of
# the ribbon RibbSetLoc[a];
# RibSetLoc[a][1][1] is first projection of the RibbSetLoc[a][1];
# RibSetLoc[a][1][2] is second projection of the RibbSetLoc[a][1];
# Position ( list , obj ) returns the position of the first occurrence
# of obj in list.
# Positions ( list , obj ) returns the number of occurrences of obj in list;
#-----

```

```

ArrayTi := function (n, kmax)

  local G, a, b, i, j, l, s, Cic, cc, RibSetLoc,
    L, L1, L2, L3, pos1, pos2, pos3 ;

  G := SymmetricGroup(n);
  L := [];
  Cic := [];

  # Construction of ribbon graphs
  RibSetLoc := RibbSetFunction(n);
  l := Length ( RibSetLoc );

  for i in [ 2 .. kmax] do
    # Construction of Ti
    Add ( Cic, Orbit ( G, CycleFromList([1 .. i]), OnPoints) );

    # Construction of the k-1 lists of listsLpt;
    # Lpt[i-1] is a list of 3 matrices Lpt[i-1][p] initialized at 0;
    # Each Lpt[i-1][p] defines the operator Ti^p with action of
    # Ti in the slot p
    Add(L, []);
    for j in [1 .. 3] do
      Add( L[i-1], NullMat ( l , l ) );
    od;
  od;

  for a in [ 1 .. l ] do
    for i in [2 .. kmax] do
      # empty the Li
      L1 := [];
      L2 := [];
      L3 := [];
      cc := Cic[i-1];
      for s in cc do
        pos1 := Position (RibSetLoc, Set(Orbit(G, [s*RibSetLoc[a][1][1],
          RibSetLoc[a][1][2]], OnPairs))) ;
        pos2 := Position (RibSetLoc, Set(Orbit(G, [RibSetLoc[a][1][1] ,
          s*RibSetLoc[a][1][2]], OnPairs))) ;
        pos3 := Position (RibSetLoc, Set(Orbit(G, [s*RibSetLoc[a][1][1],
          s*RibSetLoc[a][1][2]], OnPairs))) ;
        Add( L1 , pos1);
        Add( L2 , pos2);
        Add( L3 , pos3);
      od;
      for b in [ 1 .. l ] do
        L[i-1][1][ b, a ] := Length ( Positions ( L1 , b ) ) ;
        L[i-1][2][ b, a ] := Length ( Positions ( L2 , b ) ) ;
        L[i-1][3][ b, a ] := Length ( Positions ( L3 , b ) ) ;
      od;
    od;
  od;
  return L ;
end ;

```

Code3: Characters. The following function returns the table list of characters according to Lassalle formulae [67].

```

%%gap
# Returns the list of characters of T2 (on the left) and T3 (on the right)
# via Lassalle formulae. These characters are the eigenvalues of T's.

CharactersEigenvalues_of_Top := function( m )
  local i, j, k, p, L, L2, L3, sum2, sum3;
  p := Partitions( m ) ;
  sum2 := 0;
  sum3 := 0;
  L2 := [];
  L3 := [];
  L := [];
  for i in [ 1..Length(p) ] do
    sum2 := 0;
    sum3 := 0;
    for j in [1..Length(p[i])] do
      for k in [1..p[i][j]] do
        sum2 := sum2 - j + k ;
        sum3 := sum3 +(k-j)*(k-j);
      od;
    od;
    sum3:= sum3 - Factorial( m )/( Factorial( m-2 ) *2 ) ;
    Add( L2, sum2 );
    Add( L3, sum3 );
    Add( L, [ L2[i], L3[i] ] );
  od;
  return L;
end;

```

Code4: Primes. The two functions returns either a couple (a_1, a_2) or a triple (a_1, a_2, a_3) of prime numbers are used in the construction of the total Hamiltonian and insure that it cannot vanishes outside of the required values. This follows the sufficient conditions explained in Section 3.3.

```

%%gap
# The code produces a couple that makes the QM Hamiltonian not vanishing
# unless the triples (R1,R2,R3) = (R1',R2',R3')

CouplePrime := function(m)
  local p1, L;
  p1 := NextPrimeInt( m*(m-1) );
  L := [];
  Add(L, p1);
  Add(L, NextPrimeInt( (p1+1)*m*(m-1)));
  return L ;
end;

# The code produces a tripe that makes the QM Hamiltonian not vanishing
# unless the triples (R1,R2,R3) = (R1',R2',R3')

CouplePrime2 := function(n)
  local p1, p2, p3, L;
  p1 := NextPrimeInt( n*(n-1) );
  p2 := NextPrimeInt( Int( (n/3)*(n-1)*(3 + 2*p1*(n-2))) ) ;
  p3 := NextPrimeInt( Int( (n/3)*(n-1)*(1+p2)*( 3 + 2*p1*(n-2) ) ) ) ;

```

```

L := [];
Add(L, p1);
Add(L, p2 );
Add(L, p3);
return L ;
end;

```

Code5: Matrices and null spaces. We are now in position to address the nullspace of the multiple actions of T_2 and T_3 . The following code constructs the stack of matrices (note that, in comments, we also give instructions to construct the total Hamiltonian calling prime numbers) made of 3 (resp. 6) matrices determined by three Young Diagram R, S, T , given $kmax = 2$ (resp. $kmax = 3$). After the construct, it returns the null space of the resulting matrix.

```

%%gap
MatForNullVectors := function(m, kmax, R, S,T)
    local a, l, chi , M1, M2, M3, M5, M4, M6, Arr, Id;

    # for the total Hamiltonian version uncomment the following
    #local a, l, chi , M1, M2, M3, M5, M4, M6, Arr, Id, c, c2, Ham;

    l:=Length(RibbSetFunction(m)); # cardinality of the ribbon set
    chi := CharactersEigenvalues_of_Top(m);

    # for the Hamiltonian uncomment the following
    #c := CouplePrime(m);
    #c2 := CouplePrime2(m);

    #Constructing the matrices
    Arr := ArrayTi(m , kmax );

    if kmax > 3 then
        Print ("kmax >3");
        return 0;
    fi;

    if kmax = 2 then
        M1 := Arr[1][1] - chi[R][1] * IdentityMat ( l );
        M3 := Arr[1][2] - chi[S][1] * IdentityMat ( l );
        Append(M1, M3);

        # For the Hamiltonian uncomment the following
        # Ham := M1 + c[1]*M3 ;

        M5 := Arr[1][3] - chi[T][1] * IdentityMat ( l );
        Append(M1, M5);

        # For the Hamiltonian uncomment the following
        #Ham := Ham + c[2]*M5 ;
    fi;

    if kmax = 3 then
        M1 := Arr[1][1] - chi[R][1] * IdentityMat ( l );
        M2 := Arr[2][1] - chi[R][2] * IdentityMat ( l );
        Append(M1, M2);
    fi;
end;

```

```

# For the Hamiltonian uncomment the following
# Ham := M1 + c2[1]*M2 ;

M3 := Arr[1][2] - chi[S][1] * IdentityMat ( 1 );
Append(M1, M3);
M4 := Arr[2][2] - chi[S][2] * IdentityMat ( 1 );
Append(M1, M4);

# For the Hamiltonian uncomment the following
# Ham := Ham + c2[2]*(M3 + c2[1]*M4) ;

M5 := Arr[1][3] - chi[T][1] * IdentityMat ( 1 );
Append(M1, M5);
M6 := Arr[2][3] - chi[T][2] * IdentityMat ( 1 );
Append(M1, M6);

# For the Hamiltonian uncomment the following
# Ham := Ham + c2[3]*(M5 + c2[1]*M6) ;
fi;

return NullspaceIntMat(TransposedMat (M1) );
# For the total Hamiltonian uncomment the following
#return NullspaceMat(TransposedMat ( Ham) ) ;
end;

```

REFERENCES

- [1] Jan Ambjorn, Bergfinnur Durhuus, and Thordur Jonsson, *Three-dimensional simplicial quantum gravity and generalized matrix models*, Mod. Phys. Lett. A **6** (1991), 1133–1146.
- [2] N. Amburg, H. Itoyama, Andrei Mironov, Alexei Morozov, D. Vasiliev, and R. Yoshioka, *Correspondence between feynman diagrams and operators in quantum field theory that emerges from tensor model*, Eur. Phys. J. C **80** (2020), no. 5, 471.
- [3] Frank Arute et al., *Quantum supremacy using a programmable superconducting processor*, Nature **574** (2019), no. 7779, 505–510.
- [4] Remi C. Avohou, Joseph Ben Geloun, and Nicolas Dub, *On the counting of $o(n)$ tensor invariants*, Adv. Theor. Math. Phys. **24** (2020), no. 4, 821–878.
- [5] Vijay Balasubramanian, Bartłomiej Czech, Klaus Larjo, and Joan Simon, *Integrability versus information loss: A simple example*, J. High Energy Phys. **11** (2006), 001.
- [6] Gennadii Vladimirovich Belyi, *On galois extensions of a maximal cyclotomic field*, Mathematics of The USSR-Izvestiya **14** (1980), 247–256.
- [7] Joseph Ben Geloun, *On the counting tensor model observables as $u(n)$ and $o(n)$ classical invariants*, PoS **CORFU2019** (2020), 175.
- [8] Joseph Ben Geloun and Sanjaye Ramgoolam, *Counting tensor model observables and branched covers of the 2-sphere*, Ann. Inst. H. Poincaré D Comb. Phys. Interact. **1** (2014), no. 1, 77–138.
- [9] ———, *Tensor models, kronecker coefficients and permutation centralizer algebras*, J. High Energy Phys. **11** (2017), 092.
- [10] ———, *Quantum mechanics of bipartite ribbon graphs: Integrality, lattices and kronecker coefficients*, (2020), <https://arxiv.org/abs/2010.04054>.
- [11] Rajsekhar Bhattacharyya, Storm Collins, and Robert de Mello Koch, *Exact multi-matrix correlators*, J. High Energy Phys. **03** (2008), 044.
- [12] Rajsekhar Bhattacharyya, Robert de Mello Koch, and Michael Stephanou, *Exact multi-restricted schur polynomial correlators*, J. High Energy Phys. **06** (2008), 101.
- [13] Giulio Bonelli, Alessandro Tanzini, and Maxim Zabzine, *On topological m -theory*, Adv. Theor. Math. Phys. **10** (2006), no. 2, 239–260.
- [14] Thomas William Brown, P. J. Heslop, and S. Ramgoolam, *Diagonal free field matrix correlators, global symmetries and giant gravitons*, J. High Energy Phys. **04** (2009), 089.

- [15] Peter Burgisser, Matthias Christandl, and Christian Ikenmeyer, *Nonvanishing of Kronecker coefficients for rectangular shapes*, Adv. Math. **227** (2011), 2082–2091.
- [16] Peter J. Cameron, *Combinatorics: Topics, techniques, algorithms*, Cambridge University Press, 1994 (reprinted 1996).
- [17] Sylvain Carrozza, *Large n limit of irreducible tensor models: $o(n)$ rank-3 tensors with mixed permutation symmetry*, J. High Energy Phys. **06** (2018), 039.
- [18] Sylvain Carrozza and Adrian Tanasa, *$o(n)$ random tensor models*, Lett. Math. Phys. **106** (2016), no. 11, 1531–1559.
- [19] Erick Castro and Itzhak Roditi, *A combinatorial matrix approach for the generation of vacuum feynman graphs multiplicities in ϕ^4 theory*, J. Phys. A **51** (2018), no. 39, 395202.
- [20] Henri Cohen, *A course in computational algebraic number theory*, Graduate Texts in Mathematics, Springer Science & Business Media, 2000.
- [21] Stefan Cordes, Gregory W. Moore, and Sanjaye Ramgoolam, *Lectures on 2-d yang-mills theory, equivariant cohomology and topological field theories*, Nucl. Phys. B Proc. Suppl. **41** (1995), 184–244.
- [22] Steve Corley, Antal Jevicki, and Sanjaye Ramgoolam, *Exact correlators of giant gravitons from dual $n=4$ sym theory*, Adv. Theor. Math. Phys. **5** (2002), 809–839.
- [23] Sylvie Corteel, Alain Goupil, and Gilles Schaeffer, *Content evaluation and class symmetric functions*, Adv. Math. **188** (2004), 315–336.
- [24] Robert De Mello Koch, David Gossman, Nirina Hasina Tahiridimbisoa, and Augustine Larweh Mahu, *Holography for tensor models*, Phys. Rev. D **101** (2020), no. 4, 046004.
- [25] Robert de Mello Koch, David Gossman, and Laila Tribelhorn, *Gauge invariants, correlators and holography in bosonic and fermionic tensor models*, J. High Energy Phys. **09** (2017), 011.
- [26] Robert de Mello Koch and Sanjaye Ramgoolam, *From matrix models and quantum fields to hurwitz space and the absolute galois group*, (2010), <https://arxiv.org/abs/hep-th/1002.1634>.
- [27] ———, *A double coset ansatz for integrability in ads/cft*, J. High Energy Phys. **06** (2012), 083.
- [28] ———, *Strings from Feynman Graph counting : without large n* , Phys. Rev. D **85** (2012), 026007.
- [29] B. de Wit, J. Hoppe, and H. Nicolai, *On the quantum mechanics of supermembranes*, Nucl. Phys. B **305** (1988), 545.
- [30] Nicolas Delporte and Vincent Rivasseau, *The tensor track v: Holographic tensors*, 17th Hellenic School and Workshops on Elementary Particle Physics and Gravity, 4 2018.
- [31] Pablo Diaz, *Tensor and matrix models: a one-night stand or a lifetime romance?*, J. High Energy Phys. **06** (2018), 140.
- [32] ———, *Backgrounds from tensor models: A proposal*, Phys. Rev. D **103** (2021), no. 6, 066010.
- [33] Pablo Diaz and Soo-Jong Rey, *Invariant operators, orthogonal bases and correlators in general tensor models*, Nucl. Phys. B **932** (2018), 254–277.
- [34] ———, *Orthogonal bases of invariants in tensor models*, J. High Energy Phys. **02** (2018), 089.
- [35] Michael J. Duff, *Classical and quantum supermembranes*, Class. Quant. Grav. **6** (1989), 1577–1598.
- [36] Dennis R. Estes, *Eigenvalues of symmetric integer matrices*, J. Number Theory **42** (1992), no. 3, 292–296.
- [37] Valentin Féray, *Proof of Stanley’s conjecture about irreducible character values of the symmetric group*, Ann. Comb. **13** (2010), 453–461.
- [38] William Fulton, *Young tableaux*, London Mathematical Society Student Texts, vol. 35, Cambridge University Press, 1997.
- [39] The GAP Group GAP4, *Gap – groups, algorithms, and programming, version 4.11.0*, Lattices Algorithms and Applications (2022), <https://www.gap-system.org/>.
- [40] Joaquim Gomis, Kiyoshi Kamimura, and Paul K. Townsend, *Non-relativistic superbranes*, J. High Energy Phys. **11** (2004), 051.
- [41] R. Goodman and N.R. Wallach, *Symmetry, representations, and invariants*, Graduate Texts in Mathematics, Springer New York, 2009.
- [42] Rajesh Gopakumar, *What is the simplest gauge-string duality?*, (2011), <https://arxiv.org/abs/1104.2386>.
- [43] K. Krishna Gopala, Patrick Labelle, and Vasilisa Shramchenko, *Enumeration of n -rooted maps using quantum field theory*, Nucl. Phys. B **936** (2018), 668–689.
- [44] David J. Gross and Washington Taylor, *Two-dimensional qcd is a string theory*, Nucl. Phys. B **400** (1993), 181–208.
- [45] Alexandre Grothendieck, *Esquisse d’un programme*, 1984.
- [46] S. S. Gubser, Igor R. Klebanov, and Alexander M. Polyakov, *Gauge theory correlators from noncritical string theory*, Phys. Lett. B **428** (1998), 105–114.

- [47] Razvan Gurau, *The complete $1/n$ expansion of colored tensor models in arbitrary dimension*, Ann. Henri Poincaré **13** (2012), 399–423.
- [48] ———, *Random tensors*, Oxford University Press, 2017.
- [49] Morton Hamermesh, *Group theory and its application to physical problems*, Dover Books on Physics, Dover, 2003.
- [50] Petr Horava, *Topological rigid string theory and two-dimensional qcd*, Nucl. Phys. B **463** (1996), 238–286.
- [51] ———, *Membranes at quantum criticality*, J. High Energy Phys. **03** (2009), 020.
- [52] Christian Ikenmeyer, Ketan Mulmuley, and Michael Walter, *On vanishing of Kronecker coefficients*, Comput. Complexity **26** (2017), 949–992.
- [53] H. Itoyama, A. Mironov, and A. Morozov, *Cut and join operator ring in tensor models*, Nucl. Phys. B **932** (2018), 52–118.
- [54] ———, *Tensorial generalization of characters*, J. High Energy Phys. **12** (2019), 127.
- [55] Hiroshi Itoyama and Reiji Yoshioka, *Generalized cut operation associated with higher order variation in tensor models*, Nucl. Phys. B **945** (2019), 114681.
- [56] Gareth Aneurin Jones, *Regular embeddings of complete bipartite graphs: classification and enumeration*, Proc. Lond. Math. Soc. **101** (2010), 427–453.
- [57] Garreth Kemp and Sanjaye Ramgoolam, *Bps states, conserved charges and centres of symmetric group algebras*, J. High Energy Phys. **01** (2020), 146.
- [58] Yusuke Kimura, *Multi-matrix models and noncommutative frobenius algebras obtained from symmetric groups and brauer algebras*, Commun. Math. Phys. **337** (2015), no. 1, 1–40.
- [59] ———, *Noncommutative frobenius algebras and open-closed duality*, (2017), <https://arxiv.org/abs/1701.08382>.
- [60] Yusuke Kimura and Sanjaye Ramgoolam, *Branes, anti-branes and brauer algebras in gauge-gravity duality*, J. High Energy Phys. **11** (2007), 078.
- [61] ———, *Enhanced symmetries of gauge theory and resolving the spectrum of local operators*, Phys. Rev. D **78** (2008), 126003.
- [62] Igor R. Klebanov, Alexey Milekhin, Fedor Popov, and Grigory Tarnopolsky, *Spectra of eigenstates in fermionic tensor quantum mechanics*, Phys. Rev. D **97** (2018), no. 10, 106023.
- [63] Igor R. Klebanov, Fedor Popov, and Grigory Tarnopolsky, *Tasi lectures on large n tensor models*, PoS **TASI2017** (2018), 004.
- [64] Vaclav Kotousevec, *Oeis foundation inc. entry a000108 in the on-line encyclopedia of integer sequences*, <https://oeis.org/A279819>.
- [65] Chethan Krishnan and K. V. Pavan Kumar, *Exact solution of a strongly coupled gauge theory in $0+1$ dimensions*, Phys. Rev. Lett. **120** (2018), no. 20, 201603.
- [66] S.K. Lando, R.V. Gamkrelidze, V.A. Vassiliev, and A.K. Zvonkin, *Graphs on surfaces and their applications*, Encyclopaedia of Mathematical Sciences, Low dimensional topology II, Springer Berlin Heidelberg, 2013.
- [67] Michel Lassalle, *An explicit formula for the characters of the symmetric group*, Math. Ann. **340** (2007), 383–405.
- [68] Arjen K. Lenstra, Hendrik W. Lenstra, and László Miklós Lovász, *Factoring polynomials with rational coefficients*, Math. Ann. **261** (1982), 515–534.
- [69] Hai Lin, Oleg Lunin, and Juan Martin Maldacena, *Bubbling ads space and $1/2$ bps geometries*, J. High Energy Phys. **10** (2004), 025.
- [70] Dino Lorenzini, *Smith normal form and laplacians*, J. Combin. Theory Ser. B **98** (2008), 1271–1300.
- [71] Juan Martin Maldacena, *The large n limit of superconformal field theories and supergravity*.
- [72] Laurent Manivel, *On the asymptotics of Kronecker coefficients*, J. Algebraic Combin. **42** (2014), 999–1025.
- [73] Paolo Mattioli and Sanjaye Ramgoolam, *Permutation centralizer algebras and multi-matrix invariants*, Phys. Rev. D **93** (2016), no. 6, 065040.
- [74] Daniele Micciancio, *Basic algorithms*, Lattices Algorithms and Applications.
- [75] Ketan Mulmuley and Milind A. Sohoni, *Geometric Complexity Theory I: An Approach to the P vs. NP and Related Problems*, SIAM J. Comput. **31** (2001), 496–526.
- [76] Francis D. Murnaghan, *On the representations of the symmetric group*, Amer. J. Math. **59** (1937), 437–488.
- [77] ———, *The analysis of the Kronecker product of irreducible representations of the symmetric group*, Amer. J. Math. **60** (1938), 761–784.
- [78] Tadasi Nakayama, *On some modular properties of irreducible representations of a symmetric group, I*, Jpn. J. Math. **18** (1941), 89–108.

- [79] Igor Pak and Greta Panova, *On the complexity of computing Kronecker coefficients*, *Comput. Complexity* **26** (2017), 1–36.
- [80] Igor Pak, Greta Panova, and Ernesto Vallejo, *Combinatorics and complexity of Kronecker coefficients*, *Workshop Summary* (2015), <https://simons.berkeley.edu/sites/default/files/docs/1899/slidespanova.pdf>.
- [81] Igor Pak, Greta Panova, and Damir Yeliussizov, *On the largest Kronecker and Littlewood-Richardson coefficients*, *J. Combin. Theory Ser. A* **165** (2019), 44–77.
- [82] G. S. Paroanu, *Recent progress in quantum simulation using superconducting circuits*, *J. Low Temp. Phys.* **175** (2014), 633.
- [83] Jurgis Pasukonis and Sanjaye Ramgoolam, *Quivers as calculators: Counting, correlators and riemann surfaces*, *J. High Energy Phys.* **04** (2013), 094.
- [84] Arun Ram, *Dissertation, chapter 1: Representation theory*, 2004, <http://math.soimeme.org/~arunram/Preprints/dissertationChapt1.pdf>.
- [85] Sanjaye Ramgoolam, *Permutations and the combinatorics of gauge invariants for general n* , *PoS CORFU2015* (2016), 107.
- [86] Vincent Rivasseau, *The tensor track, iv*, *PoS CORFU2015* (2016), 106.
- [87] Leila Schneps, *The Grothendieck Theory of Dessins D'Enfants*, *Lecture note series*, Cambridge University Press, 1994.
- [88] Alexander Schrijver, *Theory of linear and integer programming*, *Wiley Series in Discrete Mathematics & Optimization*, Wiley, 1998.
- [89] Barry Simon, *Representations of finite and compact groups*, American Mathematical Society, 1991.
- [90] Richard P. Stanley, *Positivity problems and conjectures*, *Mathematics: frontiers and perspectives*, American Mathematical Society, Providence, RI, (2000), 295–319.
- [91] ———, *A conjectured combinatorial interpretation of the normalized irreducible character values of the symmetric group*, *arXiv: Combinatorics* (2006), <https://arxiv.org/abs/math/0606467>.
- [92] Richard P. Stanley and Sergey Fomin, *Enumerative combinatorics*, *Cambridge Studies in Advanced Mathematics*, vol. 2, Cambridge University Press, 1999.
- [93] Wikipedia, *On Murnaghan-Nakayama rule*, https://en.wikipedia.org/wiki/Murnaghan%E2%80%93Nakayama_rule.
- [94] Edward Witten, *Anti-de sitter space and holography*, *Adv. Theor. Math. Phys.* **2** (1998), 253–291.
- [95] ———, *An syk-like model without disorder*, *J. Phys. A* **52** (2019), no. 47, 474002.

JOSEPH BEN GELOUN, Laboratoire d'Informatique de Paris Nord UMR CNRS 7030, Université Sorbonne Paris Nord, 99, avenue J.-B. Clement, 93430 Villetaneuse, France. International Chair in Mathematical Physics and Applications, ICMPA–UNESCO Chair, 072 B.P. 50 Cotonou, Benin.
E-mail : bengeloun@lipn.univ-paris13.fr

SANJAYE RAMGOOLAM, School of Physics and Astronomy, Centre for Research in String Theory, Queen Mary University of London, London E1 4NS, United Kingdom. School of Physics and Mandelstam Institute for Theoretical Physics, University of Witwatersrand, Wits, 2050, South Africa.
E-mail : s.ramgoolam@qmul.ac.uk