



Article

Fédération Equestre Internationale eventing: Fence-level risk factors for falls during the cross-country phase (2008–2018)

Bennet, Euan D, Cameron-Whytock, Heather and Parkin, Tim

Available at <https://clock.uclan.ac.uk/47719/>

Bennet, Euan D, Cameron-Whytock, Heather orcid iconORCID: 0000-0003-0760-2584 and Parkin, Tim (2023) Fédération Equestre Internationale eventing: Fence-level risk factors for falls during the cross-country phase (2008–2018). Equine Veterinary Journal, 55 (3). ISSN 0425-1644

It is advisable to refer to the publisher's version if you intend to cite from the work.
doi10.1111/evj.13863

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the [policies](#) page.

Fédération Equestre Internationale eventing: Fence-level risk factors for falls during the cross-country phase (2008–2018)

Euan David Bennet¹  | Heather Cameron-Whytock²  | Tim D. H. Parkin¹ 

¹Bristol Veterinary School, University of Bristol, Langford Campus, Bristol, UK

²School of Animal Rural and Environmental Science, Nottingham Trent University, Brackenhurst Campus, Southwell, UK

Correspondence

Euan David Bennet, School of Biodiversity, One Health, and Veterinary Medicine, University of Glasgow, Garscube Campus, Glasgow, G61 1QH, UK.
Email: euane.bennet@glasgow.ac.uk

Funding information

Federation Equestre Internationale

Abstract

Background: The equestrian discipline of eventing tests athletes' and horses' skill over three phases: dressage, jumping and cross-country. Falls during cross-country can be particularly serious and result in serious or fatal injury for both horse and athlete. Cross-country course and fence design are crucial contributory factors to safety.

Objectives: To provide descriptive statistics and identify fence-level risk factors for horses competing in Fédération Equestre Internationale (FEI) events worldwide.

Study Design: Retrospective cohort study.

Methods: Data were collected for every horse start worldwide in all international (CI), championship (CH), Olympics (OG) and World Equestrian Games (WEG) eventing competitions between January 2008 and December 2018 and univariable logistic regression, followed by multivariable logistic regression were applied. The final model was built in a stepwise bi-directional process, with each step assessed by the Akaike information criterion.

Results: Risk factors were identified at the fence level covering aspects of fence design and course design. Ten fence types were at increased odds of a fall occurring compared with square spread fences, and seven types were at reduced odds. Fences with an approach downhill (odds ratio [OR] 1.35, 95% confidence interval [CI] 1.19–1.52), with landing into water (OR 1.82, CI 1.62–2.01), frangible devices (OR 1.28, CI 1.15–1.41) and later elements of combined obstacles (OR 1.33 CI 1.25–1.42 for the second element, OR 1.21 CI 1.10–1.32 for later elements) were associated with increased risk of falls occurring.

Main Limitations: Although the dataset covers every international competition worldwide, it does not include national-level competitions.

Conclusions: It is recommended that the most challenging fences are placed near the beginning of the course, and not in downhill or water settings. The complexity of individual elements in combined fences should be reduced. Adopting evidence-based course design is a crucial intervention for reducing the incidence of horse falls and associated serious and fatal injuries to horse and human athletes.

KEYWORDS

eventing, falls, horse, safety

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Equine Veterinary Journal* published by John Wiley & Sons Ltd on behalf of EVJ Ltd.

1 | INTRODUCTION

The Fédération Equestre Internationale (FEI) equestrian discipline of eventing, also known as three-day eventing, is a multi-phase competition that tests multiple aspects of horse and athlete skill. Each event consists of three phases: dressage, jumping and cross-country.¹ In protecting both horse and athlete welfare, significant focus must be given to the cross-country phase of eventing, as falls during cross-country can have very serious consequences up to and including the death of both horse and athlete.^{2,3} So-called 'rotational falls' are a particular concern, defined as when the horse somersaults as a result of hitting a fence. The horse's hindquarters end up significantly higher than its front end and it will land on the landing side of the fence.⁴ Despite the sport's long history, it was not until 1999 that safety in eventing achieved global attention, when five athlete fatalities in the UK that year prompted major reviews of safety.⁵ The International Eventing Safety Committee concluded in the year 2000 that 'everything should be done to prevent horses falling'.⁶ Since then, studies to identify risk factors have primarily focussed on course-level factors during the cross-country phase, as well as behavioural factors on the part of the horse and athlete.⁷⁻¹¹ Until recently, the most recent season for which risk factors for horse and athlete falls during cross-country had been published was the 2002 season. A study of horse-, athlete- and course-level risk factors covering the period 2008–2018 published in 2021 was the first peer-reviewed publication since 2008.¹² Factors at the level of event and horse previously reported to be associated with increased risk of falls include higher event level, longer course length or more fences, individual history of falls and poor performance in the dressage phase.^{7,12} Other factors previously reported include the presence of water for takeoff or landing, approach and landing gradients, angle of fence and approach speed.^{9,10}

The Barnett report—an audit covering the period 2008–2014—was published by the FEI in 2016 with statistical analysis of falls and investigation of factors contributing to rotational falls in particular, but the report was not published in a peer-reviewed journal.¹³ This report showed that 14.5% of riders who had a rotational fall were seriously or fatally injured, compared with 4.3% of riders who had a nonrotational fall. A report published by the FEI in 2020 found that in the period 2009–2020, 16.7% of riders who had a rotational fall were seriously or fatally injured, compared with 4.8% of riders who had a nonrotational fall.¹⁴

Rule changes have been implemented throughout the last two decades relating to course design, fence composition and competition format, but to date no academic studies have examined the impact of such changes. One prominent example of a rule change was the introduction of frangible fences, that is fences that are designed to break and/or deform to help prevent somersault falls that are induced by contact of the horse with the fence.^{15,16} The first type of frangible device introduced was the frangible pin, first used in eventing in 2002.¹⁷ More types of frangible devices including MIM clips, which enable a wider range of fence types to be designed as collapsible, were introduced later and their use has become more popular since 2015. Anecdotally, there is a perception that frangible fences have reduced the number of rotational falls.

The FEI reported in 2021 that the percentage of competition starts resulting in a fall was 5.94% during the period 2005–2008, 5.37% during the period 2009–2013 and 5.38% during 2014–2020.¹⁸ The same FEI report also showed that in the period 2008–2017, over 90% of falls during FEI eventing competitions occurred at fences during cross-country. In the period 2018–2020, this proportion fell to just under 80% of falls, although there was a change in how fall locations were classified in 2018. Until 2008 it was permitted for athletes who became unseated to simply remount and continue in competition. From 2009 onwards, either a horse fall or an athlete becoming unseated resulted in automatic elimination. Thus, it was the case for prior studies, and remains the case in this study, that the primary focus on understanding and aiming to reduce the incidence of falls should be at fences during cross-country.

This article presents the results of a multivariable model incorporating risk factors at the level of the fence, course and event. The goal of this is to understand which risk factors contribute to increased odds of a horse falling or athlete being unseated at a particular fence or element of combination fence. The main hypothesis was that some combination of fence, course and event-level risk factors—including fence design, location within the course and factors such as approach gradient or presence of water—would contribute to the likelihood of a horse falling or athlete being unseated at a given fence. The focus on the data at the fence level, and use of an outcome definition 'one or more falls at this fence', allows a greater understanding of how fence and course design impacts horse and athlete safety.

2 | MATERIALS AND METHODS

The dataset used was the FEI's Global Eventing Database. A form of the database is publicly available online¹⁹—the authors had access to the complete dataset for this study in collaboration with the FEI. The data used in this study consist of every FEI eventing competition held between 1 January 2008 and 31 December 2018. The database is substantial and multifaceted—it contains detailed information about each competition, along with specific information about results, fences, horses, athletes and falls.

Falls are defined for the athlete as 'when he/she is separated from the horse in such a way as to necessitate remounting', and for the horse as 'when at the same time, both its shoulder and quarters have touched either the ground or the obstacle and the ground or when it is trapped in a fence in a way that it is unable to proceed without assistance or is liable to injure itself'.²⁰ Under these definitions, horse falls and athlete falls are mutually exclusive events. A detailed fall report form is completed for every fall, so that the circumstances of the fall are recorded as part of the FEI database.

This study investigated the fence- and course-level potential risk factors associated with falls of either type. The data were modelled at the fence level—that is, every fence at every event was included in the study—and the deleterious outcome studied was a binary variable indicating whether or not there had been at least one fall (athlete or horse) at a fence. Note that 842 fences (10.2% of those that had any

TABLE 1 Potential risk factors included for consideration in the multivariable logistic regression model

Risk factor	Categorisation	Notes
Year	Categorical	Collapsed into two categories: 2008–2015 and 2016–2018
Event level	Categorical	1*, 2*, 3* or 4*. Olympics and World Equestrian Games were included in 4*
Event format	Binary	Long format (CCI) or short format (CIC)
Fence number	Continuous	Scaled such that a unit increase in the model variable corresponds to an increase of five in fence number
Fence element	Categorical	If a fence is part of a combination, indicates at which element the fall happened
Fence type	Categorical	Fence type according to course design guidelines (reference)
Optional route	Binary	If the fence is part of an optional route
Fence is frangible	Binary	If the fence is frangible, that is, has some moving parts which are designed to yield when collided with
Fence is portable	Binary	
Take off from water	Binary	
Landing into water	Binary	
Associated with water	Binary	N.B. there is an incomplete overlap between 'takeoff from', 'landing into' and 'associated with' water
Approach to jump	Categorical	Describes the ground level before the jump
Landing after jump	Categorical	Describes the ground level after the jump
Number of cross-country starters	Continuous	Scaled such that a unit increase of the model variable represents an increase of 10
Course length	Categorical	
Course level	Categorical	N.B. there is an incomplete overlap between course level and event level (above)
Effort count of entire course	Continuous	The total number of jumping efforts—combined fences are counted as one jumping effort
Number of individual fences on entire course	Continuous	The total number of unique fences—usually greater than the corresponding effort count
Total jumping efforts made at this event	Continuous	The number of fences multiplied by the field size of cross-country starters

Note: Categorisation shows the form chosen after testing both continuous and categorical variants, for those risk factors which were originally continuous.

TABLE 2 The event level categorisations used in this study were those in place up until 2018

Categorisation 2018 and earlier Olympics and World Equestrian Games Special category	Categorisation 2019 onwards Olympics and World Equestrian Games Special category
CCI4*	CCI5*-L (long)
CCI3*	CCI4*-L (long)
CIC3*	CCI4*-S (short)
CCI2*	CCI3*-L (long)
CIC2*	CCI3*-S (short)
CCI*	CCI2*-L (long)
CIC*	CCI2*-S (short)
<i>New introductory level</i>	CCI* (unified) Not compulsory for qualifications

Note: In 2019, the FEI redesignated all event levels. This table shows the old and new categories to aid readers in interpreting the event levels used in the study. Note that these categories include sub-designation of short or long format (CCI or CIC respectively in the pre-2019 system)—event format was included in the present study as a separate risk factor.

falls) had more than one fall occur, meaning the risk of falls may be underestimated for some individual fences in the final model.

Multivariable logistic regression models were constructed in a bespoke code written in R version 4.1.2 (R Foundation for Statistical Computing).²¹ Potential risk factors included in this study, along with category definitions, are shown in Table 1. Note that the variable 'event level' was included in this study in the form that it took during the time period studied. Events were assigned a star rating from 1* to 4*, with 4* representing the most challenging competitions. In 2019, the categories were altered—see Table 2 for an explanation of the old and new event level systems. Risk factors included in continuous form were also examined in categorical form, with the best fitting form as assessed using the Akaike information criterion (AIC) included in the final model. Variables were assessed for collinearity during initial data exploration. The first stage of modelling examined each risk factor in turn in a univariable logistic regression model, with a maximum *p* value of 0.20 used to select candidates for the final model. Multivariable mixed-effect logistic regression models were constructed using a stepwise bidirectional process (R function 'stepAIC') with each step assessed using the AIC, until the best-fitting models were identified.^{22,23}

Fence type	Fences	Fences with at least one fall (%)	Fence-type definitions
Total	204 399	8253 (4.0%)	
A0	1850	97 (5.2%)	A: post and rails
A1	8893	428 (4.8%)	
A2	2142	90 (4.2%)	
A3	3475	111 (3.2%)	
A4	936	37 (4%)	
B1	5653	229 (4.1%)	B: palisade
B2	1879	85 (4.5%)	
C1	6713	228 (3.4%)	C: square spread
C2	10 761	356 (3.3%)	
C3	17 345	585 (3.4%)	
D1	2816	80 (2.8%)	D: ascending spread
D2	5117	136 (2.7%)	
D3	19 021	466 (2.4%)	
D4	5868	172 (2.9%)	
E1	9467	400 (4.2%)	E: brush
E2	12 923	462 (3.6%)	
E3	4306	199 (4.6%)	
E4	3612	252 (7%)	
E5	464	17 (3.7%)	
E6	2992	86 (2.9%)	
F1	24 133	1173 (4.9%)	F: round
F2	17 614	565 (3.2%)	
G1	2172	157 (7.2%)	G: corner
G2	1801	122 (6.8%)	
G3	7394	600 (8.1%)	
H1	5871	283 (4.8%)	H: Trakehner
H2	180	12 (6.7%)	
H3	253	13 (5.1%)	
J1	67	3 (4.5%)	J: step
J2	1707	197 (11.5%)	
J3	2208	119 (5.4%)	
J4	3884	108 (2.8%)	
J5	4977	147 (3%)	
K	330	15 (4.5%)	K: water ('splash')
L	3722	161 (4.3%)	L: ditch
Other	1301	43 (3.3%)	
Unknown	552	19 (3.4%)	

TABLE 3 Descriptive statistics of falls by fence type, in FEI eventing competitions between 2008 and 2018

Variables rejected at the univariable and multivariable stages were subsequently tested for confounding in the final model.²⁴ Biologically plausible combinations of risk factors were tested for second-order interaction and included for assessment in the final model. The final single-level model was tested for goodness-of-fit using the Hosmer–Lemeshow test.²² Any potential impact of event-level clustering was assessed by the mixed-effect model which included event as a random effect. Power calculations indicated that logistic regression models would have 80% power to detect odds

ratios of 1.03 or higher, with 95% confidence, for variables in continuous form. For variables in binary categorical form, the model had 80% power to detect odds ratios of 1.07 or higher, with 95% confidence.

3 | RESULTS

Table 3 shows the descriptive statistics of falls by fence type (diagrams of each fence type are shown in Figure 1) for the full cohort of




FIGURE 1 Diagrams of all fence types in use for the cross-country phase of FEI eventing competitions between 2008 and 2018. This figure is adapted from the FEI fence-types diagram document.²⁵

fences as recorded in the Global Eventing Database. The proportion of fences of each type at which there was at least one fall ranged from 2.4% for fence type D3 (covered ascending spread) to 11.5% for fence type J2 (step down into water).

Of 202 771 horse starts between 1 January 2008 and 31 December 2018, 190 429 started the cross-country phase. Of these, 10 519 (5.2%) had a fall recorded—henceforth, unless specified, ‘fall’ refers to either an athlete being unseated or a horse fall. Of these falls, 9358 (89.0%) occurred at a fence during the cross-country stage. At fence level, there were 204 399 unique fences used in 6450 unique FEI competitions during the time period. These data represent a total of approximately 6100 000 individual jumping efforts, where one jumping effort is one horse attempting one fence. Note that to account for potential modifications year to year, the ‘same’ fence at the ‘same’ event in different years was regarded as being unique. The 9358 recorded falls occurred at 8253 fences—4.0% of all unique

fences. The study cohort used for analysis was the 204 399 unique fences with each fence as a unit of observation. Cases were defined as the 8253 fences at which there was at least one fall recorded. Table 4 shows the final multivariable model. The univariable model results are shown in Table S1.

At event level, compared with competitions in 2016, 2017 and 2018 combined, fences in competitions between 2008 and 2015 were at increased odds of being associated with a fall (odds ratio 1.11, [95% confidence interval 1.06–1.17]). Fences in events at 3* or 4* level were more likely to have been associated with a fall than fences in 1* and 2* events, at odds ratio 1.09 (1.03–1.15) for 3* events and odds ratio 1.59 (1.42–1.79) for 4* events. An increase in the number of cross-country starters was associated with increased odds of a fall occurring, with field sizes at or above the 75th percentile (42 horse starts) at odds ratio 2.00 (1.95–2.05) compared with field sizes at or below the 25th percentile (12 horse starts).

TABLE 4 Multivariable model results for the outcome 'at least one fall at this fence'

Risk factor	Cases (%)	Controls (%)	Odds ratio	95% Confidence interval	p
Event year					
2016–2018 ^a	2428 (3.7%)	63 360 (96.3%)	1.00	-	-
2008–2015	5825 (4.2%)	132 786 (95.8%)	1.11	1.06–1.17	<0.001
Event level					
Level 1 or 2 ^a	6067 (3.9%)	150 821 (96.1%)	1.00	-	-
Level 3	1811 (4.2%)	41 721 (95.8%)	1.09	1.03–1.15	0.004
Level 4	375 (9.4%)	3604 (90.6%)	1.59	1.42–1.79	<0.001
Number of cross-country starters					
Per additional 10 horses	Median = 24 Min = 0 ^b	IQR = 30 Max = 142	1.26	1.25–1.27	<0.001
Fence number					
Per additional five fences	Median = 12 Min = 1	IQR = 10 Max = 38	1.07	1.05–1.09	<0.001
Element of combined fence					
Not combined ^a	3513 (3.4%)	101 275 (96.6%)	1.00	-	-
Element A	1915 (4.6%)	40 054 (95.4%)	1.15	1.08–1.22	<0.001
Element B	2203 (5.1%)	40 692 (94.9%)	1.33	1.25–1.42	<0.001
Element C or other	622 (4.2%)	14 125 (95.8%)	1.21	1.10–1.32	<0.001
Fence type					
C ^a (square spread)	1231 (3.4%)	35 441 (96.6%)	1.00	-	-
A0 (post and rails)	97 (5.2%)	1753 (94.8%)	1.32	1.06–1.64	0.01
A1	428 (4.8%)	8465 (95.2%)	1.15	1.02–1.30	0.02
A2–4	238 (3.6%)	6315 (96.4%)	0.91	0.79–1.06	0.2
B1 (palisade)	229 (4.1%)	5424 (95.9%)	1.02	0.88–1.18	0.8
B2	85 (4.5%)	1794 (95.5%)	1.49	1.19–1.88	<0.001
D1 or D4 (ascending spread)	252 (2.9%)	8432 (97.1%)	0.84	0.73–0.97	0.01
D2	136 (2.7%)	4981 (97.3%)	0.73	0.61–0.88	<0.001
D3	466 (2.4%)	18 555 (97.6%)	0.69	0.62–0.78	<0.001
E1 (brush)	400 (4.2%)	9067 (95.8%)	0.96	0.85–1.08	0.5
E2, E3, E5 or E6	764 (3.7%)	19 921 (96.3%)	0.90	0.82–0.99	0.04
E4	252 (7%)	3360 (93%)	2.02	1.75–2.34	<0.001
F1 (round)	1173 (4.9%)	22 960 (95.1%)	1.00	0.92–1.10	>0.9
F2	565 (3.2%)	17 049 (96.8%)	0.77	0.69–0.85	<0.001
G1 (corner)	157 (7.2%)	2015 (92.8%)	1.91	1.59–2.30	<0.001
G2	122 (6.8%)	1679 (93.2%)	2.09	1.72–2.55	<0.001
G3	600 (8.1%)	6794 (91.9%)	2.44	2.20–2.72	<0.001
H1 (Trakehner)	283 (4.8%)	5588 (95.2%)	1.49	1.30–1.71	<0.001
H2	12 (6.7%)	168 (93.3%)	2.00	1.09–3.67	0.03
H3	13 (5.1%)	240 (94.9%)	1.98	1.13–3.50	0.02
J1, J3, J4 or J5 (step)	377 (3.4%)	10 759 (96.6%)	0.78	0.68–0.89	<0.001
J2	197 (11.5%)	1510 (88.5%)	1.16	0.97–1.40	0.1
K (water)	15 (4.5%)	315 (95.5%)	0.38	0.22–0.64	<0.001
L (ditch)	161 (4.3%)	3561 (95.7%)	1.10	0.89–1.36	0.4
Approach gradient					
Level or up ^a	7201 (3.9%)	176 813 (96.1%)	1.00	-	-
Down	1052 (5.2%)	19 333 (94.8%)	1.35	1.19–1.52	<0.001

(Continues)

TABLE 4 (Continued)

Risk factor	Cases (%)	Controls (%)	Odds ratio	95% Confidence interval	p
Landing gradient					
Level ^a	5701 (3.7%)	147 884 (96.3%)	1.00	-	-
Down	2139 (5.7%)	35 367 (94.3%)	1.42	1.32–1.54	<0.001
Up	413 (3.1%)	12 895 (96.9%)	0.80	0.70–0.93	0.002
Landing into water					
No ^a	7044 (3.7%)	184 779 (96.3%)	1.00	-	-
Yes	1209 (9.6%)	11 367 (90.4%)	1.82	1.65–2.01	<0.001
Jump associated with water					
No ^a	6198 (3.5%)	168 850 (96.5%)	1.00	-	-
Yes	2055 (7%)	27 296 (93%)	1.46	1.34–1.59	<0.001
Jump is on optional route					
No ^a	8142 (4.1%)	189 654 (95.9%)	1.00	-	-
Yes	111 (1.7%)	6492 (98.3%)	0.23	0.19–0.28	<0.001
Fence is frangible					
No ^a	7678 (3.9%)	187 758 (96.1%)	1.00	-	-
Yes	575 (6.4%)	8388 (93.6%)	1.28	1.15–1.41	<0.001
Fence is portable					
No ^a	4470 (4.3%)	98 770 (95.7%)	1.00	-	-
Yes	3783 (3.7%)	97 376 (96.3%)	0.94	0.89–0.99	0.02
Interactions terms					
Approach down × landing down			0.78	0.67–0.92	0.002
Jump associated with water × landing up			1.40	1.09–1.78	0.008

Note: Cases were fences at which a fall of any kind was recorded. Risk factors with a *p* value of less than 0.05 were retained in the final model.

^aReference category among categorical variable levels.

^bTwo competitions in the database had zero cross-country starters recorded.

At fence level, higher fence numbers (i.e., further from the start of the course) were associated with increased odds of a fall occurring at that fence. Fences at or above the 75th percentile (fence number 17) were at odds ratio 1.14 (1.10–1.19) compared with fences at or below the 25th percentile (fence number 7). Fences that were part of combined obstacles—that is, jumping efforts with multiple individual fences—were more likely to be associated with a fall than isolated, noncombined fences. Compared with fences that were not part of a combined obstacle, element A of a combined fence was associated with an odds ratio of 1.15 (1.08–1.22), element B was at odds ratio 1.33 (1.25–1.42) and element C or later were at odds ratio 1.21 (1.10–1.32) for a fall to occur at that fence.

Compared with fences that had a level or uphill approach, fences with a downhill approach were associated with an increased odds of a fall, with odds ratio 1.35 (1.19–1.52). Similarly, in comparison to fences which had a level landing gradient, fences with a downward landing were associated with an increased odds of a fall (1.42 [1.32–1.54]), while fences with an uphill landing were associated with reduced odds (0.80 [0.70–0.93]). Fences that had a landing into water were associated with an increased odds of a fall compared with fences without a water landing, at odds ratio 1.82 (1.65–2.01). Fences that were defined as being ‘associated with water’ (which included fences

with take-off from and landing into water, and fences with neither factor) were associated with an increased odds of a fall, at odds ratio 1.46 (1.34–1.59). Fences that were part of an optional route for a particular obstacle were at greatly reduced odds of a fall (0.23 [0.19–0.28]). Fences that were frangible were associated with a greater odds of a fall (1.28 [1.15–1.41]). Fences that were portable were at reduced odds of a fall than permanent fences (0.94 [0.89–0.99]).

Seventeen fence types were found to have significant associations with the likelihood of falls occurring. The reference fence type used was C—square spread, one of the more common fence types. Compared with type C fences, type G3 fences—corner with solid walls and solid top (2.44 [2.20–2.72]), type G2—corner with ‘post and rail’ walls and solid top (2.09 [1.72–2.55]), type E4—brush with ditch in front (2.02 [1.75–2.34]), type H2—Trakehner with cross rails on top (2.00 [1.09–3.67]) were at least twice as likely to be associated with a fall, and type K fences—a water obstacle (colloquially known as a ‘splash’) with no physical fence—were at most reduced odds of a fall occurring (0.38 [0.22–0.64]).

Two second-order interaction terms were retained in the final model. Fences for which the approach and landing gradients were both downhill were at a total odds ratio of 1.50 (1.05–2.15) of a fall occurring, compared with fences with level gradients at approach and

landing. Fences that were associated with water and that had an upwards gradient on the landing were at a total odds ratio of 1.64 (1.02–2.63) for a fall occurring, compared with fences that were not associated with water and that had a level landing gradient.

No confounding was detected between retained risk factors and those rejected at any stage of model-building, with none of the model estimates of risk factors retained in the final model changing by more than 10% upon the inclusion of any of the rejected risk factors. The inclusion of event as a random effect accounted for 18% of the variance in the final model and altered the model coefficients of one risk factor by more than 10%—the odds ratio of ‘fence is portable’ in the mixed-effects model was 0.94 instead of 0.93 in a fixed effects-only model. No evidence of a lack of fit for the final model was found with the Hosmer–Lemeshow goodness-of-fit test, which returned a *p* value of 0.3.

4 | DISCUSSION

Risk factors at event-level and fence-level were found to be statistically significantly associated with the likelihood of a fall (of a horse or athlete) occurring at each individual fence. A summary of the fence-level results is shown in Table 5.

Later years included in the data were associated with lower odds—this could be related to rule changes over that time period such as alterations to event format and course design including fence design, or changes to the minimum eligibility requirements for qualification.

Event levels at 3* and 4* must naturally include longer courses with more fences, and more challenging obstacles.²⁰ Therefore, it should perhaps be expected that fences in those events are more likely to have falls occur at them compared with fences in 1* and 2* events, even accounting for the fact that better quality horses/combinations are competing at the higher levels. It is nevertheless important for athletes stepping up to higher levels, to know how much greater the risk is for their new level of competition. Larger field sizes were more likely to result in falls occurring at the fence level simply because there would be more opportunity for falls (i.e. more jumping efforts on the day) compared with smaller field sizes. It could also be the case that competitions with more competitors were more likely to have poorer ground conditions, for example, damage to the footing either side of fences, in particular where the ground was softer after wet weather. It was reported by Murray et al.¹⁰ that competitions where the ground conditions were soft/heavy were more likely to have falls occur compared with where there were firm ground conditions.

Several of the results of this study are consistent with those reported in earlier studies. Associations between increased likelihood of falls and (i) fences later in the course; (ii) elements (in particular mid and final elements) of combined fences; and (iii) certain fence types were reported by Singer et al.⁷ That study was a case-control study which found in a multivariable model that ascending spread fences (fence type D in the present study) were less likely to be associated with falls than fences that were not ascending spread type. That study also found that fences with a ditch in front were more likely to be associated with falls compared with fences without a ditch in front.

TABLE 5 A summary of fence-level risk factors identified in the final multivariable model shown in Table 4

Fence-level factor	Reference category	Increased odds of fall	Reduced odds of fall
Approach gradient	Level or uphill	Downhill, 1.35 (1.19–1.52)	
Landing gradient	Level	Downhill, 1.42 (1.32–1.54)	Uphill, 0.80 (0.70–0.93)
Landing into water	No	Yes, 1.82 (1.65–2.01)	
Associated with water	No	Yes, 1.46 (1.34–1.59)	
On optional route	No		Yes, 0.23 (0.19–0.28)
Frangible fence	No	Yes, 1.28 (1.15–1.41)	
Portable fence	No		Yes, 0.94 (0.89–0.99)
Approach downhill and landing downhill	No and No	Yes and Yes, 1.50 (1.05–2.15)	
Associated with water and landing uphill	No and No	Yes and Yes, 1.64 (1.02–2.63)	
Element of combined fence	Not combined	Element A, 1.15 (1.08–1.22) Element B, 1.33 (1.25–1.42) Element C, 1.21 (1.10–1.32)	
Fence type	C (square spread)	A0, 1.32 (1.06–1.64) A1, 1.15 (1.02–1.30) B2, 1.49 (1.19–1.88) E4, 2.02 (1.75–2.34) G1, 1.91 (1.59–2.30) G2, 2.09 (1.72–2.55) G3, 2.44 (2.20–2.72) H1, 1.49 (1.30–1.71) H2, 2.00 (1.09–3.67) H3, 1.98 (1.13–3.50)	D, 0.84 (0.73–0.97) D2, 0.73 (0.61–0.88) D3, 0.69 (0.62–0.78) E, 0.90 (0.82–0.99) F2, 0.77 (0.69–0.85) J, 0.78 (0.68–0.89) K, 0.38 (0.22–0.64)

Note: Odds ratios are reported in the format odds ratio (95% confidence interval). Fence-type diagrams are shown in Figure 1.

Another case-control study demonstrated an association between water approach/landing and increased likelihood of falls.⁹ An association between jumps with drop landings and increased likelihood of falls has also previously been detected.^{8,9}

Fences with higher numbers—that is, located later in the course—were generally more likely to have falls occur at them compared with fences earlier on the course. This could be the result of fatigue for both athlete and horse—the longer the course, and more obstacles that they have overcome, the more likely that fatigue could contribute to a mistake being made. Additionally, fences later in competitions would naturally have fewer jumping efforts made over them compared with fences earlier in the competition, due to the attrition of retirals/falls/eliminations for some competitors earlier in the course. This means that the odds ratio reported here is likely to be an underestimate of the true odds ratio for fences located later in the course. Fences that were part of combined obstacles were more likely to have falls occur at them than isolated, noncombined fences. Fences that were described as being element B—that is, the second fence in the combination—of a combined obstacle were at the highest odds ratio compared with fences that were not combined. This reflects the additional complexity of combined obstacles—horse and athlete have to carefully approach each stage, and in the event of a successful but slightly misjudged jump over element A, for example, they could have little time to recover before attempting to jump element B.

The setting around a fence was found to have a significant impact on the likelihood of a fall occurring at the fence, in several aspects of the course design. A downward slope on the approach to or landing from a jump was associated with increased odds of a fall. When approaching down a hill, controlling the centre of balance of the horse is more challenging and precisely identifying the point of take-off more difficult. As for landing, it is possible that a downwards slope after a jump could require significant adjustment of horses' bodies in order to continue in their stride, thus making such a jump more difficult than on the flat ground. It could also be more likely that a horse could slip on a downwards landing compared with a flat landing. Fences that have their landing into water or otherwise are associated with water have the extra difficulty of the ground not being visible to the horse or athlete—as well as any other challenges that the water would add, for example to pacing and positioning on approach and while jumping, as well as potential alterations to the kinematics of the horse's movement.^{26,27}

Fences on optional routes—found to be at reduced odds of a fall occurring—are by design less challenging than those on the direct route through an obstacle. However, the data showing which branch of a fence containing an optional route was chosen by competitors were not available. For example, a fall that occurred at an optional fence would be recorded for that specific fence, but the number of successful jumping efforts (controls) at the same fence was unknown. This is unique to optional route fences because specific route choice is not recorded for successful jumping efforts. Fences that are frangible rather than solid are perhaps more likely to be misjudged as the athlete thinks they 'can get away with' clipping it and may approach the fence with less caution. It could also be that course designers

intentionally build more challenging fences when they know they will be including a frangible device. Portable fences are perhaps smaller and slightly less challenging compared with permanent fences.

Some fences are designed to be very challenging for horses and athletes, and this was reflected in the likelihood of falls occurring at certain fence types. The fence type with the highest proportion of falls—11.5% of J2 fences had at least one fall occur—was not statistically significant in the final model. At the univariable stage, J2 fences (step down into water) were associated with increased odds of a fall occurring compared with C type fences (square spread), at odds ratio 3.74 (3.20–4.38), and *p* value <0.001. This implies that other fixed effects that were retained in the final model account for some of the risk associated with J2 fences. Further investigation revealed that 86% of type J2 fences were in the category 'yes' for the risk factor 'landing into water'. Consequently, in the final model, the variable 'landing into water' was retained while type J2 fences were excluded. One curious aspect of this is that according to the FEI fence design document,²⁵ type J2 fences are described as a 'step down into water'. It might reasonably be expected that 100% of such fences would be recorded as having a landing into water, but in the available data 14% of J2 fences were not recorded as such. Accurate data recording is critical for future studies and risk management and this should be a key message to those responsible for reporting and recording these data.

The headline FEI statistics on falls, along with the similarities between prior work which was completed using a selected cohort of case-control data from the 2000, 2001 and 2002 seasons,^{7–11} and this study which covers the full cohort of data from 2008–2018 indicate that it is difficult to conclude that some aspects of eventing cross-country course design have become safer—at least in terms of reducing falls—since the International Eventing Safety Committee (IESC) reported its findings in 2000. When considering the Barnett report,¹³ this study has identified some of the same risk factors including event level, fence types A1, E4, G1, G3, downhill landings, association with water and frangible fences.

The results from this work are the first step towards building a 'risk profile' or 'score' for each cross-country course and could contribute to further grading of cross-country phases of events within different levels of competition, helping to inform athletes as to the expected difficulty of the course on which they are about to compete. Course risk profiles can be used to support the development of horses and riders and be included in qualification criteria to progress to higher event levels. These results also motivate a discussion about whether safety could be a higher priority in course design. It would not be desirable to look at these results and say, for example, that jumps in or out of water, corner and Trakehner fences should no longer be used. Rather, it should be considered whether it might be possible to design around these more challenging fences. For example, it would be more appropriate to ensure that very challenging fence types generally are not over represented in the second half of cross-country courses such that the effect of fatigue (identified as a potential explanation for falls at later fences in this study and previously⁷) is not exacerbated by a very difficult fence

design. Where feasible, the inclusion of more optional routes on cross-country courses would contribute to reducing the overall 'difficulty rating' of the course. An awareness of the risk factors identified here can inform course design—for example, through policy and course design documents—that aim to reduce the incidence of athlete and horse falls, while also maintaining the level of challenge that stakeholders would expect to see. Participants in equestrian sports recognise that there is inherent risk involved, especially when riding at speed over solid obstacles during cross-country. An increased focus on safety for both horse and athlete with the goal of minimising inherent risks as far as possible will also positively impact public perception of the sport, and bolster the social licence of eventing in the public eye.²⁸ This is a particularly important time for stakeholders to focus on the social licence to operate, both in the broader context of equestrian sports and for the particular case of eventing.²⁹

More than two decades after the IESC report urged 'everything should be done to prevent horses from falling', an improved understanding of the true level of risk posed by a particular set of fences on a specific course could form the focus of further risk reviews. Grading of courses, based on the risk profile of all fences, would be a useful next step to help inform athletes about the level of risk to which they would be exposing themselves and their horses. In combination with validated horse and athlete risk profiling, course grading would reduce the risk of serious injury associated with this challenging Olympic sport.

AUTHOR CONTRIBUTIONS

Euan David Bennet contributed to study design, study execution and data analysis and interpretation. Euan David Bennet had full access to all the data in the study and is responsible for data integrity and accuracy of the analysis. Heather Cameron-Whytock contributed to study execution and data analysis and interpretation. Tim D. H. Parkin contributed to study design, and data analysis and interpretation. All authors contributed to the preparation of the manuscript and gave final approval to the manuscript.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge the support of the FEI, which funded this research and thank Göran Akerström and Caterina Termini for their help in interpreting parts of the GED.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1111/evj.13863>.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the FEI. Restrictions apply to the availability of these data, which were used under licence for this study. Data are available from the authors with the permission of the FEI.

ETHICAL ANIMAL RESEARCH

Not applicable: data provided by a sports regulator were analysed.

ORCID

Euan David Bennet  <https://orcid.org/0000-0001-9049-1920>

Heather Cameron-Whytock  <https://orcid.org/0000-0003-0760-2584>

Tim D. H. Parkin  <https://orcid.org/0000-0003-3566-9030>

REFERENCES


1. FEI website [cited 2022 3 May]. Available from: <https://inside.fei.org/fei/disc/eventing>
2. O'Brien D. Look before you leap: what are the obstacles to risk calculation in the equestrian sport of Eventing? *Animals*. 2016;6:13. <https://doi.org/10.3390/ani6020013>
3. Eventing obituary tracker [cited 2022 3 May]. Available from: <https://www.horsetalk.co.nz/2012/04/02/eventing-safety/>
4. FEI glossary [cited 2022 3 May]. Available from: https://inside.fei.org/system/files/Glossary_0.pdf
5. News coverage [cited 2022 3 May]. Available from: <https://www.independent.co.uk/news/death-of-fifth-rider-shocks-eventing-into-crisis-reforms-1120793.html>
6. IESC report. 2000 [cited 2022 3 May]. Available from: <https://inside.fei.org/system/files/2000%20Safety%20report.pdf>
7. Singer ER, Saxby F, French NP. A retrospective case-control study of horse falls in the sport of horse trials and three-day eventing. *Equine Vet J*. 2003;35(2):139–45.
8. Murray JK, Singer ER, Saxby F, French NP. Factors influencing risk of injury to horses during eventing. *Veterinary Record*. 2004;154:204–8.
9. Murray JK, Singer ER, Morgan KL, Proudman CJ, French NP. Risk factors for cross-country falls at one-day events and at two-/three-day events. *Vet J*. 2005;170:318–24.
10. Murray JK, Singer ER, Morgan KL, Proudman CJ, French NP. The risk of a horse-and-rider partnership falling on the cross-country phase of eventing competitions. *Equine Vet J*. 2006;38(2):158–63.
11. Singer ER, Barnes J, Saxby F, Murray JK. Injuries in the event horse: training versus competition. *Vet J*. 2008;2008(175):76–81.
12. Bennet ED, Cameron-Whytock H, Parkin TDH. Fédération Equestre Internationale eventing: risk factors for horse falls and unseated riders during the cross-country phase (2008–2018). *Equine Vet J*. 2021; 1–10. <https://doi.org/10.1111/evj.13522>
13. FEI Barnett report. 2016 [cited 2022 3 May]. Available from: <https://inside.fei.org/system/files/Eventing%20Audit%20-%20Charles%20Barnett%20-%20Final%20Report%2026.07.16.pdf>
14. FEI risk management report 2009–2020 [cited 2022 3 May]. Available from: <https://inside.fei.org/system/files/2020%20Statistics%2020.01.2021.pdf>
15. FEI frangible fence standards [cited 2022 3 May]. Available from: <https://inside.fei.org/sites/default/files/FAQ%20Frangible%20Devices%2030%20Sept%202021.pdf>
16. Foreman MH, Engsborg JR, Foreman JH. Point mass impulse-momentum model of the equine rotational fall. *Comparative Exercise Physiology*. 2019;15(3):157–65.
17. Frangible pins [cited 2022 3 May]. Available from: <https://www.horseandhound.co.uk/tag/frangible-pins>
18. FEI risk management report 2009–2021. [cited 2022 3 May]. Available from: <https://inside.fei.org/system/files/2021%20Statistics%2024.01.2022.pdf>
19. FEI database [cited 2022 3 May]. Available from: <https://data.fei.org/default.aspx>
20. FEI eventing rules [cited 2022 3 May]. Available from: <https://inside.fei.org/fei/disc/eventing/rules>

21. R Foundation [cited 2022 3 May]. Available from: <https://www.r-project.org/foundation/>
22. Hosmer DW, Lemeshow S. Applied logistic regression. 2nd ed. New York: John Wiley and Sons Inc.; 2000.
23. Akaike H. Information theory and an extension of the maximum likelihood principle. In: Petrov BN, Csaki F, editors. Second international symposium on information theory. Budapest, Hungary: Akademiai Kaido; 1973. p. 267–81. Reprinted (1992) in: Breakthroughs in statistics, Volume I. eds. Kotz, S. and Johnson, N.L. Springer, New York, pp. 599–624.
24. Dohoo I, Martin W, Stryhn H. Veterinary epidemiologic research. Charlottetown, Prince Edward Island, Canada: AVC Inc.; 2003.
25. FEI fence type diagrams [cited 2022 3 May]. Available from: <https://inside.fei.org/system/files/2021%20Fence%20Diagrams%2017.03.2021.pdf>.
26. McCrae P, Bradley M, Rolian C, Leguillette R. Water height modifies forelimb kinematics of horses during water treadmill exercise. *Comparat Exercise Physiol.* 2020;17(1):91–8.
27. Scott R, Nankervis K, Stringer C, Westcott K, Marlin D. The effect of water height on stride frequency, stride length and heart rate during water treadmill exercise. *Equine Vet J.* 2010;42(s38):662–4.
28. Heleski C, Stowe CJ, Fiedler J, Peterson ML, Brady C, Wickens C, et al. Thoroughbred racehorse welfare through the lens of social license to operate with an emphasis on a U.S. perspective. *Sustainability.* 2020;12(5):1706. <https://doi.org/10.3390/su12051706>
29. 'Horse sport must act to stay in Olympics: "we are guests at these Games"' [cited 2022 20 Jun]. Available from: <https://www.horseandhound.co.uk/news/equestrian-sport-olympics-decisions-776843>.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Bennet ED, Cameron-Whytock H, Parkin TDH. Fédération Equestre Internationale eventing: Fence-level risk factors for falls during the cross-country phase (2008–2018). *Equine Vet J.* 2023;55(3):463–73. <https://doi.org/10.1111/evj.13863>



VETCT

WOULD YOU LOVE...

A dark room to view your radiographs?
A few more hours in the day?

Well the good news is, VetCT can give you **BOTH!**

Our team of specialist equine radiologists can read your radiographs for you.

Send them while you are still horse-side and receive a fully annotated radiology report in your inbox by the time you're back at the practice.

Contact us today
+44 (0) 1223 080453