



Full length article

Spatial assessment of the attributable burden of disease due to transportation noise in England

Calvin Jephcote^{a,1}, Sierra N. Clark^{b,1}, Anna L. Hansell^{a,c}, Nigel Jones^d, Yingxin Chen^a, Claire Blackmore^a, Katie Eminson^a, Megan Evans^a, Xiangpu Gong^c, Kathryn Adams^a, Georgia Rodgers^b, Benjamin Fenech^{b,c,*}, John Gulliver^{a,c,2,*}

^a Centre for Environmental Health and Sustainability, University of Leicester, Leicester, UK

^b Noise and Public Health, Radiation Chemical and Environmental Hazards, Science Group, UK Health Security Agency, UK

^c NIHR Health Protection Research Unit in Environmental Exposures and Health at the University of Leicester, UK

^d Extriium, UK

ARTICLE INFO

Handling Editor: Adrian Covaci

Keywords:

Road-traffic noise
Aircraft noise
Railway noise
Exposure-response relationship
Geospatial analysis
Environmental burden of disease
Environmental health

ABSTRACT

Background: Noise pollution from transportation is one of the leading contributors to the environmental disease burden in Europe. We provide a novel assessment of spatial variations of these health impacts within a country, using England as an example.

Methods: We estimated the burden of annoyance (highly annoyed), sleep disturbance (highly sleep disturbed), ischemic heart disease (IHD), stroke, and diabetes attributable to long-term transportation noise exposures in England for the adult population in 2018 down to local authority level (average adult population: 136,000). To derive estimates, we combined literature-informed exposure-response relationships, with population data on noise exposures, disease, and mortalities. Long-term average noise exposures from road, rail and aircraft were sourced from strategic noise mapping, with a lower exposure threshold of 50 dB (decibels) L_{den} and L_{night} .

Results: 40 %, 4.5 % and 4.8 % of adults in England were exposed to road, rail, and aircraft noise exceeding 50 dB L_{den} . We estimated close to a hundred thousand (~97,000) disability adjusted life years (DALY) lost due to road-traffic, ~13,000 from railway, and ~17,000 from aircraft noise. This excludes some noise-outcome pairs as there were too few studies available to provide robust exposure-response estimates. Annoyance and sleep disturbance accounted for the majority of the DALYs, followed by strokes, IHD, and diabetes. London, the South East, and North West regions had the greatest number of road-traffic DALYs lost, while 63 % of all aircraft noise DALYs were found in London. The strategic noise mapping did not include all roads, which may still have significant traffic flows. In sensitivity analyses using modelled noise from all roads in London, the DALYs were 1.1x to 2.2x higher.

Conclusion: Transportation noise exposures contribute to a significant and unequal environmental disease burden in England. Omitting minor roads from the noise exposure modelling leads to underestimation of the disease burden.

1. Introduction

Noise from road traffic, rail, and aviation transport affects millions of people in Europe (EEA, 2020) and are major contributors to the overall environmental disease burden (Hanninen et al., 2014; EEA, 2020). In

England, the Environmental Noise Regulations 2006 (as amended) requires, on a five year cycle, the determination, through noise mapping, of exposure to environmental noise from major sources of road, rail and aircraft, and in urban areas (known as agglomerations) (HM Government, 2006). The Environmental Noise (England) Regulations 2006

* Corresponding authors at: 23 Stephenson Street, UK Health Security Agency, Birmingham B2 4BH, UK (B. Fenech). School of Geography, Geology & The Environment, University of Leicester, Leicester LE1 7RH, UK (J. Gulliver).

E-mail addresses: benjamin.fenech@ukhsa.gov.uk (B. Fenech), jgulliver@sgul.ac.uk (J. Gulliver).

¹ Joint first authors.

² Joint senior authors.

<https://doi.org/10.1016/j.envint.2023.107966>

Received 17 January 2023; Received in revised form 14 April 2023; Accepted 7 May 2023

Available online 13 May 2023

0160-4120/© 2023 Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

transpose the provisions of the European Commission's Environmental Noise Directive (END, 2002/49/EC) (HM Government, 2006). Strategic noise mapping undertaken in 2017 revealed that ~ 11.5 million people are exposed to noise above 55 dB L_{den} (day-evening-night noise) from roads (DEFRA, 2019a,c) and ~ 1.5 million from railways (DEFRA, 2019a,b), respectively. England is also a major hub for commercial aviation, with one of Europe's busiest airports (Heathrow) situated within the London urban area (UECNA, 2020). 55 dB L_{den} is a historic noise threshold used for reporting strategic noise mapping data in accordance with the END requirements. The health-based guidelines for noise published by the World Health Organization (WHO) Regional Office for Europe were 53 dB L_{den} for road and 54 dB L_{den} for railway noise, respectively (WHO 2018). The equivalent guideline for aircraft noise exposure is 45 dB L_{den} , however currently aircraft noise exposure is not routinely modelled down to this level in England.

Long-term exposure to environmental noise has been shown to adversely impact psychological and physiological health and wellbeing (Guski et al. 2017; Hegewald et al. 2020; WHO 2018). Noise exposure at night can disturb sleep patterns, with downstream short and long-term consequences on daytime cognitive alertness, memory consolidation, quality of life and health (Basner and McGuire 2018). Exposures during daytime and night-time are also associated with cardiometabolic diseases such as incident cases of ischemic heart disease (IHD), stroke, and diabetes, with the majority of the epidemiological evidence related to road-traffic sources (Roswall et al. 2021; Sakhvidi et al. 2018; van Kempen et al. 2018; Vienneau et al. 2019). Mechanistic studies point to sustained stress responses in the body and thus repeated stimulation of the sympathetic nervous and endocrine systems as etiological pathways (Eriksson and Pershagen 2018; Munzel et al. 2014; Munzel et al. 2021a; Munzel et al. 2021b). Furthermore, noise impaired sleep is thought to be an important pathway for adverse cardiometabolic impacts through elevation of blood pressure and stress hormones, and induced changes to appetite and glucose dysregulation (Basner and McGuire 2018; Eriksson and Pershagen 2018; Munzel et al. 2021a).

Comparative burden of disease assessments provide important input into health decision-making and planning processes, research prioritization, and funding (Ezzati et al. 2004). For such comparisons to be made across health outcomes, risk factors, and locations, standardized metrics are needed. The Disability Adjusted Life Year (DALY) is one such metric that combines the years of life lost due to premature death and years of life lived in less than full health as a result of disease (WHO 2020). Several studies have quantified the attributable DALYs lost, or other measures of health burden, due to transportation noise exposures, largely within Europe at the city, regional or country level (Eriksson et al., 2017; Hanninen et al., 2014; Hegewald et al., 2021; Khomenko et al., 2022; Mueller et al., 2017; Murphy and Faulkner, 2022; Sohrabi and Khreis, 2020; Stassen et al., 2008; EEA, 2020; WHO Regional Office for Europe, 2011). The European Environment Agency (EEA) estimated that among a total of 33 European countries in 2017, 453,000 DALYs were lost to annoyance (being highly annoyed), 437,000 to sleep disturbance (being highly sleep disturbed), and 156,000 to ischaemic heart disease (IHD), as a result of environmental noise exposures. These estimates do not highlight the unequal distribution of attributable disease burdens within countries, such as within England, as a result of geographic variations in noise exposures and underlying disease incidence and mortality.

Using a more granular and geographically flexible approach, our study quantified, compared, and mapped the burden of annoyance (highly annoyed), sleep disturbance (highly sleep disturbed), IHD, stroke, and diabetes attributable to transportation noise exposures within England at the national, regional, and local authority level. Specifically, we estimated Population Attributable Fractions (PAF), the number and percentage of the population highly annoyed and sleep disturbed, and DALYs lost, in the adult population in England ($n = 42,738,500$) in 2018 due to long-term average noise exposures ≥ 50 dB L_{den} and L_{night} derived from strategic noise maps of road, rail, and

aircraft sources.

2. Material and Methods

2.1. Environmental burden of disease approach

We calculated the burden of disease attributable to noise exposures from road, rail, and civil airports in England at the national, regional, and local level using established epidemiological methodologies (Murray et al. 2003). To compare the burden of disease across noise sources and health outcomes, we used a standard metric, the Disability Adjusted Life Year (DALY). Key pieces of information needed for these calculations were:

- Noise exposures and population distributions
- Population data on disease occurrence, mortality rates, and life expectancy
- Disability weights (i.e., weighting factor that reflects the relative severity of a disease/health state)
- Exposure-response relationships (ERR) between health outcomes and noise exposures

Due to limitations in the availability of public data on exposure, demographic, and health at the spatial resolution required for our assessment, we made several assumptions on temporal stability, which are discussed in more detail in the following sections. Our estimates are based on population demographic and health data for the year 2018, and population exposure distributions based on noise mapping carried out for the year 2012.

2.2. Geography and population

We conducted the analysis in England and looked at trends at the national, regional, and local authority level. Geographic boundaries for nine regions and 314 Local Authority Districts (LAD) were obtained from the UK Office of National Statistics (licensed under the Open Government Licence v.3.0; contains OS data © Crown copyright and database right) (ONS Geography 2021) (Fig. 1). Regional populations ranged in size from ~ 2.5 to 9 million people, while LAD populations ranged in size from ~ 2.2 thousand to 1.1 million people. As the majority of the evidence of the health effects of noise is from cohort studies with adult subjects, we limited the analysis to the adult population normally resident in England (20 + years, $n = 42,738,500$) in 2018 (NOMIS 2019).

2.3. Road, railway, and aircraft traffic noise exposure data

We utilized the 2012 noise exposure datasets for road, rail, and aircraft sources generated by the 'Round 2 (II)' strategic noise mapping to fulfil the requirements of the Environmental Noise (England) Regulations 2006 (ENR) (HM Government, 2006). For road traffic, exposure was assessed via a single national exposure dataset produced by combining:

- exposure to major sources (defined as a trunk road, or a motorway, or a principal or classified road that has more than three million vehicle passages a year), for receptors outside agglomerations (DEFRA, 2014a), with
- exposure to all motorways and classified (A roads) roads, for receptors living within agglomerations.

The datasets also included noise exposure from:

- Major railways: railways that record >30,000 train passages per year.




Fig. 1. Regions and Local Authority Districts (LADs) in England. Local Authority District boundaries (2020) were sourced from the Office for National Statistics Open Geography portalx (ONS Geography 2021) licensed under the Open Government Licence v.3.0 (contains OS data © Crown copyright and database right 2022). The geographic boundaries for the United Kingdom, England, and Regions within England were sourced from the Ordnance Survey website (Contains OS data © Crown copyright and database right 2022) (Ordnance Survey 2022).

- Major airports: civil airports, which record >50,000 movements per year (a movement being a take-off or a landing), excluding those purely for training purposes on light aircraft.

Modelled noise exposures are 'A' Frequency Weighted. The 'A' weighting is a standard weighting of the audible frequencies designed to reflect the response of the human ear to noise (weighted between 20 Hz and 20 kHz). L_{den} , also known as the day-evening-night noise indicator, is the annual average A-weighted equivalent noise level (L_{eq}) over a whole day, but with a penalty of + 10 dB for night-time noise (23:00–07:00) and + 5 dB for evening noise (19:00–23:00). L_{night} is the annual average A-weighted equivalent noise level over the 8-hour night period of 23:00 to 07:00 h, also known as the night noise indicator (WHO 2018).

The 'Round 2 (II)' strategic noise maps of England and the associated residential population exposure assessment, were produced by Extrium environmental consultants under contract to the UK Department for Environment, Food & Rural Affairs (DEFRA). The 'Round 2 (II)' noise mapping data production and exposure assessment process is known to be based on input data which had a high degree of currency consistency. This included road and rail transport movements; Ordnance Survey topographic data, e.g., the locations of buildings, noise barriers and transport infrastructure; Ordnance Survey address data and the 2011 Census. Source specific exposures were modelled for the façade of a building that faces the nearest major noise source, which is theoretically the point of maximum exposure. This was undertaken by intersecting residential building data features (vector polygons) with noise levels which are structured in 10 m × 10 m grid datasets. The maximum noise

grid level is then attributed to each residential building and ultimately to the people assigned to each residential building following the residential population distribution process. Population level exposures were then calculated for the 2011 Census Output Area (COA) communities, which are typically formed of 300 residents living in 125 households. Residential buildings were identified from postal addresses, categorised into 1 dBA increments of L_{den} and L_{night} noise levels ranging from 50 dB to ≥ 80 dB, and then population counts were evenly assigned to the households. In 2011, there were 171,372 COAs in England that perfectly nested into the 314 Local Authority Districts operating in 2020. The COA population counts by exposure bands were then aggregated to form population exposure distributions (%) by LAD.

We then estimated LAD population exposures to noise in 2018 by applying the noise mapping exposure distributions to population counts recorded in 2018 by LAD. Doing this, we assumed that (i) the contribution and spatial distribution of major noise sources remained temporally stable between 2012 and 2018, and (ii) exposure distributions were the same for all age groups. These assumptions of uniform internal population dynamics were imposed by the decision to use the 2012 'Round 2 (II)' rather than the 2017 'Round 3 (III)' strategic noise maps of England. Furthermore, by using the 'Round 2 (II)' noise level data and the exposure assessment model, we had access to the underlying noise exposures in 1 dB bands and data starting consistently at 50 dB for all noise indicators. END exposure data in the public domain a) is in 5 dB bands, and b) starts from 55 dB L_{den} . Whilst road and railway noise exposure distributions are expected to be relatively stable over a period of six years, our assumptions may have introduced uncertainties for the aircraft noise estimates, due to growth in the aviation sector (Department for Transport 2022), replacement of older noisier aircraft types with more modern quieter ones (Civil Aviation Authority 2021c) and potential changes to flightpaths (Civil Aviation Authority 2021b) during this period. The population data was sourced from NOMIS, the Official Labour Market Statistics service provided by the Office for National Statistics (ONS) (NOMIS 2019) and was based on adult persons normally resident in English Local Authority Districts in 2018.

2.4. Health outcomes

Following a systematic review of reviews (Chen et al. 2022), we selected health outcomes based on the strength of the epidemiological and mechanistic evidence (Eriksson and Pershagen 2018) of the health effects of noise. The health outcomes included: annoyance (highly annoyed (HA)) (Fenech et al. 2022; Guski et al. 2017), sleep disturbance (highly sleep disturbed (HSD)) (Basner and McGuire 2018; Smith et al. 2022), IHD (also referred to as coronary heart disease) (van Kempen et al. 2018; Vienneau et al. 2019), stroke (Roswall et al. 2021; van Kempen et al. 2018), and diabetes (Sakhvidi et al. 2018; Vienneau et al. 2019). We did not quantify cognitive impairment in children (Clark et al. 2021) as our study focused on the burden of disease among adults.

2.4.1. Primary exposure response relationships (ERRs)

In cases where there were more than one recent systematic review/meta-analysis proposing an ERR for a health outcome and exposure pair, we considered the chronology of the publication and data (preference given to reviews with the most up-to-date evidence) and whether the evidence came from a published peer-reviewed paper versus a conference paper (preference given to peer-reviewed publications). We also only considered relative risks (RR) that were statistically significant (95 % confidence intervals around the central estimate did not cross zero) and which were associated with a specific traffic source (road, rail, or aircraft as opposed to 'total noise'). As such we selected ERRs for IHD and stroke from the WHO-commissioned systematic review by van Kempen et al (van Kempen et al. 2018) which considered studies published up until 2015, and for diabetes by Sakhvidi et al (Sakhvidi et al. 2018) which considered studies published up until 2017. It should be noted that the ERR for incident stroke in van Kempen et al is derived

from only one large cohort study, and so as described in Section 2.4.2, we conduct a sensitivity analysis with an alternative ERR derived from pooling multiple cohorts from Scandinavian countries (Roswall et al. 2021). We used ERRs for sleep disturbance (HSD) from Smith et al (Smith et al. 2022), which updated the WHO-commissioned systematic review (Basner and McGuire 2018) by including studies published until 2021. Finally we used ERRs for annoyance (HA) from the WHO-commissioned systematic review by Guski et al (Guski et al. 2017) which included studies published up until 2014. Within their review, Guski et al identified a subgroup of studies from Asia and the Alpine valleys in Austria with several study characteristics that could influence the results, including the range of noise exposure levels, housing characteristics linked to ventilation and air conditioning, geographical terrain, and the annoyance scale cut-offs used to determine the highly annoyed category. As such, they proposed two ERRs for road-traffic noise: one using the full dataset of studies and another with Asian and Alpine studies excluded. For our study based in England, we used the ERR which excluded Asian and Alpine studies. ERRs can be found in Table 1.

Due to a lack of evidence and/or statistically significant relative risk estimates, we did not calculate attributable burden of disease for railway noise and IHD, stroke, and diabetes, as well as aircraft noise and stroke and diabetes. Though as the epidemiological evidence base develops and strengthens, it is possible that significant and robust relative risk estimates for these exposure-outcome pairs may be identified in the future.

2.4.2. Secondary ERRs for sensitivity analyses

We conducted a series of sensitivity analyses with alternative ERRs which either reflected an update to the evidence base or the consideration of study context and inclusion in ERRs. As such, we conducted the following sensitivity analyses:

- IHD and road-traffic noise: ERR proposed by Vienneau et al (Vienneau et al. 2019) of 1.02 [95 % CI: 1.00–1.04] RR increase per 10 dB L_{den} . Results were published in a conference paper and included studies published up until 2019.
- Stroke and road-traffic noise: ERR proposed by Roswall et al (Roswall et al. 2021) of 1.06 [95 % CI 1.03–1.08] Hazard Ratio (HR) per 10 dB L_{den} . Results were based on pooling data from nine large-scale Scandinavian cohorts (135,951 participants).
- Annoyance (HA) and road-traffic noise (a): ERR proposed by Guski et al (Guski et al. 2017) which used the full WHO dataset and included studies published up until 2014.
- Annoyance (HA) and road-traffic (b) and railway noise: ERRs proposed by Fenech et al (Fenech et al. 2022) which updates the evidence from the WHO-commissioned systematic reviews by including studies published up until 2022. Results are published in a conference paper.

2.5. Population attributable fractions (PAF)

To estimate the proportion of all new cases of disease, or other adverse condition, in a population that is attributable to a specific exposure (Calentano et al. 2019; Murray et al. 2003) we calculate Population Attributable Fractions (PAF). To calculate a PAF, the exposure distribution within a population, as well the relative risk of disease due to exposure, must be known. Therefore, we used Equation 1 to calculate PAFs for IHD, stroke, and diabetes due to noise exposure for the adult population within each LAD in 2018:

$$PAF = \frac{\sum_{i=1}^n p_i \times (RR_i - 1)}{\sum_{i=1}^n p_i \times (RR_i - 1) + 1}$$

Where i represents a noise level in 1 dB increments; n is the total number of noise levels within the defined valid range for the PAF calculation; p_i represents the proportion of the population exposed to

Table 1
Parameters used to estimate the burden of disease attributable to transportation noise for the adult population in England in 2018.

	Health outcome	Noise metric	ERR source	ERR function/relative risk estimate [95 % confidence interval]	ERR lower	ERR upper	Disability weight [95 % confidence interval if available]
Road	Highly annoyed	L _{den}	(Guski et al. 2017) *	%HA = 116.4304–4.7342 × L _{den} + 0.0497 × L _{den} ² <i>Excluding Asian and Alpine studies</i>	40 dB	80 dB	0.02 (WHO Regional Office for Europe 2011)
	Highly sleep disturbed	L _{night}	(Smith et al. 2022) **	%HSD = 31.18323 – 1.47351 × L _{night} + 0.01851 × L _{night} ²	40 dB	65 dB	0.07 (WHO 2009)
	Ischemic heart disease	L _{den}	(van Kempen et al. 2018)	1.08 [1.01–1.15] per 10 dB	53 dB	80 dB	0.405 (WHO 2018)
	Stroke	L _{den}	(van Kempen et al. 2018)	1.14 [1.03–1.25] per 10 dB	50 dB	70 dB	0.522 [0.377–0.707] (Salomon et al. 2015)
	Diabetes mellitus	L _{den}	(Sakhvidi et al. 2018)	1.07 [1.02–1.12] per 5 dB	50 dB	80 dB	0.049 [0.031–0.072] (Global Burden of Disease Collaborators 2017)
Railway	Highly annoyed	L _{den}	(Guski et al. 2017)	%HA = 38.1596 – 2.05538 × L _{den} + 0.0285 × L _{den} ²	40 dB	85 dB	0.02 (WHO Regional Office for Europe 2011)
	Highly sleep disturbed	L _{night}	(Smith et al. 2022) **	%HSD = 63.56140 – 3.00711 × L _{night} + 0.03717 × L _{night} ²	40 dB	65 dB	0.07 (WHO 2009)
Aircraft	Highly annoyed	L _{den}	(Guski et al. 2017)	%HA = – 50.9693 + 1.0168 × L _{den} + 0.0072 × L _{den} ²	40 dB	75 dB	0.02 (WHO Regional Office for Europe 2011)
	Highly sleep disturbed	L _{night}	(Smith et al. 2022) **	%HSD = 17.07421 – 1.12624 × L _{night} + 0.02502 × L _{night} ²	40 dB	65 dB	0.07 (WHO 2009)
	Ischemic heart disease	L _{den}	(van Kempen et al. 2018)	1.09 [1.04–1.15] per 10 dBA	47 dB	75 dB	0.405 (WHO 2018)

ERR: Exposure response relationship; ERR lower: Lowest noise level at which the ERR is considered valid; ERR upper: Highest noise level above which the risk stays constant.

*WHO commissioned systematic review derived two ERR curves for highly annoyed due to road-traffic noise exposure. One curve utilizing the full WHO dataset and another excluded Asian and Alpine studies (Guski et al. 2017).

**Smith et al presented multiple curves for HSD. We used the ‘combined estimate’ where noise was explicitly mentioned in the question (Smith et al. 2022).

noise level *i*; and *RR_i* is the relative risk increase in the health outcome at noise level *i*. As relative risks in the noise epidemiological literature are often represented as 10 dB or 5 dB increment increases in L_{den} or L_{night}, we first scaled our literature informed relative risks (RR) to 1 dB noise increases assuming a linear relationship. In the absence of a consensus for the theoretical minimum risk exposure level associated with cardiometabolic diseases, for each exposure-outcome pair we assigned relative risk increases starting from a lower noise threshold level based on the noise ranges reported within each review/meta-analysis, or from the information contained in the individual studies within each review (Table 1). This approach ensured that we do not extrapolate relationships outside their validity range. For IHD, we assigned the lower ERR threshold to reflect the weighted average of the lowest noise levels measured in the studies, based on WHO guidance (WHO 2018). For the stroke ERR, 50 dB L_{den} reflects the noise level at the bottom 5th percentile (rounded from 49 to 50 dB L_{den}) within the cohort study (Sorensen et al. 2011; Sorensen et al. 2014) used in the WHO commissioned systematic review (van Kempen et al. 2018). For the diabetes ERR, 50 dB L_{den} reflects the noise level at the bottom 5th percentile (rounded from 49 to 50 dB L_{den}) within the study that had the majority weight (Sorensen et al. 2013) in the meta-analysis by (Sakhvidi et al. 2018). If the assigned lower ERR threshold started below 50 dB L_{den} or L_{night} (i.e., the noise level cut-off of our exposure data (see Section 2.3)), then the burden of disease attributable to exposures between that level and 49 dB would not be captured in the results. Furthermore, our noise exposure data had an upper threshold of 80 dB L_{den} and L_{night}; if actual exposures exceeded this limit, they were capped at 80 dB.

2.6. Percentage of the population highly annoyed and sleep disturbed

The percentage of the population highly annoyed and highly sleep disturbed within LADs was estimated directly from the quadratic exposure–response function equations provided by Smith et al and Guski et al (Guski et al. 2017; Smith et al. 2022). We calculated the number of

highly annoyed and highly sleep disturbed adults by multiplying the number of adults within each 1 dB noise band above 50 dB L_{den} (HA) and L_{night} (HSD) by the percentage of highly annoyed and highly sleep disturbed individuals at the corresponding noise level.

2.7. Burden of disease

Our main measure of disease burden was the Disability Adjusted Life Year (DALY) (Equation 2). The DALY simultaneously considers the reduced health state due to disability before death (Years of Life Lived with Disability (YLD)) and the decline in life expectancy due to death (Years of Life Lost (YLL)). We estimated DALYs for LADs represented as total DALYs as well as DALY rates per 100,000 people to adjust for local population sizes.

$$DALY = YLL + YLD \quad (2).$$

We calculated YLL from IHD, stroke, and diabetes separately for males and females and by 5-year age bands within each LAD by multiplying the number of disease-specific mortalities by the life expectancy at age of death. We then aggregated the age and gender specific estimates to achieve a single LAD-level estimate. We sourced a 3-year average life expectancy LAD dataset from the UK Office for National Statistics for the years 2017–2019 (Office for National Statistics, 2020a). Life expectancy information was not available for the City of London. Therefore, regional life expectancies for London were used for this LAD. We estimated the number of disease-specific mortalities by combining data on annual mortality counts for the year 2018 (Office for National Statistics 2019) and disease-specific mortality fractions. Local records of mortalities by their primary cause were not publicly available beyond 2013; therefore, the fraction of mortalities for each health outcome were calculated over a 5-year period from 2009 to 13 (Office for National Statistics 2015). We defined the underlying cause of death using codes provided by the International Classification of Diseases Tenth Revision (ICD-10) (Appendix A).

We calculated the YLD from IHD, stroke, and diabetes within each LAD by multiplying the annual prevalence rates of disease by the disease-specific disability weights (Kim et al. 2022; WHO 2020). Similar to the WHO approach, we calculated prevalence-based YLD (WHO 2020) and therefore did not include information on average duration of disease into the calculation. Disease prevalence rates come from the Quality and Outcomes Framework reporting (NHS Digital 2022), which was sourced from the UK Office for Health Improvement and Disparities 'Fingertips' service (Office for Health Improvement & Disparities 2022). Prevalence records were obtained for 191 Clinical Commissioning Groups (CCG) in England, the highest spatial resolution of data that is currently available. CCG datasets were resampled to 314 LADs using population-weighted interpolation, this approach calculates average rates based on the intersections of population counts at postcode points and the two administrative boundaries. Prevalence data for the years 2011–2018 were additionally averaged to minimise the influence of any unusual temporal anomalies. Disability weights allow non-fatal disability as a result of disease/health states to be measured under a common unit, ranging between 0 (no disability) and 1 (full disability). The disability weights that we used for IHD (DW: 0.405), annoyance (DW: 0.02) and sleep disturbance (DW: 0.07) were published and used in WHO noise guidelines (WHO 2009; WHO Regional Office for Europe 2011), and additionally used to estimate noise attributable disease burdens in Europe (EEA, 2020) (Table 1). As used in the Global Burden of Disease studies, we used a DW for stroke relating to long-term consequences of severe cases (DW: 0.522) and for diabetes we used a DW relating to uncomplicated cases (0.049) (Global Burden of Disease Collaborators 2017; Salomon et al. 2015). Alternative DWs for each health outcome were considered and the impact on DALYs was quantified in sensitivity analyses (National Institute for Public Health and the Environment (RIVM) 2018; Salomon et al. 2015) (Appendix A).

2.8. Noise attributable burden of disease

As a final step, the noise attributable DALYs were estimated by multiplying the PAF (i.e., disease attribution to noise) by the DALY (i.e., disease burden). The noise attributable DALYs were estimated with respect to each noise source (road, railway, aircraft) and health outcome separately at the LAD level. The calculation procedure for annoyance and sleep disturbance followed a slightly different process, whereby the number of adults highly annoyed and sleep disturbed were multiplied by the corresponding disability weight to estimate the attributable DALYs. For annoyance and sleep disturbance, only the impacts on morbidity contribute to the DALY. We estimated attributable DALYs at the LAD level, and then aggregated up to the regional and national levels to provide results at varying geographical scales.

2.9. Sensitivity analysis of alternative road-traffic noise exposure distributions: Case-study applied to the London region

Our main analysis quantified the attributable burden of disease for the adult population in England exposed above 50 dB L_{den} and L_{night} from road, railway, and aviation (commercial airports) sources, based on noise modelling conducted to fulfil the ENR strategic noise mapping requirements. However, we are likely under-estimating the attributable burden of disease due to the (i) restricted exposure range (>50 dB) and because (ii) exposures from "minor" roads (i.e., collector, residential and local access roads) are not counted. Many collector roads can still carry significant volumes of traffic, meaning that a) dwellings exposed to arterial and busy collector roads will experience higher exposure than that indicated by strategic noise mapping, and b) dwellings only exposed to collector and local roads could have exposures within the exposure range (50 – 80 + dB) but may be excluded from strategic noise mapping estimates. Furthermore, the 50 dB threshold does not account for low-level exposures for some of the population living in relatively quiet areas. Therefore, we conducted a sensitivity analysis where the

attributable burden of disease was estimated with an alternate exposure distribution derived from all road sources (CNOSSOS-all roads model) and with a wider exposure range, starting at 40 dB L_{den} and 35 dB L_{night} . We conducted this analysis for the London region only, due to the availability of data. The exposure estimates were created by the Centre for Environmental Health and Sustainability (CEHS) at the University of Leicester, using a road-transport noise model for 2013 in accordance to the European Commissions 'Common framework for noise assessment' (CNOSSOS) (Kephalopoulos et al. 2012).

The CNOSSOS noise propagation algorithms were implemented in PostgreSQL via the PostGIS v2.1 extension, following the protocol described by Gulliver et al. (Gulliver et al. 2015). Annual Average Daily Traffic (AADT) counts and traffic speeds are considered along with information relating to the surface roughness of land cover, building heights, wind profiles and annual average temperatures. Roads are divided into 10 m segments and ray paths are drawn to the receptor locations accounting for the angle of view, source distance, and façade reflections. The AADT counts used here are available for the entire UK road-network in 2013 and were modelled by (Morley and Gulliver 2016). The noise contributions from major (motorways and a-roads) roads within 1 km and minor roads within 100 m of the receptor location are modelled as separate components. The estimated noise level at each receptor is the sum of sound propagation from every road in the network, including all public-accessible minor and local roads. The CNOSSOS-all roads model was run for 140,793 address locations in London, which were selected by assigning population-weighted postcode centroids to the nearest building. The 33 LADs in London typically contain 4,266 postcodes (SD = 1,355), which on average house 58 residents (SD = 44). Noise levels are modelled at the loudest façade, identified by calculating the AADT count of the nearest road inverse to the roads distance.

2.10. Software

We used Microsoft Excel workbooks (Version 2022) for the burden of disease calculations, the open-source statistical computing language and environment R (R Core Team 2022) for generating figures/plots, and ArcGIS software by Esri (Version 10.5.1) for mapping.

3. Results

3.1. Population noise exposures in England

40 % of the adult population in England were exposed to road-traffic noise that exceeded 50 dB L_{den} , however, there was a large degree of spatial variation across LAD (Fig. 2). 4.8 % of the population was exposed to aircraft noise above 50 dB L_{den} , though the spatial distribution was particularly skewed as 95 % of LADs had less than 20 % of populations exposed. The highest exposures to aircraft noise from major airports were found near London. Lastly, 4.5 % of the population in England were exposed to railway noise from mainlines that exceeded 50 dB L_{den} , though there was less variation between LADs (Range: 0 – 21 %), compared with road-traffic and aircraft exposures. There were similar variations and spatial distributions for night-time noise (Appendix A). We also found that 27 % and 2.9 % of the adult population were exposed to road and railway noise above the WHO guideline levels of 53 dB L_{den} (road) and 54 dB L_{den} (railway), respectively. However, we could not estimate the % of the population exposed above the L_{night} guideline levels for all sources (road: 45 dB; rail: 45 dB; aircraft 40), or for L_{den} aircraft noise (45 dB), as the guideline levels fall below the lowest exposure threshold of our data (50 L_{den} and 50 L_{night} dB).

3.2. Noise attributable burden of disease in England

Road-traffic accounted for the majority of the DALYs lost in England in 2018, accounting for close to 97,000 DALYs/yr (Table 2). This was




Fig. 2. Spatial variation in the percentage (%) of the population exposed to road-traffic, railway, and aircraft noise from major sources above 50 dB (L_{den}) across local Authority Districts (LADs) in England, based on strategic noise mapping carried out in 2012.

Table 2
Attributable burden of annoyance, sleep disturbance, ischemic heart disease, stroke, and diabetes mellitus due to road-traffic, railway, and aircraft noise exposures above 50 dB L_{den} and L_{night} in England. Estimates are for the adult population (20+) in 2018.

	Road-traffic Central estimate [95 % confidence interval]*	Railway traffic Central estimate [95 % confidence interval]*	Aircraft traffic Central estimate [95 % confidence interval]*
Highly annoyed (HA)			
% of population	3.9 %	0.7 %	1.2 %
Number of people	1,662,157	295,766	524,321
Total DALYs/yr	33,243	5,915	10,486
DALYs per 100,000 people/yr	78	14	25
Highly sleep disturbed (HSD)			
% of population	0.9 % [0.6 % – 1.2 %]	0.2 % [0.2 % – 0.3 %]	0.2 % [0.1 % – 0.2 %]
Number of people	382,333 [236,040 – 521,004]	101,815 [66,788 – 127,132]	65,455 [47,155 – 83,625]
Total DALYs/yr	26,763 [16,523 – 36,470]	7,127 [4,675 – 8,899]	4,582 [3,301 – 5,854]
DALYs per 100,000 people/yr	63 [39–85]	17 [11–21]	11 [8–14]
Ischemic heart disease (IHD)			
PAF (%)	1.5 % [0.2 % – 2.7 %]	–	0.2 % [0.1 % – 0.4 %]
Total DALYs/yr [95 % CI]	11,556 [1,427–21,942]	–	1,970 [876–3,273]
DALYs per 100,000 people/yr [95 % CI]	27 [3–51]	–	5 [2–8]
Stroke			
PAF (%)	3.8 % [0.8 % – 6.6 %]	–	–
Total DALYs/yr [95 % CI]	18,592 [2,926–41,093]	–	–
DALYs per 100,000 people/yr [95 % CI]	44 [7–96]	–	–
Diabetes Mellitus			
PAF (%)	4.2 % [1.2 % – 7.2 %]	–	–
Total DALYs/yr [95 % CI]	6,686 [1,291–16,301]	–	–
DALYs per 100,000 people/yr [95 % CI]	16 [3–38]	–	–

Total DALYs/yr: Total number of Disability Adjusted Life Years lost per year; YLL: Years of Life Lost; YLD: Years of Life Lived with Disability; PAF (%): Population Attributable Fraction percentage (%); % of pop: Percentage of the population.

* The 95 % confidence intervals (CI) around the central burden of disease estimates of IHD, stroke, and diabetes were based on the combined uncertainty reported for the ERR functions, disease prevalence, disability weights, and life expectancy. While the 95 % CIs around the central estimate for sleep disturbance was based solely on the uncertainty estimate of the ERR function as the corresponding disability weight did not have a 95 % CI. We did not have uncertainty estimates for the ERRs or disability weights to be able to construct a 95 % confidence interval around the central burden of disease estimate for annoyance.

followed by aircraft (~17,000/yr) and then railway (~13,000/yr) noise exposures. Across health outcomes, annoyance accounted for the largest number of total DALYs/yr from road-traffic (33,243/yr) and aircraft (10,486/yr) exposures, while sleep disturbance accounted for the most from railway noise exposures (7,127/yr) (Table 2). From road-traffic noise exposures, we estimated 18,592 attributable DALYs/yr lost from strokes, 11,556 from IHDs, and 6,686 from diabetes. The attributable DALYs for stroke were higher than for IHD due to a combination of the magnitude of the relative risk estimates and the higher disability weight. The lower number of attributable DALYs for diabetes compared with IHD and stroke was driven by a lower disability weight (IHD: 0.405; stroke: 0.522; diabetes: 0.049) and annual mortality rates in the population (comparative statistics can be found at (British Heart Foundation, 2020)).

When we tested the sensitivity of our estimates for a range of disability weights, we found that the total DALYs/yr for road-traffic noise varied between 7,040 – 11,556 for IHDs, 5,043 – 18,592 for strokes, and 2,763 – 13,003 for diabetes. By applying alternative disability weights for annoyance and sleep disturbance (National Institute for Public Health and the Environment (RIVM) 2018), the attributable DALYs/yr were reduced to 16,622 and 6,691, respectively. See Appendix A for more information.

3.3. Noise attributable burden of disease at the regional and local authority level

London had the greatest number of DALYs/yr lost attributable to road (~20,000/yr), railway (~5,000/yr) and aircraft (~11,000/yr) noise exposures compared with all other regions in England (Fig. 3 - upper panel). In fact, London had six times greater road-traffic DALYs

compared to the region with the lowest levels (North East). The regional differences were particularly marked for aircraft noise exposures as the London region accounted for 63 % of all aircraft noise DALYs in England. The South East and North West of England also had relatively high number of total attributable DALYs/yr due to road, railway, and aircraft noise, while the North East had the lowest, compared with other regions.

When the influence of population size was removed from the road-traffic attributable DALYs by expressing them as rates per 100,00 people (Appendix A - Table A7), London still had the largest burden for annoyance (116 per 100,000 people) and sleep disturbance (101 per 100,000 people). However, the DALY rates for London were on par with the North West with respect to diabetes (London: 21, North West: 21) and in fact lower than the North West for stroke (London: 39, North West: 63) and IHD (London: 28, North West: 43). Not only did the absolute number of DALYs, and the rates, vary between regions, but so too did the proportional contributions of each health outcome to the total DALYs within each region. Notably, the proportional contribution of road-traffic IHD and stroke DALYs to the total DALYs within London was noticeably lower than the proportional contributions within other regions (Fig. 3 - lower panel).

There was a high-degree of variation in the DALY rates (per 100,000 people/yr) attributable to road-traffic noise exposure across LADs for annoyance (Interquartile range (IQR): 39–139), sleep disturbance (IQR: 29–109), IHD (IQR: 15–46), stroke (IQR: 28–75), and diabetes (IQR: 8–28). This variation also showed distinct spatial patterns across the country. We found the highest DALY rates clustered within and around London, the North West, and in the eastern part of the West Midlands (Fig. 4). For railway noise, the highest DALY rates were observed in local authorities within and around London, and those located along the major North-South railway routes in England (Fig. 5). For aircraft noise,




Fig. 3. Attributable Disability Adjusted Life Years (DALYs) lost due to road, railway, and aircraft traffic noise exposures above 50 dB across regions in England. Estimates are for the adult population (20 +) in 2018. Note that no robust ERRs were identified for railway noise and IHD, stroke, and diabetes, and for aircraft noise and stroke and diabetes.

Attributable DALYs due to road traffic noise exposure




Fig. 4. Attributable Disability Adjusted Life Years (DALY) lost per 100,000 people/yr due to road-traffic noise exposures above 50 dB within Local Authority Districts (LAD) in England. Estimates are for the adult population (20+) in 2018. Inset map is Greater London.

local authorities within and around London had by far the highest DALY rates. In particular Hounslow and Richmond near Heathrow airport (Fig. 6). DALYs shown by noise source, health outcome, and LAD are given in Appendix B, C and D.

3.4. Sensitivity analysis of alternate exposure response relationships (ERR)

The estimated burden of disease from IHDs and strokes attributable to road-traffic noise exposure was very minimally impacted by the application of alternative ERRs proposed by Roswall et al (stroke) and Vienneau et al (IHD) (Roswall et al. 2021; Vienneau et al. 2019) (details can be found in the Appendix A). We did, however, find that the attributable DALYs from annoyance (HA) was impacted by the choice of ERR. By applying alternative road-traffic ERRs, we found that the number of DALYs/yr were 1.4x (Guski et al. 2017) and 1.3x (Fenech et al. 2022) times higher than our main estimate. For railway noise, the

alternative ERR for annoyance resulted in a smaller difference (1.15x times higher than our main estimate) (details in Appendix A).

3.5. Sensitivity analysis of an alternative noise exposure distribution for London

As a sensitivity analysis for the noise exposure modelling, the attributable burden of disease for London was estimated using an alternative noise exposure distribution calculated by the *CNOSSOS-All roads* model (see Section 2.9 for details on model). *CNOSSOS-All roads* uses estimates of traffic flows/speeds from all roads and provides data down to a lower exposure threshold (40 dB L_{den} , 35 dB L_{night}) compared with the England-wide *ENR strategic mapping*, as illustrated in Fig. 7.

Within London we estimated a higher number of attributable DALYs/yr with *CNOSSOS-All roads* compared with *ENR strategic mapping*. The relative increase in attributable DALYs/yr varied between health outcomes, ranging from 1.1x (IHD) to 2.2x (HSD) times higher (Table 3).

Attributable DALYs due to railway noise exposure

Highly annoyed (HA)

Highly sleep disturbed (HSD)




Fig. 5. Attributable Disability Adjusted Life Years (DALY) lost per 100,000 people/yr due to railway traffic noise exposures above 50 dB within Local Authority Districts (LAD) in England. Estimates are for the adult population (20 +) in 2018. Inset map is Greater London.

Sleep disturbance had the highest relative increase in DALYs compared with other health outcomes, likely because of the following factors: the lower ERR noise threshold (40 dB); the shape and shift in the L_{night} exposure distribution; and because the lower noise cut-off for L_{night} differed more between the models (35 to 50 dB L_{night}) compared with L_{den} (40 to 50 dB L_{den}). Based on these results, we presume that we are likely underestimating the road-traffic noise attributable burden of disease in England using exposure modelling from strategic noise mapping.

4. Discussion

While European studies have shown that environmental noise is a major contributor to the overall environmental disease burden (Hanninen et al., 2014; EEA, 2020; WHO Regional Office for Europe, 2011), detailed spatial assessments within countries are still lacking. Using recent and scientifically robust epidemiological evidence, we quantified, compared, and mapped the burden of annoyance, sleep disturbance, IHD, stroke, and diabetes attributable to long-term transportation noise exposures for adults in England at the national, regional, and local authority level. We estimated that overall, approximately a hundred thousand (~97,000) DALYs were lost in England in 2018 due to road-traffic, ~13,000 from railway, and ~17,000 from aircraft long-term averaged noise exposures above 50 dB. The magnitude of the attributable disease burden varied significantly across regions and LADs.

Compared to Global Burden of Disease study estimates of other types of pollution in England in 2018, our road noise attributable DALYs (~97,000 DALYs) were lower than for ambient $\text{PM}_{2.5}$ pollution (~269,000 DALYs), but higher than ambient ozone (~7,000 DALYs) and occupational noise (~29,000 DALYs) (Global Burden of Disease Collaborative Network 2020). However, we note these comparisons between risk factors should be made with caution due to differences in methodologies, assumptions, and the state of the evidence. Furthermore, whilst our estimates of the overall noise attributable burden of disease was significant, it may still be under-estimated in England due to limitations in modelling noise exposures designed specifically for strategic noise mapping (i.e., only including roads with very high traffic volumes and only modelling above 50 dB L_{den} and L_{night}). Our noise attributable DALY estimates at a national level for England were largely in line with similar estimates produced for the United Kingdom (England plus Scotland, Wales and Northern Ireland) by the European Environment

Agency (EEA) using END mapping (EEA, 2020). The available public information from the EEA does not provide information on within-country variations, so comparisons at a regional and local district level were not possible.

One of the main drivers of spatial variations in noise attributable disease burdens is non-uniform exposure to noise. This is particularly evident for aviation, as we found that London had 63 % of all attributable aircraft noise DALYs in England. Between 2012 and 2019, air traffic at Heathrow, Gatwick and London City was approximately 40 % of the total UK air traffic (Department for Transport 2022). Variations in noise attributable disease burdens are also impacted by the distribution of the population. For example, the London region had the highest total attributable DALYs due to road-traffic noise, but accounting for population size, the rates of road-traffic attributable DALYs (per 100,000 people) in the North West region were on par with London. Furthermore, spatial variations in local disease burden (British Heart Foundation, 2021b; Steel et al. 2018) also contribute to spatial contrasts of noise attributable DALYs across England. This is evident in the fact that the proportional contribution of road-traffic IHD and stroke DALYs to the total DALYs within London was noticeably lower than the proportional contributions within other regions. This is because the underlying prevalence and mortality rates of IHD and stroke were generally lower in London compared with many other areas in England (comparative statistics can be found at (British Heart Foundation. 2021a,b) (also see Tables A8 and A9 in Appendix A), possibly because of the younger age demographic in London (median age 35.3yrs in London compared with 39.5 – 43.9 in other regions) (Office for National Statistics, 2020b).

The comparative environmental burden of disease concept facilitates comparisons across geographies, health outcomes, and risk factors, but comparisons should be made with care when the methodological assumptions and data inputs are not aligned across studies. For example, the epidemiological evidence on noise and health is developing and strengthening (Clark et al. 2020; Fenech et al. 2022; Persson Waye and van Kempen 2021; van Kamp et al. 2020), and therefore choices about which health outcomes to include, and their associated exposure response relationships will vary by study, and this can affect the comparability between studies. For example, a noise burden of disease study for Hesse, Germany (Hegewald et al. 2021) included depressive disorders based on relative risk estimates from a large study conducted around Frankfurt Airport (Seidler et al. 2017). We excluded depression

Attributable DALYs due to aircraft noise exposure

Highly annoyed (HA)

Highly sleep disturbed (HSD)




Fig. 6. Attributable Disability Adjusted Life Years (DALY) lost per 100,000/yr people due to aircraft traffic noise exposures above 50 dB within Local Authority Districts (LAD) in England. Estimates are for the adult population (20+) in 2018. Inset map is Greater London.

and anxiety from our assessment, however given that the evidence base is developing, we recommend that these health outcomes are kept under review for future updates. We also recommend that the shape of the exposure–response relationships for all health outcomes, and any potential threshold effects which would influence the choice of a counterfactual, also remain under continual review. There is limited but growing evidence of the shape of the exposure–response relationships and whether any potential thresholds of effect exist (Fu et al. 2022; Thacher et al. 2022; Vienneau et al. 2022). Another example is that the WHO-commissioned systematic review for annoyance from road traffic proposed two exposure response relationships (ERRs): a global one and one excluding Asian and Alpine studies. For our main analysis we used the latter, as did the aforementioned study in Hesse, Germany (Hege-wald et al. 2020), whereas other European studies used the global ERR (Murphy and Faulkner, 2022; EEA, 2020). Within England, we found

that the two curves led to a relative 1.4x difference in attributable road-traffic annoyance DALYs/yr. Another source for discrepancy is that some studies have applied ERRs from one noise source to another (e.g., using a road-traffic noise and IHD ERR for railway and aircraft exposures (EEA, 2020)) where information has been limited. We took a more conservative approach and only quantified the burden of disease if a source-specific statistically significant association was given in a review or meta-analysis of the epidemiological evidence. Our decision to use the same lower exposure cut-off of 50 dB for L_{den} and L_{night} from the strategic mapping means that the relative differences between the L_{den} -based YLD and the YLD due to sleep disturbance are not comparable to other assessments taking the more conventional approach of using a 5 dB lower cut-off for night-time noise. Lastly, similar to the WHO approach for DALY calculations (WHO 2020), we used disease prevalence estimates to derive YLD with respect to IHD, stroke, and diabetes,




Fig. 7. Distribution of the number of adults exposed to road-traffic noise levels (L_{den}) in London based on ENR *strategic mapping* and CNOSSOS – *All roads* modelling. Vertical blue line indicates the threshold at the 50 dB L_{den} band. The numbers on the Y-axis are in units of thousands.

as incidence-based data was not available at the geographic resolution of our study. This could impact comparability with studies using the incidence-based calculation approach (Park et al. 2019; von der Lippe et al. 2020; Wagner et al. 2015).

For our main analysis we used exposure estimates from strategic noise mapping carried out to satisfy the requirements of the Environmental Noise (England) Regulations 2006 (as amended) (implementation of the European Environmental Noise Directive in England). Although a third round of strategic mapping was undertaken in 2017, we used noise mapping data from the second round (2012) for reasons already described in Section 2.3. For road and railway noise, the spatial distribution of exposure is relatively stable over time. The population within the > 55 dB L_{den} contour in 2017 was approximately 10 % higher than in 2012 for road and railway noise (DEFRA, 2014b; 2019). This increase is likely to have been due to a combination of changes in the population exposed and changes in noise emissions taking place between 2012 and 2017. Our analysis includes the former effect but not the latter. For aircraft noise, airports and air traffic control organisations in the UK sometimes carry out short-term trials of changes to the airspace structure (and hence flight paths) (Civil Aviation Authority 2023). More generally, the ongoing recovery of the aviation sector from the impacts due to Covid, and the upcoming large-scale changes to UK airspace (Civil Aviation Authority 2018) mean that any retrospective estimates of spatial variations in disease burden attributable to aviation noise should not be considered representative of future years, and regular assessment updates may be warranted (subject to availability of exposure data and up to date ERRs).

The availability of strategic noise mapping data greatly facilitates burden of disease assessments at national, regional and local levels; the downside is that not all roads, railways and airports may be included in the modelling, and there may be issues with the chosen exposure thresholds giving an incomplete picture of population exposure in the range where health effects are known to occur. As we showed with a sensitivity analysis for London, these two factors are likely to lead to under-estimation of the attributable disease burden, and demonstrates the importance of modelling all roads, and using a lower threshold than 50 dB. A similar result, but with even greater differences, was found for the Hessian population in Germany (Hegewald et al. 2021). Hegewald et al compared estimated DALYs due to road-traffic noise using exposures from END strategic mapping (exposures ≥ 55 dB) and ‘PLUS-Mapping’ which included all roads and noise levels as low as 40 dB. They found that DALYs estimated with PLUS-Mapping for cardiovascular diseases and depressive disorders were $\sim 4.2x$ times higher than estimated with END strategic mapping. Although the absolute relative risk estimates for

cardiometaabolic health outcomes below 50 dB L_{den} can be low (van Kempen et al. 2018; WHO 2018), a large proportion of the population in England is likely to be exposed to road traffic noise within the 40 to 50 dB L_{den} range (DEFRA, 2000, 2013). Our sensitivity analysis in London used the CNOSSOS methodology (Kephalopoulos et al. 2012), whereas the strategic modelling used the UK’s national calculation methodology (Calculation of Road Traffic Noise (CRTN) (Department of Transport et al. 1988)). We have not assessed the differences between the approaches in terms of exposure estimates, but we expect this to not significantly impact results when aggregated to our minimum unit of assessment (local authority district).

The sensitivity analyses of the ERRs showed that there was minimal difference in attributable DALYs for IHD and stroke when using the main and alternative ERRs. This was ascribed to the fact that while our main relative risk estimates were larger in magnitude than the alternatives, the lower thresholds for the alternative ERRs were set lower down on the exposure range, at 40 dB L_{den} (informed by the valid noise ranges in the included studies) resulting in similar DALY estimates. When setting lower ERR thresholds, it is important to consider the valid noise range from which each ERR was derived as it can influence the magnitude of the slope coefficient but also so as not to extrapolate relationships out of range of the data. Additionally, we conducted these comparisons for the population exposed to noise above > 50 dB (L_{den}), though if populations exposed to lower noise levels were to be included in the estimations (i.e., between 40 and 50 dB), then we might expect a widening of these differences. This may be particularly relevant for health impacts associated with night-time exposures, such as sleep disturbance, as a larger proportion of the population is exposed within the 40–50 dB L_{night} range. Lastly, the choice of ERR for annoyance had an impact on attributable DALYs, as evidenced by the differences observed when applying the road-traffic curves using the full WHO dataset versus curves excluding Asian and Alpine studies (Fenech et al. 2022; Guski et al. 2017). The WHO ENG guidelines recommended that data and ERRs derived in a local context should be used whenever possible (WHO 2018). As there has been no recent large-scale socio-acoustic study for road and railway noise in the UK (last one was in 1984) (Fenech et al. 2022), we used the WHO aggregate curve excluding Asian and Alpine studies for the main analysis, but we conducted sensitivity analyses with the full WHO curves (original (Guski et al. 2017) and updated curves (Fenech et al. 2022)) to provide additional estimates and to be transparent about the impacts of our methodological decisions. A socio-acoustic study on attitudes to aviation noise was carried out in England in 2014–15, however an ERR expressed in terms of an annual L_{den} is not yet published from this study (Civil Aviation Authority 2021a).

Table 3
Comparison of the impact of modelled noise exposure distributions on the road-traffic attributable burden of disease in London.

Health outcome	Road-traffic: strategic mapping (ENR)*	Road-traffic: all roads (CNOSSOS-All roads)**	Relative increase (CNOSSOS-All roads compared with ENR)
	≥50 dBA L _{den} ≥50 dBA L _{night}	≥40 dBA L _{den} ≥35 dBA L _{night}	
Highly annoyed (HA)	% of population Number of adults Total DALYs/yr	5.8 % 388,389 7,768	8.0 % 534,467 10,689 1.38x higher
Highly sleep disturbed (HSD)	% of population Number of adults Total DALYs/yr	1.4 % 96,501 6,755	3.2 % 213,723 14,961 2.21x higher
Ischemic Heart Disease (IHD)	PAF Total DALYs/yr	2.3 % 1,886	2.5 % 2,057 1.09x higher
Stroke	PAF Total DALYs/yr	5.5 % 2,623	6.9 % 3,330 1.27x higher
Diabetes mellitus	PAF Total DALYs/yr	5.9 % 1,381	7.3 % 1,704 1.23x higher

*ENR: Noise exposures derived from modelling to fulfil the ENR Round 2 strategic mapping requirements (main analysis).

**CNOSSOS-All roads: Noise exposures created by the Centre for Environmental Health and Sustainability at the University of Leicester in accordance with CNOSSOS-EU modelling framework (Gulliver et al. 2015).

4.1. Strengths and limitations

While the majority of noise burden of disease studies derive estimates at a national or city-level, we generated high spatial resolution estimates across 314 Local Authority Districts across England. Our calculations took into account spatial variations in noise exposures, population distributions, and underlying disease prevalence and mortality across the country. We made estimates for a range of health outcomes, including annoyance, sleep disturbance, and ischemic heart disease. Given the strengthening of the epidemiological evidence, we also made estimates for strokes and diabetes, which have largely been uncharacterised in other studies (Eriksson et al., 2017; Khomenko et al., 2022; Murphy and Faulkner, 2022; EEA, 2020; WHO Regional Office for Europe, 2011). We utilised recent synthesised scientific evidence to derive our burden of disease estimates (Guski et al. 2017; Sakhvidi et al. 2018; Smith et al. 2022; van Kempen et al. 2018) and undertook a comprehensive set of sensitivity analyses to show the impact of using alternative exposure–response relationships and disability weights. Finally, we also showed that there can be a substantial impact (under-estimation) on the estimated attributable burden of disease from road-

traffic noise when the exposure range is limited and when traffic flows on collector or local roads are not taken into account in the modelling.

Our study has several potential limitations. We quantified the attributable burden of disease assuming the population was exposed to transportation noise sources in isolation. There are still uncertainties on the actual health burden in the event of co-exposure to multiple sources of transport noise, and there may be a risk of double counting. While the evidence is still limited, the EEA estimated the potential for double counting for annoyance and sleep disturbance from the combined effects of multiple sources in the 2020 report (EEA, 2020) to be a maximum of 13 % for annoyance, 16 % for sleep disturbance, and negligible for IHD, particularly if the PAFs are below 0.10 (Houthuijs et al. 2018). Conversely, a limited but growing evidence base is investigating whether the combined risks from multiple sources may be larger than the sum of the individual risks (Seidler et al. 2019; Thacher et al. 2021; Thacher et al. 2022). As the evidence base develops, future work should consider how to incorporate these potential combined effects into noise burden of disease assessments. Due to these current uncertainties, we recommend that our burden of disease estimates should not be combined across noise sources.

Due to a lack of evidence and/or statistically significant relative risk estimates, we did not calculate attributable burdens of disease for railway noise and IHD, stroke, and diabetes, as well as aircraft noise and stroke and diabetes. Though as the epidemiological evidence base develops and strengthens, it is possible that significant relative risk estimates for these exposure-outcome pairs may be identified in the future. We did approximate the total DALYs associated with each transportation source by summing across health outcomes, though this approach did not take into account comorbidities in the DALY calculation. Given the uncertainty in these estimates, we reported the total DALYs to the nearest thousand.

We used yearly IHD, stroke, and diabetes disease prevalence data which were based on general practice (GP) reporting within the National Health Service (NHS) Quality and Outcomes Framework (QOF). The QOF prevalence data were based on the number of patients recorded as having disease on the practice register over the total practice list size (Office for Health Improvement & Disparities 2022). These data are limited however, as people who have not been formally diagnosed would not be included in the estimate and it is also possible that there are variations across practices in how conditions are diagnosed (though this is likely to be random). Furthermore, the prevalence rates for IHD and stroke are given with respect to people of all ages registered with the GP, and Diabetes for people aged 17+, which means that we may be under-estimating the noise attributable burden of disease for adults (20 +) in England as the underlying disease prevalence rates include a population of children and adolescents who are at lower risk. Furthermore, we may be under-estimating the most in London as it has the lowest regional median age in England (Office for National Statistics, 2020b).

As discussed in Section 2.3, we applied population noise exposure distributions from strategic mapping undertaken in 2012 to the population distribution (i.e., population at risk) within LAD in 2018. We thus, assumed that the spatial distribution and magnitude of the noise exposure was stable between these time points. Furthermore, while the availability of strategic noise mapping data greatly facilitates burden of disease assessments at national, regional and local levels, the downside is that only major sources are captured and the lower exposure threshold of 50 dB L_{den} and L_{night} mean that we are likely under-estimating the total noise attributable burden of disease in England (as we showed in our Sensitivity Analysis in Section 3.5). Lastly, while people spend the majority of their time indoors, we were unable to account for geographical differences in the ingress of outdoor noise into indoor environments due to variations in building characteristics and ventilation practices across the country.

5. Conclusions

We produced the first geographical assessment of the burden of disease due to noise for England. Our study shows that transportation noise exposure, particularly from road-traffic sources, is responsible for a significant disease burden in England and this varies unequally across regions and LADs. Our work provides useful nationwide information for identifying areas with varying burdens of disease due to different noise sources and for setting priorities in environmental health research, policies, and interventions. Quantifying the disease burden is the first step in building an environmental, social and economic case for action that takes into account the views and expertise of national and local government, environmental health practitioners, acoustics specialists, industry, the third sector and affected citizens. The epidemiological evidence on noise and health continues to develop, and disease burden estimates are likely to change as more data from good quality longitudinal studies becomes available. Lastly, we showed that exposure data derived from strategic noise mapping models can lead to underestimates in the attributable burden of disease if not all transport sources are accounted for and if lower levels of noise exposure are omitted.

CRediT authorship contribution statement

Calvin Jephcote: Data curation, Formal analysis, Methodology, Software, Writing – original draft, Writing – review & editing. **Sierra N. Clark:** Data curation, Formal analysis, Visualization, Methodology, Writing – original draft, Writing – review & editing. **Anna L. Hansell:** Methodology, Investigation, Writing – review & editing. **Nigel Jones:** Conceptualization, Data curation, Project administration, Supervision, Writing – original draft, Writing – review & editing. **Yingxin Chen:** Formal analysis, Writing – review & editing. **Claire Blackmore:** Formal analysis, Writing – review & editing. **Katie Eminson:** Formal analysis, Writing – review & editing. **Megan Evans:** Formal analysis, Writing – review & editing. **Xiangpu Gong:** Formal analysis, Writing – review & editing. **Kathryn Adams:** Data curation, Writing – review & editing. **Georgia Rodgers:** Writing – review & editing. **Benjamin Fenech:** Conceptualization, Investigation, Methodology, Project administration, Resources, Supervision, Writing – original draft, Writing – review & editing. **John Gulliver:** Conceptualization, Investigation, Methodology, Project administration, Software, Supervision, Visualization, Writing – original draft, Writing – review & editing.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

The epidemiological, population, and health data are all in the public domain and cited throughout the paper. The ENR strategic noise exposure data is available in the public domain aggregated in 5 dB bands with lower thresholds of 55 dB L_{den} and 50 dB L_{night} .

Acknowledgements

We thank the Department for Environment, Food and Rural Affairs (Defra) for granting us permission to use the ENR strategic noise dataset for this assessment. We thank Dr. Sophie Hamilton for providing insight into methodological aspects of noise and health meta-analyses. The research was supported by National Institute for Health Research (NIHR) Health Protection Research Unit in Environmental Exposures and Health, a partnership between UK Health Security Agency, the Health and Safety Executive and the University of Leicester. The views expressed are those of the author(s) and not necessarily those of the

NHS, the NIHR, the Department of Health and Social Care, Defra or the UK Health Security Agency. While finishing the paper, SC's affiliation changed to Air Quality and Public Health, RCE, Science group, UK Health Security Agency and JG's affiliation changed to Population Health Research Institute, St George's University of London, Cranmer Terrace, London, SW17 0RE. The joint first authors contributed equally to the paper and have the right to list their own names first in the reference list for funding applications or on CVs. The same applies for the joint senior authors.

Funding declarations

Funding was provided by Public Health England (now the UK Health Security Agency) as part of a contract delivered by Extriium in conjunction with the University of Leicester to develop a "Toolkit for Environmental Noise".

Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.envint.2023.107966>.

References

- Basner, M., McGuire, S., 2018. WHO Environmental Noise Guidelines for the European Region: A Systematic Review on Environmental Noise and Effects on Sleep. *Int. J. Environ. Res. Public Health* 15, 519.
- British Heart Foundation. Death rates - Heart and circulatory disease in context. <https://www.bhf.org.uk/what-we-do/our-research/heart-and-circulatory-diseases-in-numbers/heart-and-circulatory-disease-in-context>; 2020.
- British Heart Foundation. Death rates - by local authority. <https://www.bhf.org.uk/what-we-do/our-research/heart-and-circulatory-diseases-in-numbers/death-rates-by-local-authority>; 2021a.
- British Heart Foundation. Diagnosed prevalence by local authority. <https://www.bhf.org.uk/what-we-do/our-research/heart-and-circulatory-diseases-in-numbers/incidence-and-prevalence-incidence-by-local-authority>; 2021b.
- Calentano, D., Szklo, M., Gordis, L., 2019. More on Risk: Estimating the potential for prevention. *Gordis Epidemiology*, sixth ed.
- Chen, Y., Blackmore, C., Eminson, K., Gong, X., Hansell, 2022. Systematic review of meta-analyses for noise. *Inter-noise*.
- Civil Aviation Authority. Airspace structure trials: How trials are used to assess potential changes to airspace. <https://www.caa.co.uk/consumers/guide-to-aviation/airspace/airspace-structure-trials/#>; 2023.
- Civil Aviation Authority. CAP1711: Airspace Modernisation Strategy. <https://publicapps.caa.co.uk/modalapplication.aspx?appid=11&mode=detail&id=8960>; 2018.
- Civil Aviation Authority. CAP1506: Survey of Noise Attitudes 2014: Aircraft Noise and Annoyance, Second Edition. <https://publicapps.caa.co.uk/modalapplication.aspx?appid=11&mode=detail&id=7744>; 2021a.
- Civil Aviation Authority. CAP1616: Airspace change: Guidance on the regulatory process for changing the notified airspace design and planned and permanent redistribution of air traffic, and on providing airspace information. <https://publicapps.caa.co.uk/modalapplication.aspx?appid=11&mode=detail&id=8127>; 2021b.
- Civil Aviation Authority. Heathrow Airport 2019 Summer Noise Contours and Noise Action Plan Contours. https://www.heathrow.com/content/dam/heathrow/web/common/documents/company/local-community/noise/reports-and-statistics/reports-noise-action-plan-Contours/LHR_2019_contour_report.pdf; 2021c.
- Clark, C., Crumpler, C., Notley, A.H., 2020. Evidence for Environmental Noise Effects on Health for the United Kingdom Policy Context: A Systematic Review of the Effects of Environmental Noise on Mental Health, Wellbeing, Quality of Life, Cancer, Dementia, Birth, Reproductive Outcomes, and Cognition. *Int. J. Environ. Res. Public Health* 17, 393.
- Clark, C., Head, J., Haines, M., van Kamp, I., van Kempen, E., Stansfeld, S.A., 2021. A meta-analysis of the association of aircraft noise at school on children's reading comprehension and psychological health for use in health impact assessment. *J. Environ. Psychol.* 76,101646.
- DEFRA. National Noise Incidence Survey (England and Wales). 2000.
- DEFRA. Focussed Noise Monitoring Study 2013 (FNM2013). 2013.
- DEFRA. Noise Action Plan: Roads (Including Major Roads). 2014a.
- DEFRA. Noise action plans: large urban areas, roads and railways (2014). 2014b.
- DEFRA. Noise Action Plan: Agglomerations (Urban Areas); 2019a.
- DEFRA. Noise Action Plan: Railways; 2019b.
- DEFRA. Noise Action Plan: Roads; 2019c.
- DEFRA. Noise action plans: large urban areas, roads and railways (2019). 2022.
- Department of Transport; Welsh Office; HMSO. Calculation of Road Traffic Noise (CRTN). 1988.
- Department for Transport, 2022. Civil Aviation Authority. Aviation statistics: data tables (AVI). <https://www.gov.uk/government/statistical-data-sets/aviation-statistics-data-tables-avi>.

- EEA. Environmental noise in Europe - 2020. Luxembourg: European Environment Agency; 2020.
- Eriksson, C., Bodin, T., Selander, J., 2017. Burden of disease from road traffic and railway noise – a quantification of healthy life years lost in Sweden. *Scand. J. Work Environ. Health* 43, 519–525.
- Eriksson, C., Pershagen, G. Biological mechanisms related to cardiovascular and metabolic effects by environmental noise. World Health Organization; 2018.
- Ezzati, M.; Lopez, A.D.; Rodgers, A.; Murray, C.J. Comparative quantification of health risks: global and regional burden of disease attributable to selected major risk factors ed²: World Health Organization; 2004.
- Fenech, B.; Clark, S.N.; Rodgers, G. An update to the WHO 2018 Environmental Noise Guidelines exposure response relationships for annoyance from road and railway noise. Inter-noise; 2022.
- Fu, W.N., Liu, Y.F., Yan, S.J., Wen, J., Zhang, J., Zhang, P., Zou, L., 2022. The association of noise exposure with stroke incidence and mortality: A systematic review and dose-response meta-analysis of cohort studies. *Environ. Res.* 215.
- Global Burden of Disease Collaborative Network. Global Burden of Disease Study 2019 (GBD 2019) Results. Seattle: Institute of Health Metrics and Evaluation (IHME) 2020.
- Global Burden of Disease Collaborators, 2017. Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 390, 1211–1259.
- Gulliver, J., Morley, D., Vienneau, D., Fabbri, F., Bell, M., Goodman, P., Beevers, S., Dajnak, D., Kelly, F.J., Fecht, D., 2015. Development of an open-source road traffic noise model for exposure assessment. *Environ. Modell. Softw.* 74, 183–193.
- Guski, R., Schreckenberg, D., Schuemer, R., 2017. WHO Environmental Noise Guidelines for the European Region: A Systematic Review on Environmental Noise and Annoyance. *Int. J. Environ. Res. Public Health* 14, 1539.
- Hanninen, O., Knol, A.B., Jantunen, M., Lim, T.A., Conrad, A., Rappolder, M., Carrer, P., Fanetti, A.C., Kim, R., Buekers, J., Torfs, R., Iavarone, I., Classen, T., Hornberg, C., Mekel, O.C.L., Grp, E.W., 2014. Environmental Burden of Disease in Europe: Assessing Nine Risk Factors in Six Countries. *Environ. Health Perspect.* 122, 439–446.
- Hegewald, J., Schubert, M., Freiberg, A., Starke, K.R., Augustin, F., Riedel-Heller, S.G., Zeeb, H., Seidler, A., 2020. Traffic Noise and Mental Health: A Systematic Review and Meta-Analysis. *Int. J. Environ. Res. Public Health* 17, 6175.
- Hegewald, J., Schubert, M., Lochmann, M., Seidler, A., 2021. The burden of disease due to road traffic noise in Hesse, Germany. *Int. J. Environ. Res. Public Health* 18, 9337.
- HM Government. The Environmental Noise (England) (Amendment) Regulations 2018. <https://www.legislation.gov.uk/uksi/2018/1089/contents/made>; 2006.
- Houthuijs, D., Swart, W., van Kempen, E. Implications of environmental noise on health and wellbeing in Europe: Based on data from the second (2012) and third (2017) round of noise assessment in the framework of the European Noise Directive. 2018.
- Kephalopoulos, S., Paviotti, M., Anfosso-Lédée, F., 2012. Common Noise Assessment Methods in Europe (CNOSSOS-EU). European Commission.
- Khomenko, S., Cirach, M., Barrera-Gómez, J., Pereira-Barboza, E., Iungman, T., Mueller, N., Foraster, M., Tonne, C., Thondoo, M., Jephcote, C., 2022. Impact of road traffic noise on annoyance and preventable mortality in European cities: A health impact assessment. *Environ. Int.* 162, 107160.
- Kim, Y.-E., Jung, Y.-S., Ock, M., Yoon, S.-J., 2022. DALY Estimation Approaches: Understanding and Using the Incidence-based Approach and the Prevalence-based Approach. *J. Prev. Med. Public Health* 55, 10–18.
- Morley, D.W., Gulliver, J., 2016. Methods to improve traffic flow and noise exposure estimation on minor roads. *Environ. Pollut.* 216, 746–754.
- Mueller, N., Rojas-Rueda, D., Basagaña, X., Cirach, M., Cole-Hunter, T., Dadvand, P., Donaire-Gonzalez, D., Foraster, M., Gascon, M., Martinez, D., 2017. Health impacts related to urban and transport planning: A burden of disease assessment. *Environ. Int.* 107, 243–257.
- Munzel, T., Gori, T., Babisch, W., Basner, M., 2014. Cardiovascular effects of environmental noise exposure. *Eur. Heart J.* 35, 829.
- Munzel, T., Sorensen, M., Daiber, A., 2021a. Transportation noise pollution and cardiovascular disease. *Nat. Rev. Cardiol.* 18, 619–636.
- Munzel, T., Steven, S., Hahad, O., Frenis, K., Oelze, M., Kroll-Schon, S., Daiber, A., 2021b. Environmental traffic noise triggers stress reactions, oxidative stress, inflammation and vascular dysfunction - comparison of studies in mice and men. *Free Radic. Biol. Med.* 177 (Supplement 1), S56.
- Murphy, E., Faulkner, J.P., 2022. Estimating the harmful effects of environmental transport noise: An EU study. *Sci. Total Environ.* 811, 152313.
- Murray, C.J., Ezzati, M., Lopez, A.D., Rodgers, A., Vander Hoorn, S., 2003. Comparative quantification of health risks conceptual framework and methodological issues. *Popul. Health Metr.* 1, 1.
- National Institute for Public Health and the Environment (RIVM). Study on methodology to perform environmental noise and health assessment. Netherlands: RIVM. <https://www.rivm.nl/bibliotheek/rapporten/2018-0121.pdf>; 2018.
- NHS Digital. Quality and Outcomes Framework (QOF): Disease prevalence and care achievement rates. <https://digital.nhs.uk/data-and-information/data-tools-and-services/data-services/general-practice-data-hub/quality-outcomes-framework-qof>; 2022.
- NOMIS. Mid-year population estimates. <https://www.nomisweb.co.uk/articles/1165.aspx>; 2019.
- Office for Health Improvement & Disparities. Fingertips | Public health data. <https://fingertips.phe.org.uk/profile/cardiovascular>; 2022.
- Office for National Statistics. Deaths by sex, age group, cause and local authority, England, deaths registered 1993 to 2013. in: Office for National Statistics, ed. <http://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/causesofdeath/adhocs/004352deathsbysexagegroupcauseandlocalauthorityenglanddeathregistered1993to2013>; 2015.
- Office for National Statistics. Deaths by LSOA in England and Wales, mid-year 2000/01 to 2017/18. in: Office for National Statistics, ed. <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/adhocs/10772deathsbylowerlayersuperoutputarealsoenglandandwalesmidyear2001to2018>; 2019.
- Office for National Statistics. Life expectancy for local areas of the UK between 2001 to 2003 and 2017 to 2019. <https://www.ons.gov.uk/releases/lifeexpectancyforlocalareasoftheuk2001to2003to2017to2019>; 2020a.
- Office for National Statistics. UK population summary by age and geography, mid-2018. <https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationestimates/adhocs/11516ukpopulationsummarybyageandgeographymid2018>; 2020b.
- ONS Geography. Open Geography portal: Local Authority Districts (May 2020) Boundaries UK BFE. https://geoportal.statistics.gov.uk/datasets/9d86e7bcf2864343ab72e2914756b86d_0/explore?location=55.154076%2C-3.313872%2C6.36; 2021.
- Ordnance Survey. Boundary-Line data. <https://beta.ordnancesurvey.co.uk/products/boundary-line>; 2022.
- Park, B., Park, B., Lee, W.K., Kim, Y.E., Yoon, S.J., Park, H., 2019. Incidence-Based versus Prevalence-Based Approaches on Measuring Disability-Adjusted Life Years for Injury (vol 34, e69, 2019). *J. Korean Med. Sci.* 34.
- Persson Waye, K.; van Kempen, E. Non-auditory effects of noise: an overview of the state of the science of the 2017-2020 period. ICIBEN. Stockholm; 2021.
- R Core Team. R: A language and environment for statistical computing. Vienna; 2022.
- Roswall, N., Pyko, A., Ogren, M., Oudin, A., Rosengren, A., Lager, A., Poulsen, A.H., Eriksson, C., Segerson, D., Rizzuto, D., Andersson, E.M., Aasvang, G.M., Engstrom, G., Jorgensen, J.T., Selander, J., Christensen, J.H., Thacher, J., Leander, K., Overvad, K., Eneroth, K., Mattisson, K., Barregaard, L., Stockfelt, L., Albin, M., Ketzel, M., Simonsen, M.K., Spanne, M., Raaschou-Nielsen, O., Magnusson, P.K.E., Tiittanen, P., Molnar, P., Ljungman, P., Lanki, T., Lim, Y.H., Andersen, Z.J., Pershagen, G., Sorensen, M., 2021. Long-Term Exposure to Transportation Noise and Risk of Incident Stroke: A Pooled Study of Nine Scandinavian Cohorts. *Environ. Health Perspect.* 129, 1665.
- Sakhvidi, M.J.Z., Sakhvidi, F.Z., Mehrparvar, A.H., Foraster, M., Dadvand, P., 2018. Association between noise exposure and diabetes: A systematic review and meta-analysis. *Environ. Res.* 166, 647–657.
- Salomon, J., Haagsma, J.A., Davis, A., de Noordhout, C.M., Polinder, S., Havelaar, A.H., Cassini, A., Devleeschauwer, B., Kretzschmar, M., Speybroeck, N., Murray, C.J.L., Vos, T., 2015. Disability weights for the Global Burden of Disease 2013 study. *Lancet Glob. Health* 3, E712–E723.
- Seidler, A., Hegewald, J., Seidler, A.L., Schubert, M., Wagner, M., Droge, P., Haufe, E., Schmitt, J., Swart, E., Zeeb, H., 2017. Association between aircraft, road and railway traffic noise and depression in a large case-control study based on secondary data. *Environ. Res.* 152, 263–271.
- Seidler, A., Hegewald, J., Seidler, A.L., Schubert, M., Zeeb, H., 2019. Is the Whole More Than the Sum of its Parts? Health Effects of Different Types of Traffic Noise Combined. *Int. J. Environ. Res. Public Health* 16, 1665.
- Smith, M.G., Cordoza, M., Basner, M., 2022. Environmental Noise and Effects on Sleep: An Update to the WHO Systematic Review and Meta-Analysis. *Environ. Health Perspect.* 130, 76001.
- Sohrabi, S., Khreis, H., 2020. Burden of disease from transportation noise and motor vehicle crashes: Analysis of data from Houston, Texas. *Environ. Int.* 136, 105520.
- Sorensen, M., Andersen, Z.J., Nordsborg, R.B., Becker, T., Tjonneland, A., Overvad, K., Raaschou-Nielsen, O., 2013. Long-Term Exposure to Road Traffic Noise and Incident Diabetes: A Cohort Study. *Environ. Health Perspect.* 121, 217–222.
- Sorensen, M., Hvidberg, M., Andersen, Z., Nordsborg, R., Lillelund, K., Jakobsen, J., Tjonneland, A., Overvad, K., Raaschou-Nielsen, O., 2011. Road Traffic Noise and Stroke: A Prospective Cohort Study. *Eur. Heart J.* 32, 737–744.
- Sorensen, M., Luhdorf, P., Ketzel, M., Andersen, Z.J., Tjonneland, A., Overvad, K., Raaschou-Nielsen, O., 2014. Combined effects of road traffic noise and ambient air pollution in relation to risk for stroke? *Environ. Res.* 133, 49–55.
- Stassen, K.R., Collier, P., Torfs, R., 2008. Environmental burden of disease due to transportation noise in Flanders (Belgium). *Transp. Res. Part D-Transp. Environ.* 13, 355–358.
- Steel, N., Ford, J.A., Newton, J.N., 2018. Changes in health in the countries of the UK and 150 English Local Authority areas 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 392, 1647–1661.
- Thacher, J.D., Poulsen, A.H., Hvidtfeldt, U.A., Raaschou-Nielsen, O., Brandt, J., Geels, C., Khan, J., Münzel, T., Sørensen, M., 2021. Long-term exposure to transportation noise and risk for type 2 diabetes in a nationwide cohort study from Denmark. *Environ. Health Perspect.* 129, 127003.
- Thacher, J.D., Poulsen, A.H., Raaschou-Nielsen, O., Hvidtfeldt, U.A., Brandt, J., Christensen, J.H., Khan, J., Levin, G., Münzel, T., Sørensen, M., 2022. Exposure to transportation noise and risk for cardiovascular disease in a nationwide cohort study from Denmark. *Environ. Res.* 211, 113106.
- UECNA. Statistics: Main European Airport. <https://www.uecna.eu/wp-content/uploads/2021/10/UECNA-STATS-EUROPE-Synthesis-til-2020-with-2007-En-Cours-2.pdf>; 2020.
- van Kamp, I., Simon, S., Notley, H., Baliatsas, C., van Kempen, E., 2020. Evidence Relating to Environmental Noise Exposure and Annoyance, Sleep Disturbance, Cardio-Vascular and Metabolic Health Outcomes in the Context of IGCB (N): A Scoping Review of New Evidence. *Int. J. Environ. Res. Public Health* 17, 3016.
- van Kempen, E., Casas, M., Pershagen, G., Foraster, M., 2018. WHO Environmental Noise Guidelines for the European Region: A Systematic Review on Environmental Noise and Cardiovascular and Metabolic Effects: A Summary. *Int. J. Environ. Res. Public Health* 15, 379.

- Vienneau, D., Eze, I.C., Probst-Hensch, N., Roosli, M., 2019. Association between transportation noise and cardio-metabolic diseases: an update of the WHO meta-analysis. 23rd International Congress on Acoustics . Aachen, Germany.
- Vienneau, D., Saucy, A., Schaffer, B., Fluckiger, B., Tangermann, L., Stafoggia, M., Wunderli, J.M., Roosli, M., Grp, S.S., 2022. Transportation noise exposure and cardiovascular mortality: 15-years of follow-up in a nationwide prospective cohort in Switzerland. *Environ. Int.* 158, 106974.
- von der Lippe, E., Devleeschauwer, B., Gourley, M., Haagsma, J., Hilderink, H., Porst, M., Wengler, A., Wyper, G., Grant, I., 2020. Reflections on key methodological decisions in national burden of disease assessments. *Arch. Public Health* 78, 1-14.
- Wagner, R.G., Ibinda, F., Tollman, S., Lindholm, L., Newton, C.R., Bertram, M.Y., 2015. Differing Methods and Definitions Influence DALY estimates: Using Population-Based Data to Calculate the Burden of Convulsive Epilepsy in Rural South Africa. *PLoS One* 10, e0145300.
- WHO. Night Noise Guidelines for Europe. <https://www.euro.who.int/en/health-topics/environment-and-health/noise/publications/2009/night-noise-guidelines-for-europe>; 2009.
- WHO. Environmental Noise Guidelines for the European Region. <https://www.euro.who.int/en/publications/abstracts/environmental-noise-guidelines-for-the-european-region-2018>; 2018.
- WHO. WHO methods and data sources for global burden of disease estimates 2000-2019. Geneva: World Health Organization; 2020.
- WHO Regional Office for Europe. Burden of disease from environmental noise: Quantification of healthy life years lost in Europe. World Health Organization; 2011.