

## THESIS / THÈSE

### MASTER IN BIOLOGY

#### Unusual organization of the rRNA genes in the Rickettsia genus

ROMEDENNE , Magali

*Award date:*  
1995

*Awarding institution:*  
University of Namur

[Link to publication](#)

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

0 1 DEC. 1995



**FACULTES UNIVERSITAIRES N.-D. DE LA PAIX  
NAMUR  
FACULTE DES SCIENCES**

---

**Unusual organization of the rRNA genes  
in the *Rickettsia* genus**

Mémoire présenté pour l'obtention du grade  
de Licencié en Sciences  
biologiques

Romedenne Magali

Décembre 1995

**Facultés Universitaires Notre-Dame de la Paix**  
**FACULTE DES SCIENCES**  
Rue de Bruxelles 61 - 5000 NAMUR  
Tél. 081/72.41.11 - Telex 59222 Facnam-b - Telefax 081/72.44.20

**Unusual organization of the rRNA genes  
in the *Rickettsia* genus**

ROMEDENNE Magali

**Summary**

Dans une première partie du travail, nous décrivons l'organisation des gènes codant pour le peptide déformylase (PDF) et méthionyl-tARN<sup>fMet</sup> formyltransférase (MTF) dans la bactérie *Rickettsia prowazeki*. L'analyse de la séquence de clones ainsi que l'analyse par PCR de l'ADN génomique grâce aux amorces, correspondant à chacun des deux gènes, confirme que les gènes *fms* et *fmt* sont étroitement liés dans le génome de *Rickettsia prowazeki*.

Dans un second temps, nous continuons de démontrer l'organisation inhabituelle des gènes codant pour les ARN ribosomiaux déjà observée dans *Rickettsia prowazeki*. Nous montrons dans cette étude que *fmt* est aussi localisé en amont du gène *rrl* dans le génome du *R. sibirica*, *R. parkeri*, *R. rickettsii*, *R. amblyommii* tout comme dans l'agent ELB. Ces résultats suggèrent que le réarrangement génomique des gènes codant pour les ARN ribosomiaux précèdent la divergence des espèces TG (Typhus group) et des espèces SFG (Spotted Fever group).

Mémoire de licence en Sciences Biologiques

Décembre 1995

**Promoteur : J. VANDENHAUTE**

**Co-promoteur : S. ANDERSSON**

Nine months, two weeks and a few hours, I had just arrived in the snow of Sweden for a great adventure! Back in Belgium, I miss all these people that I met, but anyway I am happy to see again the rain of my native country and all these people who supported me in my "crazy" trip. Now, it is time to thank everybody who encourage me during this work.

First I wish to express my gratitude to Professor Charles G. Kurland who welcomed me in his department and introduced me in the scientific world.

I am extremely grateful for my "swedish" supervisor Siv Andersson. Your advices, supports and patience were really helpful for the realisation of this report. I thank you for the interesting subject.

I thank Professor J. Vandenhoute who accepted to be my supervisor.

I am grateful for the ERASMUS program which gives me the wonderful opportunity to discover another university, to meet a lot of different people, culture and languages. I specially thank Father Devos and Marita Wriegen who organised my stay in Uppsala.

In addition, I would like to thank all my friends from the Molecular Biology Department.

Ulrika, for her patience to share the same bench, desk, and PCR machine! I thank you for the precious advices and encouragement in difficult moments (N'oublie pas ton Français, ça soulage parfois!).

Asfar, for all the informations about science but also about Pakistan (food and culture!). Be careful on the snow! I really thank you for your friendship.

Ali (for advices), Haley (for all the primers), Thomas and Morten (for helping me with the computers, and that is not easy!), Anne-Sophie (for your humour, joy and patience with ALF), Jan (for your marker), Kristina (for your discrete but friendly presence) and Surrinder (unfortunately, you arrived to late but soon enough for me to enjoy your conversation).

I thank also Vildan and Anders (for the turkish music, see you for the next movie), Pedro (be careful with chocolate), Monika (I will think about you next time when I will be by my yoghourt!), Tracy and Mark (for your friendship), David and Mattias (for the BC), Angie (to take over the organization of the BC with an american accent),....

*But there is also evrybody from outside of the department who make my stay in Uppsala a great pleasure.*

*Pernilla, my great buddy (or bodygard!). I thank you for your presence at anytime (specially on sunday afternoon in Kalmar nation)....*

*I thank also all the swedish students in biology for the nice parties, cafés, folkmusic, Valborg,... And of course, I should not forget the innumerable exchange students from everywhere in the world.*

*Je remercie Natacha pour sa compagnie durant ces cinq années, pour son courage de me supporter au kot, aux TP et pendant ces très très longs blocus. Je te remercie tout particulièrement pour l'encouragement dans mon expédition suédoise.*

*Un tout grand merci à Xavier qui, en dernière minute, a eu la patience de lire et corriger ce travail.*

*Evidemment, je ne peux oublier de remercier Ialex pour ses nombreux e-mails tout au long de ces neuf mois. Ils m'ont été réconfort, soutien et humour belge (et ça fait du bien!). Merci aussi pour l'aide pour l'achèvement de ce mémoire.*

*Merci beaucoup aux membres du laboratoire de biochimie qui m'ont accueillie ces derniers jours. Je remercie surtout Marie qui, il y a plus d'un an, m'a décidée de partir à Uppsala. Ton avis a joué un grand rôle dans cette expérience innoubliable. Merci à Christine que je découvre bien tard.*

*Je n'oublie pas tous les bios et tous les autres.*

*Il y a bien sûr Sophie que je remercie pour tout, mais je ne m'étale pas, ça risquerait d'être trop long.*

*Ce travail ne pourrait être achevé sans remercier de tout coeur mes parents et ma soeur Laurence qui depuis 22 ans (bientôt 23) ont eu le courage de me supporter et soutenir. Je ne peux d'ailleurs que vous féliciter!*

*Tack*

*Thanks*

*Merci*

*Magali*

# Contents

<b>Introduction</b> .....	1
1. Structure and evolution of bacterial genomes.....	1
1.1. Universal genalogical tree.....	1
1.2. Comparative molecular data on bacterial genome.....	1
1.3. rRNA genes locus and flanking region.....	3
2. <i>Rickettsia</i> characteristics.....	4
2.1. <i>Rickettsia</i> , as a parasitic bacteria.....	4
2.2. Genomic characteristics of <i>Rickettsia</i> .....	5
2.3. rRNA operon in <i>Rickettsia prowazekii</i> .....	6
2.4. Interest to study the organisation of the rRNA genes in <i>Rickettsia</i> .....	6
<b>Goals of this project</b> .....	7
<b>Conserved organisation of the genes encoding Peptide Deformylase and Methionyl-tRNA<sup>fMet</sup> Formyltransferase in <i>Rickettsia prowazekii</i></b> .....	8
Summary.....	9
1. Introduction.....	10
2. Materials and method.....	11
2.1. Purification of plasmids.....	11
2.2. DNA sequencing.....	11
2.3. Sequence analysis.....	11

2.4. PCR amplification.....	11
3. Results.....	12
3.1. Nucleotide sequence of the <i>fms</i> gene and upstream regions.....	12
3.2. The <i>fms</i> gene is linked to the <i>fmt</i> gene in <i>Rickettsia prowazekii</i> .....	13
3.3. Comparative analysis of the <i>fms</i> and <i>fmt</i> gene products.....	13
3.4. Organization of the <i>fms</i> - <i>fmt</i> locus in bacterial genomes.....	14
4. Concluding remarks.....	15

**Rearrangement of rRNA Genes in *Rickettsia*  
Preceded the Divergence of the Typhus and the  
Spotted Fever Group *Rickettsia*.....** 16

Summary.....	17
1. Introduction.....	18
2. Materials and methods.....	19
2.1. PCR amplification and purification.....	19
2.2. DNA sequencing.....	19
2.3. Sequence analysis.....	20
3. Results.....	20
3.1. PCR amplification with primers from <i>fmt</i> and 23S rRNA.....	20
3.2. Comparative nucleotide sequence analysis.....	21
4. Concluding remarks.....	22
<b>General conclusion.....</b>	<b>24</b>

<b>Materials and methods</b> .....	25
1. Preparation of plasmids.....	25
1.1. Introduction.....	25
1.2. Components.....	25
1.3. Protocol.....	26
2. Polymerase Chain Reaction (PCR).....	28
2.1. Introduction.....	28
2.2. Components.....	28
2.3. Protocol.....	29
2.4. Control.....	30
3. Direct purification from PCR products.....	30
3.1. Introduction.....	30
3.2. Components.....	31
3.3. Protocol.....	31
4. DNA sequencing.....	32
4.1. Introduction.....	32
4.2. Auto Read Sequencing Kit (Pharmacia).....	33
4.3. Thermo Sequenase fluorescent labelled primer cycle sequencing kit (Amersham).....	37
<b>References</b> .....	39






Figure 1: Universal tree in rooted form, showing the three domains: Archaea, Bacteria and Eucarya.

# Introduction

## 1. Structure and evolution of bacterial genomes

### 1.1. Universal genealogical tree

All life forms could be accommodated in five kingdoms of organisms: the Animals, Plants, Fungi, Protistes and Monera, and in two superkingdoms with the eukaryotes and prokaryotes.

The prokaryotic cells are divided into the Bacteria and the Archaea domains. The Eukaryotes fall into the Eucarya domain (Fig. 1).

### 1.2. Comparative molecular data on bacterial genomes

Within this phylogenetic tree, the genomic evolutionary forces have had an impact on the genome size, nucleotide composition and general organisation, on the genetic code and in particular on the nature, the number and the disposition of the genes coding for the translation apparatus.

#### a. Genome size

The genome is the complete heritable set of genes of an organism. Bacterial genome sizes vary from 600 kb (*Mycoplasma genitalium*) to 12.8 Mb (in *Calothrix* strains). Organisms with values between these two extremes includes *Borrelia burgdorferi* with 950 kb, *Mycoplasma hypopneumoniae* (1.14 Mb), *E. coli* (4.7 Mb) and *Myxococcus*

*xanthus* (9.5 Mb) (Krawiec and Riley, 1990). The number of genes spans the range from 400 to 8000.

The genome of bacteria can be divided into three fractions: chromosomal DNA, plasmids and transposable elements (which are common components of the genome).

It is suggested that the larger genomes may have evolved from smaller genomes by genomic duplications as well as gene elongations, and that the smaller genomes may be the result of a reductive mode of genomic evolution (Andersson and Kurland, 1995).

#### b. Nucleotide composition

The nucleotide composition (in %) is another important genomic trait in procaryotes. It seems to be related to phylogeny with closely related species having the same G+C content. However, in the alpha proteobacteria for example, some species have high G+C contents (such as *Agrobacterium tumefaciens*) while others have low G+C contents (such as *Rickettsia prowazekii*). The genomic G+C content values vary from 25% to 75% (Muto and Osawa, 1987).

This variation in nucleotide composition among genomes can be explained by two hypotheses:

-the selectionists suggest that different environmental factors may influence the G+C content, the observed differences resulting from adaptations of bacteria to their environment (Li and Graur, 1991). For instance the thermophile bacteria have a high G+C content to stabilise the proteins at high temperature.

-the mutationists invoke biases in the mutation patterns as an explanation for differences in genomic G+C contents (Muto and Osawa, 1987). The redundancy of the genetic code permits changes in base composition which are neutral. A change in the third codon position most often do not alter the amino acid sequence of the protein and is therefore most strongly affected by biased mutation patterns (change in the second or the first position can also occur) (Andersson and Kurland, 1995).




Figure 2: Schematic representation of a typical rRNA operon.

Arrangement of rRNA	Organisms	Number of copies
16S-23S-5S	<i>E. coli</i>	7
	<i>Bacillus subtilis</i>	10
	<i>Mycoplasma capricolum</i>	2
	<i>Streptomyces coelicolor</i>	6
23S-16S-5S	<i>Vibrio harveyi</i>	1
5S and 16S-23S	<i>Mycoplasma</i>	1
	<i>hyopneumoniae</i>	
16S and 23S-5S-23S-5S	<i>Borrelia burgdorferi</i>	1
16S and 23S-5S	<i>Thermus thermophilus</i>	2

Table 1: Copy number and arrangement of the rRNA genes in some bacterial species

## THE rRNA OPERON





Figure 3: The seven rRNA operons in *E. coli*. Abbreviations (A, D, H, B, E, G, C) used to indicate the position of the different operons in the genome. 16S = rRNA small subunit, 23S = rRNA large subunit, 5S = rRNA subunit, Ile = tRNA for isoleucine, Ala = tRNA for alanine, Glu = tRNA for glutamic acid, Thr = tRNA for threonine, Asp = tRNA for aspartic acid, Trp = tRNA for tryptophane.

Changes in genomic global nucleotide composition involve differences in the frequencies of AT-rich or GC-rich codons, and thus produce compositional differences in amino acid sequence for homologous proteins.

Some bacterial proteins, tRNA or rRNA genes have a remarkably conservative amino acid composition and may be relevant to discuss the evolution of code.

### c. Architectures

Surprisingly, the genes for components of the translation apparatus are often organized similarly among different bacteria and archaea (and even eukaryotes). But there are exceptions in their organization and it is suggested that such unusual genomic organizations may contain important information about the evolutionary history of these bacteria.

#### 1.3. rRNA genes locus and flanking region

In most eubacterial species, the three rRNA genes are closely linked in the order 16S-23S-5S in one transcription unit, corresponding to the so called rRNA operon. This operon is typically composed of a promoter, a small subunit gene (SSU, 16S or *rrs*), a spacer usually containing tRNAs, a large subunit gene (LSU, 23S or *rrl*) and a spacer and a 5S gene (*rrf*). Other tRNAs may be found downstream of the 5S gene. (Fig. 2)

The rRNA operons are usually found in several copies in the genome, ranging from several hundred of copies in vertebrate genomes to a few copies in small bacterial genomes (Table 1). The different rRNA operons of *E. coli* are shown in the figure 3.

A very low rRNA copy number is often observed in obligate intracellular organisms together with a small genome size.

It is generally believed that obligate intracellular parasitic bacteria derive from extracellular free-living bacteria. The development of mechanisms for entering host cells, for multiplying, for survival and




Figure 4: Phylogenetic tree showing *R. prowazekii* within the alpha-proteobacteria.



for infecting new cells are important for a parasitic life style. During the transition to this parasitic life style, the obligate intracellular bacteria may have gained new characters and lost old characters.

The very rich medium (cytosol) could explain the reduced number of genes resulting from a degenerative mode of evolution from a larger, bacterial genome. The objective behind a small genome could also be a quick replication and division.

## **2. Rickettsia characteristics**

In this report we will study the unusual organisation of the rRNA operon in the *Rickettsia* genus.

*Rickettsia* is an obligate intracellular gram-negative eubacteria, member of the branch of the alpha-proteobacteria (purple bacteria) (Fig. 4). *Rickettsia* is interesting to study because of its small genome size (1.2 Mb), its close phylogenetic relationship to mitochondria and its obligate intracellular lifestyle.

### **2.1 *Rickettsia*, as a parasitic bacteria**

*Rickettsia* have natural arthropod hosts (ticks, mites or insects). Man and other vertebrates may be affected by these organisms, most of which are pathogenic.

The penetration of the *Rickettsia* into host cells involves the phagocytosis of the microorganism after its attachment to the cell surface. Penetration of a *Rickettsia* into its host cell is accomplished within 10-20 minutes, but the intracellular multiplication process is very slow. It escapes quickly from the phagosome into the cytosol where the parasite grows. This very rich medium produces energy and every intermediates of most pathways that *Rickettsia* needs for its growth. Some species are also capable of growth and division in the nucleus.

Species	Human disease	Principal invertebrate host	Principal vertebrate host	Geographic distribution
<u>Spotted fever group</u>				
<i>R. rickettsii</i>	Rocky Mountain Spotted fever	Several genera of ticks	rabbit, hare, dog	North and South America
<i>R. conorii</i>	Boutonneuse fever, Mediterranean fever	Several genera of ticks	rodent, dog	Mediterranean countries, Black Sea Coast of Russia, Africa
<i>R. sibirica</i>	Siberian tick typhus	Several genera of ticks	rodent	Eurasia, Asia
<i>R. akari</i>	Rickettsiapox	Mouse mite	rodent	worldwide
<i>R. australis</i>	Queensland tick typhus	<i>Ixodes</i> tick	rodent, marsupial	Australia
<i>R. parkeri</i>	No known disease	<i>Amblyomma</i>	unknown	USA
<i>R. montana</i>	No known disease	<i>Dermacentor</i> tick	unknown	USA
<i>R. amblyommi</i>	No known disease	<i>Amblyomma</i>	unknown	Missouri, USA
<i>R. rhipicephali</i>	No known disease	<i>Rhipicephalus</i>	dog, meadow vole	USA
<i>R. honei</i>	Flinder's Island spotted fever	<i>Ixodes</i> tick		Flinder's Island (Tasmania)
<i>R. helvetica</i>	No known disease	<i>Ixodes</i> tick	meadow vole	Switzerland
<i>R. massiliae</i>	No known disease	<i>Rhipicephalus</i>	dog	France, Greece
<i>R. africae</i>	African tick-bite fever	<i>Amblyomma</i>	dog, ruminant, rabbit	South Africa
<i>R. japonica</i>	Japanese spotted fever	unknown	rodent, dog	Japan

Species	Human disease	Principal invertebrate host	Principal vertebrate host	Geographic distribution
<u>Typhus group</u>				
<i>R. typhi</i>	Murine typhus	Flea, louse	rat, mouse	worldwide
<i>R. prowazekii</i>	Epidemic typhus	Body louse	human	worldwide
	Recrudescant typhus		human	worldwide
	Sylvatic typhus	Flea and louse	flying squirrel	Eastern United States
<i>R. canada</i>	No known disease	<i>Haemaphysalus</i>	rabbit, hare	Ontario, California
<i>ELB agent</i>	Murine-like typhus	Flea	opossum, dog	Texas, California
<u>Scrub typhus group</u>				
<i>R. tsutsugamushi</i>	Scrub typhus	Trombiculid mites	rodent	Japan, Eastern Asia to India, Australia, Pacific Islands
<u>Undetermined</u>				
<i>R. belii</i>	No known disease	Several genera of ticks	unknown	USA
AB bacterium	No known disease	Ladybird beetle	unknown	England

Table 2: *Rickettsia* genus

The *Rickettsia* genus is divided into three groups: The Spotted Fever Group (SFG), the Typhus Group (TG) and the Scrub Typhus Group (STG) (Table 2)

a. The Spotted Fever Group (or SFG)

The members of the SFG grow and divide in the cytosol, but they can also grow within the nucleus.

The average G+C content of the coding sequences has a value of 32 and 33 %.

*R. rickettsii*, *R. sibirica*, *R. parkeri*, *R. montana* and *R. amblyommi* will be discussed later in this study.

b. The Typhus Group (TG)

The characteristic hosts of the TG species are the insects.

The members of the Typhus Group are not able to invade the nucleus.

The G+C content has an average of 29-30 %.

*R. prowazekii* and *ELB agent* will be the object of this project.

c. The Scrub Typhus Group (STG)

The scrub typhus agent (*R. tsutsugamushi*) is very different from the species of the SFG and the TG in its morphology cell surface proteins.

It is capable to grow in the cytosol and can be found in the nucleus.

This until now unique member of the STG will not be considered in this study.

d. *R. bellii* and *AB bacterium* are not included within any of these three groups.

## 2.2. Genomic characteristics of *Rickettsia*

The small and unique chromosome of the bacterium *Rickettsia* has a size around 1.2 Mb (Eremeeva et al. 1993). Calculations based on a coding potential of 60-70%, a genome size of 1.2 Mb and an average




Figure 5: Organization of the rRNA genes in *Rickettsia prowazekii*. Thin lines indicate a noncoding region; Boxes indicate a coding region. Abbreviations used to indicate genes are as follows: rrs = rRNA small subunit gene, orf = open reading frame, fmt = methionyl-tRNA<sup>Met</sup> formyltransferase gene, rrl = rRNA large subunit gene, rrf = 5S rRNA gene.

gene length of 1 kb suggests that there may be around 800 genes in the *R. prowazekii* genome (Andersson and Sharp, 1995).

*Rickettsia* is a AT-rich bacteria with average values between 29 to 33% due to a mutational bias towards A+T instead of G+C base pairs. Protein coding and intergenic region have G+C content values of 32.5% and 21.6%, respectively. The functionally less important third codon position is more biased than the first two positions (Andersson and Sharp, 1995).

### 2.3. rRNA operon in *Rickettsia prowazekii*

In most eubacterial species, the three rRNA genes are closely linked in the order 16S-23S-5S grouped into one operon. The number of copies in bacteria varies from 1 (*Mycoplasma hypopneumoniae*) to 11 in some *Bacillus* species.

Studies of the rRNA genes in *Rickettsia prowazekii* shows that 16S rRNA gene is separated from 23S-5S rRNA gene cluster. One unidentified open reading frame (ORF) was detected downstream of the 16S rRNA gene. An ORF encoding the methionyl-tRNA<sup>fMet</sup> N-formyltransferase gene (*fnt* gene) was found upstream of the 23S rRNA gene. No tRNAs have been found upstream from 23S gene nor in the region downstream from 16S gene (Andersson et al., 1995) (Fig.5). In *Rickettsia* there is only one copy of each of the rRNA genes.

### 2.4. Interest to study the organization of the rRNA genes in *Rickettsia*

It has been shown that *R. prowazekii* has one copy of the 16S rRNA gene, one copy of the 23S-5S rRNA gene cluster (Andersson et al., 1995) and also one copy of the *tuf* gene which encodes elongation factor Tu (EF-Tu) (Syvänen et al., 1995). Since these genes are found in several copies in others bacteria, and the organization seems to be unusual in *R. prowazekii*, it is suggested that the genome of *Rickettsia* has evolved from a larger genome.

Because of their ability to recombination, duplication, inversion and deletion, the sequences which are found in multiple copies (as the rRNA genes) are interesting to study the hypothesis of a reductive evolution.

## Goals of this project

The study of the unusual organization of the rRNA genes in the genus *Rickettsia* is divided in two parts.

First, we will study the *fms-fmt* arrangement in *Rickettsia prowazekii* (*fms* encodes peptide deformylase and *fmt* encodes methionyl-tRNA<sup>Met</sup> N-formyltransferase gene).

The colocalisation of these two genes might be a conserved feature within the bacterial kingdom. Probably because of their related fonctions, the two genes *fms* and *fmt* are found linked to each other in *E. coli*, *Thermus thermophilus* and *Haemophilus influenzae*. The conservation of this linkage in these three species might be a conserved trait within the bacterial organisms (Meinzel and Blanquet, 1994).

Recently the complete sequencing of the genome of *Mycoplasma genitalium* has shown a different organisation in which they are separated from each other by 328 kb (Fraser et al., 1995).

In the present work, the *fms* gene has been sequenced in *Rickettsia prowazekii* and a PCR amplification shows the linkage of *fms* gene with the *fmt* gene which is situated upstream of the 23S rRNA gene.

Secondly, we will analyse and compare the sequences of the *fmt* gene and the sequences of the spacer between *fmt* and 23S genes for seven different species of the genus *Rickettsia*.

Noncoding regions are often altered by random mutation without any troubles in the genome. Thus, the study of the spacer is interesting as a fast clock of evolution and also for eventual diagnostic of the different species of *Rickettsia*.

**Conserved organization of the genes encoding Peptide  
Deformylase and Methionyl-tRNA<sup>fMet</sup> Formyltransferase  
in *Rickettsia prowazekii***

Magali P. C. Romedenne\*, Siv G. E. Andersson and Charles G.  
Kurland

*Department of Molecular Biology, Uppsala University, Biomedical  
Center, Box 590, S751 24 Uppsala, Sweden*

*\*Present address: Laboratory of Genetics, Facultés Universitaires  
Notre dame de la Paix, 61 rue de Bruxelles, 5000 Namur, Belgium*

Running title: *fms* and *fmt* in *Rickettsia prowazekii*

Keywords: *fms*, *fmt*, genomic organization, *Rickettsia prowazekii*



## Summary

We describe here the organization of the genes encoding peptide deformylase (*fms*) and methionyl-tRNA<sup>fMet</sup> formyltransferase (*fmt*) in *Rickettsia prowazekii*. Sequence analysis of cloned fragments as well as a PCR analysis of genomic DNA using primers from each of the two genes confirm that the *fms* and *fmt* genes are tightly linked together in the *R. prowazekii* genome.


## 1. Introduction

A characteristic feature of translation initiation in bacteria, chloroplast and mitochondria is the N-formylation of the initiator Met-tRNA<sub>f</sub><sup>Met</sup>, which is catalyzed by methionyl-tRNA<sub>f</sub><sup>Met</sup> formyltransferase (MTF) using formyltetrahydrofolate as a cofactor. After about 15-30 rounds of elongation the formyl group is removed by the peptide deformylase (PDF) and in about half of the proteins the methionine residue is removed by the methionine aminopeptidase.

Genes encoding functionally related products are sometimes organized into co-transcribed units in bacterial genomes. The genes encoding the enzymes involved in the formylation and deformylation of the Met-tRNA<sub>f</sub><sup>Met</sup>, *fmt* and *fms*, are genetically linked in several different species of diverse phylogenetic affiliations such as *Escherichia coli* (Guillone et al., 1992), *Haemophilus influenzae* (Fleishman et al., 1995) and *Thermus thermophilus* (Meinzel and Blanquet, 1994). In contrast, in the 0.6 Mb genome of *Mycoplasma genitalium* these two genes are separated from each other by 328 kb (Fraser et al., 1995). Disruption of otherwise conserved bacterial operons, such as for example the rRNA operon, has been observed in several small sized bacterial genomes and suggested to be a consequence of a reductive mode of evolution, where extensive deletion and intrachromosomal recombination events have rearranged an ancestral genome with a size and an organization that was more typical of the bacteria (Andersson and Kurland, 1995).

Since *R. prowazekii* has a small-sized genome of 1.2 Mb (Eremeeva et al., 1993) it was of interest to determine how the *fms* and *fmt* genes are arranged in this organism. It has previously been shown that the *fmt* gene is situated upstream of the 23S rRNA gene (Andersson et al., 1995). In this paper, we present evidence which suggests that the *fms* and *fmt* genes are closely linked to each other in the *R. prowazekii* genome.

# A.



# B.




Figure 1a: Schematic presentation of the region surrounding the *fms* and *fmt* genes in *R. prowazekii*. (A) small subunit rDNA region. (B) large subunit rDNA region. The location of clone 132 that was used in sequencing is indicated. Thin lines indicate a noncoding region; Boxes indicate a coding region. Abbreviations used to indicate restriction enzyme sites are as follows: B = *Bam* HI, RI = *Eco* RI, RV = *Eco* RV, H = *Hind* III, X = *Xho* I. Abbreviations used to indicate genes are as follows: rrs = rRNA small subunit gene, orf = open reading frame, *gyr A* = gyrase A gene, *fms* = peptide deformylase gene, *fmt* = methionyl-tRNA<sup>Met</sup> formyltransferase gene, rrl = rRNA large subunit, rrf = rRNA subunit.

## 2. Material and method

### 2.1. Purification of plasmids.

DNA plasmid containing the clone 132 was prepared using Qiagen Maxi Kit (KEBU, Stockholm, Sweden).

### 2.2. DNA sequencing

Autoread TM sequencing kit (Pharmacia, Uppsala, Sweden) which used Sanger's method (dideoxy chain termination method) was used for sequencing the clone 132.

For the DNA sequencing of both strands of the insert we used modified T7 DNA polymerase (sequenase) and fluorescent forward and reverse primers. The sequence of the gene was performed by following the primer walking method and the non-fluorescent primers deduced are used by addition of fluorescent ATP (Pharmacia, Uppsala, Sweden).

The product of the sequencing reaction were separated and analysed with the aid of an ALF (Automated Laser Fluorescent) DNA sequencer (Pharmacia, Uppsala, Sweden).

### 2.3. Sequence analysis

The data banks were searched for homologous sequences using the BLAST (Basic Local Alignment Search Tool) program (Altsihul et al., 1990).

The predicted protein products of the *fms* and *fnt* genes were aligned (Fig.3a and 3b) using CLUSTAL W (Higgins et al., 1992).

### 2.4. PCR amplification

The polymerase chain reaction (PCR) amplification were carried out in 50 µl reaction mixture containing 10 nmoles of dATP, dCTP, dGTP and dTTP, 50 pmoles each of indicated primer, 20 to 50 ng of

ATG GAA TTC ATC TTG ACG GTT ACA GAG AAC GGT TTT GGT AAA AGA AGC TCG GCA TAT GGT  
M E F I L T V T E N G F G K R S S A Y G  
TAT AGA ATT ACT GAT CGT GGT GGT AGT GGT ATA ATA AAT ATG GAT ATT AAC GAT AAA ACC  
Y R I T D R G G S G I I N M D I N D K T  
GGT TTA GTC GTT GGT GTT ATG CCA GTT AAA ATG GAT GAT GAG TTA ATG CTA ATC ACA AAT  
G L V V G V M P V K M D D E L M L I T N  
AGT GGT AAG TTA ATT CGC TGT AAA CTT GAG TCG GTA CGT ATT ACG GGT CGT AAC ACT AGT  
S G K L I R C K L E S V R I T G R N T S  
GGT GTA ATT CTG TTT AAA CTA GAT GAT GAT GAA AAA GTA GTA TCT GTT TCT TTA ATA GCT  
G V I L F K L D D D E K V V S V S L I A  
GAA ACT TCT GAA AGC GGA GAA GCT AGT GAA TTA GCG GAA GAG GGA TTA GAG AAT GAT GTT  
E T S E S G E A S E L A E E G L E N D V  
AAG GTA TAG TAGTGGAAGTATATGCTGATTTTATTTAAACCCATGGACATTTTTTATATATTATTCATTTAAAACCGG  
K V \*  
CTCTTTTTGCAGCGCATGACTTGATATATAGTACTAAATCAATACTAATATTAGTATTTTTAGTCTCGATCATTGAATAG  
TTATGATATTAATTAATAATGTCTTGTACTTTTCATCTTAATTAATAAGAACAATTTATGTGCTTTTTGTTGTTAAATTA  
TTTACCTATTTTTATAACCATGATGTTAATCAATGCATTTATCTGTTTGTGTTGCGTTTAGCAATATCTCATATAAAAATTC  
TAGAAAACACTAATTTTTATAAAAATAGATATCAATTTTTGAATATACAAAATATTACTTTTTAATCTTGTACTACTA  
TAATAGGTTTCAGCGAGTAATAATATTTCTACATTAATTAGATCAATCTATTTCCAATGCACATAATTATTAATCAGI  
TTTTATGTTTTCGTAACATACAACCTGTGGCTATTAATTATAGCTGTTATTTTTGTTACACACA\*CTCAATTACTTTCAAA  
CTTATTATTTTTTGTTTACCTTTTAAATGTTAGC\*AAAaGAGCTAGCCcAAAAGATTTTCATTCATATATTTTATGT\*TATC  
TGGTATGTGTAGTTAGCGCATACTCCTATAATGATATCAATTTTTATTTATCTA\*TATAGTATAGGAGAAACGATAAAATGA  
AAGT\*TTGATGTTTTCTTTGATTTTTGCTATCATTATTTCTAATGACTTTAGTATTTTTAAAAATTTTGTGTTTAAAGGAAI  
TGCATTGTACTTCAAGATAGCAGCATTGTAATATTAATTTGTTTATCTATATAATCTATGATCTTTACCATTAAACTAAA  
GTTTTAGAAAAGTATTTAAAGATTATCCCTTTATTTTTTATGATAATTGCATATGCTGCAGTTTTGTTATTGCTCGTTT  
ACTTCAGTAATAGCGGAGTTAAGAATATTA AAAAGTGTGATTTTAAACGATAAAATCCCGAGATATGATGAGATGATATAAC  
TTGATCTAATATTTAGTAGAAAATAATACAAAAATAAAGCGCTTGCTCGTACTGTTATTATAGTAAAAGTAAATATTTAI  
TCACTATTGAGTGTTTTACTCCGATTTTAGCGTAAGTAATCCTAAAAAGTTGTTAATTAGCTTTAATAGCGATAATAATA  
TCAAGGTGGTGTAATTTATGTAATGCTTGCTATTTCTGTTAAAAAGACATATATTTATATCTTATTTAATCCTCTTATTT  
TTATAAAATGATGTTTTTTTTAAGATTGATAGATAATCTATATATCTATAACTCAGAAAACCTTAAAATAACGCAAAAT  
TGTATGGATTAAATGCCATGT\*AACT\*ATAAAAATTTTAGCAATATTTATTAAGTATCTCTTTAGGTTTCTGACAAAAA  
TTCTCTTAAACGCACTATATTTCAACTAAAAATTAATGATGATAAATCGTTCTTTTATCAATTAAGTATTACATTTATTA  
TATTCAATAAATTAATCCATATATTATTTTTTTTATAGAAGAAATTTTTGAATTGGAATTTTTTATTAATTAAGATAA  
TTGCTCTTTAAAAGCACAAAAGCATATAATCTTGTTGTAAGATTTATTTTTAGTATAAAATTAATGATGTGGTTATTAAT  
GCCATTGGGTTAACCGTTTCAGTTCTTATTATAGGTATAGTGTCTATGGCTATAGGAGGAAAAATTGATAAGAAATTGAG  
CTTAAAACATAATGACTTTGAGAGTATTTTTTTCAGGCCGTTTCTATATTTTTTATTAATTTATTTTTTATAAAGTTCATTT  
TATACTAAAATTAAGTGAATGATCTATTTTTTATAGGTGAATTTTTATTTTTTAGTAAGATTTATATCTCTTAAACTATT  
TACTTCATGAGGTATGTAATAATCAAAAGATATTATAACCCAAACTGTTAATACCAATTCAGATCAATAACTACTTAATT  
TAT\*TCTATCATCTCTACAAATAATTTTTTCAATTTGTCTAGAGTTTTTTGTTATAACTTGTCTACTTCATCTCTAGAA  
CCTTTTTAA\*TCATCAATTTCTCTAGCATTGATAATGCATGATATCGCTATAATGTGTTGAGTTATTCACAAATTTCT  
CCATTATCACACTATAATATATTAAATCTTCCTTAATTGATACATATATTTTATGCTGTCTGATATTTGTTATAATCA  
TGCAATAAATGATTGAAAGTAGGTGAGCTTATTATACAACTATTACCGGAATTAGCAAGATTTACTTTCTTTGCATTTA  
ATTTTATTGTATATTATAAGTAAAAGATTAAGATTGAATT ATG TCA ATA TTC TCC ATA GTA ACC GCA CCI  
GAC GAA AGA TTA AAG CAA AAA TCT AAG CCT GTT TTA GAA TGT ACC GAT CAA ACA CGA AAA  
D E R L K Q K S K P V L E C T D Q T R K  
TTT ATG CAT GAT ATG CTT GAA ACT ATG TAC AAT GCG GAT GGT GCA GGG CTT GCG GCA GTA  
F M H D M L E T M Y N A D G A G L A A V  
CAA GTA GGA ATA TTG GCA CGT ATT TTA GTA ATT GAT ATA AAA GAA TAT GAT CCA GTA GAA  
Q V G I L A R I L V I D I K E Y D P V E  
AGA CCG AAA GAT TTT TAT CCA CTT TTT ATA GTG AAC CCT GAA ATA ATA GAA AAA TCA ACA  
R P K D F Y P L F I V N P E I I E K S T  
GAG TTA GTG ACT GCT AAT GAA GGC TGT ATC TCA TTA CCA AAG CAA CGT ATT GAG GTT ACC  
E L V T A N E G C I S L P K Q R I E V T

AGA	CCG	GAG	TCT	GTA	AAG	ATA	AGA	TAT	TTA	GAT	TAT	CAT	GGT	AAA	TCA	CAA	GAG	CTA	AAA
R	P	E	S	V	K	I	R	Y	L	D	Y	H	G	K	S	Q	E	L	K
GCA	AAT	GAT	TGG	CTT	GCA	AGA	GTT	ATT	CAG	CAT	GAG	TAT	GAT	CAT	TTA	GAA	GGT	AAG	CTT
A	N	D	W	L	A	R	V	I	Q	H	E	Y	D	H	L	E	G	K	L
ATG	GTT	GAT	TAT	TTA	AGT	AAT	TTA	AAA	CGA	GAT	GTA	GTA	CTT	CGT	AAG	CTT	AAG	AAA	CTT
M	V	D	Y	L	S	N	L	K	R	D	V	V	L	R	K	L	K	K	L
AAA	AAT	AAT	ATA	GTG	TGA	AAGTAATTTTT	ATG	GGA	ACG	CCT	GAA								
K	N	N	I	V	*		M	G	T	P	E								

Figure 1b: Nucleotide sequence of the 5' region of *fms* and the deduced amino acid sequence of the *fms* gene product. Asterisks signify stop codons and, in the spacer, asterisks signify ambiguities.

a b λ




Figure 2. PCR analysis. Primers 132+2 and ot56-7 with genomic DNA from *R. prowazekii* isolated in H. Winklers laboratory, Mobile, Alabama (lane a). Lane b are water controls. Molecular marker were based on digests of lambda DNA with the enzymes EcoR I and Hind III, producing the following sizes of bands in kb: 21.2, 5.1, 4.9, 4.2, 3.5, 2.0, 1.9, 1.584, 1.375, 0.974, 0.831, 0.563 and 0.125.

chromosomal DNA, 1.25 units of Taq polymerase (Promega Biotech, Madison, WI) and 5  $\mu$ l 10X buffer (0.5M Tris-HCl pH 8.8, 0.15M  $(\text{NH}_4)_2\text{SO}_4$ , 15mM  $\text{MgCl}_2$ , 1% Triton X-100 and 0.1% gelatin). After a hot start, 30 cycles of amplification was performed in a DNA thermal cycler (AMS Biotechnology, Taby, Sweden), with a standard profile of denaturation at 94°C for 1min, annealing at 53°C for 1 min, and extension at 72°C for 1 or 2 min. Following amplification, 5 to 10  $\mu$ l of the amplified material was fractionated by electrophoresis on 1% agarose gels.

### **3. Results**

#### **3.1. Nucleotide sequence of the *fms* gene and upstream regions.**

In order to study the arrangement of the *fms* and *fmt* gene, we selected for sequence analysis clone 132 (Fig. 1a) from a set of 1000 clones randomly isolated from a genomic library constructed in lambda Zap II (Andersson et al., 1995). The nucleotide sequence of the 3 272 base pairs long insert revealed one complete open reading frame of 525 base pairs between positions 2718 and 3243 (Fig. 1b). The overall G+C content of this ORF is 33.7%, with the first and the second codon positions having higher G+C content values (50.3% and 29.1%, respectively) than the third (21.7%). The observed position-specific differences in base frequency patterns are in perfect agreement with the results obtained from an analysis of 27 protein coding genes (Andersson and Sharp, 1995), suggesting that the open reading frame identified corresponds to an actively transcribed gene. Translation of this gene produces a protein of 175 amino acids; a search in the database for homologous sequences using the BLAST program (Altschul et al., 1990) gives strongest homology to the *E. coli* peptide deformylase, encoded by the gene *fms* (Fig. 3a)

A short open reading within the first 363 nucleotides of clone 132 was found to correspond to the previously sequenced gene *gyrA*, encoding DNA gyrase (Wood and Waite, 1994). An homology search of the intergenic region (2355 bp) of *gyrA* and *fms* failed to identify any homologs in the databanks. The average G+C content of this regions is only 23.3%, which is close to 21.4%, previously calculated from a large



```

Rp.fms -----MSIFSIVTAPDERLKQKSKPVLECTDQTRKFMHDMLETMYN-----AD
Tt.fms -----MVYPIRLYGDPVLRKARPVEDFSG- IKRLAEDMLETMFE-----AK
Ec.fms -----MSVLQVLHIPDERLRKVAKPVEEVNAEIQRIVDDMFETMYA-----EE
Mg.def MTFQPTKTWLVFDDNALINKPTEAVNFPIDEQIETCIKKMIAVVDASYDGKAQEYDIIP
Hi.def -----MTALNVLIIYPDDHLKVVCEPVTKVNDAIRKIVDDMFDTMYQ-----EK
                                     * . . . .

Rp.fms AGLAAVQVGILARILVIDIKEYDPVERPKD-----FYPLFIVNPEIIEKSTE---L-VT
Tt.fms VGLAAPQIGLSQRLFVAVEYADEPEGEEERPLRELVRRVYVVAWWRGGEPGDHLPGGTG
Ec.fms IGLAATQVDIHQRIIVIDVSENDRER-----LVLINPELLEKSSE-----TG
Mg.def IGIAANQIGYWKQLFYIHLNDLNKEKK-----CLLINPKIIDQSENKAF--LE
Hi.def IGLAAPQVDILQRIITIDVEGDKQNQ-----FVLINPEILASEGE-----TG
      * . * * * . . . . .

Rp.fms NEGCLSLPKQ---RIEVT---RPESVKIRYLDYH-GKSQELKANDWLARVIQHEYDHLE
Tt.fms GDGGVPLPARPLLRRGGAPGGAHPGGVPPGRGGPWARVGARGVHGPGPLPAZDRPSGRDPLL
Ec.fms EEGCLSIPEQ---RALVP---RAEKVKIRALDRD-GKPFLEADGLLAICIQHEMDHLV
Mg.def GEGCLSVKKQH--KGYVI---RSEWITIKGYDWFEEKKEITIKATGLFGMCLQHEFDHLQ
Hi.def EEGCLSIPGF---RALVP---RKEKVTVRALDRD-GKEFTLDADGLLAICIQHEIDHLN
      * . . . . . * *

Rp.fms KLMVDYLSNLK--RDVVLRLKLLKLNIV-----
Tt.fms APAQAQAGGLSGGQPGGARPLPEGGPGLAEGAFPGM
Ec.fms KLFMDYLSPLK--QQRIRQKVEKLDRLKARAZL---
Mg.def RFFYQRINPLN--PWFKKPEWKVINPTLKTSNG---
Hi.def ILFVDYLSPLK--RQR!KEKLIKYKKQIAKS-----
                                     *

```

Figure 3a: Multiple alignment of the deduced amino acid sequences of the *fms* gene in 5 bacterial species. Symbols beneath the aligned sequences indicate identical residues (\*) and sites with conservative replacements (.). The 34 amino acids in the 5' end of the *fms* gene in *M. genitalium* are not represented. Abbreviations used are: Rp = *R. prowazekii*, Tt = *T. thermophilus*, Ec = *E. coli*, Mg = *M. genitalium*, *fms* = *def* = peptide deformylase gene.

```

Rp.fmt -----MGTPEFAVPTLKKLIH--HEVKAVFTQQPKAKGR-GLHLAK-SPIHQLAF
Tt.fmt ----MRVAFFGTPLWA-PCWTPYASA---TRWSWWS-PSPTSPRAGAZGPPRAPWPATPR
Ec.fmt VSESLRIIFAGTPDFAARHLDALSSG-HNVVGVFTQPD RPAGR-GKKLMP-SPVKVLAE
Mg.fmt ---MFKIVFFGTSTLSKKCLEQLFYDNDFEICAVVTQPDKINHR-NNKIVP-SDVKSFCL
Hi.fmt -MKSLNIIFAGTPDFAAQHLOAILNSQ-HNVIAVYTQPDKPAGR-GKKLQA-SPVKQLAE
          **                               *

Rp.fmt EHQIPVYSPSTLRNDETINLIKKIDADIIIVVIAYGFIVPKAILEAKKYGCLNIHPSDLPR
Tt.fmt RRGFPFCARRGLGRRPSSRPCARPRRWRWRPTGSSSPRRPWTSPHTASSTSTPPSSPS
Ec.fmt EKGLPVFQPVSLRPQENQQLVAELQADVMVVVAYGLILPKAVLEMPRLGCINVHGSLLPR
Mg.fmt EKNITFFQPK--QSIKADLEKLEKADIGICVSFGQYLHQDIIDLFPNKVINLHPSKPLPL
Hi.fmt QNNIPVYQPKSLRKEEAQSELKALNADVMVVVAYGLILPKAVLDAPRLGCLNVHGSILPR
          *                               *

Rp.fmt HRGAAPLQRTIIIEGDRKSSVCIMRMDSGLDTDG DILLKEDLNLERRITLDELSNKCAHLGA
Tt.fmt TGGRPPSSGPGSSPGNGRPGSPSCAWTRAWTPVPSTPSGGRPSCRTRTPWPWGTGSGTRGW
Ec.fmt WRGAAPIQRSWAGDAETGVTIMQMDVGLDTGDMLYKLSCPITAEDTSGTLYDKLAE LGP
Mg.fmt LRGGAPLHWTIINGFKKSALSVIQLVKKMDAGPIWKQDFLVNNDWNTGDL SIYVEEHSP
Hi.fmt WRGAAPIQRSIWAGDVQGTGVTIMQMDVGLDTGDMLHKVYCDILPTETSTSLYNKLAELAP
          * * *

Rp.fmt ELLIKTLANIDN--IVPIKQSSNGITYAHKLTKAEGKINWYESAYSIDCKIRG--MNPWP
Tt.fmt SSFLRCWSASRS--SPPGPRKGRSPTHRPFPRR---RGGWTSGRAPRPSTAATGPSSSPGP
Ec.fmt QGLITTLKQLADGTAKPEVQDETLVTYAEKLSKEEARIDWLSLSAAQLERCIRA--FNPWP
Mg.fmt SFLIECTKEILN-KKGKWFQIGEP TFGLNIRKEQEHLDLNQIYKSFLNWVKG--LAPKP
Hi.fmt SALIDVLDNLENGKFIAEKQDGSQSNYAEKLSKEEAQLNWSLSAMQLERNIRA--FNPWP
          * *

Rp.fmt GAYFSYNDKI-----IKILRAEYFNYNHH-FIPGTVINN---KLEIACGSG-ILRVKKLQ
Tt.fmt GATSSTGAGG----SRPZGSAPSPGKSGPGWPGWRRARWARPRASSSS--WRS-SLR
Ec.fmt MSWLEIEGQP-----VKVWKASVIDTATN-AAPGTILEANKQGIQVATGDG-ILNLLSLQ
Mg.fmt GGWLSFEGKN-----IKIFKAKYVSKS NYKHQLGEIVNISRKGINIALKSNEIISIEKIQ
Hi.fmt IAYFSTEDKDGNAHTLKVYQAKVLP HQD--KPAGTILSADKNGIQIATVDG-VLNL LQLQ
          * * *

Rp.fmt QESKKALNIEEFLRGTN--ILKDTILKZ
Tt.fmt EGGPCPPPTGPGAMAWP--RGPGSARY-
Ec.fmt PAGKKAMSAQDLLNSRREWFVPGNRLVZ
Mg.fmt IPGKRVMEVSEIINGKHP-FVVGKCFK-
Hi.fmt SAGKKPMSAQDLLNGRAEWF TIGKVL A-

```

Figure 3b: Multiple alignment of the deduced amino acid sequences of the *fmt* gene in 5 bacterial species. Symbols beneath the aligned sequences indicate identical residues (\*) and sites with conservative replacements (.). Abbreviations used are: Rp = *R. prowazekii*, Tt = *T. thermophilus*, Ec = *E. coli*, Mg = *M. genitalium*, *fmt* = methionyl-tRNA<sup>Met</sup> formyltransferase gene

number of intergenic regions (Andersson and Sharp, 1995). This is consistent with the interpretation that the region between *gyrA* and *fms* corresponds to noncoding DNA.

### 3.2. The *fms* gene is linked to the *fmt* gene in *Rickettsia prowazekii*.

Sequence analysis of the end-fragments of clone 132 and t56 suggests that these may share a common genomic breakpoint (Fig. 1a). To determine whether *fms* is indeed linked to *fmt* in the genome, we amplified by PCR the region in between these two genes by using a primer from the upstream region of the *fms* gene, 132+2 (ATTAGCAAGATTTACTTTCTT) together with a primer from within the *fmt* gene, ot56-7 (CACTGCCACAAGCGATT). According to sequence data we expect a genomic fragment of 1 361 base pairs to be amplified in this reaction. Indeed, a fragment of the expected size was obtained (Fig. 2), confirming that the *fms* gene is located immediately upstream of the *fmt* gene in *R. prowazekii*.

### 3.3. Comparative analysis of the *fms* and *fmt* gene products.

The amino acid sequences of the peptide deformylase (PDF) and methionyl-tRNA<sup>Met</sup> formyltransferase (MTF) from five different bacterial species (including *Rickettsia prowazekii* sequences) have been analysed. Three of these belong to the Proteobacteria; *Rickettsia prowazekii* to the alpha proteobacteria, *Haemophilus influenza* and *Escherichia coli* to the gamma proteobacteria. *Mycoplasma genitalium* belongs to the low G+C Gram-positive lineage. *Thermus thermophilus* has been placed in a lineage near to the green non-sulfur bacteria (Olsen et al., 1994).

The sequence alignments of the *fms* and *fmt* gene products were performed using CLUSTAL W (Higgins et al., 1992) and are presented in Figs. 3a and 3b. The two proteins show approximately the same degree of functional conservation, as inferred from the percent identity values in a direct comparison of the *R. prowazekii* and *E. coli* homologs; 48.5% and 39.8% for PDF and MTF, respectively.




Figure 4. Organization of the region surrounding the genes *fms* and *fmt* in five bacterial species. Abbreviations used to indicate genes are as follows: *fms* = peptide deformylase gene, *fmt* = methionyl-tRNA<sup>Met</sup> formyltransferase gene, *gyr A* = gyrase A gene, 16S = rRNA small subunit, 23S = rRNA large subunit, 5S = rRNA subunit, *orf* = open reading frame, *gmk* = 5' guanylate kinase gene and *mob 13* = mobilisation protein gene.

The peptide deformylase from *M. genitalium* is about 41 amino acids longer at the N-terminal end than its homologs in the other species. Interestingly, there is a Methionine codon at amino acid position 34 that if used as the initiation codon would produce a protein of approximately the same length as observed in the other species. The gene encoding PDF in *M. genitalium* is located in the opposite direction of its surrounding genes and is overlapping with as much as 269 bases at its 5' end with the *gmk* gene and 23 bases with its other neighbouring gene. The overlap with the *gmk* (5'guanylate kinase) gene is such that the third codon is shared by the two genes. The G+C content values at the first two positions are more similar to the *gmk* gene upstream of the Met-34 codon, but more similar to the *fms* gene downstream of the Met-34 codon. A translation initiation start site at codon position 34 in *M. genitalium* would produce a protein that is only 8 amino acids longer than its homologs. Most of the gaps observed in the alignment appear to be caused by the *T. thermophilus* sequence which has a few extra amino acids at a number of different positions. The region between positions 60-69 appear to be particularly well conserved among these five bacterial species.

The *fmt* gene product is of approximately the same length in all species, but with only a few positions being conserved among all of the species analysed.

#### 3.4. Organization of the *fms-fmt* locus in bacterial genomes.

The organization of the *fms* and *fmt* genes as well as their genomic neighbourhoods is presented in Fig. 4. Here it is shown that *fms* and *fmt* are genetically linked in *R. prowazekii*, *E. coli*, *T. thermophilus* and *H. influenzae*, but that the two genes are separated from each other in *M. genitalium*.

The conservation of gene order appears to be restricted to the *fms-fmt* locus since no similarities between any of the species can be observed with respect to their flanking sequences. It is interesting to note however that the *fms-fmt* locus is located immediately downstream of the rRNA operon in *H. influenzae* but immediately upstream of the 23S-5S rRNA gene cluster in *R. prowazekii*.

#### 4. Concluding remarks.

In this report, we have shown that in *R. prowazekii* the *fms* gene which encodes peptide deformylase (PDF) is positioned immediately upstream of the *fmt* gene which encodes methionyl-tRNA<sub>f</sub><sup>Met</sup> formyltransferase (MTF). These two enzymes are involved in the formylation and subsequent deformylation of the formyl group of the initiator Met-tRNA<sub>f</sub><sup>Met</sup> during protein synthesis. That genes encoding functionally related enzymes have also been organized into the same operons appear to be a common theme in bacterial evolution. Colocalisation of *fms* and *fmt* has been demonstrated in organisms as different as *H. influenzae* (Fleischman et al., 1995), *E. coli* (Guillone et al., 1992) and *T. thermophilus* (Meinzel and Blanquet, 1994). Since it seems unlikely that *fms* and *fmt* should have become linked to each other independently in different bacterial lineages, the colocalization of these two genes probably represents the organization of these two genes in a bacterial ancestor. The sole exception observed so far is *M. genitalium* where *fms* and *fmt* are several hundred kb apart on the chromosome. Since the evolution of *M. genitalium* is believed to have been associated with a large number of deletion and rearrangement events, the disruption of the *fms* and *fmt* linkage probably represents a recent, derived event. Despite the fact that *R. prowazekii* similarly has a small genome size and is thought to have undergone a reductive mode of evolution as reflected in low copy numbers and rearranged operon structures for the rRNA and elongation factor genes (Andersson et al., 1995; Syvänen et al., 1995), the ancestral arrangement of *fms* and *fmt* has not been disrupted. Thus, the rearranged 23S-5S rRNA gene cluster is flanked by an operon that is highly preserved in structure.

**Rearrangement of rRNA Genes in *Rickettsia*  
Preceded the Divergence of the Typhus and the  
Spotted Fever Group *Rickettsia***

Magali P. C. Romedenne<sup>1\*</sup>, Diane Stothardt<sup>2</sup>, Siv G. E. Andersson<sup>1</sup>  
and Charles G. Kurland<sup>1</sup>

<sup>1</sup>*Department of Molecular Biology, Uppsala University, Biomedical  
Center, Box 590, S751 24 Uppsala, Sweden*

<sup>2</sup>*Department of Ecology and Evolutionary Biology, University of  
Arizona,  
Tucson, AZ 85721, USA*

*\*Present address: Laboratory of Genetics, Facultés Universitaires  
Notre dame de la Paix, 61 rue de Bruxelles, 5000 Namur, Belgium*

Running title: rRNA genes in *Rickettsia*

Keywords: genomic organization, molecular evolution, *Rickettsia*

## Summary

An unusual arrangement of the rRNA genes has previously been observed in *Rickettsia prowazekii*, where the 23S rRNA gene is preceded by *fmt* which encodes methionyl-tRNA<sup>fMet</sup> formyltransferase (Andersson et al., 1995). In this study we show that *fmt* is located upstream of *rrl* also in the genomes of *R. sibirica*, *R. parkeri*, *R. rickettsii*, *R. amblyomii* as well as of the ELB agent. These findings suggest that the genomic rearrangement of the rRNA genes preceded the divergence of the typhus and the spotted fever group rickettsia.



## 1. Introduction

Rickettsia are small, gram-negative bacteria that live in close association with eukaryotic cells (Weiss and Moulder, 1984). They are members of the family *Rickettsiaceae*, which includes the three tribes *Rickettsiacea*, *Ehrlichiaea* and *Wolbachia*. The genus *Rickettsia* can be further divided into the spotted fever group (SFG), the typhus group (TG) and the scrub typhus group (STG).

Members of the spotted fever group *Rickettsia* are distributed worldwide. These bacteria grow and divide primarily within the host cell cytoplasm, but are capable of multiplying within the cell nucleus as well. Many species are pathogens causing diseases such as for example Siberian Tick Typhus (*R. sibirica*), Rickettsialpox (*R. akari*) and Queensland Tick Typhus (*R. australis*). The typhus group *Rickettsia* grow and divide exclusively within the host cell cytoplasm and include species such as *R. prowazekii* and *R. typhi*, the etiological agents of epidemic and murine typhus, respectively. The scrub typhus group of *Rickettsia* is composed of several heterogeneous strains of *R. tsutsugamushi*, the causative agent of scrub typhus. A few species can not easily be classified into either of these three categories. The ELB agent for example was first classified into the typhus group rickettsia based on the observation that it causes a disease similar to murine typhus in human populations (Azad et al., 1992; Williams et al., 1992). However, sequence comparisons of the 17 kDa antigen gene and the 16S rRNA gene have subsequently shown that this species may be more closely related to the spotted fever group rickettsia than to the typhus group rickettsia (Azad et al., 1992; Williams et al., 1992).

Features in common of all the different species of *Rickettsia* are their intracellular growth environment, their small genome sizes (1.2 Mb) (Eremeeva et al., 1994) and their low genomic G+C content values (29-33%). It is generally believed that the *Rickettsia* have evolved from free-living bacteria with larger size genomes (Andersson and Kurland, 1995). From this perspective it seems plausible that the single copy number of the rRNA genes and the disruption of the traditional rRNA operon structure as observed in *R. prowazekii* is the result of a degenerate mode of evolution (Andersson et al., 1995). In this study, we examine the genus *Rickettsia* for the occurrence and

distribution of the unusual arrangement of rRNA genes previously observed in *R. prowazekii* (Andersson et al., 1995). We report here the sequence of the intergenic region between *fmt* and the 23S rRNA gene in six different Rickettsia species.

## **2. Materials and methods**

### **2.1. PCR amplification and purification**

The polymerase chain reaction (PCR) amplification were carried out in 100 µl reaction mixture containing 20 nmoles of dATP, dCTP, dGTP and dTTP, 100 pmoles each of indicated primer, 40 to 100 ng of chromosomal DNA, 2.5 units of Taq polymerase (Promega Biotech, Madison, WI) and 10 µl 10X buffer (0.5M Tris-HCl pH 8.8, 0.15M (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 15mM MgCl<sub>2</sub>, 1% Triton X-100 and 0.1% gelatin). After a hot start, 30 cycles of amplification was performed in a DNA thermal cycler (AMS Biotechnology, Taby, Sweden), with a standard profile of denaturation at 94°C for 1min, annealing at 53°C for 1 min, and extension at 72°C for 1 or 2 min. Following amplification, 5 to 10 µl of the amplified material was fractionated by electrophoresis on 1% agarose gels.

The PCR products were purified by using Wizard PCR Preps DNA purification System (Promega). The concentration of the final product was measured by running a 5 µl on a 1% agarose gel side by side with lambda DNA of known concentration.

### **2.2. DNA sequencing**

Thermo Sequenase fluorescent labelled primer cycle sequencing kit (Amersham) which used dideoxy chain termination method was used to sequence PCR products. Cycle sequencing uses repeat cycles of thermal denaturation, annealing and elongation/termination.

For the DNA sequencing we used Thermo sequenase (Amersham) and fluorescent primers.




Figure 1a: PCR analysis. Primers ot56+2 and ot56-2 with genomic DNA from *R. prowazekii* (lane b), *R. sibirica* (lane d), *R. parkerii* (lane f). Lanes a, c and e are water controls. Molecular marker were based on digests of lambda DNA with the single enzyme *Hind* III, producing the following sizes of bands in kb: 23.1, 9.4, 6.6, 4.4, 2.3, 2.0 and 0.6.




Figure 1b: PCR analysis. Primers ot56+2 and ot56-2 with genomic DNA from *R. prowazekii* (lane b), ELB agent (lane d), *R. rickettsii* (lane f), *R. montana* (lane h) and *R. amblyomii* (lane j). Lanes a, c, e, g and i are water controls. Molecular marker were based on digests of lambda DNA with the single enzyme *Hind* III, producing the following sizes of bands in kb: 23.1, 9.4, 6.6, 4.4, 2.3, 2.0 and 0.6.

The product of the sequencing reaction were separated and analysed with the aid of an ALF (Automated Laser Fluorescent) DNA sequencer (Pharmacia, Uppsala, Sweden).

### 2.3. Sequence analysis

The alignment of the *fmt* gene and the spacer were performed by using CLUSTAL W (Higgins et al., 1992) as well as the tree deduced following the Neighbor-joining method.

The Ks and Ka were calculated with the Li's method (Li, W-H., 1993)

## 3. Results

### 3.1. PCR amplifications with primers from *fmt* and 23S rRNA

To study the genomic neighbourhoods of the *fmt* gene and the 23S rRNA gene we performed PCR reactions using one primer from within the *fmt* gene, ot56+2 (CTAAAGCAGAAGGAAAAATT) and one primer from within the *rrl* gene, ot56-2 (GCTTCTAGTGCCAAGGCATC). This primer pair yielded a product of 1.0 kb in size when genomic DNA from *R. prowazekii* was used to drive the PCR reaction (Fig. 1), as previously observed (Andersson et al., 1995). Successful PCR amplifications were also obtained when these primers were used in PCR reactions together with genomic DNA from the following species: *R. sibirica*, *R. parkeri*, *R. rickettsii*, *R. montana*, *R. ablyommii* and the ELB agent (Fig. 1). This suggests that the *fmt* and the 23S rRNA genes are near to each other in all of these genomes. Some size variation of the amplified fragments was observed, with genomic DNA from the ELB agent producing a PCR product of only 500 bp in length.

	1	2	3	4	5	6	7	8
1 P1;Rsib <i>fmt</i>	0.0	1.3	0.5	1.0	1.3	1.6	8.9	6.6
2 P1;Rrick <i>fmt</i>	5.0	0.0	0.7	1.3	1.5	1.8	9.2	6.9
3 P1;Rpark <i>fmt</i>	1.3	5.0	0.0	0.5	0.7	1.0	8.4	6.0
4 P1;Ramb <i>fmt</i>	0.0	5.0	1.3	0.0	1.3	1.6	8.1	5.8
5 P1;ELB <i>fmt</i>	3.9	9.1	5.3	3.9	0.0	1.8	7.5	5.2
6 P1;Rmont <i>fmt</i>	1.3	6.1	2.6	1.3	5.2	0.0	9.5	7.2
7 P1;Rp <i>fmt</i>	26.5	22.6	26.5	26.5	33.1	27.6	0.0	4.2
8 P1;Rty <i>fmt</i>	23.2	20.5	23.2	23.2	29.2	25.3	11.7	0.0

Table 1: Evolutionary distances based on the *fmt* nucleotide sequences. Substitutions at synonymous sites ( $K_a$ ) are shown above the diagonal. Substitutions at nonsynonymous sites ( $K_s$ ) are shown below the diagonal.

```

b_fmt      YFSYNDKIIKILEAEYLNADHHCTSGTVISDKLEIACGSGILRVKKLQQESKKALNIEEF
ck_fmt     YFSYNDKIIKILEAEYLNADHHFTSGTVISDKLEIACGSGILRVKKLQQESKKALSIEEF
rk_fmt     YFSYNDKIIKILEAEYLNADHHFTSGTVISDKLEIACGSGILRVKKLQQESKKALNIEEF
b_fmt      YFSYNDKIIKILEAEYLNADHHFTAGTVISDKLEIACGSGILRVKKLQQESKKALNIEEF
b_fmt      YFSYNDKIIKILEAEYLNADHHFTSGTVISDKLEIACGSGILRVKKLQQESKKALNIEEF
nt_fmt     YFSYNDKIIKILEAEYLNAAHHFTSGTLISDKLEIACGSGILRVKKLQQESKKALNIEEF
ow_fmt     YFSYNDKIIKILRAEYFNYNHHFIPGTVINNKLEIACGSGILRVKKLQQESKKALNIEEF
r_fmt      YFSYNDKIIKILEAEYLNYNHHFIPGTVISNKLEIACGSGILRVTKLQQESKKALNIEAF
***** ** * ** * * ***** ***** ** *

b_fmt      LRGTNILKDTVVKZ
ck_fmt     LRGTNILKDTVVKZ
rk_fmt     LRGTNILKDTVVKZ
b_fmt      LRGTNILKDTVVKZ
b_fmt      LRGTNIL-DTILKZ
nt_fmt     LRGTNILKDTVVKZ
ow_fmt     LRGTNILKDTILKZ
r_fmt      LRGTNILKDTILKZ
***** ** .***

```

Figure 2. Multiple alignment of the deduced amino acid sequences of the *fmt* gene in eight *Rickettsia* species. Rsib = *R. sibirica* ; Rrick = *R. rickettsii* ; Rpark = *R. parkeri* ; Ramb = *R. amblyomii* ; ELB = *ELB agent* ; Rmont = *R. montana* ; Rprow = *R. prowazekii* ; Rty = *R. typhi*. Symbols beneath the aligned sequences indicate identical residues (\*) and sites with conservative replacements (.).

### 3.2. Comparative nucleotide sequence analysis

The identity of the PCR products obtained were verified by a complete sequence analysis. The sequence data obtained corresponded to 219 basepairs of the 3' end of the *fmt* gene in addition to a spacer region of 303 to 886 base pairs ending 3 base pairs upstream of the 23S rRNA gene. The length of the 3' end of the *fmt* gene was found to be identical in all species, with the exception of the ELB agent in which 3 bases had been deleted. The deletion was found to correspond to exactly one codon and was therefore not associated with any shift in reading frame.

The alignment of the amino acid sequences of the partial *fmt* gene products is presented in Fig. 2. Based on this alignment we have calculated the number of synonymous ( $K_S$ ) and nonsynonymous ( $K_A$ ) nucleotide substitutions in the *fmt* gene among the eight species analysed (Table 1). As expected, for most pairs the  $K_S$  values are higher than the  $K_A$  values. The observed substitution frequencies clearly show that the two members of the typhus group rickettsia (*R. prowazekii* and *R. typhi*) are distinct from the members of the spotted fever group rickettsia. On the average, the nonsynonymous and synonymous substitution frequencies are 7.4% and 25.6%, respectively, between the two groups of species. For comparison, it can be noted that the synonymous substitution frequencies are less than 5% among the different species within the spotted fever group rickettsia. The substitution frequencies observed for the ELB agent suggest that it is more closely related to the spotted fever group rickettsia ( $K_S < 0.09$ ,  $K_A < 0.02$ ) than to the typhus group rickettsia ( $K_S > 0.29$ ,  $K_A > 0.05$ ).

The lengths and the G+C content values of the spacer regions between *fmt* and *rml* are presented in Table 2. In *R. prowazekii* the G+C content of this spacer sequence is 22.5%, close to an average value of 21.4% calculated from a large number of *R. prowazekii* intergenic regions (Andersson and Sharp, 1995). This suggests that the region in between *fmt* and *rml* corresponds to noncoding DNA, as also supported by the lack of homologous sequences in the databanks. The other six species analysed were found to have slightly higher G+C content values ranging from 26.4% (ELB agent) to 28.7% (*R. montanta*). The three

Species	Length (bp)	G+C content (%)
SFG		
<i>R. sibirica</i>	854	28.5
<i>R. rickettsii</i>	886	28.7
<i>R. parkii</i>	857	28.1
<i>R. ambirica</i>	844	27.8
<i>R. montana</i>	845	27.8
TG		
<i>ELB-agent</i>	303	26.4
<i>R. prowazekii</i>	716	22.5

Table 2: Size and G+C content of the spacer region in seven Rickettsia species.



1 vs. 2:	DIST = 0.0235;	p = 0.0176;	q = 0.0059;	length = 851
1 vs. 3:	DIST = 0.0082;	p = 0.0070;	q = 0.0012;	length = 854
1 vs. 4:	DIST = 0.0308;	p = 0.0225;	q = 0.0083;	length = 844
1 vs. 5:	DIST = 0.0284;	p = 0.0213;	q = 0.0071;	length = 845
1 vs. 6:	DIST = 0.0660;	p = 0.0528;	q = 0.0132;	length = 303
1 vs. 7:	DIST = 0.2420;	p = 0.1399;	q = 0.1021;	length = 715
2 vs. 3:	DIST = 0.0234;	p = 0.0187;	q = 0.0047;	length = 854
2 vs. 4:	DIST = 0.0380;	p = 0.0285;	q = 0.0095;	length = 841
2 vs. 5:	DIST = 0.0380;	p = 0.0297;	q = 0.0083;	length = 842
2 vs. 6:	DIST = 0.0627;	p = 0.0462;	q = 0.0165;	length = 303
2 vs. 7:	DIST = 0.2416;	p = 0.1404;	q = 0.1011;	length = 712
3 vs. 4:	DIST = 0.0296;	p = 0.0225;	q = 0.0071;	length = 844
3 vs. 5:	DIST = 0.0296;	p = 0.0237;	q = 0.0059;	length = 845
3 vs. 6:	DIST = 0.0660;	p = 0.0528;	q = 0.0132;	length = 303
3 vs. 7:	DIST = 0.2420;	p = 0.1413;	q = 0.1007;	length = 715
4 vs. 5:	DIST = 0.0178;	p = 0.0154;	q = 0.0024;	length = 843
4 vs. 6:	DIST = 0.0596;	p = 0.0464;	q = 0.0132;	length = 302
4 vs. 7:	DIST = 0.2325;	p = 0.1317;	q = 0.1008;	length = 714
5 vs. 6:	DIST = 0.0660;	p = 0.0528;	q = 0.0132;	length = 303
5 vs. 7:	DIST = 0.2392;	p = 0.1371;	q = 0.1021;	length = 715
6 vs. 7:	DIST = 0.1294;	p = 0.0734;	q = 0.0559;	length = 286

Table 3: Evolutionary distances based on the spacer sequence between *fmt* and *rfl*. Abbreviations are as followed: 1 = *R. sibirica*, 2 = *R. ricketsii*, 3 = *R. parkeri*, 4 = *R. amblyommi*, 5 = *R. montana*, 6 = *ELB agent*, 7 = *R. prowazekii*, DIST = percentage of divergence, p = rate of transition (A<->G; C<->T), q = rate of tranversion, Length = number of sites used in comparison.

Rsib.\_spac GAGTATTATTATGGGTAAAGATAAAAAAGAAGTTAGGTTTAAATGATAATTGAGACGTTAA  
Rrick.\_spa GAGTATTATTATGGGTAAAGATAAAAAAGAAGTTAGGTTTAAATGATAATTGAGACGTTAA  
Rpark.\_spa GAGTATTATTATGGGTAAAGATAAAAAAGAAGTTAGGTTTAAATGATAATTGAGACGTTAA  
Ramb.\_spac GAGTATTATTATGGGTAAAGATAAAAAAGAATTTAGGTTTAAATGATAATTGAGACGTTAA  
Rmont.\_spa GAGTATTATTATGGGTAAAGATAAAAAAGAAGTTAGGTTTAAATGATAATTGAGACGTTAA  
ELB.\_space -----  
Rprow.\_spa -----GAATTTATTAGTTATAAAGCTAAAGT

Rsib.\_spac GGAGTTTACCTCAAAGTTTGAAAAGAATAGTGATATTTATTCCTTTAAAAAGGAAAGTAAT  
Rrick.\_spa GGAGTTTACCTCAAAGTTTGAAAAGAATAGTGATATTTATTCCTTTAAAAAGGAAAGTAAT  
Rpark.\_spa GGAGTTTACCTCAAAGTTTGAAAAGAATAGTGATATTTATTCCTTTAAAAAGGAAAGTAAT  
Ramb.\_spac GGAGTTTACCTCAAAGCTTGAAAAGAATAGTGATATTTATTCCTGTA AAAAGGAAAGTAAT  
Rmont.\_spa GGAGTTTACCTCAAAGCTTGAAAAGAATAGTGATATTTATTCCTGTA AAAAGGAAAGTAAT  
ELB.\_space -----  
Rprow.\_spa TCAGTTTAAATTGTAATTTTCGAGTTTACATATAATTCATAAGAATTTGTAAA-AAAGTAAT

Rsib.\_spac TAATATAGTAG-----ATGATGAAGTAGAG  
Rrick.\_spa TAATATGGTAACAAAGCTTTGATAGCATTTATAAGCATCTATAGTAGATGATGAAGTAGAG  
Rpark.\_spa TAATATAGTAGTA-----GATGATGAAGTAGAG  
Ramb.\_spac TAATATAGTAG-----ATGATGAAGTAGAG  
Rmont.\_spa TAATATAGTAG-----ATGATGAAGTAGAG  
ELB.\_space -----  
Rprow.\_spa TAG-ATATTAG-----ATGATGAGGTAGAA

Rsib.\_spac GATCAAACCTAAGCTACTTACGGAAACTCAATTTCTTGTGCAAAAAACTCCACATGGTAAA  
Rrick.\_spa GATCAAACCTAAGCTACTTACGGAAACTCAATTTCTTGTGCAAAAAACTCCACATGGTAAA  
Rpark.\_spa GATCAAACCTAAGCTACTTACGGAAACTCAATTTCTTGTGCAAAAAACTCCACATGGTAAA  
Ramb.\_spac GATCAAACCTAAGCTATTTATGGAAACTCAATTTCTTGTGCAAAAA-CTCCACATGGTAAA  
Rmont.\_spa GATCAAACCTAAGCTACTTACGGAAACTCAATTTCTTGTGCAAAAAACTCCACATGGTAAA  
ELB.\_space -----  
Rprow.\_spa GATTAAATTGAGATTTTACAG-----AATTTTGTGAAGAAAAATTTCCACTCTAAA

Rsib.\_spac ACATCTCAATATGAAATTAGCGGCTTTATTAACAAAGCCGGCGCACATATAAACGATCCA  
Rrick.\_spa ACATCTCAATATGAAATTAGCGGCTTTATTAACAAAGCCGGCGAACATATAAACGATCCA  
Rpark.\_spa ACATCTCAATATGAAATTAGCGGCTTTATTAACAAAGCCGGCGCACATATAAACGATCCA  
Ramb.\_spac ACATCTCAATATGAAATTAGCGGCTTTATTAACAAAGCCGGCGAACATATAAACGATCCA  
Rmont.\_spa ACATCTCAATATGAAATTAGCAGCTTTATTAACAAAGCCGACGAACATATAAACGATCCA  
ELB.\_space -----  
Rprow.\_spa G-----TCTGAAATTAGTAGCTTGGTT-----GTTAATAAACAT-TGCACAACCTCA

Rsib.\_spac AAGCAAATAAAAAGAAATTTCAAAATTTATACGCTCGTGCCGTATTTGCTGATTTACAGAAA  
Rrick.\_spa AAACAAATAAAAAGAAATTTCAAAATTTATACGCTCATGCCGTATTTGCTGATTTACAGAAA  
Rpark.\_spa AAGCAAATAAAAAGAAATTTCAAAATTTATACGCTCATGCCGTATTTGCTGATTTACAGAAA  
Ramb.\_spac AAGCAAATAAAAAGAAATTTCAAAATTTATACACTTATGCTGCATTTGCTGATTTACAGAAA  
Rmont.\_spa AAGCAAATAAAAAGAAATTTCAAAATTTATACGCTCATGCTGCATTTGCTGATTTACAGAAA  
ELB.\_space -----  
Rprow.\_spa AAATAGATAAAAAGCAATTTTAAAAATTTATATACTTATGATGAATCTGCTAATTTACAGAAA

Rsib.\_spac AATATTTTAGGTAAATTTGATAAGAACGGTAATTTAAGTATAACTAAAGATTTTGCAATTT  
Rrick.\_spa AATATTTTAGGTAAATTTGATAAGAACGGTAATTTAAGTATAACTAAAGATTTTGCAATTT  
Rpark.\_spa AATATTTTAGGTAAATTTGATAAGAACGGTAATTTAAGTATAACTAAAGATTTTGCAATTT  
Ramb.\_spac AATATTTTAGGTAAATTTGATAAGAACGGTAATTTAAGTATAACTAAAGATTTTGCAATTT  
Rmont.\_spa AATATTTTAGGTAAATTTGATAAGAACGGTAATTTAAGTATAACTAAAGATTTTGCAATTT  
ELB.\_space -----  
Rprow.\_spa AATGTTTGAATC-----TAAGATAATCATTTAAT-----AGTTTCAAATC

```

Rsib._spac      GATGAAAATCCAGTCTTTAGAAAAGACGGTAAATTATTATGGGTTAAAACCTCAAAGCGGT
Rrick._spa     GATGAAAATCCAGTCTTTAGAAAAGACGGTAGATTATTATGGGTTAAAACCTCAAAGCGGT
Rpark._spa     GATGAAAATCCAGTCTTTAGAAAAGACGGTAAATTATTATGGGTTAAAACCTCAAAGCGGT
Ramb._spac     GATGAAAATCCAGTCTTTAGAAAAGACGGTAAATTATTATGGGTTAAAACCTCAAAGCGGT
Rmont._spa     GATGAAAATCTAGTCTTTAGAAAAGACGGTAAATTATTGTTTGTAAAACCTCAAAGCGGT
ELB._space     -----
RproW._spa     G-----TCTAGTCTTTAGAAAA-----TAAAACTACAGCTTG

Rsib._spac      TTGATAAATAGAGAAGGAGAAATGACAGCAGTAAATCATATTTTACAAAAAGGCAGATTT
Rrick._spa     TTGATAAATAGAGAAGGAGAAATGACAGCAGTAAATCATATTTTACAAAAAGGCAGATTT
Rpark._spa     TTGATAAATAGAGAAGGAGAAATGACAGCAGTAAATCATATTTTACAAAAAGGCAGATTT
Ramb._spac     TTGATAAATAGAGAAGGAGAAATGACAGCAGTAAATCATGTTTTACAAAAAGGCAGATTT
Rmont._spa     TTGATAAATAGAGAAGGAGAAATGACAGCAGTAAATCATGTTTTACAAAAAGGCAGATTT
ELB._space     -----
RproW._spa     TTAATAAAAAGAAAATGTATACTGAAAACAGTAAATCATATTTTACAAAAGAAGTAGATTT

Rsib._spac      TTGTTTGAAGATTTAGGCGGTAAGGTAGATACAAAATTAGGGACCACAAATCTAAACAAA
Rrick._spa     TTGTTTGAAGATTTAGGCGGTAAGGTAGAT---AAATTAGGGACTACAAATCTAAACAAA
Rpark._spa     TTGTTTGAAGATTTAGGCGGTAAGGTAGATACAAAATTAGGGACCACAAATCTAAACAAA
Ramb._spac     TTGTTTGAAGATTTAGGCGGTAAGGTAGATACAAAATTAGGGACCACAAATCTAAACAAA
Rmont._spa     TTGTTTGAAGATTTAGGCGGTAAGGTAGATACAAACATTAGGGACCACAAATCTAAACAAA
ELB._space     -----TTAGGAACTGAAAATCTAAACAAA
RproW._spa     T-----GAAGATTTAAGTGA-----ATATAAAATCGTAGACTGTAA-TCTAAGCACA
                                *      **      **      *      *      *      *      *

Rsib._spac      AAGCCGACAAAGCCTACTTTAAAAAGAGGTTAAAATAACTTTAAGCCTTCTCATAAGAAA
Rrick._spa     AAGCCGACAAAGCCTACTTTAAAAAGAGGTTAAAATAACTTTAAGCCTTCTCATAAGAAA
Rpark._spa     AAGCCGACAAAGCCTACTTTAAAAAGAGGTTAAAATAACTTTAAGCCTTCTCATAAGAAA
Ramb._spac     AAGTCGACAAAGCCTACTTTAAAAAGAGGTTAAAATAACTTTAAGCCTTCTCATAAGAAA
Rmont._spa     AAGCCGACAAAGCCTACTTTAAAAAGAGGTTAAAATAACTTTAAGCCTTCTCATAAGAAA
ELB._space     AAGCCGACAAAGCCTACTTTAAAAAGAGGTTAAAATAACTTTAAGCCTTCTCATAAGAAA
RproW._spa     TA-----TTTTCTTTAAAAAGAGGTTAAAATGAGTTTAAAGATACTACATCAAAA
                                *      *      *      *      *      *      *      *      *      *      *      *

Rsib._spac      TAAAAATAAAAATTATTTGAAATTAGTGTTTGACAAGCTTTAAAAATATAGATATAAAAC-
Rrick._spa     TAAAAATAAAAATTATTTGAAATTAGTGTTTGACAAGCTTTAAAAATATAGATATAAAAT-
Rpark._spa     TAAAAATAAAAATTATTTGAAATTAGTGTTTGACAAGCTTTAAAAATATAGATATAAAAC-
Ramb._spac     TAAAAATAAAAATTATTTGAAATTAGTGTTTGACAAGCTTTAAAAATATAGATATAAAAT-
Rmont._spa     TAAAAATAAAAATTATTTGAAATTAGTGTTTGACAAGCTTTAAAAATATAGATATAAAAGT-
ELB._space     TAAAAATAAAAATTATTTGAAATTAGTGTTTGACAAGCTTTAAAAATATAGATATAAAAT-
RproW._spa     TAAAAATAAAAATTATTTGAAATTAGTGTTTGACAAGCTTTAAAAATATAGATATAAAGATT
*****

Rsib._spac      CATCACACAAAGTTTGAACGTA AAACTTTGTAGCTGTTTAAAAAGTTGATAAAGCTAAA
Rrick._spa     CATCACACAAAGTTTGAACGTA AAACTTTGTAGCTGTTTAAAAAGTTGGTAAAGCTAAA
Rpark._spa     TATCACGCAAGTTTGAACGTA AAACTTTGTAGCTGTTTAAAAAGTTGATAAAGCTAAA
Ramb._spac     CATCACACAAAGTTTGAACGTA AAACTTTGTAGCTGTTTAAAAAGTTGATAAAGCTAAA
Rmont._spa     CACCACACAAAGTTTGAACGTA AAACTTTGTAGCTGTTTAAAAAGTTGATAAAGCTAAA
ELB._space     CATCACGCAAGTTTGAACGTA AAACTTTGTAGCTGTTTAAAAAGTTGATAAAGCTAAA
RproW._spa     CATCACACAAAGTTTAAAAATTTAAACTTTGTAGCTGTTTAAAAAGTTGATAAAGCTAAA
                                *      *      *      *      *      *      *      *      *      *

Rsib._spac      AACTAAATACACCGCGATAATAAAAAATTTATATCGTGAGTAGATGCTGAAAATAAGAAAA
Rrick._spa     AACTAAATACACCGCGATAATAAAAAATTTATATCGTGAGTAGATGCTGAAAATAAGAAAA
Rpark._spa     AACTAAATACACCGCGATAATAAAAAATTTATATCGTGAGTAGATGCTGAAAATAAGAAAA
Ramb._spac     A-CTAAATACACCGCGATAATAAAAAATTTATATCGTGAGTAGATGCTGAAAAA-----
Rmont._spa     AACTAAATACACCGCGATAATAAAAAATTTATATCGTGAGTAGATGCTGAAAA-----
ELB._space     AACTAAATACACCGCGATAATAAAAAATTTATATCGTGAGTAGATACTGAAAA-----
RproW._spa     A-CTAAATACACCGCGATAATAA-----TATATCGTGAGTAGATACTGAAAA-----
                                *      *      *      *      *      *      *      *      *      *

Rsib._spac      ATAATTGTGGAATTAATTCTGTATGCATTGTGTAATCTCAAGTTAAAAGC
Rrick._spa     ATAATTGTGGAATTAATTCTGTATGCATTGTGTAATCTCAAGTTAAAAGC
Rpark._spa     ATAATTGTGGAATTAATTCTGTATGCATTGTGTAATCTCAAGTTAAAAGC
Ramb._spac     -TAATTGTGGAATTAATTCTGTATGCATTGTGTAATCTCAAGTTAAAAGC
Rmont._spa     -TAATTGTGGAATTAATTCTGTATGCATTGTGTAATCTCAAGTTAAAAGC
ELB._space     -TAATTGTGGAATTAATTCTGTATGCATTGTGTAATCTCAAGTTAAAAGC
RproW._spa     -TAATTGTGGAATTAATTCTGTATGCATTGTGTAATCTCAAGTTAAAAGC
*****

```

Figure 3: Multiple alignment of the nucleotide sequences of the spacer regions between *fnt* and *rrl* in seven Rickettsia species. Rsib = *R. sibirica*; Rrick = *R. rickettsii*; Rpark = *R. parkerii*; Ramb = *R. amblyomii*; ELB = *ELB agent*; Rmont = *R. montana*; RproW = *R. prowazekii*. Symbols beneath the aligned sequences indicate identical residues (\*).




Figure 4: Relationship among the *Rickettsia* inferred from the nucleotide sequence of the spacer region between *fmt* and *rriI*. The phylogeny was estimated by the neighbor-joining method applied to pairwise sequence differences corrected for multiple replacements. The number at each node indicates the percentage of bootstraps in which the taxa to the right were clustered.

most closely related spotted fever group rickettsia, *R. sibirica*, *R. rickettsii* and *R. parkerii* were observed to have virtually identical G+C content values within the spacer regions (from 28.1% to 28.7%).

The alignments of the spacer sequences are presented in Fig. 3. Here it can be seen that the reduced size of the PCR product obtained with genomic DNA from the ELB agent can be entirely explained by a contiguous deletion of almost 500 base pairs within the spacer region. The deletion extends from the region immediately next to the *fmt* termination codon to 500 base pairs into the spacer region. The 303 base pairs of the remaining spacer region show strong homology to the other *Rickettsia* sequences (Table 3). The corrected nucleotide distance values among the different species of the spotted fever group rickettsia are very small, less than 5%, in accordance with the low  $K_S$  values obtained within this group of species (<5%). An average distance value of 23.9% is observed between *R. prowazekii* and the five representatives of the spotted fever group rickettsia (*R. parkerii*, *R. sibirica*, *R. montana* and *R. amblyomii*). This value is similar to the average substitution frequency at synonymous sites for the same pairs of species (25.6%).

Phylogenetic trees based on these distance values are presented in Fig. 4. Neighbour joining as well as parsimony analysis produce three main clusters; *R. prowazekii* and the ELB agent, *R. parkerii* and *R. sibirica*, *R. montana* and *R. amblyomii*. The low bootstrap values associated with *R. rickettsii* suggests that the branching pattern of this species can not be determined with certainty in this analysis.

#### **4. Concluding remarks**

It has been shown previously that the *fmt* gene encoding methionyl-tRNA<sup>Met</sup> formyltransferase is situated upstream of the 23S-5S rRNA cluster in *R. prowazekii* (Andersson et al., 1995). Here we have shown that the same organization is found in *R. sibirica*, *R. rickettsii*, *R. parkerii*, *R. amblyomii*, *R. montana* and ELB agent.

Analyses of the alignment of the *fmt* gene and of the spacer between *fmt* and 23S rRNA genes show a large difference between the species from the spotted fever group (SFG) (*R. sibirica*, *R. rickettsii*, *R. parkerii*,

*R. amblyommi* and *R. montana*) and the species from the typhus group (TG) (*R. prowazekii*). Small difference among the spotted fever group has been found.

The ELB agent which was classified into the typhus group because of its disease similar to murine typhus, seems to be closer to the spotted fever group. Anyway the phylogenetic tree deduced from distance values produce a cluster with ELB agent together with *R. prowazekii* (TG). Because of the small number of positions being compared (the length of the ELB agent is 303bp), it is difficult to evaluate the significance of the tree obtained. Analyses of longer spacers or genes may help to deduce a phylogenetic tree of the *Rickettsia* genus. Studies of the other species from the *Rickettsia* genus may also be helpful specially for *R. typhi* which is from the typhus group and for *R. akari* and *R. australis* which are close to ELB agent (the three bacteria are often called AAE group).

Since the rearrangement of the rRNA operon is conserved in the seven species, it is believed that the reorganization of the 16S-23S-5S gene cluster has occurred before the diversification of the genus *Rickettsia*.

## General conclusion

Deletions and recombination events between duplicated genetic sequences lead to genomic shrinkage as well as to rearrangements of genomic neighborhoods (Krawiec and Riley, 1990). Genomic rearrangements are frequent events in bacteria, but free living bacterial populations are quite resistant to the inroads of these variations. In contrast, mitochondrial populations have quite small and highly variable genome (Gray, 1992). We expect the intracellular lifestyle to provide an evolutionary setting within which the genomes of endoparasitic bacteria would tend to have characteristics that are intermediate between those of free living bacteria and organelles.

Deletions and rearrangements of the ribosomal RNA genes in the *Rickettsia* genus were to be expected because genomic rearrangements are often associated with repeated sequences, which serve as sites of intrachromosomal homologous recombination.

It has been shown in this work that the highly preserved arrangement of the *fms* gene which encodes peptide deformylase (PDF) and the *fmt* gene encoding methionyl-tRNA<sup>Met</sup> formyltransferase (MTF) has been found immediately upstream of the rearranged 23S-5S rRNA gene cluster.

Furthermore, we show that the unusual organisation of the rRNA genes are found in six other species of the *Rickettsia* genus.

All these observations suggest that the rearrangement of rRNA genes in *Rickettsia* preceded the divergence of the typhus group and the spotted fever group.

## Materials and methods

### 1. Preparation of Plasmids

(QIAGEN Plasmid Kit) (KEBO, Stockholm, Sweden)

#### 1.1. Introduction

The Clone 132 sequenced was one of thousand clones selected randomly for universal and reverse sequencing from lambda Zap library.

The lambda Zap II phage is designed with pBluescript plasmid DNA, flanking the foreign insert in the phage linear genome. *E. coli* is infected by lambda Zap II phage and by a helper phage. The helper phage recognises the flanking region of the insert in the lambda DNA. The single strand phagemid circularizes and contains the clone 132. Then the phagemid infect another *E. coli* cells (XLI) and the cells containing the double strand plasmid will be ampiciline resistant. The plasmid is stored at -80°C in LB with 30% of glycerol.

The clone 132 which contains the *fms* gene was produced and purified by the folowing protocol.

#### 1.2. Components

Resuspension Buffer : 100 ug/ml RNase A  
50 mM Tris-HCl  
10 mM EDTA (pH 8)

Lysis Buffer: 200 mM NaOH



1% SDS

Neutralisation Buffer : 3.0 M KAc (pH 5.5)

Equilibration Buffer : 750 mM NaCl  
50 mM MOPS  
15% ethanol (pH 7.0)  
0.15% Triton X-100

Wash Buffer : 1.0 M NaCl  
50 mM MOPS  
15% ethanol (pH 7.0)

Elution Buffer: 1.25 M NaCl  
50 mM Tris-HCl  
15% ethanol (pH 8.5)

TE: 10 mM Tris-HCl  
1mM EDTA (pH 8.0)

5x LB Broth (per liter) : 500 g NaCl  
(Stock solution) 50 g Bacto-Tryptone  
25 g Yeast Extract

1 x LB (working solution) : 200 ml 5x LB  
800 ml dd H<sub>2</sub>O  
10 ml 20% glucose  
1 ml Ampiciline (100ug/ml)

LB/Ampiciline plate: 14g agar  
1 liter 1x LB

### 1.3. Protocol

Streak transformes cell within a LB/Ampiciline plate and incubate overnight at 37°C. Colonies who appear on the plate, contain the double stranded plasmid with the cloned DNA insert.

Take a single isolated colony and started a culture in 20 ml of working solution. When the mixture becomes cloudy, start an overnight culture with 2ml of the mixture and 200ml of 1x LB.

Centrifuge at 4,000 RPM during 10 minutes. Add 10 ml of Resuspension Buffer to the bacterial pellet and 10 ml of Lysis Buffer. Mix gently.

After incubation at room temperature for 5 minutes, the lysis should occur, and the solution appears viscous. Add 10 ml of Equilibration Buffer. Place the solution on ice for 20 minutes. Mix the solution and centrifuge it at 4°C for 30 minutes at 10,000 RPM.

The supernatant should be clear after centrifugation. If not a second centrifugation is performed.

Add 10 ml of Equilibration Buffer to a Qiagen-tip 500. Let the liquid going out without applying pressure. Apply the clear supernatant onto the column, let the liquid enter the resin by gravity flow.

Wash two times with 30 ml of Wash Buffer .

Elute the DNA by adding 15 ml of Elution Buffer and collect it in 30 ml glass tube (corex).

Add 0.7 volumes of isopropanol for precipitation of DNA. Centrifuge directly at 10,000 RPM for 30 minutes at 4°C. Remove carefully the supernatant.

Wash carefully the DNA pellet with 15 ml of cold 70% ethanol. Discard the ethanol and let dry for 5 minutes and dissolve the pellet in a small volume of TE.

The DNA concentration was measured by running it and a aliquot of lambda DNA of known concentration on a 1% agarose gel.

## 2. Polymerase Chain Reaction (PCR)

### 2.1. Introduction

The goal of this technique is to amplify short segments between two defined primers.

First the DNA is denatured. Two short primers (around 20 bases) are annealed with the two single-stranded DNA, at a complementary site.

The enzyme Taq DNA polymerase is used to synthesize a complementary single-strand from the 3' end of each primer.

The cycle of 3 steps is repeated 20 to 30 times to have an increasing of the number of copies.

PCR is required to amplify the spacer between *fmt* gene and 23S gene for *R. prowazakii*, *R. rickettsii*, *R. sibirica*, *R. montana*, *R. parkii*, *ELB-agent* and *R. ambilyommi*.

We use two primers, one from the *fmt* gene (5'-CTAAAGCAGAAGGAAAAATT-3') (position 575) and one primer from the beginning of the the 23S gene (5'-GCTTCTAGTGCCAAGGCATC-3').

These segments will be sequenced later.

A PCR of *R. prowazaki* genomic DNA was also performed in order to amplify the locus containing both *fms* and *fmt* genes.

The amplification by PCR is using two primers : one from the *fms* gene (5'-ATTAGCAAGATTTACTTTCTT-3') and the second one from the *fmt* gene (5'-CACTGCCACAAGCGATTTCTAGTTT-3').

### 2.2. Components

Taq DNA Polymerase in Storage Buffer containing :

- 50% glycerol
- 50mM Tris-HCl (pH8.0)
- 100mM NaCl
- 0.1mM EDTA
- 1mM DTT
- 1.0% Triton X-100

10X Thermophilic Buffer (Magnesium Free) :  
500mM KCl  
100mM Tris-HCl (pH 9.0)  
1%Triton X-100

Magnesium Chloride 25mM

dNTP 2.5 mM : 5  $\mu$ l of 100 mM of dATP  
5  $\mu$ l of 100 mM of dCTP  
5  $\mu$ l of 100 mM of dGTP  
5  $\mu$ l of 100 mM of dTTP  
180  $\mu$ l H<sub>2</sub>O

### 2.3. Protocol

The reaction buffer was: 135  $\mu$ l of sterilised water  
25  $\mu$ l of 10 x Thermophilic Buffer  
15  $\mu$ l of MgCl<sub>2</sub>  
25  $\mu$ l of dNTP (2.5 mM)  
12.5  $\mu$ l of each primers (25 pmoles).

The total reaction volume was 225  $\mu$ l.

Two aliquots of 90  $\mu$ l of this reaction are prepared in two tubes. Add in one tube 5  $\mu$ l of water as a control and add 5  $\mu$ l of genomic DNA in the second tube.

Add one drop of mineral oil.

Hot Start at 94° C for two minutes, then 80° C for three minutes. Add 5  $\mu$ l of diluted Taq DNA polymerase (2.5 units).

The cycle started at 94° C for one minute, followed by one minute at 53° C and finished at 72° C for three minutes for 30 cycles.

The program finished by ten minutes at 72° C.

## 2.4. Control

A control of the PCR product on a 1% agarose gel was performed to observe eventual contaminations and to estimate the yield of the PCR.

### Components :

5 x TBE (stock solution):      54 g Tris base  
   27.7 g boric acid  
   20 ml 0.5 M EDTA (pH 8)

0.5 M EDTA (pH 8) :    35 g of disodium ethylen diamine tetraacetate  
   160 ml H<sub>2</sub>O Adjust to pH to 8  
   Adjust the volume to 200 ml

The components of the agarose gel 1% :

0.5 g of agarose  
50 ml 0.5 x TBE  
5 µl EtBr (ethidiumbromide)

10 µl of each reaction were loaded together with 2 µl of dye.  
The samples were electrophoresed at 100 volts for thirty minutes.

## **3. Direct Purification from PCR Products**

(Wizard PCP Preps DNA purification System) (Promega)

### 3.1. Introduction

Before sequencing, it is important to purify the PCR products to remove the mineral oil and the excess of components. It is necessary to remove all the oil before charging the minicolumn. After washing the minicolumn, the DNA was collected in TE.

### 3.2. Components

Direct Purification Buffer: 50 mM KCl  
10 mM Tris-HCl (pH 8.8)  
1.5 mM MgCl<sub>2</sub>

Magic PCR Preps Resin (Promega)

Magic Minicolumn (Promega)

TE buffer : 10 mM Tris-HCl (pH7.5)  
1 mM EDTA

Chloroform 100%

Isopropanol 80%

### 3.3. Protocol

Add 100 µl of water and 50 µl of chloroform to each completed PCR reaction. Mix carefully and spin one minute at 12,000 RPM.

Remove 150 µl from the upper layer of the sample.

Add 100 µl Direct Purification Buffer, vortex.

Add 1 ml of magic PCR Preps resin and vortex briefly 3 times over a one minute period.

Prepare one Magic Minicolumn for each PCR product and connect it to a 2 ml syringe.

Apply the DNA mix into the barrel and push slowly the plunger, after that, place 2 ml of isopropanol 80% and push gently the liquid out to wash the minicolumn.

Remove the syringe and place the minicolumn into a 1.5 ml microcentrifuge tube. Spin the minicolumn for twenty seconds at 12,000 RPM to remove all the liquid out.

Let the minicolumn for five minutes at room temperature to dry the resin.

Add 50  $\mu$ l of warm TE buffer (or water) to the Magic Minicolumn and wait ten minutes. Centrifuge the minicolumn for twenty seconds at 12,000 RPM to elute the DNA fragment.

## **4. DNA sequencing**

### **4.1. Introduction**

The Automated Laser Fluorescent ALF DNA sequencer from Pharmacia LKB has been conceived in accordance with Sanger's method (chain termination technique).

Sequencing requires hybridization of the single stranded DNA to be sequenced with a short complementary primer.

The DNA polymerase starts at the 3' hydroxyl end of the primer and synthesises a complementary copy of the single strand DNA.

DNA sequencing by enzymatic synthesis employs four different reaction mixtures, each contains all four deoxyribonucleoside triphosphates (dNTP) and one of the four dideoxyribonucleoside triphosphate (ddNTP). The ddNTP has no 3' hydroxyl group and so, when incorporated, causes the chain termination.

When the sequencing reaction is fractionated by electrophoresis on gel, the pattern of bands shows the distribution of synthesised DNA.

The use of fluorescent primers or fluore dATP are required for the analyse of the fluorescent curves by the A.L.F. DNA sequencer which will translate them into a DNA sequence, exported as a computer file.

## 4.2. Auto Read Sequencing kit (Pharmacia)

Auto Read Sequencing kit is designed for use with Automated Laser Fluorescent ALF DNA sequencer from Pharmacia LKB.

This method is to sequence a clone by using fluorescent M13 Universal Primer and fluorescent M13 Reverse Primer or primers no fluorescent in combination with Fluore-dATP Labelling mix.

### *4.2.1. Sequencing with fluorescent primer*

The reversal and universal fluorescent primers start elongation at specific priming sites allowing sequencing of both strands.

#### 4.2.1.1. Components

A mix : 5 uM ddATP, 1 mM dATP, 1 mM dCTP, 1 mM dTTP, 1 mM c7dGTP, 50 mM NaCl and 40 mM Tris-HCl (pH 7.6)

C mix : 5 uM ddCTP, 1 mM dATP, 1 mM dCTP, 1 mM dTTP, 1 mM c7dGTP, 50 mM NaCl and 40 mM Tris-HCl (pH 7.6)

G mix : 5 uM ddGTP, 1 mM dATP, 1 mM dCTP, 1 mM dTTP, 1 mM c7dGTP, 50 mM NaCl and 40 mM Tris-HCl (pH 7.6)

T mix : 5 uM ddTTP, 1 mM dATP, 1 mM dCTP, 1 mM dTTP, 1 mM c7dGTP, 50 mM NaCl and 40 mM Tris-HCl (pH 7.6)

T7 DNA polymerase : 8 units/ul in 25 mM Tris-HCl (pH 7.5), 0.25 M NaCl, 5 mM DTT and 50% glycerol

M13 Universal Primer : 5'-fluorescein-d(CGACGTTGTAAAACGACGGCCAGT)-3' in aqueous solution, 1.5 uM (2.1 pmol/ul; 0.55 A<sub>260</sub> unit/ml)

M13 Reverse Primer : 5'-fluorescein-d(CAGGAAACAGCTATGAC)-3' in aqueous solution, 2.1 uM (2.1 pmol/ul; 0.42 A<sub>260</sub> unit/ml)

Annealing Buffer : 1 M Tris-HCl (pH 7.6) and 100 mM MgCl<sub>2</sub>



Extension Buffer : 304 mM citric acid, 324 mM DTT and 40 mM MnCl<sub>2</sub> (pH 7.5)

Enzyme Dilution Buffer : 20 mM Tris-HCl (pH 7.5), 5 mM DTT, 100 ug/ml BSA and 5% glycerol

DMSO : dimethyl sulfoxide 100%

Stop solution : 100% deionized formamide and Dextran Blue 2000 (5 mg/ml)

1M NaOH for denaturation

#### 4.2.1.2. Protocol

##### a. Denaturation

Adjust the concentration of the template so that 20  $\mu$ l contains 5 to 10 ug of DNA.

Add 5  $\mu$ l of 1M NaOH. Incubate at 65° C for five minutes. Spindown the tube briefly.

The mixture is loaded on a microspin column and centrifuge at 12,000 RPM during two minutes.

Two aliquots (F and R) of 10  $\mu$ l of the solution containing denatured template DNA are prepared in a 1.5 ml microcentrifuge tube.

##### b. Annealing of primers (Forward and Reverse)

Add 2  $\mu$ l of annealing buffer in both tube (F and R).

Add 2  $\mu$ l of fluorescent M13 Universal Primer in one tube (F) and 2  $\mu$ l of fluorescent M13 Reverse Primer in the second tube (R).

The total volume in each tube is 14  $\mu$ l. Vortex and centrifuge gently. Preheat the annealing reaction mixtures at 65° C for five minutes. and then incubate immediately in a water bath at 37° C for ten minutes. Remove the tubes and leave them at room temperature for five minutes. Spindown the content of the tubes briefly.

c. Termination :

Preliminaries :

- For sequencing with a double stranded templates the T7 DNA polymerase is diluted in Enzyme dilution Buffer to a concentration of 6 to 8 units per 2  $\mu$ l.
- Pipette 2.5  $\mu$ l of the A mix, C mix, G mix, T mix respectively in four wells of a microsample plate. Keep it on ice.

Termination reaction :

In a 1.5 ml microcentrifuge tube add for each annealing reaction :

- 1  $\mu$ l of extension buffer
- 3  $\mu$ l of DMSO
- 2  $\mu$ l of diluted T7 DNA polymerase.

Remove 6  $\mu$ l of this mixture and add it in each annealing reactions (F and R), the total volume in each tube being now 20  $\mu$ l.

Pipette 4.5  $\mu$ l of this mixture into each of the four wells of the microsample plate. Mix carefully and incubate for five minutes at 37°C.

Add 5  $\mu$ l of Stop Solution to each reaction and mix gently.  
Heat the reactions at 95° C for two minutes and then keep them on ice.

Load 6 to 8  $\mu$ l into the appropriate wells of a sequencing gel.

#### 4.2.2. Sequencing with no fluorescent primer

This technique uses Fluore-dATP Labelling Mix in combination with Auto Read Sequencing kit and an Automated Laser Fluorescent ALF DNA sequencer from Pharmacia.

This fluorescein-labelled dATP is incorporated during the labelling reaction.

##### 4.2.2.1. Protocol

###### a. Denaturation and Annealing reactions

The denaturation reaction and annealing reaction are the same than B.1.2.a. and B.1.2.b. with a concentration of the primer between 10 to 100 pmoles.

###### b. Labelling reaction

To the tube containing the annealed template and the primer add 2  $\mu$ l of fluor-dATP and 2  $\mu$ l of diluted T7 DNA polymerase (6-8 units/2  $\mu$ l). Mix gently.

Incubate the reaction mixtures at 37°C for ten minutes .

###### c. Termination

After incubation, add 1  $\mu$ l of Extension Buffer and 3.5  $\mu$ l of DMSO to each labelling reaction. Mix gently

Pipette 3  $\mu$ l of each of the A mix, C mix, G mix, and T mix into four wells of a microsample plate. Add 5.4  $\mu$ l of the reaction mixture to each of the appropriate wells. Mix carefully.

Incubate for five minutes at 37° C. Then stop the reactions by adding 6  $\mu$ l of Stop Solution.

Load 6 to 8  $\mu$ l into the appropriate wells of the sequencing gel.

### 4.3. Thermo Sequenase fluorescent labelled primer cycle sequencing kit (Amersham)

The Thermo Sequenase is a thermostable DNA polymerase and then many cycles can be performed without the need to add new enzyme. Uniform sequence band patterns are generated and are easier to read.

#### *4.3.1. Components*

A reagent            Tris-HCl (pH9.5), magnesium chloride, Tween 20, Nonidet P-40, 2-mercaptoethanol, dATP, dCTP, dGTP, dTTP, ddATP, thermostable pyrophosphatase and Thermo Sequenase DNA polymerase.

C reagent            Same composition with ddCTP

G reagent            Same composition with ddGTP

T reagent            Same composition with ddTTP

Formamide  
loading dye        Formamide, EDTA and methyl violet

Fluorescent primers (5pmol/ul)

Template DNA (0.5-5ug)

#### *4.3.2. Protocol*

For each template DNA, prepare four microcentrifuge tubes containing respectively 2  $\mu$ l of the A reagent, C reagent, G reagent and T reagent. Store on ice until required.

In a 1.5ml microtube add 5  $\mu$ l of template DNA, 19  $\mu$ l of water and 2  $\mu$ l of the fluorescent primers. Mix carefully.

Add 6  $\mu$ l of this mixture into each tubes containing the dNTP. Vortex. Add one drop of mineral oil and centrifuge the tubes at high speed for 20 seconds.

Start the PCR program by a pre-denaturation at 94°C for 2 minutes followed by 25 cycles : denaturation for 30 seconds at 95°C, annealing for 30 secondes at 60°C and elongation for 1 minute at 72°C.

When the program is completed, add 3  $\mu$ l of Formamide loading dye to stop the elongation. Mix the reactions and centrifuge the tubes at high speed for 20 seconds.

Remove the 11  $\mu$ l or each reaction from under the mineral oil, and put them on the wells of a microsample plate.

Warm the plate at 95°C for two minutes and place it directly on ice. Load 6 to 8  $\mu$ l Of the reactions into the appropriate wells of the gel.

## References

- Ahern, H., 1994.** *Does a molecular clock tick for bacteria?* ASM news 60: 600-603.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W. and Lipman, D. J. 1990.** *Basic local alignment search tool.* J. Mol. Biol. 215:403-410.
- Amman, R. I., Ludwig, W., and Schleifer, K-H. 1995.** *Phylogenetic identification and in situ detection of individual microbial cells without cultivation.* Microbial. Rev. 59: 143-169.
- Andersson, S. G. E. and Kurland C. G. 1995.** *Genomic evolution drives the evolution of the translational system.* Biochem. Cell Biol. In press.
- Andersson, S. G. E. and Sharp, P. M. 1995.** *Codon usage and base composition in Rickettsia prowazekii.* J. Mol. Evol. Submitted.
- Andersson, S. G. E., Zomorodipour, A., Winkler, H. H. and Kurland, C. G. 1995.** *Unusual organization of the rRNA genes in Rickettsia prowazekii.* J. Bacteriol. 177: 4171-4175.
- Azad, A. F., Sacci, J. B., Jr., Nelson, W. M., Dasch, G. A., Schmidtman, E. T., and Carl, M. 1992.** *Genetic characterization and transovarial transmission of a typhus-like rickettsia found in cat fleas.* Proc. Natl. Acad. Sci. 89: 43-46.
- Baylis, H. A., and Bibb, M. J., 1988.** *Organisation of the ribosomal RNA genes in Streptomyces coelicolor A3(2).* Mol. Gen. Genet. 211: 191-196.

**Chen, X, and Frinch, L. R., 1989.** *Novel arrangement of rRNA genes in Mycoplasma gallisepticum: separation of the 16S gene of one set from the 23S and 5S genes.* J. Bacteriol. 171: 2876-2878.

**De Rijk, P., Van de Peer, Y., Van den Broeck, I., De Wachter, R., 1995.** *Evolution according to large ribosomal subunit RNA.* J. Mol. Evol. 41: 366-375.

**Drancourt, M., and Raoult, D., 1994.** *Taxonomic position of the Rickettsiae: current knowledge.* FEMS Microbiol. Rev. 13: 13-24.

**Eremeeva, M. E., Roux, V. and Raoult, D. 1993.** *Determination of genome size and restriction pattern polymorphism of Rickettsia prowazekii and Rickettsia typhi by pulsed field gel electrophoresis.* FEMS Microbiol. Lett. 112: 105-112.

**Fleischman, R. D., Adms, M. D., White, O., Clayton, R. A., Kirkness, E. F., Kerlavage, A. R., Tomb, J-F., Dougherty, B. A., Merrick, J. M., McKenney, K., Sutton, G., FitzHugh, W., Fields, C., Gocayne, J. D., Scott, J., Shirley, R., Liu, L-I, Glodek, A., Kelley, J. M., Weidman, J. F., Phillips, C. A., Spriggs, T., Hedblom, E., Cotton, M. D., Utterback, T. R., Hanna, M. C., Nguyen, D. T., Saudek, D. M., Brandon, R. C., Fine, L. D., Fritchman, J. L., Fuhrmann, J. L., Geoghagen, N. S. M., Gnehm, C. L., McDonald, L. A., Small, K. V., Fraser, C. M., Smith, H. O., Venter, C. 1995.** *Whole-genome random sequencing and assembly of Haemophilus influenzae.* Rd.Science 269: 496-512.

**Fraser, C. M., Gocayne, J. D., White, O., Adms, M. D., Clayton, R. A., Fleischman, R. D., Bult, C. J., Kerlavage, A. R., Sutton, G., Kelley, J. M., Fritchman, J. L., Weidman, J. F., Small, K. V., Sandusky, M., Fuhrmann, J., Nguyen, D., Utterback, T. R., Saudek, D. M., Phillips, C. A., Merrick, J. M., Tomb, J-F., Dougherty, B. A., Bott, K. F., Hu, P-C., Lucier, T. S., Peterson, S. N., Smith, H. O., Hutchison III, C. A., Venter, C. 1995.** *The minimal gene complement of Mycoplasma genitalium.* Science 270: 390-403.

**Goffeau, A. 1995.** *Life with 482 genes.* Science 270: 445-446.

**Gray, M. W., 1993.** *Origin and evolution of organelle genomes.* Current Opinion in genetics and development 3: 1-7.

**Guillon J-M., Mechulam, Y, Schmitter, J-M., Blanquet, S., and Fayat, G. 1992.** *Disruption of the gene for Met-tRNA<sup>fMet</sup> Formyltransferase severely impairs growth of Escherichia coli.* J. of Bacteriol. 174: 4294-4301.

**Higgins, D. G., Bleasby, A. J. and Fuchs, R. 1992.** *ClustalW: improved software for multiple sequence alignment.* CABIOS 8: 189-191.

**Krawiec, S. and Riley, M. 1990.** *Organisation of the bacterial chromosome.* Microbiol. Rev. 54: 502-539.

**Kurland C. G., 1993.** *Evolution of mitochondrial genomes and the genetic code.* BioEssays 14, 10: 709-714.

**Li, W-H., 1993.** *Unbiased estimation of the rates of synonymous and nonsynonymous substitution.* J. Mol. Evol. 36: 96-99.

**Meinzel, T. and Blanquet, S. 1994.** *Characterisation of the Thermus thermophilus locus encoding peptide deformylase and Methionyl-tRNA<sup>fMet</sup> Formyltransferase.* J. Bacteriol. 176: 7387-7390.

**Muto, A. and Osawa, S. 1987.** *The guanine and cytosine content of genomic DNA and bacterial evolution.* Proc. Natl. Acad. Sci. 84: 166-169.

**Ochman, H., and Wilson, C. 1987.** *Evolution in bacteria: evidence for a universal substitution rate in cellular genomes.* J. Mol. Evol. 26: 74-86.

**Ohashi, N., Fukuhara, M., Shimada, M., Tamura, A., 1995.** *Phylogenetic position of Rickettsia tsutsugamushi and the relationship among its antigenic variants by analyses of the 16S rRNA gene sequences.* FEMS Micro. Letters 125: 299-304.

**Olson, G. J., Woese, C. R., and Overbeek, R. 1994.** *The winds of (Evolutionary) change: breathing new life into microbiology.* J. Bacteriol. 176: 1-6.



**Sanger, F., Nicklen, S., and Coulson A. R., 1977.** *DNA sequencing with chain-terminating inhibitors.* Proc. Natl. Acad. Sci. 74: 5463-5467

**Stothard, D. R., Clark, J. B. and Fuerst, P. A., 1994.** *Ancestral divergence of Rickettsia belii from the spotted fever and typhus group of Rickettsia and antiquity of the genus Rickettsia.* Inter. J. of Syst. Bacteriol. 44, 4/ 798-804.

**Stothard, D. R., 1995.** *The evolution history of the genus Rickettsia as inferred from 16S and 23S ribosomal RNA genes and the 17 Kilodalton cell surface antigen gene.* Dissertation, Ohio State University, Columbus, OH.

**Syvänen, A-C., Amiri, H., Jamal, A., Andersson, S. G. E., and Kurland, C. G., 1995.** *Disposition of elongation factor genes reflects the genomic evolution of Rickettsia prowazekii.* PNAS submitted.

**Tamura, A., 1988.** *Invasion and intracellular growth of Rickettsia tsutsugamushi.* Microbiol. Sci. 5: 228-232.

**Taschke, C., Klinkert, M-Q., Wolters, J., and Hermann, R., 1986.** *Organisation of the ribosomal RNA genes in Mycoplasma hyopneumoniae: the 5S rRNA gene is separated from the 16S and 23S rRNA genes.* Mol. Gen. Genet. 205: 428-433.

**Weisburg W. G., Dobson, M. E., Samuel, J. E., Dasch, G. A., Mallavia, L. P., Baca, O., Mandelco, L., Sechrest, J. E., Weis, E., Woese, C. R., 1989.** *Phylogenetic diversity of the Rickettsiae.* J. Bacteriol. 171: 4202-4206.

**Weiss, E. and Moulder, J. M., 1984.** In Bergeys's manual of Systematic Bacteriology, N.R. Kreig and J.G. Holt, eds. (Williams and Wilkins, Baltimore) p.687.

**Williams, S. G., Sacci, J. B., Jr., Schriefer, M. E., Andersen, E. M., Fujioka, K. K., Sorvillo, F. J., Barr, A. R., and Azad, A. F., 1992.** *Typhus and typhus-like rickettsiae associated with opossums and their fleas in Los Angeles County, California.* J. Clin. Microbiol. 30°: 1758-1762.

**Winkler H. H., 1995.** *Rickettsia prowazekii*, ribosomes and slow growth. Trends in Microbiol. 3: 196-198.

**Winkler H. H., 1990.** *Rickettsia* species (as organisms). Annu. Rev. Microbiol. 44: 131-153.

**Woese, C. R., 1994.** There must be a prokaryote somewhere: microbiology's search for itself. Microbiol. Rev. 58: 1-9.

**Wood, D. O. and Waite, R. T. 1994.** Sequence analysis of the *Rickettsia prowazekii* *gyrA* gene. Gene 151: 191-196.