

University of Groningen

## Adalimumab (Humira®) Induced Recurrent Peritonsillar Abscess in A Patient Received Three Different Anti-TNF Therapies

Vural, Sevilay; Kuşdoğan, Mikail; Kaya, Hasan Burak; İkiz, Venhar; Albayrak, Levent

*Published in:*  
Indian Journal of Otolaryngology and Head and Neck Surgery

*DOI:*  
[10.1007/s12070-022-03203-0](https://doi.org/10.1007/s12070-022-03203-0)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2023

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Vural, S., Kuşdoğan, M., Kaya, H. B., İkiz, V., & Albayrak, L. (2023). Adalimumab (Humira®) Induced Recurrent Peritonsillar Abscess in A Patient Received Three Different Anti-TNF Therapies: A Case Report. *Indian Journal of Otolaryngology and Head and Neck Surgery*, 75, 1013–1015. Advance online publication. <https://doi.org/10.1007/s12070-022-03203-0>

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



# Adalimumab (Humira®) Induced Recurrent Peritonsillar Abscess in A Patient Received Three Different Anti-TNF Therapies: A Case Report

Sevilay Vural<sup>1,2</sup> · Mikail Kuşdoğan<sup>2</sup> · Hasan Burak Kaya<sup>2</sup> · Venhar İkiz<sup>2</sup> · Levent Albayrak<sup>2</sup>

Received: 2 July 2022 / Accepted: 23 September 2022 / Published online: 6 December 2022  
© The Author(s) 2022

## Abstract

Anti-tumor necrosis factor agents are widely used in treating ankylosing spondylitis, but they increase the risk of infection by suppressing the immune response. Therefore, physicians should be careful about recurrent infections in patients under anti-tumor necrosis factor agents.

**Keywords** Anti-TNF · Adalimumab · Etanercept · Infliximab · Peritonsillar abscess · Deep infection

## Abbreviations

AS	Ankylosing spondylitis
TNF- $\alpha$	Tumor necrosis factor alpha
PTA	Peritonsillar abscess
ED	Emergency Department

## Introduction

Ankylosing spondylitis (AS) is a chronic inflammatory disease mainly affecting the sacroiliac joints and spine. Tumor necrosis factor alpha (TNF- $\alpha$ ) plays an essential role in the pathogenesis of AS and many other inflammatory diseases, which can be used as a target point during the treatment. If

the disease activity continues despite the first-line therapies, anti-TNF drugs can be used for AS [1]. The most widely used anti-TNF drugs are infliximab, etanercept, and adalimumab. Peritonsillar abscess (PTA) is the most common deep infection of the head and neck, which generally occurs as a complication of acute tonsillitis. There are no reported numbers in the literature regarding PTA prevalence or recurrence rates in immunosuppressed patients.

## Case Report

A 31-year-old non-smoker male patient admitted to the emergency department (ED) with sore throat, dysphagia, and trismus for two days. His vitals were stable. The physical examination revealed a left-sided peritonsillar abscess without any tonsillitis findings (Fig. 1). His medical history was significant for AS, and receiving anti-TNF treatment for nine years. He had been prescribed etanercept during the first two years and infliximab during the second two-year period. His current treatment was 40 mg Adalimumab (Humira®) for the last five years. It was his third ED admission due to PTA. He had been treated for PTA twice in a different medical center and was hospitalized once during the current year. He denied any similar and/or recurrent medical condition under other anti-TNF therapies. The laboratory tests showed that WBC: 9020 /mm<sup>3</sup>, neutrophil ratio: 68.2%, and CRP: 9.2 mg/L. The patient was hospitalized by the otolaryngology clinic for further treatment. Abscess drainage was performed, and antibiotic therapy was initiated. No additional

✉ Sevilay Vural  
s.vural@umcg.nl  
Mikail Kuşdoğan  
dr\_mk7784@hotmail.com  
Hasan Burak Kaya  
hasanburakkaya@hotmail.com  
Venhar İkiz  
vnhr.ikiz@gmail.com  
Levent Albayrak  
drleventalbayrak@gmail.com

<sup>1</sup> Department of Emergency Medicine, University Medical Center Groningen, Hanzeplein 1, 9713 GZ Groningen, The Netherlands

<sup>2</sup> Department of Emergency Medicine, Yozgat Bozok University, Yozgat, Turkey



**Fig. 1** The examination revealing a left-sided peritonsillar abscess without any tonsillitis findings

pathology such as lymphoma or infectious mononucleosis was found during the differential diagnostics.

Two and a half months after the discharge, he admitted to our ED again with similar complaints. The vitals and laboratory results were similar to the previous admission. He was diagnosed with PTA for the fourth time in two years. The patient was discharged with an antibiotic prescription after the abscess drainage was performed in the ED and referred to his rheumatologist.

## Discussion

The most common bacterium isolated in PTA are *Streptococcus* and *Fusobacterium*, but many abscesses have a mixed profile [2, 3]. Primary treatment strategies are the initiation of appropriate antibiotic therapy and percutaneous evacuation of the abscess if needed. “Recurrence” term for PTA can be defined as a new episode of PTA in  $\geq 30$  days from the initial PTA [4]. The frequency of recurrent PTA in the normal population is 5.15–16%, and even much higher in younger populations aged 13–18 [5, 6]. As early recurrence can be linked to inadequate treatment, extra-peritonsillar spread of infection, or autoimmune deficiency, late recurrence is more likely related to autoimmune deficiency or prior history of tonsillitis or pharyngitis [6]. Unfortunately, the literature does not answer to the question, “How

many times can a PTA recur.” Our patient with AS receiving anti-TNF treatment had recurrent PTA attacks (four times), requiring surgical interventions and hospitalizations (two times) in the two-year time period. Although his anti-TNF regimen was the usual suspect, neither the patient himself stated his emergency visits or interventions to his rheumatologist during his follow-up, nor was there any executed rheumatology referral or consultation. Since none of PTAs accompanied by tonsillitis, tonsillectomy was not considered. In addition, the microorganisms causing PTAs were not determined because no culture study was performed. The large-scale studies on the relationship between anti-TNF and infection focused predominantly on tuberculosis activation and opportunistic infections [7, 8]. We could not be sure if PTA episodes were related to any opportunistic microorganism since no culture study was conducted.

As we assume that his repetitive medical condition was associated with the anti-TNF agent, we investigated the drug regimens he received. He had used etanercept and infliximab, and he has been treated with adalimumab for the last five years. All three are the most commonly used anti-TNF agents. He had four PTA during active use of adalimumab. However, the findings comparing the risk of infection in patients treated with etanercept, adalimumab, and infliximab showed no significant difference between the three drugs [9]. In a meta-analysis, significant increases were found in anti-TNF drug use by 20% for any infection, 40% for severe infection, and 250% for tuberculosis reactivation [7]. Nevertheless, we did not encounter any reports or studies including or specifically addressing PTA under anti-TNF treatment.

## Conclusion

Anti-TNF drugs should also be questioned in specific infection types and repetitive infection patterns that suggest the possibility of immunosuppression in the EDs.

**Acknowledgements** None.

**Funding** No funding was received for the study.

## Declarations

**Conflict of Interest** The authors declare that they have no conflict of interest.

**Funding Sources/Disclosures** None declared.

**Ethical Approval** Not applicable.

**Informed Consent** Informed consent was obtained from the patient of the case report.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

1. Haibel H, Sieper J (2010) Editorial review: how early should ankylosing spondylitis be treated with a tumor necrosis factor-blocker? *Curr Opin Rheumatol* 22:388–392. <https://doi.org/10.1097/BOR.0b013e32833aaf93>
2. Mazur E, Czerwińska E, Korona-Główniak I, Grochowalska A, Koziół-Montewka M (2015) Epidemiology, clinical history and microbiology of peritonsillar abscess. *Eur J Clin Microbiol Infect Dis* 34:549–554. <https://doi.org/10.1007/s10096-014-2260-2>
3. Tsai YW, Liu YH, Su HH (2018) Bacteriology of peritonsillar abscess: the changing trend and predisposing factors. *Braz J Otorhinolaryngol* 84:532–539. <https://doi.org/10.1016/j.bjorl.2017.06.007>
4. Wolf M, Even-Chen I, Kronenberg J (1994) Peritonsillar abscess: repeated needle aspiration versus incision and drainage. *Ann Otol Rhinol Laryngol* 103:554–557. <https://doi.org/10.1177/000348949410300709>
5. Wang YP, Wang MC, Lin HC, Chou P (2014) The impact of prior tonsillitis and treatment modality on the recurrence of peritonsillar abscess: a nationwide cohort study. *PLoS ONE* 9:e109887. <https://doi.org/10.1371/journal.pone.0109887>
6. Ali SA, Kovatch KJ, Smith J, Bellile EL, Hanks JE, Hoff PT (2019) Implication of *Fusobacterium necrophorum* in recurrence of peritonsillar abscess. *Laryngoscope* 129:1567–1571. <https://doi.org/10.1002/lary.27675>
7. Minozzi S, Bonovas S, Lytras T, Pecoraro V, González-Lorenzo M, Bastiampillai AJ et al (2016) Risk of infections using anti-TNF agents in rheumatoid arthritis, psoriatic arthritis, and ankylosing spondylitis: a systematic review and meta-analysis. *Expert Opin Drug Saf* 15:11–34. <https://doi.org/10.1080/14740338.2016.1240783>
8. Xu Z, Xu P, Fan W, Yang G, Wang J, Cheng Q et al (2017) Risk of infection in patients with spondyloarthritis and ankylosing spondylitis receiving antitumor necrosis factor therapy: A meta-analysis of randomized controlled trials. *Exp Ther Med* 14:3491–3500. <https://doi.org/10.3892/etm.2017.5003>
9. Galloway JB, Hyrich KL, Mercer LK, Dixon WG, Fu B, Ustianowski AP et al (2011) Anti-TNF therapy is associated with an increased risk of serious infections in patients with rheumatoid arthritis especially in the first 6 months of treatment: updated results from the British Society for Rheumatology Biologics Register with special emphasis on risks in the elderly. *Rheumatology (Oxford)* 50:124–131. <https://doi.org/10.1093/rheumatology/keq242>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.