

University of Groningen

Clinical significance of mitochondrial DNA content in acute promyelocytic leukaemia

Pereira-Martins, Diego A.; Coelho-Silva, Juan L.; Weinhauser, Isabel; Franca-Neto, Pedro L.; Silveira, Douglas R.; Ortiz, Cesar; Moreira-Aguiar, Amanda; Lima, Marinus M.; Koury, Luisa C.; de Melo, Raul A.

Published in:
British Journal of Haematology

DOI:
[10.1111/bjh.18510](https://doi.org/10.1111/bjh.18510)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2023

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Pereira-Martins, D. A., Coelho-Silva, J. L., Weinhauser, I., Franca-Neto, P. L., Silveira, D. R., Ortiz, C., Moreira-Aguiar, A., Lima, M. M., Koury, L. C., de Melo, R. A., Gloria, A. B., Fagundes, E. M., Lino, B. K., Pagnano, K., Bittencourt, R., Nunes, E., Traina, F., Figueiredo-Pontes, L., Keating, A., ... Lucena-Araujo, A. R. (2023). Clinical significance of mitochondrial DNA content in acute promyelocytic leukaemia. *British Journal of Haematology*, 200(2), 170-174. Advance online publication. <https://doi.org/10.1111/bjh.18510>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.












Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

SHORT REPORT

Clinical significance of mitochondrial DNA content in acute promyelocytic leukaemia

Diego A. Pereira-Martins^{1,2,3,4}  | Juan L. Coelho-Silva^{1,3,4} | Isabel Weinhäuser^{2,3,4} | Pedro L. Franca-Neto¹ | Douglas R. Silveira⁵  | César Ortiz^{3,4} | Amanda Moreira-Aguiar¹ | Marinus M. Lima¹ | Luisa C. Koury^{3,4} | Raul A. de Melo⁶ | Ana B. Glória⁷ | Evandro M. Fagundes⁷ | Bruno K. Lino⁸  | Katia Pagnano⁸  | Rosane Bittencourt⁹ | Elenaide Nunes¹⁰ | Fabiola Traina^{3,4} | Lorena Figueiredo-Pontes^{3,4} | Armand Keating¹¹ | Martin S. Tallman¹² | Raul C. Ribeiro¹³  | Richard Dilon¹⁴  | Arnold Ganser¹⁵ | Miguel A. Sanz^{16,17} | Nancy Berliner¹⁸ | Peter Valk¹⁹ | Bob Löwenberg¹⁹ | Tiziana Ottone^{20,21}  | Nelida I. Noguera^{20,21}  | Maria T. Voso^{20,21}  | Francesca Paoloni²¹ | Paola Fazi²² | Emanuele Ammatuna² | Gerwin Huls² | Jan Jacob Schuringa² | Eduardo M. Rego^{3,4,23}  | Antonio R. Lucena-Araujo¹ 

¹Department of Genetics, Federal University of Pernambuco, Recife, Brazil

²Department of Hematology, Cancer Research Centre Groningen, University Medical Centre Groningen, University of Groningen, Groningen, the Netherlands

³Department of Medical Imaging, Haematology, and Oncology, Ribeirão Preto Medical School, University of São Paulo, Ribeirão Preto, SP, Brazil

⁴Center for Cell Based Therapy, São Paulo Research Foundation, Ribeirão Preto, SP, Brazil

⁵Myeloid Leukaemia Genomics and Biology Group, School of Cancer and Pharmaceutical Sciences, King's College London, London, UK

⁶Department of Internal Medicine, University of Pernambuco, Recife, Brazil

⁷Hematology Division, Federal University of Minas Gerais, Belo Horizonte, Brazil

⁸Hematology and Hemotherapy Center, University of Campinas, Campinas, Brazil

⁹Hematology Division, Federal University of Paraná, Curitiba, Brazil

¹⁰Hematology Division, Federal University of Rio Grande do Sul, Porto Alegre, Brazil

¹¹Princess Margaret Cancer Centre, Toronto, Ontario, Canada

¹²Leukemia Service, Memorial Sloan-Kettering Cancer Center/Weill Cornell Medical College, New York City, New York, USA

¹³Department of Oncology, St. Jude Children's Research Hospital, Memphis, Tennessee, USA

¹⁴Department of Medical and Molecular Genetics, King's College London School of Medicine, London, UK

¹⁵Department of Hematology, Hemostasis, Oncology, and Stem Cell Transplantation, Hannover Medical School, Hannover, Germany

¹⁶Department of Hematology, Valencia University Medical School, Valencia, Spain

¹⁷CIBERONC, Instituto Carlos III, Madrid, Spain

¹⁸Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA

¹⁹Department of Hematology, Erasmus University Medical Center, Rotterdam, The Netherlands

²⁰Department of Biomedicine and Prevention, University of Tor Vergata, Rome, Italy

²¹Santa Lucia Foundation, I.R.C.C.S., Neuro-Oncohematology, Rome, Italy

²²Gruppo Italiano Malattie Ematologiche Maligne dell'Adulto (GIMEMA), GIMEMA Foundation, Rome, Italy

²³Hematology Division, LIM31, Faculdade de Medicina, University of São Paulo, São Paulo, Brazil

Diego A. Pereira-Martins and Juan L. Coelho-Silva contributed equally for this study.

© 2022 British Society for Haematology and John Wiley & Sons Ltd.

Correspondence

Antonio R. Lucena-Araujo, Department of Genetics, Federal University of Pernambuco, Av. Prof. Moraes Rego, 1235, Recife, PE 50670-901, Brazil.
 Email: araujoarl@gmail.com

Funding information

Conselho Nacional de Desenvolvimento Científico e Tecnológico, Grant/Award Number: 303479/2018-3; Fundação de Amparo à Pesquisa do Estado de São Paulo, Grant/Award Number: 2013/08135-2, 2015/09228-0, 2016/23191-4 and 2017/23117-1

Summary

Although a growing body of evidence demonstrates that altered mtDNA content (mtDNAC) has clinical implications in several types of solid tumours, its prognostic relevance in acute promyelocytic leukaemia (APL) patients remains largely unknown. Here, we show that patients with higher-than-normal mtDNAC had better outcomes regardless of tumour burden. These results were more evident in patients with low-risk of relapse. The multivariate Cox proportional hazard model demonstrated that high mtDNAC was independently associated with a decreased cumulative incidence of relapse. Altogether, our data highlights the possible role of mitochondrial metabolism in APL patients treated with ATRA.

KEYWORDS

anthracycline-based chemotherapy, ATRA, mtDNA content, oxidative phosphorylation

INTRODUCTION

Being in use in the clinics for more than three decades, all-*trans* retinoic acid (ATRA)-based treatment (in combination with chemotherapy or arsenic trioxide, ATO) represents the first-line therapy for standard-risk acute promyelocytic leukaemia (APL). Consequently, APL remains the most curable subtype of acute myeloid leukaemia (AML) to date. In addition to the degradation of the PML/RARA fusion protein,¹ ATRA modulates the mitochondria-mediated cellular metabolism and promotes a metabolic shift from glycolysis to oxidative phosphorylation.² Considering that mitochondrial alterations potentially represent a metabolic vulnerability in leukaemic cells,³ metabolic changes driven by ATRA could be clinically relevant.

Being an important part of the metabolic machinery, the biosynthesis of mitochondrial DNA (mtDNA) has been associated with increased mitochondrial biogenesis and dependence on oxidative phosphorylation.⁴ In addition, studies in solid tumours suggested that alterations in mtDNA content (mtDNAC) could have clinical implications.^{5,6} Yet, whether altered mtDNAC has an impact on clinical outcomes of patients with haematological malignancies (in particular APL and non-APL AML) remains largely unknown. Here, we hypothesize that APL patients with high mtDNAC, and consequently a high oxidative phosphorylation-driven metabolism, would be better responder to ATRA-based therapies.⁷ To test this hypothesis, we determined the mtDNAC in bone marrow (BM) samples of patients treated with ATRA and anthracycline-based chemotherapy as well as patients treated with the state-of-the-art combination ATRA+ATO.

METHODS

Overall, we enrolled 156 patients treated with ATRA and chemotherapy. Details of treatment protocols are published elsewhere.⁸ A second cohort was composed of 22 patients treated according to the APL 0406 protocol.⁹ All samples

were obtained at diagnosis. For comparison purposes, we included peripheral blood (PB) samples from 293 age- and sex-adjusted healthy volunteers with no history of haematological disorders. Additionally, we analysed CD34⁺ cells isolated from BM ($n = 3$), cord blood ($n = 9$), and apheresis ($n = 4$) samples from healthy donors. The study adhered to the tenets of the Declaration of Helsinki and informed consent was obtained from all patients or their relatives. The local Research Ethics Board of each participating centre approved the study. mtDNAC was quantified using quantitative real-time polymerase chain reaction and the results were normalized and expressed as fold-change relative to a reference DNA. Details can be found in the Supplemental data. Descriptive analyses (frequency and central tendency and dispersion measurements) were performed for patients' characteristics at baseline. Details of the statistical analysis and clinical endpoints are described in the supplemental data.

RESULTS AND DISCUSSION

First, we compared the mtDNAC between healthy (CD34⁺ and PB cells) and APL samples. **Figure 1A** shows that patients with APL had a higher-than-normal mtDNAC. Multiple comparison tests indicated no difference between healthy CD34⁺ and PB cells ($p = 0.956$), while the relative mtDNAC was significantly lower in CD34⁺ cells compared to APL ($p = 0.005$). **Table S1** summarizes the central tendency and dispersion of mtDNAC measurements of all the enrolled subjects. Restricting our analyses to APL, we noticed that the relative mtDNAC of APL patients ranged between 0.1 and 39. To evaluate the impact of mtDNAC on clinical outcomes without seeking an optimal cut-point, we used values of mtDNA higher than the 95th percentile of healthy PB samples (i.e., ≥ 1.63 ; **Figure 1A**, dashed line) to define APL patients with high mtDNAC. Consequently, patients that presented values within the range of normal control samples (< 1.63) were classified as normal mtDNAC. To account




FIGURE 1 Relative quantification of mtDNA in CD34⁺ and peripheral blood cells from healthy donors and bone marrow cells from patients with acute promyelocytic leukaemia (APL) treated with ATRA + ATO (A + A) and ATRA + chemotherapy (A + C) (A). The relative quantification of mtDNA was performed by quantitative real-time polymerase chain reaction relative to a single-copy nuclear gene (*PKLR* and *HBB*). The horizontal bars represent the median value of mtDNA (fold-change). Patient survival. The probability of disease-free survival (B) and cumulative incidence of relapse (C) in APL patients treated with A + C according to the mtDNA (entire cohort). Cumulative incidence of relapse in APL patients with low-risk of relapse (D) treated with A + C according to the mtDNA. Survival curves were estimated using the Kaplan–Meier method, and the log-rank test was used for comparison. Cumulative incidence curves for non-relapse death and relapse with or without death were constructed to reflect time to relapse and time to non-relapse death as competing risks. Time to relapse and time to non-relapse death was measured from the date of complete remission. Forest plot representing the multivariate Cox proportional hazard model for the cumulative incidence of relapse in APL patients treated with A + C according to the mtDNA (entire cohort) (E) and in low-risk patients (F). Confounding variables included in the model were mtDNA (high mass versus normal mass), risk of relapse (high-risk versus low-risk),¹⁰ and white blood cells counts (continuous variable).

for the potential variability in mtDNA between different sources, we measured the mtDNA of 22 paired BM and PB samples from APL patients collected at the time of diagnosis. Our analysis showed a strong correlation between BM and PB measured mtDNA of the same patient (Pearson correlation coefficient, $r = 0.78$, 95% confidence interval, CI: 0.54–0.9; Figure S1), suggesting that mtDNA detected in BM and PB could be used interchangeably with little bias.

Analysis of baseline characteristics indicated significantly lower mtDNA in *FLT3*-ITD mutated (median: 1.62, 95%CI: 1.6–4.4) compared to *FLT3*-wild-type patients (median: 2.7, 95%CI: 3.7–6.9) ($p = 0.03$; Figure S1). No other differences were found between patients with normal and high mtDNA (Table S2). Patient follow-up was last updated in May 2022. All patients treated with the combination ATRA+ATO achieved complete haematological remission with no deaths or relapses until the last follow-up. Consequently, univariate or multivariate regression

analyses were performed only for patients treated with ATRA and chemotherapy. With a median follow-up of 40 months (95% CI: 34–47 months), the estimated 5-year overall survival rate was 79% (95% CI: 72–84%). Overall, 129/156 (83%) patients achieved complete haematological remission. mtDNA had no impact on complete remission ($p = 0.518$) or overall survival ($p = 0.69$). Yet, patients with normal mtDNA had significantly lower 5-year disease-free survival (DFS) (61%, 95% CI: 46%–82%) compared to patients with high mtDNA (86%, 95% CI: 78%–95%) ($p = 0.035$; Figure 1B). Considering non-relapse mortality as a competing cause of failure, the 5-year cumulative incidence of relapse (CIR) for patients with normal and high mtDNA was 35% (95%CI: 16%–49%) and 10% (95%CI: 2%–17%), respectively ($p = 0.0082$; Figure 1C). After stratifying patients into low- and high-risk of relapse,¹⁰ we observed that the prognostic impact of mtDNA on CIR was applicable to the low-risk group (high mtDNA: 3%,

95%CI: 1%–7% versus normal mtDNAC: 26%, 95% CI: 5%–43%) ($p = 0.0018$; **Figure 1D**), but not to the high-risk group (high mtDNAC: 34%, 95% CI: 12%–51% versus normal mtDNAC: 58%, 95% CI: 14%–80%) ($p = 0.41$). Finally, the multivariate Cox proportional hazard model indicated that high mtDNAC was independently associated with decreased CIR in the entire cohort (**Figure 1F**) and in patients assigned to the low-risk group (**Figure 1F**).

In summary, we show that patients with higher-than-normal mtDNAC had better clinical outcomes (DFS and CIR) regardless of tumour burden. In addition, mtDNAC was useful to sub-stratify low-risk patients and identify those with a higher probability of relapse. Despite our promising data, these findings are only applicable to patients treated with ATRA and chemotherapy due to the limited sample size of patients treated with ATRA+ATO. Yet, it is conceivable that the greater and more sustained antileukemic effect of the combination ATRA+ATO¹¹ may overcome characteristics at diagnosis associated with an adverse outcome,¹² abrogating the need and the use of mtDNAC as prognostic marker in APL. Therefore, perhaps more important than providing a new prognostic marker for APL, we are interested to determine how to integrate the metabolic profile of leukaemic blasts into clinical decision making. Unfortunately, most studies of the mitochondrial genome in acute leukaemia are related to mtDNA mutations,^{13,14} and very few studies have explored these findings in a clinical context.^{15,16} Most of our knowledge regarding mtDNAC in cancer comes from studies conducted in solid tumours and, although altered amounts of mtDNAC have been widely reported, there is no consensus on how it affects tumour development. Both high and low mtDNAC have been associated with tumour progression and worse prognosis.^{5,6} These inconsistencies may be a result of different types of tumours, tissue sites, and treatments, whereby acute leukaemias are no exception. Similarly in acute leukaemias, unpublished data from our group indicate contrasting results in non-APL AML patients treated with cytotoxic chemotherapy compared to the results presented herein (Pereira-Martins et al., manuscript in progress). Altogether, it is possible that different types of tumours (including APL and non-APL AML) present different bioenergetic requirements. Particularly in non-APL AML, it remains to be explored whether its genetic heterogeneity dictate different metabolic requirements.

AUTHOR CONTRIBUTIONS

D.A.P-M, J.L.C-S, I.W. and P.L.F-N, performed experiments, collected, analysed, and interpreted data; and drafted the manuscript. D.R.S. performed the statistical analyses, interpreted the data, and drafted the manuscript. A.M-A., M.M.L., C.O., performed experiments data; and drafted the manuscript. L.C.K., R.A.M. A.B.G., E.M.F., B.K.L., K.P., R.B., E.N., F.T., L.L.F-P, A.K., M.S.T., R.C.R., R.D., A.G., M.A.S., N.B., P.V., B.L., T.O., N.I.N., M.T.V., F.P., P.F., E.A. G.H., obtained patient samples, updated the clinical data, and drafted the manuscript. D.A.P-M, J.L.C-S, J.J.S., E.M.R., A.R.L-A conceived and designed the study and reviewed the

manuscript. A.R.L-A. gave the final approval of the version to be submitted.

ACKNOWLEDGEMENTS

This investigation was supported by *Conselho Nacional de Desenvolvimento Científico e Tecnológico* (CNPq; grant # 303479/2018-3) and *Fundação de Amparo à Pesquisa do Estado de São Paulo* (FAPESP, Grant #2013/08135-2). D.A.P.M. (Grant #2017/23117-1), J.L.C.-S. (Grant #2016/23191-4) and I.W. (Grant #2015/09228-0) received a fellowship from FAPESP. I.W. and D.A.P-M were sponsored by the Abel Tasman Talent Program of the Graduate School of Medical Sciences of the University of Groningen/University Medical Center Groningen, The Netherlands.












CONFLICT OF INTEREST

The authors have no competing financial interests.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ORCID

Diego A. Pereira-Martins  <https://orcid.org/0000-0002-3302-4311>
 Douglas R. Silveira  <https://orcid.org/0000-0001-5298-9592>
 Bruno K. Lino  <https://orcid.org/0000-0002-6155-3253>
 Katia Pagnano  <https://orcid.org/0000-0001-7975-0805>
 Raul C. Ribeiro  <https://orcid.org/0000-0002-9956-5647>
 Richard Dillon  <https://orcid.org/0000-0001-9333-5296>
 Tiziana Ottone  <https://orcid.org/0000-0003-1936-3592>
 Nelida I. Noguera  <https://orcid.org/0000-0001-7682-862X>
 Maria T. Voso  <https://orcid.org/0000-0002-6164-4761>
 Eduardo M. Rego  <https://orcid.org/0000-0003-1567-4086>
 Antonio R. Lucena-Araujo  <https://orcid.org/0000-0003-1694-8958>

REFERENCES

- Zhu J, Gianni M, Kopf E, Honore N, Chelbi-Alix M, Koken M, et al. Retinoic acid induces proteasome-dependent degradation of retinoic acid receptor alpha (RARalpha) and oncogenic RARalpha fusion proteins. *Proc Natl Acad Sci*. 1999;96(26):14807–12. <http://www.pnas.org/cgi>, <https://doi.org/10.1073/pnas.96.26.14807>
- Tournaire F, Musinovic H, Gouranton E, Astier J, Marcotorchino J, Arreguin A, et al. All-trans retinoic acid induces oxidative phosphorylation and mitochondria biogenesis in adipocytes. *J Lipid Res* [Internet] 2015;56(6):1100–9. <https://linkinghub.elsevier.com/retrieve/pii/S002227520317223>
- Panina SB, Pei J, Kirienko N V. Mitochondrial metabolism as a target for acute myeloid leukemia treatment. *Cancer Metab* [Internet] 2021;9(1):17. <https://doi.org/10.1186/s40170-021-00253-w>. <https://cancerandmetabolism.biomedcentral.com/articles>
- Basak NP, Banerjee S. Mitochondrial dependency in progression of acute myeloid leukemia. *Mitochondrion* [Internet] 2015;21:41–8. <https://linkinghub.elsevier.com/retrieve/pii/S1567724915000173>
- He X, Qu F, Zhou F, Zhou X, Chen Y, Guo X, Li J., Huang Q., Yang Y., Lyu Z., Zhang H., Xing J. High leukocyte mtDNA content contributes to poor prognosis through ROS-mediated immunosuppression in hepatocellular carcinoma patients. *Oncotarget* [Internet] 2016;7(16):22834–45.
- Dang S, Qu Y, Wei J, Shao Y, Yang Q, Ji M, et al. Low copy number of mitochondrial DNA (mtDNA) predicts worse prognosis in early-stage laryngeal cancer patients. *Diagn Pathol* [Internet] 2014 5;9:28.

7. Wierenga ATJ, Cunningham A, Erdem A, Lopera NV, Brouwers-Vos AZ, Pruis M, et al. HIF1/2-exerted control over glycolytic gene expression is not functionally relevant for glycolysis in human leukemic stem/progenitor cells. *Cancer Metab* [Internet]. 2019 27;7(1):11. <https://doi.org/10.1186/s40170-019-0206-y>. <https://cancerandmetabolism.biomedcentral.com/articles/>
8. Rego EM, Kim HT, Ruiz-Argüelles GJ, Undurraga MS, Uriarte Mdel R, Jacomo RH, et al. Improving acute promyelocytic leukemia (APL) outcome in developing countries through networking, results of the International Consortium on APL. *Blood* [Internet] 2013;121(11):1935–43.
9. Lo-Coco F, Avvisati G, Vignetti M, Thiede C, Orlando SM, Iacobelli S, et al. Retinoic acid and arsenic trioxide for acute promyelocytic leukemia. *N Engl J Med* [Internet] 2013;369(2):111–21.
10. Sanz MA, Lo Coco F, Martín G, Avvisati G, Rayón C, Barbui T, et al. Definition of relapse risk and role of nonanthracycline drugs for consolidation in patients with acute promyelocytic leukemia: a joint study of the PETHEMA and GIMEMA cooperative groups. *Blood* [Internet] 2000;96(4):1247–53.
11. Platzbecker U, Avvisati G, Cicconi L, Thiede C, Paoloni F, Vignetti M et al. Improved outcomes with retinoic acid and arsenic trioxide compared with retinoic acid and chemotherapy in non-high-risk acute promyelocytic leukemia: final results of the randomized Italian-German APL0406 trial. *J Clin Oncol* [Internet] 2017;35(6):605–12.
12. Cicconi L, Divona M, Ciardi C, Ottone T, Ferrantini A, Lavorgna S, Alfonso V, Paoloni F, Picocchi A, Avvisati G, Ferrara F, di Bona E, Albano F, Breccia M, Cerqui E, Sborgia M, Kropp MG, Santoro A, Levis A, Sica S, Amadori S, Voso MT, Mandelli F, Lo-Coco F PML-RAR α kinetics and impact of FLT3-ITD mutations in newly diagnosed acute promyelocytic leukaemia treated with ATRA and ATO or ATRA and chemotherapy. *Leukemia* [Internet] 2016;30(10):1987–92.
13. Piccoli C, Ripoli M, Scrima R, Stanziale P, Di Ianni M, Moretti L, et al. MtDNA mutation associated with mitochondrial dysfunction in megakaryoblastic leukaemic cells. *Leukemia* [Internet] 2008;22(10):1938–41.
14. Yao Y-G, Ogasawara Y, Kajigaya S, Molldrem JJ, Falcão RP, Pintão M-C, et al. Mitochondrial DNA sequence variation in single cells from leukemia patients. *Blood* [Internet] 2007;109(2):756–62.
15. Silkjaer T, Nørgaard JM, Aggerholm A, Ebbesen LH, Kjeldsen E, Hokland P, Nyvold CG Characterization and prognostic significance of mitochondrial DNA variations in acute myeloid leukemia. *Eur J Haematol* [Internet] 2013;90(5):385–96.
16. Kang M-G, Kim Y-N, Lee JH, Szardenings M, Baek H-J, Kook H, Kim HR, Shin MG Clinicopathological implications of mitochondrial genome alterations in pediatric acute myeloid leukemia. *Ann Lab Med* [Internet] 2016; 36(2):101–10.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Pereira-Martins DA, Coelho-Silva JL, Weinhäuser I, Franca-Neto PL, Silveira DR, Ortiz C, et al. Clinical significance of mitochondrial DNA content in acute promyelocytic leukaemia. *Br J Haematol*. 2022;00:1–5. <https://doi.org/10.1111/bjh.18510>