

University of Groningen

Identification of annual routines and critical stopover sites of a breeding shorebird in the Yellow Sea, China

Wu, Yang; Lei, Weipan; Zhu, Bingrun; Xue, Jiaqi; Miao, Yuanxiang; Zhang, Zhengwang

Published in:
 Avian Research

DOI:
[10.1016/j.avrs.2022.100068](https://doi.org/10.1016/j.avrs.2022.100068)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2022

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Wu, Y., Lei, W., Zhu, B., Xue, J., Miao, Y., & Zhang, Z. (2022). Identification of annual routines and critical stopover sites of a breeding shorebird in the Yellow Sea, China. *Avian Research*, 13, Article 100068. <https://doi.org/10.1016/j.avrs.2022.100068>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Identification of annual routines and critical stopover sites of a breeding shorebird in the Yellow Sea, China



Yang Wu^{a,1}, Weipan Lei^{a,b,1}, Bingrun Zhu^{a,c}, Jiaqi Xue^a, Yuanxiang Miao^d, Zhengwang Zhang^{a,*}

^a Key Laboratory for Biodiversity Science and Ecological Engineering, College of Life Sciences, Beijing Normal University, Beijing, 100875, China

^b Demonstration Center for Experimental Life Sciences & Biotechnology Education, College of Life Sciences, Beijing Normal University, Beijing, 100875, China

^c Conservation Ecology Group, Groningen Institute for Evolutionary Life Sciences (GELIFES), University of Groningen, PO Box 11103, 9700, CC, Groningen, the Netherlands

^d School of Life Science and Technology, Shanghai Tech University, Shanghai, 201210, China

ARTICLE INFO

Keywords:

Bohai bay

EAAF

Lianyungang

Migration

Recurvirostra avosetta

ABSTRACT

Pied Avocets (*Recurvirostra avosetta*) are common migratory shorebirds in the East Asian–Australasian Flyway. From 2019 to 2021, GPS/GSM transmitters were used to track 40 Pied Avocets nesting in northern Bohai Bay to identify annual routines and key stopover sites. On average, southward migration of Pied Avocets started on 23 October and arrived at wintering sites (mainly in the middle and lower reaches of the Yangtze River and coastal wetlands) in southern China on 22 November; northward migration started on 22 March with arrival at breeding sites on 7 April. Most avocets used the same breeding sites and wintering sites between years, with an average migration distance of 1124 km. There was no significant difference between sexes on the migration timing or distance in both northward and southward migration, except for the departure time from the wintering sites and winter distribution. The coastal wetland of Lianyungang in Jiangsu Province is a critical stopover site. Most individuals rely on Lianyungang during both northward and southward migration, indicating that species with short migration distances also heavily rely on a few stopover sites. However, Lianyungang lacks adequate protection and is facing many threats, including tidal flat loss. We strongly recommend that the coastal wetland of Lianyungang be designated as a protected area to effectively conserve the critical stopover site.

1. Introduction

Identifying migration routines and the stopover sites of shorebirds is critical for understanding their life cycle and for conservation management (Battley et al., 2012; Giunchi et al., 2019). Such a task is essential for the East Asian–Australasian Flyway (EAAF), which is the most species-rich flyway globally but also has the most species in decline due to the loss of natural wetlands over the past several decades (Kirby et al., 2008; Murray et al., 2014; Studds et al., 2017).

Tracking the annual cycle of shorebirds can uncover their migration strategy, reveal potential stopover sites, and even show some behaviour adaptive to the natural tidal flat lost in the flyways (Lisovski et al., 2016; Piersma et al., 2021). Many efforts have been made on the EAAF, focusing mainly on long-distance migration or threatened species. For example, tracking the Great Knots (*Calidris tenuirostris*) provided further

evidence that they used the Yellow Sea as the major stopover sites on both northward and southward migration and showed some short-stay sites probably for rest only before arriving at the Yellow Sea (Ma et al., 2013; Lisovski et al., 2016). The tracking of the Red Knot (*Calidris canutus piersma*) showed that ‘long jumpers’ behaved as ‘skippers’. Rather than staging at a single site for multiple weeks halfway during migration, they made several stops of up to a week. Recent reductions in the extent of suitable habitat in the Yellow Sea area (Murray et al., 2014; Piersma et al., 2016) may have contributed to the current pattern (Piersma et al., 2021). Tracking of Whimbrel (*Numenius phaeopus*) indicated that different populations have different reliance on the Yellow Sea, which might be related to their different population trends (Kuang et al., 2020).

However, less effort has been put into tracking the annual cycle of short-distance or least concerned species, despite their greater importance in ecosystem function and service provision (Inger et al., 2015).

* Corresponding author.

E-mail address: zzw@bnu.edu.cn (Z. Zhang).

¹ These authors contributed equally to this work.

The neglect of common species hinders our knowledge of their migration route, breeding/wintering area, and the potential risk they may face during their life cycle. Population collapse can occur even in species with a billion individuals (Bucher, 1992).

Pied Avocets (*Recurvirostra avocetta*) are common shorebirds that occur across most regions of Asia, Europe, and Africa (IUCN, 2020). Males and females do not show strong sexual size dimorphism, and the breeding system is seasonally monogamous, with both sexes performing egg incubation and chick brooding (Pierce et al., 2020). Historically, Pied Avocets are common breeding waterbirds on the east coast of England during the early 19th century but disappeared in the mid-19th century due to the loss of natural habitat and overharvest of their eggs by people. They were reintroduced one hundred years later with the reconstruction of habitats and other methods in the 1940s (Hill, 1988). In the EAAF, the population of Pied Avocets is estimated to be 100,000 (Wetlands International, 2020). Although the overall population trend is not clear, a local study found that the winter peak in Shenzhen Bay/Deep Bay in southern China has been decreasing in recent years (Choi et al., 2020), which means we need a focus on this common but neglected species.

The migration of the Pied Avocet has not been well studied, although some migration routes were revealed by banding in Europe (Hotker and West, 2005; Chambon et al., 2018), which is especially poorly documented in Asia. Their traditional breeding range was distributed from Central Asia to Southeast Siberia and Northeast China. However, in recent years, Pied Avocets have been documented breeding in areas farther south, for example, in the Bohai Bay of the Yellow Sea (Lei et al., 2021), in Chongming Dongtan in the East China Sea (Wanjuan Ke, personal communication), and in Hong Kong (Allcock et al., 2012).

Tracking avocets can also provide insight into sex-specific migration strategies. The migration timing and wintering area are different between sexes and are common in shorebirds. Migration timing (arrival at or departure from breeding/wintering grounds) is supposed to relate to the breeding system. Males usually arrive at the breeding area earlier than females in shorebirds to occupy the territory. Males that do not provide incubation or brooding leave earlier from the breeding area than females (Meissner, 2006). Males and females have different winter distributions that are supposed to be related to sexual size dimorphism (Nebel, 2005) or resource partitioning (Duijns et al., 2014). Studying the sex differences in shorebirds in terms of migration strategies can be helpful for detecting the factors linked to intraspecific variation, explaining their distribution pattern, and finding conservation gaps.

The wetlands of eastern China support a large non-breeding population of Pied Avocets (Cao et al., 2009). However, to the best of our knowledge, the annual routines and critical stopover sites of Pied Avocets have not been revealed. In this study, we tracked the avocets breeding in the Yellow Sea using satellite transmitters. We aim to identify their annual routines and stopover sites. We also explored sex-differential migration in this species. We will advance our understanding of migration ecology and fill the conservation knowledge gap for a common shorebird species in the EAAF.

2. Methods

2.1. Study area and data collection

We captured Pied Avocets (hereafter “avocets”) at two sites in northern Bohai Bay, Yellow Sea: (i) Nanpu salt pans, and (ii) Yuetuo Island, both in Tangshan, Hebei Province, China (Fig. 1). Nanpu salt pans (118.1–118.3° E, 39.0–39.2° N) are an artificial habitat for waterbirds in the EAAF (Lei et al., 2018). Yuetuo Island (118.8–118.9° E, 39.1–39.2° N) is an artificial sand island located 50 km east of the Nanpu salt pans. More than 500 breeding pairs were recorded at the study sites (Wu, 2019–2021, unpublished results). Avocets arrived at the study sites in early March and started breeding in late April (Lei et al., 2021). They were captured using walk-in traps (0.8 m × 0.8 m × 0.6 m) or clap nets ($N_{2019} = 27$, $N_{2020} = 13$) during the incubation period.

Captured individuals were banded with a metal ring on the left tibia and colour flags (blue over yellow) on the right tibia with a unique code. A blood sample of 10–80 µL was collected from the brachial vein with a cotton swab and stored in a centrifuge tube at –20 °C for subsequent molecular sexing (Van der Velde et al., 2017).

In 2019–2020, we captured 40 adult avocets (22 males, 18 females) and outfitted them with tracking devices using a leg-loop harness (Verhoeven et al., 2020). Two types of transmitters were used: (1) HQPG2009P ($N = 38$, 9 g solar GPS/GSM, Hunan Global Messenger Technology Ltd., Changsha City, China) was programmed to record one location every 2–12 h, and (2) HQPG 1206 ($N = 2$, 7 g solar GPS/GSM) was programmed to record one location every 3–12 h.

To understand the present situation of critical stopover sites (see Results), we visited Lianyungang city, Jiangsu Province in 2020 to search for the 17 tagged avocets known to be present there during that autumn. In addition, we counted avocets from 31 October to 1 November in Lianyungang.

2.2. Tracking data

We retrieved the tracking data via the service platform of HQXS (<http://www.hqxs.net>), and the locations of transmitters with accuracy classes A, B, and C were retained (representing precisions of 5 m, 10 m, and 200 m, respectively).

Location was defined as the GPS point of transmitters; site was defined as the area of a bird staged for a while. Two avocets that lost signal within three months after release were excluded. Two avocets lost their signals in November 2019 during their southward migration. From July 2019 to June 2021, 36 individuals completed at least a southward or northward migration episode (single trip from wintering to breeding area or from breeding to wintering area, Appendix Table S1). There were 46 complete northward episodes from 35 individuals (19 males, 16 females) and 47 complete southward episodes from 35 individuals (18 males, 17 females).

Many avocets had second brood in southern Bohai Bay and Laizhou Bay of Bohai sea, after their reproduction in northern Bohai Bay. Thus, we take the farthest area south of the Bohai Sea (36.6° N) as the southern boundary of the breeding area. Most individuals would not travel long distances after arriving at the middle and lower reaches of the Yangtze River. Thus, we take the northern boundary of the middle and lower reaches of the Yangtze River (31° N) as the northern boundary of the wintering area. Based on the annual movements of the tracked avocets, we defined that the breeding and wintering areas were above 36.6° N and below 31° N, respectively. We considered an individual to be stationary when the recorded transmitter showed a speed of ~0 km/h (Zhu et al., 2021). Thus, the remaining stationary locations fell between these two latitudes and were considered stopping areas (Chan et al., 2019). The heatmap was used to describe the avocet breeding, wintering, and stopping areas in QGIS 3.20.0. Based on geographical location, we categorized the habitat of stationary locations into two types: (i) inland wetlands, and (ii) coastal wetlands.

During migration, we assumed that the last sites where birds remained within 50 km for more than 48 h were considered stopover sites (Kuang et al., 2020). The departure and arrival dates at the breeding area and wintering areas were determined by their last site. During southward migration, the last site in the breeding area with a duration of staying more than 48 h within 50 km was treated as the start of southward migration. The time of the last location on this site was the departure date. If tracked individuals left their breeding area and then flew back, the last time was used as the departure time. The first site in the wintering area with a duration of staying more than 48 h within 50 km was treated as the last site of southward migration. The time of the first location on this site was the arrival date. When the migration tracking signal was lost over 24 h but less than 10 d, the starting date of migration was defined as the medium date between the date of the last location in the breeding area and the first location during migration. Individuals




Fig. 1. Annual distribution of tracked Pied Avocets. The heatmap represents low densities to high densities of locations from pale to dark.

whose signal was lost over 10 d were excluded. The departure and arrival dates of northward migration were calculated similarly.

We used the R package 'geosphere' (Hijmans et al., 2017) to calculate the distance between the last location of all breeding/stopover sites before migration and the first location when arriving at the next stoping/wintering sites. The migration distance is the sum of the distance between the breeding sites, wintering sites, and all stopover sites on the route.

Tracked individuals with more than one repeated southward/northward migration episode (and individuals with incomplete episodes but had locations in breeding areas or wintering areas for more than one year) were collected to calculate breeding/wintering site fidelity at the individual level. We distinguished birds wintered in inland wetlands or coastal wetlands. Individuals were considered to have changed their wintering area only when they changed their wintering sites from inland wetlands to coastal wetlands or vice versa. Wintering site fidelity was determined by the wintering area change percentage between two years. A lower percentage indicates higher site fidelity. The breeding site fidelity was calculated similarly.

2.3. Statistical analysis

The linear mixed model (LMM) from the R package 'lme4' (Bates et al., 2015) was used to explore whether migration distance or duration differed between sexes and seasons, with individuals as random effects. We also used LMM to explore whether departure and arrival times of southward and northward migration differed between the sexes and years, with individuals as random effects. The statistical significance of fixed effects was calculated with the R package 'lmerTest' (Kuznetsova et al., 2017). Based on the wintering location of avocets, Fisher's exact test was used to explore whether males and females differed in the winter distribution. The *t*-test was used to explore whether the arrival and departure times of Lianyungang during northward migration differed between sexes. Analysis was conducted in R (R Core Team, 2020). Data are presented as the mean \pm standard deviation (SD).

3. Results

3.1. Annual routines, wintering and stopover sites

During southward migration, most individuals departed from Bohai Bay, but one departed from Liaodong Bay, Liaoning Province, in 2019 and 2020, and one departed from Hulun Buir, the Inner Mongolia Autonomous Region, in 2020 (Figs. 1 and 2), because they moved northwards after breeding in Bohai Bay. Tracked birds wintered in two areas of southern China: inland wetlands in the middle and lower reaches of the Yangtze River (hereafter "MLYR") and coastal wetlands in southeastern China (Figs. 1 and 2). Most individuals chose MLYR as their wintering sites (22 males, 7 females), and other individuals wintered in coastal wetlands (4 males, 8 females). Seven (all females) wintered in both areas and shifted from inland wetlands to coastal wetlands in the same season. There was a significant difference between sexes in the winter distribution. Males prefer inland wetlands, and females prefer coastal wetlands ($p = 0.001$; Appendix Table S1). In addition, one individual wintered on Taiwan Island in 2020, and two wintered in Yancheng coast wetland, Jiangsu Province, which is north of the wintering areas of other avocets. The coordinates of these areas are described in Fig. 1.

The change rates of both wintering area and breeding sites between years were small. In the 17 avocets with wintering tracking data both in 2019 and 2020 (Appendix Table S1), 14 wintered in inland areas, 3 wintered in a coastal area in 2019, and 11.76% of individuals (2 in 17) changed wintering areas in 2020 (both switched from inland to coastal wetlands). Twenty-one individuals captured in Bohai Bay in 2019 revisited their breeding sites in 2020. In 2021, 21 of 24 individuals (87.5%) were still bred at previous breeding sites. Thus, on average, 8.7% of individuals changed their breeding sites in three years.

During both southward and northward migration, most stopover sites were located along the coastal wetlands, while a few stopover sites were in the inland wetlands (Fig. 1).

The coastal wetland of Lianyungang (Lianyungang as follows) is the most important stopover site for avocets. More than 60% of tracked individuals staged in Lianyungang during southward migration, with an average of 36 days of stopover duration (Table 1). During the northward migration, more than 40% of tracked individuals staged in Lianyungang, with an average of 18 days of stopover duration. In the autumn of 2020, we surveyed Lianyungang and observed 39,700 avocets on the tidal flat within two days, up to 40% of the population in the EAAF (Appendix Fig. S1).

3.2. Migration timing and distance

On average, avocets started southward migration on 23 October (Table 2). The majority reached the wintering grounds on 22 November. Males and females did not differ significantly in the departure time from the breeding sites ($p = 0.59$), the time of arrival at the wintering sites ($p = 0.062$), or the duration ($p = 0.098$). On average, avocets started northward migration on 22 March, and the majority reached the breeding sites on 7 April. The males departed from the wintering grounds earlier than females ($p = 0.048$), but both sexes did not differ significantly in the time of arrival at the breeding sites ($p = 0.59$). Thus, the duration of northward migration of males was slightly longer than the duration of northward migration of females but not significantly different ($p = 0.126$). For tracked birds staged in Lianyungang, males arrived at Lianyungang earlier than females ($p = 0.013$) in northward migration, but their departure time from here was similar ($p = 0.88$), which means that males stayed longer than females.

The great circle distances between the wintering and breeding sites ranged from 80 to 2350 km with an average of 1124 km, with no significant differences between males and females ($p = 0.086$; Table 2). However, the migration duration of males was longer than the migration duration of females ($p = 0.048$). Avocets migrated longer distances ($p < 0.001$) and shorter durations ($p = 0.002$) during northward migration (1256 ± 471 km and 15.61 ± 17.12 d) than southward migration (1019 ± 268 km and 30.13 ± 36.52 d).

4. Discussion

4.1. Wide-range wintering area

Although the tracked birds were caught in a small breeding area (within 50 km) in Bohai Bay, they showed a broad wintering area with a maximum ca. 1000 km apart from each other, from the inland freshwater environments (MLYR) to coasts in southern China. Most individuals wintered in MLYR, while a few individuals overwintered in coastal wetlands. This result is inconsistent with those about only 15% (14,590) and 32% (31,971) of the estimated population of avocets overwintered on the inland wetlands at the beginning of this century (Cao et al., 2009) and in recent years (Tao et al., 2017), respectively. This difference may arise from the research method, i.e., the counting data of these two studies reflected the general wintering population, but the tracking data of this study were based on the breeding population in Bohai Bay, which may prefer the inland wintering area. Thus, more tracking data from other breeding areas or wintering areas are needed. In western Europe, most avocets tend to use coastal wetlands during the wintering period (Moreira, 1995; Hotker and West, 2005). Shorebirds that habituated both coastal and inland wetlands could also be found in Black-tailed Godwit (*Limosa limosa*; Estrella and Masero, 2010; Lourenço et al., 2010) and Dunlin (*Calidris alpina*; Abad-Gómez et al., 2020). Compared to coastal habitats, inland freshwater habitats where pathogens and parasites are generally more abundant may increase investment in the immune system (Piersma, 2003; Abad-Gómez et al., 2020). Avocets that spend more energy resisting pathogens may have a negative impact on subsequent




Fig. 2. Annual migration routes of Pied Avocets. The northernmost blue line represents individual H51. In 2020 and 2021, H51 bred in Hulun Buir, north of the Bohai Sea.

Table 1

The number of tracked Pied Avocets staged in Lianyungang during four migration periods.

Year	Migration period	Tracked individuals staged	Proportion of staged individuals (%)	Average duration of staging (d)	Range of duration (d)
2019	Southward	13	65	35.9 ±	1–110
2020	Southward	17	61	34.5	
2020	Northward	9	41	18.4 ±	1–52
2021	Northward	12	52	13.2	

Only individuals who staged for more than two days were calculated for the duration.

migration and breeding. Female avocets may be more constrained by such energy costs than males; thus, they prefer wintering in coastal wetlands.

Winter site conditions substantially influence the survival of migratory species (McMahon et al., 2003; Woodworth et al., 2017). Dispersed wintering patterns may positively affect population survival by diluting the habitat degeneration risk of specific wintering areas (Cresswell, 2014; Bracey et al., 2018). However, avocet also showed winter site fidelity at the individual level and showed strong breeding site fidelity at the population level; thus, they are still sensitive to habitat loss in both wintering and breeding areas.

4.2. Migration strategy

Males and females did not differ in departure time from the breeding area, and arrival time at wintering grounds may be due to small sexual size dimorphism and both sexes participating in reproduction. Males depart from wintering areas earlier than females, but they did not differ in arrival time at breeding sites, which is in line with the result of Black-tailed Godwits (Zhu et al., 2021). This finding can be explained by males

Table 2

Estimated regression parameters, standard errors, profile confidence intervals (2.5% PCL and 97.5% PCL), χ^2 values, df, and *P* values for each predictor variable to investigate the effects of season/year and sex (as fixed effects) on the migration distance, migration duration and migration timing of the Pied Avocet (*Recurvirostra avosetta*).

Parameters	Fix effects	Estimates (β)	2.5% PCL	97.5% PCL	SE	χ^2	df	<i>p</i>
Migration distance	Intercept	1316.90	1156.45	1477.15	82.28			
	Period (South)	-213.00	-325.16	-100.61	56.80	12.73	1	< 0.001
	Sex (M)	-178.20	-384.32	26.69	105.00	2.94	1	0.086
Migration duration	Intercept	5.84	-5.52	17.16	5.82			
	Period (South)	15.84	6.06	25.51	4.94	9.60	1	0.002
	Sex (M)	14.11	0.15	28.00	7.12	3.92	1	0.048
Distance of northward migration	Intercept	1278.47	1041.39	1514.61	122.04			
	Year (2021)	110.78	-77.24	289.09	91.30	1.46	1	0.228
	Sex (M)	-225.31	-520.02	66.16	150.26	2.34	1	0.126
Distance of southward migration	Intercept	1091.34	940.36	1244.54	76.03			
	Year (2020)	14.72	-141.11	162.06	76.09	0.03	1	0.859
	Sex (M)	-148.94	-307.01	8.02	80.76	3.48	1	0.062
Duration of northward migration	Intercept	12.28	3.89	20.63	4.32			
	Year (2021)	-5.25	-12.48	1.73	3.58	2.24	1	0.135
	Sex (M)	7.59	-2.59	17.64	5.19	2.19	1	0.139
Duration of southward migration	Intercept	18.55	0.98	36.17	9.06			
	Year (2020)	2.24	-7.71	12.40	5.00	0.21	1	0.644
	Sex (M)	19.30	-3.77	42.36	11.81	2.73	1	0.098
Arrival time at the breeding sites	Intercept	98.64	88.79	108.53	5.09			
	Year (2021)	0.25	-6.40	7.28	3.39	0.01	1	0.942
	Sex (M)	-3.38	-15.89	9.09	6.40	0.30	1	0.586
Arrival time at the wintering sites	Intercept	319.81	309.31	330.29	5.41			
	Year (2020)	1.37	-7.70	9.78	4.33	0.10	1	0.751
	Sex (M)	12.03	-0.60	24.61	6.46	3.50	1	0.061
Departure time from the breeding sites	Intercept	299.67	284.23	314.99	7.90			
	Year (2020)	0.68	-9.39	10.00	4.78	0.02	1	0.884
	Sex (M)	-5.43	-25.51	14.64	10.28	0.30	1	0.587
Departure time from the wintering sites	Intercept	85.61	75.80	95.45	5.07			
	Year (2021)	7.16	-2.53	17.42	5.00	2.14	1	0.143
	Sex (M)	-11.42	-22.50	-0.09	5.73	3.90	1	0.048

spending more time at stopover sites. Their arrival time at Lianyungang is earlier than the arrival time of females, but the departure time from Lianyungang is similar between the sexes.

The distance and main stopover sites of northward and southward migration routes are similar. The migration distance of avocets breeding in Bohai Bay is similar to some inexperienced individuals on the French Atlantic coast in a ringing study (Chambon et al., 2018). Compared to other common migration shorebirds in the EAAF (e.g., Bar-tailed Godwit *Limosa lapponica*, Red Knot, Great Knot, and Whimbrel) that fly over 10,000 km (Lisovski et al., 2016; Kuang et al., 2020; Piersma et al., 2021), avocets can be classified as short-distance migration species. Furthermore, our results indicated that short-distance migratory species also rely on the stopover site during migration, which should not be neglected.

The migration distance of avocets ranged from 80 km to more than 2300 km, showing a 30-fold variation. Some individuals migrated from breeding areas to wintering areas or vice versa with a non-stop flight. Avocets can fly more than 1000 km within a single day, but only a small proportion of individuals use such a strategy. Overall, avocets showed great diversity in their migration behaviour. They can finish the migration without rest or spend over three months at stopover sites. The migration strategy of avocets can be described as a continuum, from the “cost minimization strategy” (for the individuals stopping for a long time in stopover sites) to the “time minimization strategy” (for the individuals with non-stop migration), and more research about such diversity strategies can fill the knowledge gap of individual specifications of shorebird migration.

4.3. Importance of Lianyungang

Both northward and southward migration routes showed the “umbrella” pattern, which highlighted the importance of Lianyungang in the migration path of the population of avocets in East Asia (Fig. 3). Our survey suggested that Lianyungang supported nearly 40% of the avocet population in the EAAF. Another report from Spoon-billed Sandpiper in



Fig. 3. Flocks of Pied Avocets in Lianyungang during northward migration. (Photo credits: Yongxiang Han).

China (2022) found more than 22,000 avocets in Lianyungang during southward migration in the same year. This number may be underestimated because their count focuses on the Asian Dowitcher (*Limnodromus semipalmatus*). In addition to northward migration in 2020, more than half of the tracked avocets refuel in Lianyungang during southward migration (Table 1). Lianyungang is in the middle point between the breeding sites of Bohai Bay and the wintering sites of MLYRR, which have a straight-line distance of approximately 500 km. There was no large and suitable habitat for avocet after they migrated southward across the inland Shandong Peninsula until Lianyungang (Fig. 1). Lianyungang is the first staging site for avocets to migrate from Bohai Bay and is also the last refuelling site before they fly across the peninsula to breeding sites. If avocets choose a detour flight for an extra 500 km to fly along the coastline, their single non-stop migration distance increases from 500 km to 1000 km. In addition to avocets, Lianyungang supported over 1% of the flyway populations of other 21 shorebird species, and most are long-migration distance species threatened by losing important stopover sites (Chan et al., 2019). In particular, the Asian Dowitcher, for 97.5% of its estimated global population, was recorded in 2019 in Lianyungang (Yang et al., 2021). If Lianyungang is lost, there will be a serious impact on the annual migration of avocets, as well as other birds.

Unfortunately, 27% of Lianyungang's intertidal soft-sediment habitats have been reclaimed from 2003 to 2018 (Chan et al., 2019), and a large area of tidal flat has been invaded and occupied by the salt-water cordgrass *Spartina alterniflora*. However, with only a provincial wetland park being established, Lianyungang still lacks adequate protection.

5. Conclusions

This study is the first to examine the annual routines and main stopover sites of Pied Avocets in the EAAF. As a short-distance migratory species, avocets tracked in our study spent their whole annual life cycle within China, which is different from most shorebirds in the EAAF. They connected the inland freshwater wetlands and coastal wetlands in the Yellow Sea. Thus, their population fate is closely linked with the status of both types of wetlands in China. Identifying their critical stopover sites, such as Lianyungang, and building protected areas for the areas is vital for future conservation. The international responsibilities for conserving wetlands require China to balance economic development and biodiversity conservation. We strongly recommend that the coastal wetlands of Lianyungang be enrolled in protected areas and guaranteed under law as soon as possible.

Authors' contributions

YW and WL conceived the experiments, analyzed the data, wrote the manuscript, and reviewed the drafts; BZ analyzed the data and reviewed the drafts; JX performed the experiment in the field and analyzed the

data; YM analyzed the data; ZZ conceived and designed the experiments, and reviewed drafts of the paper. All authors read and approved the final manuscript.

Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: Zhengwang Zhang reports financial support was provided by National Natural Science Foundation of China and Non-profit Foundation of Marine Environment and Ecological Conservation of CNOOC. Weipan Lei reports financial support was provided by National Natural Science Foundation of China.

Acknowledgement

We thank volunteers (Chuyu Cheng, Shunlong Liu, Peichen Lin, Xiangwan Liu, Dan Li, Luoyi Kang, Qingxin Ma, Hengjun Xiao), colleagues from Beijing Normal University, and the Global Flyway Network who joined and offered their help in Tianjin and Nanpu. We also thanks Professor Weidong Kong, Dr. Hui Lin, Yang Sun and Jing Zhang for their advice to refine the paper. We also thank the two reviewers who provided valuable and insightful comments which greatly improved the manuscript. This research was funded by the National Natural Science Foundation of China (grant number 31830089, 32270518, 31801985), and the Non-profit Foundation of Marine Environment and Ecological Conservation of CNOOC (grant number CF-MEEC/TR/2020-20).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.avrs.2022.100068>.

Ethics statement

All experimental protocols of bird capture and handling were strictly complied with the requirement of Chinese Wild Animal Protection Law and all necessary permission for bird banding was obtained.

References

- Abad-Gómez, J.M., Villegas, A., Gutiérrez, J.S., Parejo, M., Navedo, J.G., Sánchez-Guzmán, J.M., et al., 2020. Pro-inflammatory immune response is linked to wintering habitat in a migratory shorebird. *Auk* 137, ukaa046. <https://doi.org/10.1093/auk/ukaa046>.
- Allcock, J., Carey, G., Chow, G., Welch, G., 2012. Hong Kong Bird Report 2009–10. Hong Kong Bird Watching Society. <https://www.hkbws.org.hk/cms/resource-tw/bird-report-tw/file/39-2009-2010>. (Accessed 15 September 2021).
- Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear mixed-effects models using lme4. *J. Stat. Software* 67, 1–48. <https://doi.org/10.18637/jss.v067.i01>.
- Battley, P.F., Warnock, N., Tibbitts, T.L., Gill, R.E., Piersma, T., Hassell, C.J., et al., 2012. Contrasting extreme long-distance migration patterns in bar-tailed godwits *Limosa lapponica*. *J. Avian Biol.* 43, 21–32. <https://doi.org/10.1111/j.1600-048X.2011.05473.x>.
- Bracey, A., Lisovski, S., Moore, M., McKellar, A., Craig, E., Matteson, S., et al., 2018. Migratory routes and wintering locations of declining inland North American Common Terns. *Auk* 135, 385–399. <https://doi.org/10.1642/AUK-17-210.1>.
- Bucher, E.H., 1992. The causes of extinction of the passenger pigeon. In: Power, D.M. (Ed.), *Current Ornithology*. Springer, Boston, pp. 1–36. https://doi.org/10.1007/978-1-4757-9921-7_1.
- Cao, L., Tang, S., Wang, X., Barter, M., 2009. The importance of eastern China for shorebirds during the non-breeding season. *Emu* 109, 170–178. <https://doi.org/10.1071/MU08051>.
- Chambon, R., Dugravot, S., Paillisson, J.-M., Lemesle, J.-C., Ysnel, F., Gélaud, G., 2018. Partial migration in inexperienced pied avocets *Recurvirostra avosetta*: distribution pattern and correlates. *J. Avian Biol.* 49. <https://doi.org/10.1111/jav.01549>.
- Chan, Y.C., Peng, H.B., Han, Y.X., Chung, S.S.-W., Li, J., Zhang, L., et al., 2019. Conserving unprotected important coastal habitats in the Yellow Sea: shorebird occurrence, distribution and food resources at Lianyungang. *Glob. Ecol. Conserv.* 20, e00724. <https://doi.org/10.1016/j.gecco.2019.e00724>.
- Choi, C.Y., Li, J., Xue, W.J., 2020. China coastal waterbird census report 2012–2019 (English). Hong Kong Bird Watching Society. <https://cms.hkbws.org.hk/cms/en/reso>

- urce/publication/regional-publication/file/657-ccwc-report-2012-2019-eng. (Accessed 15 September 2021).
- Cresswell, W., 2014. Migratory connectivity of Palaearctic–African migratory birds and their responses to environmental change: the serial residency hypothesis. *Ibis* 156, 493–510. <https://doi.org/10.1111/ibi.12168>.
- Duijns, S., van Gils, J.A., Spaans, B., ten Horn, J., Brugge, M., Piersma, T., 2014. Sex-specific winter distribution in a sexually dimorphic shorebird is explained by resource partitioning. *Ecol. Evol.* 4, 4009–4018. <https://doi.org/10.1002/ece3.1213>.
- Estrella, S.M., Masero, J.A., 2010. Prey and prey size selection by the near-threatened Black-Tailed Godwit foraging in non-tidal areas during migration. *Waterbirds* 33, 293–299. <https://doi.org/10.1675/063.033.0304>.
- Giunchi, D., Baldaccini, N.E., Lenzone, A., Luschi, P., Sorrenti, M., Cerritelli, G., et al., 2019. Spring migratory routes and stopover duration of satellite-tracked Eurasian Teals *Anas crecca* wintering in Italy. *Ibis* 161, 117–130. <https://doi.org/10.1111/ibi.12602>.
- Hijmans, R.J., Williams, E., Vennes, C., 2017. Geosphere: Spherical Trigonometry, Version 1.5-10. <https://CRAN.R-project.org/package=geosphere>. (Accessed 12 January 2021).
- Hill, D., 1988. Population dynamics of the avocet (*Recurvirostra avosetta*) breeding in Britain. *J. Anim. Ecol.* 57, 669–683. <https://doi.org/10.2307/4932>.
- Hotker, H., West, R., 2005. Population size, population development and habitat use by Avocets in Western Europe at the end of the 20th century. *Wader Study Group Bull.* 107, 57.
- Inger, R., Gregory, R., Duffy, J.P., Stott, I., Voříšek, P., Gaston, K.J., 2015. Common European birds are declining rapidly while less abundant species' numbers are rising. *Ecol. Lett.* 18, 28–36. <https://doi.org/10.1111/ele.12387>.
- IUCN, 2020. *Recurvirostra avosetta* (pied avocet). IUCN red list threat. Species. <https://www.iucnredlist.org/species/22693712/155534228>. (Accessed 30 November 2020).
- Kirby, J.S., Stattersfield, A.J., Butchart, S.H.M., Evans, M.I., Grimmett, R.F.A., Jones, V.R., et al., 2008. Key conservation issues for migratory land- and waterbird species on the world's major flyways. *Bird. Conserv. Int.* 18, S49–S73. <https://doi.org/10.1017/S0959270908000439>.
- Kuang, F., Coleman, J.T., Hassell, C.J., Leung, K.K.-S., Maglio, G., Ke, W., et al., 2020. Seasonal and population differences in migration of Whimbrels in the East Asian–Australasian flyway. *Avian Res.* 11, 24. <https://doi.org/10.1186/s40657-020-00210-z>.
- Kuznetsova, A., Brockhoff, P.B., Christensen, R.H.B., 2017. lmerTest package: tests in Linear mixed effects models. *J. Stat. Software* 82, 1–26. <https://doi.org/10.18637/jss.v082.i13>.
- Lei, W., Masero, J.A., Piersma, T., Zhu, B., Yang, H.-Y., Zhang, Z., 2018. Alternative habitat: the importance of the Nanpu saltpans for migratory waterbirds in the Chinese Yellow Sea. *Bird. Conserv. Int.* 28, 549–566. <https://doi.org/10.1017/S0959270917000508>.
- Lei, W., Wu, Y., Wu, F., Piersma, T., Zhang, Z., Masero, J.A., 2021. Artificial wetlands as breeding habitats for shorebirds: a case study on Pied Avocets in China's largest saltpan complex. *Front. Ecol. Evol.* 9, 622756. <https://doi.org/10.3389/fevo.2021.622756>.
- Lisovski, S., Gosbell, K., Hassell, C., Minton, C., 2016. Tracking the full annual-cycle of the great Knot, *Calidris tenuirostris*, a long-distance migratory shorebird of the East Asian–Australasian flyway. *Wader Study* 123, 177–189. <https://doi.org/10.18194/ws.00048>.
- Lourenço, P.M., Kentie, R., Schroeder, J., Alves, J.A., Groen, N.M., Hooijmeijer, J.C.E.W., et al., 2010. Phenology, stopover dynamics and population size of migrating Black-Tailed Godwits *Limosa limosa limosa* in Portuguese Rice Plantations. *Ardea* 98, 35–42. <https://doi.org/10.5253/078.098.0105>.
- Ma, Z., Hua, N., Peng, H., Choi, C., Battley, P.F., Zhou, Q., et al., 2013. Differentiating between stopover and staging sites: functions of the southern and northern Yellow Sea for long-distance migratory shorebirds. *J. Avian Biol.* 44, 504–512. <https://doi.org/10.1111/j.1600-048X.2013.00213.x>.
- McMahon, B.J., Whelan, J., Bracken, F., Kavanagh, B.P., 2003. The impact of farming on over-wintering bird populations. *Tearmann – Irish J. Agri-Environ. Res.* 3, 67–76. <https://hdl.handle.net/10779/rcsi.10767518.v2>.
- Meissner, W., 2006. Timing and phenology of Curlew Sandpiper *Calidris ferruginea* on southward migration through Puck Bay, Poland. *Int. Wader Stud.* 19, 121–124.
- Moreira, F., 1995. The winter feeding ecology of Avocets *Recurvirostra avosetta* on intertidal areas. I. Feeding strategies. *Ibis* 137, 92–98. <https://doi.org/10.1111/j.1474-919X.1995.tb03224.x>.
- Murray, N.J., Clemens, R.S., Phinn, S.R., Possingham, H.P., Fuller, R.A., 2014. Tracking the rapid loss of tidal wetlands in the Yellow Sea. *Front. Ecol. Environ.* 12, 267–272. <https://doi.org/10.1890/130260>.
- Nebel, S., 2005. Latitudinal clines in bill length and sex ratio in a migratory shorebird: a case of resource partitioning? *Acta Oecol.* 28, 33–38. <https://doi.org/10.1016/j.actao.2005.02.002>.
- Pierce, R.J., Kirwan, G.M., Boesman, P.F.D., 2020. Pied Avocet (*Recurvirostra avosetta*), Version 1.0. In: del Hoyo, J., Elliott, A., Sargatal, J., Christie, D.A., de Juana, E. (Eds.), *Birds of the World*. Cornell Lab of Ornithology, New York. <https://doi.org/10.2173/bow.pieavo1.01>.
- Piersma, T., 2003. “Coastal” versus “inland” shorebird species: interlinked fundamental dichotomies between their life- and demographic histories? *Wader Study Group Bull.* 100, 5–9.
- Piersma, T., Kok, E.M.A., Hassell, C.J., Peng, H.-B., Verkuil, Y.I., Lei, G., et al., 2021. When a typical jumper skips: itineraries and staging habitats used by Red Knots (*Calidris canutus piersmai*) migrating between Northwest Australia and the New Siberian Islands. *Ibis* 163, 1235–1251. <https://doi.org/10.1111/ibi.12964>.
- Piersma, T., Lok, T., Chen, Y., Hassell, C.J., Yang, H.-Y., Boyle, A., et al., 2016. Simultaneous declines in summer survival of three shorebird species signals a flyway at risk. *J. Appl. Ecol.* 53, 479–490. <https://doi.org/10.1111/1365-2664.12582>.
- R Core Team, 2020. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna.
- Spoon-billed Sandpiper in China, 2022. Jiangsu Coastal Waterbird Census Report 2019–2020. https://ff.lingxi360.com/?fid=BDI=dmYCPWcc6&utm_bccid=LXEQrkjWjUzjyCR. (Accessed 15 September 2022).
- Studds, C.E., Kendall, B.E., Murray, N.J., Wilson, H.B., Rogers, D.I., Clemens, R.S., et al., 2017. Rapid population decline in migratory shorebirds relying on Yellow Sea tidal mudflats as stopover sites. *Nat. Commun.* 8, 14895. <https://doi.org/10.1038/ncomms14895>.
- Tao, X.D., Lei, J.Y., Richard, H., Lei, G., 2017. Report on the Coordinated Survey for Wintering Waterbird of the Central and Lower Yangtze in 2015. China Forestry Publishing House, Beijing.
- Van der Velde, M., Haddrath, O., Verkuil, Y.I., Baker, A.J., Piersma, T., 2017. New primers for molecular sex identification of waders. *Wader Study* 124, 147–151. <https://doi.org/10.18194/ws.00069>.
- Verhoeven, M.A., Loonstra, A.J., McBride, A.D., Macias, P., Kaspersma, W., Hooijmeijer, J.C., et al., 2020. Geolocators lead to better measures of timing and re-nesting in black-tailed godwits and reveal the bias of traditional observational methods. *J. Avian Biol.* 51, e02259. <https://doi.org/10.1111/jav.02259>.
- Wetlands International, 2020. Waterbird Population Estimates. <http://wpe.wetlands.org/>. (Accessed 2 December 2020).
- Woodworth, B.K., Wheelwright, N.T., Newman, A.E., Schaub, M., Norris, D.R., 2017. Winter temperatures limit population growth rate of a migratory songbird. *Nat. Commun.* 8, 14812. <https://doi.org/10.1038/ncomms14812>.
- Yang, Z., Li, J., Han, Y., Hassell, C.J., Leung, K.K.-S., Melville, D.S., et al., 2021. Coastal wetlands in Lianyungang, Jiangsu Province, China: probably the most important site globally for the Asian Dowitcher (*Limnodromus semipalmatus*). *Avian Res.* 12, 38. <https://doi.org/10.1186/s40657-021-00272-7>.
- Zhu, B., Verhoeven, M.A., Loonstra, A.H.J., Sanchez-Aguilar, L., Hassell, C.J., Leung, K.K.-S., et al., 2021. Identification of breeding grounds and annual routines of the newly discovered *bohaili* subspecies of Black-tailed Godwits. *Emu* 121, 292–302. <https://doi.org/10.1080/01584197.2021.1963287>.