

University of Groningen

Trade in vulture parts in West Africa

Daboné, Clément; Ouéda, Adama; Thompson, Lindy J.; Adjakpa, Jacques Boco; Weesie, Peter D.M.

Published in:
 Bird Conservation International

DOI:
[10.1017/S095927092100054X](https://doi.org/10.1017/S095927092100054X)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2023

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Daboné, C., Ouéda, A., Thompson, L. J., Adjakpa, J. B., & Weesie, P. D. M. (2023). Trade in vulture parts in West Africa: Burkina Faso may be one of the main sources of vulture carcasses. *Bird Conservation International*, 33, Article e8. <https://doi.org/10.1017/S095927092100054X>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Research Article

Cite this article: Daboné C, Ouéda A, Thompson LJ, Adjakpa JB, Weesie PDM (2023). Trade in vulture parts in West Africa: Burkina Faso may be one of the main sources of vulture carcasses. *Bird Conservation International*, **33**, e8, 1–11
<https://doi.org/10.1017/S095927092100054X>

Received: 14 August 2021
Revised: 17 November 2021
Accepted: 30 November 2021


Keywords:

Body part trade; Mass mortality; Poison; Vultures; West Africa

Author for correspondence:

*Clément Daboné,
Email: dabexpi@yahoo.fr

Trade in vulture parts in West Africa: Burkina Faso may be one of the main sources of vulture carcasses

Clément Daboné^{1,2*} , Adama Ouéda², Lindy J. Thompson^{3,4},
Jacques Boco Adjakpa⁵ and Peter DM Weesie⁶

¹University Centre of Tenkodogo, University Thomas Sankara, Tenkodogo, Burkina Faso; ²Laboratory of Animal Biology and Ecology, University Joseph Ki-Zerbo, Ouagadougou, Burkina Faso; ³Endangered Wildlife Trust, Midrand, 1685, South Africa; ⁴Centre for Functional Biodiversity, School of Life Sciences, University of KwaZulu-Natal, Private Bag X01, Scottsville, 3201, South Africa; ⁵Laboratory of Research in Applied Biology, University of Abomey-Calavi, Abomey-Calavi, Benin and ⁶Integrated Research on Energy, Environment and Society, Faculty of Sciences and Engineering, University of Groningen, Groningen, The Netherlands

Summary

The current catastrophic decline in African vultures is caused mainly by poisoning, and killing for consumption and belief-based use (traditional medicine). To find out more about the key threats to vultures in West Africa, we assessed the main anthropogenic causes of vulture mortality in Burkina Faso. We analysed incidents of mass vulture mortality, based on interviews conducted in 2016 with local butchers, veterinarians, foresters, and abattoir watchmen at 44 sites across the country. A total of 730 interviews revealed that poisoning was the main cause of mortality of vultures in Burkina Faso (20 of the 23 mass mortality incidents described by respondents were caused by poisoning). Poisoning was also the most lethal threat to vultures (779 out of 879 known vulture deaths were due to poisoning). According to the survey, intentional poisoning of vultures with poisoned baits (which comprised 15 out of 23 mass mortality incidents) produced the highest number of victims (577 out of 879 dead vultures). The number of vultures killed by poisoned baits was higher closer to the borders than elsewhere in Burkina Faso and we believe these recent intentional vulture poisoning events in Burkina Faso were intended to meet the growing demand for vulture body parts in West Africa. The survey showed that unintentional poisoning was the second main cause of vulture mass mortality (5 out of 23 mass mortality incidents comprising 202 of 879 dead vultures). Other important anthropogenic causes of vulture mortality included electrocution at electricity poles and motor vehicle collisions. Our results highlight the need for awareness campaigns, improved policy and legislation, and stronger commitment from governments in West Africa, to halt the trade in vultures and prevent their extirpation.

Introduction

Today, vulture populations around the world are in serious danger (Safford *et al.* 2019). The illegal use of poison continues to be the single most important threat to vultures throughout their ranges (Margalida 2012, Pantović and Andevski 2018, Pauli *et al.* 2018, Alarcón and Lambertucci 2018), especially in Africa (Ogada *et al.* 2016a). In Europe, this practice appears to be common in some countries, where it is mostly associated with human-wildlife conflicts involving predators (Berny 2007, Margalida *et al.* 2008, Guitart *et al.* 2010, Pantović and Andevski 2018). In Asia, several studies highlighted the decline of different vulture species suggesting that poisoning is the main cause (MaMing and Xu 2015, Kim *et al.* 2016, Loveridge *et al.* 2019). In both North and South America too, vultures have been killed in various poisoning incidents (Mineau *et al.* 1999, Fleischli *et al.* 2004, Pavez and Estades 2016).

In Africa, over the last 30 years, populations of eight vulture species have decreased by an average of 62% (Ogada *et al.* 2016a, Garbett *et al.* 2018). Well-known threats to vultures in Africa include unintentional and intentional poisoning, and harvesting for belief-based use (also called traditional medicine) and consumption (Ogada *et al.* 2016a, Botha *et al.* 2017, Margalida *et al.* 2019, Henriques *et al.* 2020). Additional threats to vultures include habitat degradation, motor vehicle collisions, collisions with (or electrocution on) energy infrastructure, and decreasing food availability (Ogada and Buij 2011, Ogada *et al.* 2012, 2016a). However, exposure to poison remains the main threat for these birds in Africa because it is occurring with increasing frequency and producing high mortality rates (Ogada *et al.* 2016b, Margalida *et al.* 2019, Henriques *et al.* 2020). There is a growing concern that poison and the other threats to vultures in Africa are widespread and difficult to tackle, and the scientific data needed to address these threats remain

largely lacking. Also, the vulnerability of vulture populations to each threat, and the extent of the impact of these threats on vulture populations, are not clearly defined for each country (Botha *et al.* 2017).

Unintentional poisoning of vultures often occurs in connection with farmers whose livestock are predated by carnivores, or whose crops are frequently raided by elephants, buffalo and other large herbivores (Ogada *et al.* 2012). When farmers poison these animals to mitigate conflict, the poisoned carcasses are consumed by vultures, which then become unintentional victims of poisoning. Vultures are also intentionally poisoned by poachers due to the role they play as sentinels of the presence of dead animals, alerting park rangers to illegal poaching activities (Ogada *et al.* 2016a, Margalida *et al.* 2019). More recently, poisoned baits are being used to harvest vultures for the trade in vulture body parts for belief-based use (Buij *et al.* 2016, Daboné *et al.* 2019, Henriques *et al.* 2020). Indeed, although a variety of techniques are used to kill vultures and other raptors for these purposes (Buij and Croes 2014), the use of poisoned baits is probably the most widespread method for acquiring vultures (Ogada *et al.* 2016b), often killing many vultures at a single poisoning event. In a recent case, between September 2019 and March 2020, more than 2000 Hooded Vultures *Necrosyrtes monachus* were intentionally poisoned across eastern Guinea-Bissau for belief-based use (Henriques *et al.* 2020).

Understanding threats to vultures is critical for developing effective conservation strategies, and there is a clear need to increase knowledge on these poisoning events in West Africa. In this paper, we aim to explore the extent to which poisoned baits are being used to acquire vultures for belief-based use, by using surveys to assess the anthropogenic causes of vulture mortality throughout Burkina Faso. Ultimately, we aim to highlight the drivers of vulture declines in Burkina Faso and elsewhere in West Africa. Given that toxic products such as pesticides are commonly used to poison vultures (Ogada 2014) and given their rapid effect, high efficacy (a large number of vultures may be killed in a single incident) (Richards *et al.* 2017, Margalida *et al.* 2019, Henriques *et al.* 2020) and their availability in this area, we expected that

poisoned baits would be the method most widely used to acquire vultures for belief-based use. Vultures have become rare in Burkina Faso's neighbouring countries, and there are high levels of demand for vulture body parts in countries such as Nigeria and Benin (Buij *et al.* 2016). So, given that there are cross-border relationships between communities in these countries, we expected the intentional poisoning of vultures in Burkina Faso to occur more frequently in areas closer to the borders, where large densities of vultures still occur.

Methods

Study area

Burkina Faso is a Sahelian, land-locked country in the centre of West Africa between 9°20' and 15°3'N, and 2°20'E and 5°3'W (Figure 1). The climate is Sudano-Sahelian in which two seasons alternate: a long dry season from November to May and a short rainy season from June to October (MECV 2007, Ibrahim *et al.* 2014). Land use includes farmland, open savanna, urban areas, and protected and wildlife management areas in which six vulture species occur. These are the Hooded Vulture, White-backed Vulture *Gyps africanus*, White-headed Vulture *Trigonoceps occipitalis*, Rüppell's Vulture *Gyps rueppelli*, Lappet-faced Vulture *Torgos tracheliotos* and Egyptian Vulture *Neophron percnopterus* (Weesie and Belemsobgo 1997, Borrow and Demey 2001, Thiollay 2006a, Portier 2007). Of these, the Hooded Vulture is the most closely associated with human habitations (Mundy *et al.* 1992), and it is very commensal with humans (Thiollay 2006b, 2007b, Henriques *et al.* 2017). The other vulture species are confined to protected areas (Mundy *et al.* 1992, Thiollay 2007a, Ogada and Buij 2011). We assumed that mass mortalities of vultures would be more common in areas with high vulture densities, and so we focused our survey on localities cited in the literature (Thiollay 2006a, 2007b), and on areas known to have high numbers of vultures, based on our own experience. We conducted our survey in 31 towns and 13 townships spread across the country (Figure 1).




Figure 1. The locations of our 44 survey sites, and the main protected areas. Only the names of the main towns are given here, as not all surveyed site names appear on the map.

Data collection

We used a mixture of qualitative and quantitative methods to gather and evaluate data through both structured and unstructured interviews conducted from 3 January 2016 to 30 December 2016 (Kabir 2016). In each locality, interviewees were chosen according to a nonprobability (purposeful) sampling method (Patton 2002). This involved identifying and selecting individuals (or groups of individuals) who are especially knowledgeable about or experienced with a phenomenon of interest (Cresswell and Plano Clark 2011). In our study area, vultures (especially Hooded Vultures) were historically, and still are, very close to butchers, veterinarians, foresters, and abattoir watchmen (Mundy *et al.* 1992, Weesie and Belemsobgo 1997, Thiollay 2006b). Vultures are legally protected by local people because they provide an important public service (Daboné *et al.* 2019) in terms of consuming scraps of meat at markets and abattoirs, as well as carcasses of domestic animals and wildlife. In Burkina Faso, foresters (and to some extent veterinarians) are assigned to control and safeguard protected species including vultures. The people interviewed in this study (butchers, veterinarians, foresters, and abattoir watchmen) were those whom we considered would provide informative, high-quality responses to our survey questions. They are key players for vulture conservation in our study area and they frequently witness (or are alerted of the occurrence of) mass vulture mortalities. A typical slaughterhouse workday in Burkina Faso involves, among other activities, the slaughter of animals (ovine, caprine, bovine, porcine etc.) from 05h00 to 07h00 and veterinary control, to protect public health, from 07h00 to 11h00. More than 30 butchers and at least one veterinarian and slaughterhouse watchman remain actively present on slaughterhouse sites every day. We aimed to survey 15 butchers and at least one slaughterhouse watchman, one veterinarian, and one forester over two or three successive days in each locality. Throughout the survey, we considered parameters suggested by Spradley (1979) and Bernard (2002), such as availability, willingness to participate, and the ability to communicate experiences and opinions, when identifying and selecting interviewees. Informed consent was obtained from all of the interviewees involved in the study. Moreover, once in a locality, if we learned of any incidents of vulture mass mortality that had been reported to the court or to the police, then we followed up on these events and went to the relevant police station and/or the registry office of courts where the sentence was passed. In this survey, we use the terms “incidents of mass mortalities of vultures” to refer to any event in which at least five vultures were killed. All surveys were carried out by the first author.

Structured interviews

Structured interviews, consisting of closed and open-ended questions, were conducted during slaughterhouse visits and face-to-face with the interviewees, i.e. butchers, abattoir watchmen and veterinarians, between 08h00 and 11h00 (Appendix S1 in the online supplementary material). Each interview took about 10 to 15 minutes to complete. Interviews with foresters took place in the afternoon, in their offices, where we would survey one forester (the Head of Service) representing the service. Forester surveys took between 10 and 45 minutes to complete.

Unstructured interviews

Unstructured interviews were completed with the registrar of court or the police officer who investigated each incident of

vulture mass mortality. Police minutes and court records were then requested for consultation and reference purpose. In accordance with legal provisions in Burkina Faso, these documents can be available for consultation if requested. The interview process was such that it assured interviewees anonymity and encouraged proactive participation. We began by assessing whether interviewees were aware of the incidents of vulture mass mortality referred to. We then asked interviewees to describe the circumstances under which those incidents occurred (that is, the method used, motivation, culprit, etc.).

Monitoring of incidents of vulture mass mortality

Vulture poisoning events were assessed and approved by foresters or veterinary staff by i) checking for the presence of dead vultures close to poisoned meat or carcasses, ii) checking for poison residues and the poison container or packaging, and iii) recording testimonies of people who witnessed the culprits of intentional poisonings of vultures. Dead vultures were identified by foresters and, if needed, birds were photographed to facilitate further species identification. Poisoned meat or carcasses were destroyed by burning them to prevent further vulture mortality and in some cases, vulture carcasses were sent to Ouagadougou (the capital of Burkina Faso) for further analyses to determine which toxic chemicals were used. Regrettably, according to foresters and veterinarians, these analyses were not followed up. We sent requests for information to the service referred to, but did not receive an answer.

Data analysis

We conducted all data analyses using the SPSS statistical package (IBM SPSS Statistics for Windows, Version 27.0). We used a bivariate analysis (Pearson correlation) to assess the correlation between the number of vultures killed at poisoned baits in each mass mortality incident and the distance to the nearest international border. We also assessed (using a Pearson correlation) the relationship between the abundance of vultures at each abattoir and the distance to the nearest international border, and showed that vultures were not any more abundant in localities in border areas than elsewhere in Burkina Faso.

Results

Demographics

In the 44 localities surveyed, we completed a total of 730 interviews with 660 butchers, 44 abattoir watchmen, 14 veterinarians and 12 foresters. All interviewees were men, from 17 to 82 years old and most (93%) were over 35 years old (Table 1). This skew towards men may be because in an effort to select individuals or groups of individuals with extensive knowledge or experienced with the topics on which our survey questions were focused, we targeted professions (butchers, veterinarians, foresters and abattoir watchmen) in which women are currently very rarely hired in Burkina Faso. Most interviewees (88%) had 5–25 years of experience in their professions (Table 1).

Population trends of vultures

According to 98% of the interviewees, the Hooded Vulture is the vulture species most closely associated with human habitations. The other vulture species are reportedly confined to protected

Table 1. Demographics of interviewees who took part in the structured interviews

		Number of interviewees	Percentage (%)
Profession	Butchers	660	90,41%
	Abattoir watchmen	44	6,02%
	Veterinarians	14	1,91%
	Foresters	12	1,64%
Gender	Male	730	100%
	Female	0	0%
Age	36–82	683	93,56%
	21–35	36	4,93%
	17–20	11	1,50%
Experience (years)	5–10	35	4,79%
	11–15	125	17,12%
	16–20	134	18,35%
	21–25	346	47,39%
	>25	90	12,32%
Ethnic group	Mossi	373	51,09%
	Fulani	95	13,01%
	Bissa	55	7,50%
	Gourmantché	33	4,52%
	10 other groups (comprised less than 4%)	174	23,83%

areas. Ninety percent of respondents noted a large decline in numbers of vultures in their localities. According to 70% of interviewees, this large decline is due to poaching for belief-based use. Most (60%) of the respondents reported that the recent increase in the occurrence of Hooded Vulture carcasses in their localities (Table 2) is the result of the use of poisoned baits to obtain vultures for belief-based use.

Mass mortalities of vultures

From 1 April 2010 to 12 December 2016, 23 incidents of anthropogenic mass mortalities of vultures were recorded in 15 localities in Burkina Faso, resulting in the deaths of 879 vultures, of which 317 (36%) were beheaded (Table 3). Of these 23 incidents, 22 exclusively involved Hooded Vultures, while the remaining incident resulted in the deaths of individuals from four vulture species (18 Hooded Vultures, 15 White-backed Vultures, six Rüppell's Vultures and three Lappet-faced Vultures; Table 3). Of the 23 anthropogenic mass mortalities of vultures, 20 incidents were poisoning events. Of these 20 poisoning events, 15 were intentional, involving the use of poisoned baits (resulting in the deaths of 577 vultures, or nearly 66%), and five were unintentional (202 vultures, almost 23%). In one incident, 10 vultures (just over 1%) were killed using a slingshot, and in another incident, five vultures (less than 1%) were electrocuted on a high-voltage power line. In the remaining incident, where 85 vultures (almost 10%) were killed, the cause of mortality was not determined. Our results suggest that poisoning is the main cause (89% of vulture deaths) of anthropogenic mass mortality of

vultures in Burkina Faso. So, it seems that in Burkina Faso, poisoned baits are the most widely used method of killing vultures, with the most devastating effect on vulture populations.

Intentional poisoning events

Most (70%) of our interviewees highlighted poisoned baits as the main cause of anthropogenic mass mortality of vultures (Table 2). In these deliberate poisoning events, poisoned baits were reportedly used by poachers to obtain vulture carcasses. Some of these vulture carcasses were found to be missing body parts (always the head; Table 3).

Culprits in intentional vulture poisonings

More than 75% of our survey respondents witnessed or heard about intentional poisoning perpetrated by local people. Only 19% of respondents suggested that perpetrators may be foreign nationals (Table 2). These results are corroborated by four arrests where local communities were found to be complicit in poisoning activities (Table 4). The investigations following these arrests found that the local people involved in poisoning events were incited by foreign nationals, themselves stimulated by the decreasing availability of vultures in their own countries and the increasing profitability of the regional vulture trade. For example, in November 2012, a 32-year-old local man with three Hooded Vulture carcasses was arrested near the abattoir of Bitou, a township ~39 km from Togo and ~10 km from Ghana. In 2011, 50 beheaded Hooded Vultures were found at the same abattoir (Table 3). Similarly, in December 2011, in Bogandé town in eastern Burkina Faso, ~100 km from Niger, two Burkinabe and a Nigerian woman were arrested with 71 dead Hooded Vultures intended for export to Nigeria (Table 3). This evidence emanating from arrests leads us to suspect that during the last decade, vultures were deliberately poisoned throughout Burkina Faso for export to neighbouring countries. While the number of vultures at abattoirs was not higher closer to international borders (Pearson's correlation test: $r = 0.44$, $P = 0.107$, $n = 10$), the number of vultures killed by poisoned baits showed a significant linear negative correlation with distance to the nearest border (Pearson's correlation test: $r = -0.77$, $P = 0.003$, $n = 10$). This suggests that the number of vultures killed by poisoned baits was higher closer to the borders than elsewhere in Burkina Faso.

Drivers of intentional poisoning

There seem to be two main drivers for poisoning by poachers and their accomplices. As a result of the experience gained in their close relationship with vultures, 79% of our survey respondents stated that poachers and their accomplices harvest vulture body parts for their own use, to treat a range of physical and mental diseases, or to bring good luck during gambling, competitions and contests. However, 20% of respondents suggested that vultures are poached to sell, due to the increasing profitability of the regional trade in vultures for belief-based use (Table 2). Investigations into the four incidents of intentional poisoning in which culprits were arrested and sentenced (Table 4) showed that 90% of the Hooded Vultures killed by poisoned baits (128 out of 143, Figure 2) were destined for trade in local markets (which includes trade to foreign nationals who come to Burkina Faso to buy vulture parts). When interrogating arrested culprits, it emerged that the unit cost of a whole vulture was about 5,000 to 7,500 CFA francs (~ USD 10–15). That would have yielded about 540,000 CFA (~ USD 1,080) to two poachers arrested in one of the intentional poisoning events. These investigations further

Table 2. Responses to questions addressed to the 730 interviewees who participated in the structured interviews. (“n” is the number of respondents, and “%” is the number of respondents as a percentage of the total).

		n	%
Do you know these birds? (photos of each species were shown)	Hooded Vulture	730	100
	White-backed Vulture	40	5,47
	White-headed Vulture	30	4,10
	Rüppell's Vulture	1	0,13
	Lappet-faced Vulture	0	0
	Egyptian Vulture	0	0
Have you ever seen them around here?	Hooded Vulture	714	97,80
	White-backed Vulture	10	1,36
	White-headed Vulture	6	0,82
	Rüppell's Vulture	0	0
	Lappet-faced Vulture	0	0
	Egyptian Vulture	0	0
Do you consider vultures as a constraint or benefit for you / the community?	Benefit	593	81,23
	Constraint	75	10,27
	Do not know	62	8,49
Do you agree with that statement 'Vultures are declining'?	Yes	657	90
	No	23	3,15
	Do not know	50	6,84
Can you give the reasons for the decline?	Natural causes	7	0,95
	Poaching/intentional poisoning	511	70
	Unintentional poisoning	74	10,13
	Electrocution	44	6,02
	Diseases	12	1,64
	Motor vehicle collisions	6	0,82
	Others:	8	1,09
	Do not know	68	9,31
Who are the perpetrators of this poaching/intentional poisoning?	Local people	384	75,14
	Nationals from Nigeria	52	10,17
	Nationals from Benin	33	6,45
	Nationals from Ghana	13	2,54
	Do not know	27	5,28
The uses of vultures poached/intentionally poisoned	Intended for use by the poachers themselves	404	79,06
	Intended to be traded	103	20,15
	Do not know	4	0,78
Do you sometime encounter vulture carcasses?	Yes	439	60,13
	No	291	39,86
Have you ever witnessed a "vulture mass mortality event" in your locality or elsewhere?	Yes	322	44,10
	No	408	55,89
Have you ever heard about mass mortalities of vultures in your locality or elsewhere?	Yes	423	57,94
	No	307	42,05

*In this survey, we use the terms "incidents of mass mortalities of vultures" to refer to any unfortunate event in which there are at least five vultures killed.

Table 3. Mass mortalities of vultures recorded in Burkina Faso from 2010 to 2016. In circumstances where the exact day of the incidents was not available, the first day of the month was considered.

Date	Location	Number of dead vultures					Total	Cause	V.P.A.	Ref*
		HV	WBV	RV	LFV					
06-Jun-2012	Bitou	26	0	0	0	26	intentional poisoning	–	a	
1-Aug-2013	Bitou	20	0	0	0	20	unintentional poisoning	–	a	
8-Nov-2012	Bitou	31	0	0	0	31	intentional poisoning	head	a	
1-Mar-2011	Bitou	50	0	0	0	50	intentional poisoning	head	b	
1-Dec-2011	Bogandé	71	0	0	0	71	intentional poisoning	head	c	
1-Sep-2010	Boromo	10	0	0	0	10	hunting	–	a	
1-Sep-2015	Diabo	16	0	0	0	16	unintentional poisoning	–	a	
1-Feb-2015	Dori	6	0	0	0	6	unintentional poisoning	–	a	
1-Sep-2012	Dori	70	0	0	0	70	unintentional poisoning	–	a	
1-Apr-2011	Dori	30	0	0	0	30	intentional poisoning	head	a	
13-Apr-2012	Gourgou	15	0	0	0	15	intentional poisoning	head	a	
6-Mar-2015	Kantchari	26	0	0	0	26	intentional poisoning	–	a	
4-Aug-2015	Kantchari	20	0	0	0	20	intentional poisoning	–	a	
19-Mar-2016	Kantchari	15	0	0	0	15	intentional poisoning	–	a	
1-May-2011	Kelbo	110	0	0	0	110	intentional poisoning	–	a	
1-Apr-2012	Kompianga	90	0	0	0	90	unintentional poisoning	–	a	
06-Apr-2012	Kouba	26	0	0	0	26	intentional poisoning	–	d	
13-Sep-2012	Ouagadougou	5	0	0	0	5	electrocution	head	a	
12-Dec-2016	Parc Arly-Pendjari	18	15	6	3	42	intentional poisoning	–	a	
9-Dec-2012	Tenkodogo	25	0	0	0	25	intentional poisoning	head	a	
1-Mar-2011	Tenkodogo	30	0	0	0	30	intentional poisoning	head	a	
1-Apr-2010	Tougouri	60	0	0	0	60	intentional poisoning	head	a	
1-Mar-2015	Zabré	85	0	0	0	85	not determined	–	a	

Ref = Reference, where a = *This study*; b = AIB (2012); c = Nabaloum (2012a); d = Nabaloum (2012b). HV = Hooded Vulture; WBV = White-backed Vulture; RV = Rüppell's Vulture; LFV = Lappet-faced Vulture. V.P.A. = Vulture's body parts amputated.

Table 4. Intentional poisoning events in which culprits were arrested. ("n" = number of Hooded Vulture killed, ND = not determined).

Date	locality	n	Public officer alerted	in charge of the investigation	Court	Culprit identity	Use of dead vultures
08-nov-2012	Bitou	31	forester	Bitou police station	Court of Tenkodogo	Local people	ND
11-Dec-2011	Bogandé	71	forester	Bogandé police station	Court of Ouagadougou	Local people and Nigerian	Exported to Nigeria
06-Apr-2012	Kouba	26	forester	Koubri police station	Court of Ouagadougou	Local people	Exported to Nigeria
13-Apr-2012	Gourgou	15	forester and veterinarian	Tenkodogo police station	Court of Tenkodogo	Local people	ND

revealed that in some West African countries such as Nigeria, the unit cost of a whole Hooded Vulture ranged from 50,000 to 75,000 CFA (~ USD 100–150).

Poisoned baits: Poachers' operating method

Testimonies of people who witnessed the intentional poisoning of vultures, and the confessions of the arrested culprits stated that

poachers used highly toxic pesticides (or local traditional tobacco) to poison vultures. They attracted the birds using meat which was sprinkled with poison or local traditional tobacco, and left on abattoir roofs or on the ground where vultures gather to feed. After consuming this poisoned bait, vultures are intoxicated and unable to fly. Poachers either harvest the whole bird or take only the head. This practice is reportedly done clandestinely at abattoirs, after butchers, veterinarians and abattoir watchmen have left.

Unintentional poisoning of vultures

Almost 10% of our survey respondents reported unintentional poisoning as one of the main causes of anthropogenic mass mortality of vultures (Table 2). They reported that it was very common to find the carcasses of poisoned vultures close to the carcasses of poisoned animals which were killed to 'protect' livestock and crops. For example, in Kompianga in April 2012, a farmer used Furadan (a very toxic unregistered carbamate pesticide) to kill 12 crop-raiding pigs, which resulted in the unintentional poisoning of 90 Hooded Vultures (Table 3) as well as four dogs, six cats, and 10 Pied Crows *Corvus albus*.

Other causes of mass mortality of vultures

Electrocution (6% of respondents), motor vehicle collisions (less than 1% of respondents), and disease (more than 1% of respondents) were the other causes of mass mortality of vultures reported by interviewees (Table 2). According to our survey respondents, Hooded Vultures are often electrocuted by high-voltage power lines, especially during the rainy season. In September 2013, five vultures were electrocuted close to the abattoir of Ouagadougou (Table 3). Motor vehicle collisions occur when vultures that are feeding on the carcasses of road-killed animals are hit unintentionally by vehicles.

Discussion

Intentional poisoning of vultures in Burkina Faso for belief-based use

We found that the number of vultures killed by mass-poisoning in Burkina Faso increased with proximity to the country's borders, suggesting that the recent recurrence of intentional vulture poisoning events in Burkina Faso was intended to meet the growing demand for vulture body parts in West Africa. These poisoning events were likely driven by the decreasing availability of vultures in other West African countries (Thiollay 2007b, Mullié *et al.* 2017, Nosazeogie *et al.* 2018) and by the lucrative regional trade in vulture body parts for belief-based use (Buij *et al.* 2016). Thus, Burkina Faso appears to be one of the preferred areas for harvesting whole vultures for illegal trade in West Africa. The results of this study and the relatively high demand for vulture body parts in Nigeria and Benin (Nikolaus 2011, Buij *et al.* 2016), show that these two countries are important links in the trade network for vulture body parts in West Africa. Previous studies have established that vulture parts, their eggs and nest materials are sourced from an area spanning Burkina Faso to Chad to supply the demand in Nigeria and Benin (Mander *et al.* 2007, Nikolaus 2011, Buij *et al.* 2016, Craig *et al.* 2018).

The economic value of the trade in vulture body parts in West Africa

Some of the vulture carcasses found at poisoning scenes in Burkina Faso were missing heads, which is the body part most used by traditional healers, and which carries a higher price per unit than any other body part (McKean *et al.* 2013, Awoyemi 2014). This suggests that the trade in vulture body parts for belief-based use is driving vulture poisoning in Burkina Faso. In this study, the local value of a vulture harvested in Burkina Faso was 5,000 to 7,500 CFA

francs (~ USD 10-15), but vultures could be sold for 6-10 times more than that in Nigeria. We estimate that an inculcated Nigerian woman would have earned between 3,350,000 and 5,325,000 CFA francs (~ USD 6,700-10,650) for the vultures that she harvested once in Nigeria (Nabaloum 2012a). The high profits yielded by the trade in vulture body parts might make this threat difficult to tackle given the relatively impoverished conditions of the people in this region.

The urgent need to increase enforcement of legislation in this regard

To reduce and eventually to halt the intentional poisoning of vultures for trade for belief-based use, we recommend and encourage a review of current legislation and enforcement actions that could lead to the successful prosecution of cases involving poisoned vultures. In Burkina Faso, all vulture species have been protected since 1960 by several legislative decrees of which the most recent is the N° 1996-061-PRES/PM/MEE/MATS/MEFP/MCIA/MTT of 11 March 1996 concerning the regulation of the exploitation of Burkina Faso's wildlife (<http://faolex.fao.org/docs/pdf/bkf4885.pdf>; consulted on 10 November 2018). According to this decree, the hunting, capture, possession, and trade of vultures are banned and any person infringing this provision shall be liable to sentences ranging from 6 to 36 months and/or to a fine of 100,000 to 1,000,000 CFA (~ USD 200-2,000). For example, on 12 January 2012, three culprits in Ouagadougou (Burkina Faso) received sentences ranging from 3 to 18 months and a fine of 3,275,000 CFA (~ USD 6,550) (Nabaloum 2012a). As a result of recent convictions in Burkina Faso, poisoning for belief-based use has been scarcer (B. Douamba, Head of Forestry Services, pers. comm.). However, poisoning events have spread to other West African countries where large densities of vultures still occur. One example of this is the intentional poisoning of over 2,000 Hooded Vultures across eastern Guinea-Bissau, between September 2019 and March 2020, to obtain vulture heads for belief-based use (Henriques *et al.* 2020). In certain areas of West Africa such as the western region of The Gambia, the Ziguinchor department of southwestern Senegal, and in Guinea-Bissau, Hooded Vultures still occur in high densities (Jallow *et al.* 2016, Henriques *et al.* 2017, Mullié *et al.* 2017). This suggests that conservation actions and strategies need to be collaborative and coordinated between West African countries to reduce and eventually halt vulture poisoning and the trade in vulture parts for belief-based use in this region.

Susceptibility of Hooded Vultures to intentional poisoning

Of the 15 incidents of intentional poisoning of vultures recorded in our study, 96% of dead vultures were Hooded Vultures ($n = 577$). This is perhaps surprising in light of the results of McKean (2004), Boakye *et al.* (2019), and Mashele *et al.* (2021), who suggested that poisoning of vultures for belief-based use is indiscriminate of species. However, in West Africa, the other vulture species are now confined to protected areas and the Hooded Vulture is the only vulture species that is seen close to human settlements. It is also the most common and widespread vulture throughout West Africa (Thiollay 2007b). This widespread distribution and the close commensal relationship with people in West Africa may have further exposed the Hooded Vulture to the threat of poisoning




Figure 2. The intended uses of Hooded Vultures killed in four incidents of intentional poisoning in which culprits were arrested and sentenced by Burkina Faso's police and the courts ($n = 143$ recorded deaths, Table 4).

for belief-based use. Corroborating this, Saidu and Buij (2013) found that in West African markets, 90% of the vulture parts found were from Hooded Vultures.

Mass poisoning of African vultures and continent-wide population declines

The belief-based use of vulture body parts is very common in West Africa and has existed for many years (Buij *et al.* 2016), but with the use of poisoned bait to harvest vultures (a method which produces the highest number of victims in a single occasion), numbers of vultures are expected to decline rapidly in this area. Indeed, according to Monadjem *et al.* (2018) and Murn and Botha (2018), vultures are supremely adapted to scavenging and are therefore vulnerable to the threat of meat baits that are laced with poison. Furthermore, vultures generally show gregarious feeding behaviour, which increases their vulnerability to poison (Houston 1996, Ogada *et al.* 2012). For example, between 2012 and 2014, 11 poaching-related vulture poisoning incidents recorded in seven African countries killed 2,044 vultures (Ogada *et al.* 2016b). More recently, 537 vultures were reported dead close to the Botswana and Zimbabwe border due to the consumption of three poached elephant carcasses laced with poison (Africa Geographic 2019). Several studies (Hille and Collar 2011, Ogada and Buij 2011, Mullié *et al.* 2017) suggest that poisoning is currently the main threat to vultures in Africa, resulting in population declines (Ogada and Keesing 2010, Virani *et al.* 2011, Ogada *et al.* 2016b, Di Vittorio *et al.* 2018, Garbett *et al.* 2018). This decline assessed at about 80% or more for seven vulture species, over three generations (Ogada *et al.* 2016a, Mullié *et al.* 2017) could lead some species to the brink of extinction if it is not mitigated (Monadjem *et al.* 2018, Murn and Botha 2018).

Cultural value of vultures and impact of trade for belief-based use

In Europe and the USA, vultures have been reported to (rarely) attack and sometimes kill livestock (if food is lacking and/or if the animals are already in a weakened state: immobile, sick, weak, or abandoned lambs at, or soon after, birth) (Avery and Cummings 2004, Margalida *et al.* 2011, 2014, Duriez *et al.* 2019, Lambertucci *et al.* 2021). In South Africa too, following a recent drought and presumed food shortages in some areas, there have been reports of vultures killing livestock (L. J. Thompson pers. obs.). According to Lambertucci *et al.* (2021), this behaviour is rare and does not imply that vultures are efficient predators or hunters which may represent a significant threat to livestock production. In West Africa, vultures are considered harmless to livestock, and people value vultures for the critical ecological services they provide (Ogada *et al.* 2012). For these reasons, people consider vultures to be sacred and they provide protection to vultures in some areas of Burkina Faso (Daboné *et al.* 2019). At the same time, the widespread harvesting of vultures for belief-based use is apparently in contrast to the protection that is usually afforded to vultures in this area. Our study shows that the increasing profitability of the regional trade in vulture body parts is probably the main driver of poisoning, undermining the local cultural value of vultures. The intentional poisoning targeted at vultures for trade for belief-based use appears to be a recent practice in West Africa (Buij *et al.* 2016, Daboné *et al.* 2019, Henriques *et al.* 2020). In Burkina Faso, the first incidents of these intentional poisonings were reported in 2010, when vulture populations were already decreasing. In their West African range, from the 1970s to 2004, the number of Hooded Vultures declined from 84 to 46 individuals every 100 km, a decrease of 45% (Thiollay 2006a, 2006b).

Table 5. Non-exhaustive list of pesticides which are not registered for use by the Sahelian Pesticides Committee (CSP) but which are commonly used in Burkina Faso.

N°	Pesticides	Active ingredients	Pesticide class
1	Adwuma Wura	Glyphosate* (480 g/l)	Herbicide
2	Atrazine	Atrazine	ND
3	Benaxone super	Paraquat* (270 g/l)	Herbicide
4	Buta plus	ND	ND
5	Butaclor	ND	ND
6	Calthio E	Endosulfan* 25%, thirame 25%	ND
7	Conti-sul	ND	Insecticide
8	Ervextra 750cl	ND	ND
9	Focogeb	ND	ND
10	Furadan 5G	Carbofuran*5%	Insecticide
11	Gramoquat	Paraquat* (200 g/l)	Insecticide
12	Lambda super	Lambda-cyhalothrine* 25	Insecticide
13	Lamda best 2.5 EC	ND	ND
14	Rilof	ND	ND
15	Sunphosate G 7578	ND	ND
16	Tango 500	ND	ND

Active ingredient mentioned in the PAN (Pesticide Action Network) International List of Highly Hazardous Pesticides: PAN List of HHPs*
 ND = not determined

The intentional poisonings that have occurred over recent decades have probably hastened the decline of vultures which was already well underway in Burkina Faso and in other West African countries.

Availability of toxic substances used for poisoning

Beyond these targeted poisonings, the unintentional poisoning of vultures has probably contributed to the large decline of vultures in West Africa. Indeed, based on the results of this survey, only five unintentional poisoning events killed at least 202 Hooded Vultures. According to the veterinarians we surveyed, the underlying factor is the indiscriminate and widespread use of pesticides, rodenticides and veterinary drugs that are cheap to buy, and easily accessed across the highly porous borders in West Africa. The veterinary drugs referred to by veterinarians in our survey were (in no way exhaustive and for reference only): Benzyl benzoate, Ivermectin, Ivomec, Limoxin-200 LA, Oxytetracycline, Peni Strepto, trypamidium, Diminazeme aceturate, Isometamidium chloride, and Veriben. Furthermore, many of pesticides commonly used in this area are not registered by the Sahelian Pesticides Committee (CSP; effective since 1994 in member-states of CILSS including Burkina Faso) (Table 5) and yet it is common to find them openly for sale in markets. According to Thiollay (2006a), this widespread use of pesticides and rodenticides in West Africa has led to vulture declines. If we are to significantly reduce unintentional (often illegal) poisoning of vultures, there is an urgent need to review, develop and significantly increase the enforcement of appropriate legislation to control, ban or restrict the sale, storage, distribution, use and

disposal of toxic chemicals involved in the indiscriminate killing of wildlife.

Conclusion

West Africa is a key site for the conservation of African vultures because, despite the collapse of vulture populations highlighted in this area 30–40 years ago (Thiollay 2006b, Ogada *et al.* 2016a), in some areas, vultures still occur in appreciable densities (Jallow *et al.* 2016, Henriques *et al.* 2017). However, the rapid rise in reports of vultures killed using poisoned baits is alarming and presents a serious conservation challenge for vultures in this region. Elsewhere, as in Europe, suitable conservation actions have been implemented and vultures are increasing in number and recolonising their old ranges (Safford *et al.* 2019). In West Africa, although local actions are emerging, a lot more needs to be done by governments, NGOs, research institutions, and the media, to ensure sustainable vulture conservation. Priority actions include: a) restricting access to and use of hazardous toxic substances which may be used to poison wildlife including vultures; b) training of personnel (foresters and veterinarians) to detect and respond to wildlife poisoning incidents; c) conducting research on vultures to update population trends, breeding success and distribution in protected areas and adjacent communities, and d) compiling and enacting a regional action plan for sustainable vulture conservation in West Africa.

Acknowledgements. This project is a joint initiative between the University Joseph Ki-Zerbo in Burkina Faso and the University of Groningen in The Netherlands. The NATURAMA Foundation in Burkina Faso also provided support and assistance. We would also like to thank people interviewed in this

study (butchers, veterinarians, foresters, abattoir watchmen, police officers and court registrars) for their participation and for allowing us to undertake this work in their locality. This work was supported by the Science and Society Group in the Faculty of Sciences and Engineering, University of Groningen, The Netherlands.

Supplementary Materials. To view supplementary material for this article, please visit <http://doi.org/10.1017/S095927092100054X>.

References

- Africa Geographic (2019) Catastrophic breaking news: 537 vultures found poisoned in dark day for Botswana conservation. <https://africageographic.com/stories/catastrophic-breaking-news-537-vultures-found-poisoned-in-dark-day-for-botswana-conservation/> [accessed 09 April 2021].
- AIB (2012) *Un homme interpellé à Bitou avec trois cadavres de vautours*. Agence de l'Information du Burkina, aOuaga.com, 9 November. <http://news.aouaga.com/h/2333.html> [accessed 20 June 2018].
- Alarcón, P. A. and Lambertucci, S. A. (2018) Pesticides thwart condor conservation. *Science* **360**: 612.
- Avery, M. L. and Cummings, J. L. (2004) Livestock depredations by black vultures and golden eagles. *Sheep and Goat Res. J.* **19**: 58–63.
- Awoyemi, S. (2014) Vulture declines in West Africa: investigating the scale and (socioeconomic) drivers of the trade in vulture parts for traditional medicine. Placement Report, MPhil in Conservation Leadership (Examination no. 92), Department of Geography, University of Cambridge, UK. <https://www.4vulture.s.org/2014/10/20/significant-contributions-to-document-the-impact-of-the-trade-in-vulture-parts-for-traditional-medicine-in-africa-published/> [accessed 20 June 2020].
- Bernard, H. R. (2002) *Research methods in anthropology: Qualitative and quantitative approaches*. 3rd edition. Walnut Creek, CA, USA: Altamira Press.
- Berny, P. (2007) Pesticides and the intoxication of wild animals. *J. Vet. Pharmacol. Ther.* **30**: 93–100.
- Boakye, M. K., Wiafe, E. D. and Ziekah, M. Y. (2019) Ethnomedicinal use of vultures by traditional medicinal practitioners in Ghana. *Ostrich* **90**: 111–118.
- Borrow, N. and Demey, R. (2001) *The birds of Western Africa*. London, UK: C. Helm.
- Botha, A. J., Andevski, J., Bowden, C. G. R., Gudka, M., Safford, R. J., Tavares, J. *et al.* (2017) *Multi-species Action Plan to conserve African-Eurasian vultures*. Abu Dhabi, United Arab Emirates: Coordinating Unit of the CMS Raptors MOU. (CMS Raptors MOU Technical Publication No. 5. CMS Technical Series No. 35).
- Buij, R. and Croes, B. M. (2014) Raptors in northern Cameroon, December 2005–December 2010. *Bull. Afr. Bird Club* **2**: 26–63.
- Buij, R., Nikolaus, G., Whytock, R., Ingram, D. J. and Ogada, D. L. (2016) Trade of threatened vultures and other raptors for fetish and bushmeat in West and Central Africa. *Oryx* **50**: 606–616.
- Craig, C. A., Thomson, R. L. and Santangeli, A. (2018) Communal farmers of Namibia appreciate vultures and the ecosystem services they provide. *Ostrich* **89**: 211–220.
- Cresswell, J. W. and Plano Clark, V. L. (2011) *Designing and conducting mixed method research*. 2nd edition. Thousand Oaks: Sage.
- Daboné, C., Buij, R., Oueda, A., Adjakpa, J. B., Guenda, W. and Weesie P. D.M. (2019) Impact of human activities on the reproduction of the Hooded Vultures, *Necrosyrtes monachus* in Garango area (Burkina Faso). *Ostrich* **90**: 53–61.
- Di Vittorio, M., Hema, E. M., Dendi, D., Akani, G. C., Cortone, G., López-López, P., Amadi, M., Hoinssoudé Ségniabeto, G., Battisti, C. and Luiselli, L. (2018) The conservation status of west African vultures: an updated review and a strategy for conservation. *Vie Milieu-Life Environ.* **68**: 33–43.
- Duriez, O., Descaves, S., Gallais, R., Neouze, R., Fluhr, J. and Decante, F. (2019) Vultures attacking livestock: a problem of vulture behavioural change or farmers' perception? *Bird Conserv. Internatn.* **29**: 437–453.
- Fleischli, M. A., Franson, J. C., Thomas, N. J., Finley, D. L. and Riley, W. (2004) Avian mortality events in the United States caused by anticholinesterase pesticides: a retrospective summary of National Wildlife Health Center records from 1980 to 2000. *Arch. Environ. Contam. Toxicol.* **46**: 542–550.
- Garbett, R., Herremans, M., Maude, G., Reading, R. P. and Amar, A. (2018) Raptor population trends in northern Botswana: a re-survey of road transects after 20 years. *Biol. Conserv.* **224**: 87–99.
- Guitart, R., Sachana, M., Caloni, F., Croubels, S., Vandenbroucke, V. and Berny, P. (2010) Animal poisoning in Europe. Part 3: wildlife. *Vet. J.* **183**: 260–265.
- Henriques, M., Lecoq, M., Monteiro, H., Regalla, A., Granadeiro, J. P. and Catry, P. (2017) Status of birds of prey in Guinea-Bissau: first assessment based on road surveys. *Ostrich* **88**: 101–111.
- Henriques, M., Buij, R., Monteiro, H., Sá, J., Wambar, F., Tavares, J. P., Botha, A., Citegetse, G., Lecoq, M., Catry, P. and Ogada D. (2020) Deliberate poisoning of Africa's vultures. *Science* **370**: 304.
- Hille, S. M. and Collar, N. J. (2011) Status assessment of raptors in Cape Verde confirms a major crisis for scavengers. *Oryx* **45**: 217–224.
- Houston, D. C. (1996) The effect of altered environments on vultures. Pp. 327–336 in D. Bird, E. Varland and J. J. Negro, eds. *Raptors in human landscapes*. London, UK: Academic Press.
- Ibrahim, B., Polcher, J., Karambiri, H., Yacouba, H. and Ribstein P. (2014) Changes in rainfall regime over Burkina Faso under the climate change conditions simulated by 5 regional climate models. *Clim. Dyn.* **42**: 1363–1381
- Jallow, M., Barlow, C. R., Sanyang, L., Dibba, L., Kendall, C., Bechard, M. and Bildstein K. L. (2016) High population density of the Critically Endangered Hooded Vulture *Necrosyrtes monachus* in Western Region, The Gambia, confirmed by road surveys in 2013 and 2015. *Malimbus* **38**: 23–28.
- Kabir, S. M. S. (2016) *Basic guidelines for research: An introductory approach for all disciplines*. Chittagong, Bangladesh: Book Zone Publication.
- Kim, S., Park, M.-Y., Kim, H.-J., Shin, J. Y., Ko, K. Y., Kim, D.-G., Kim, M., Kang, H.-G., So, B. and Park, S.-W. (2016) Analysis of insecticides in dead wild birds in Korea from 2010 to 2013. *Bull. Environ. Contam. Toxicol.* **96**: 25–30.
- Lambertucci, S. A., Margalida, A., Speziale, K.L., *et al.* (2021) Presumed killers? Vultures, stakeholders, misperceptions, and fake news. *Conserv. Sci. Practice* **3**: e415: 1–10.
- Loveridge, R., Ryan, G. E., Phearun, S., Gray-Read, O., Mahood, S. P., Mould, A., Harrison, S., Crouthers, R., Ko, S., Clements, T., Eames, J. C. and Pruvot, M. (2019) Poisoning causing the decline in South-East Asia's largest vulture population. *Bird Conserv. Internatn.* **29**: 41–54.
- MaMing, R. and Xu, G. (2015) Status and threats to vultures in China. *Vulture News* **68**: 3–24.
- Mander, M., Diederichs, N., Ntuli, L., Mavundla, K., Williams, V. and McKean, S. (2007) *Survey of the trade in vultures for the traditional health industry in South Africa*. Unpublished report, pp. 54.
- Margalida, A. (2012) Baits, budget cuts: a deadly mix. *Science* **338**: 192.
- Margalida, A., Heredia, R., Razin, M. and Hernández, M. (2008) Sources of variation in mortality of the Bearded Vulture *Gypaetus barbatus* in Europe. *Bird Conserv. Internatn.* **18**: 1–10.
- Margalida, A., Campión, D. and Donazar, A. J. (2011) European vultures' altered behaviour. *Nature* **480**: 457.
- Margalida, A., Campión, D. and Donazar, A. J. (2014) Vultures vs livestock: Conservation relationships in an emergent human-wildlife conflict. *Oryx* **48**: 172–176.
- Margalida A., Ogada, D. and Botha, A. (2019) Protect African vultures from poison. *Science*, **365**: 1089–1090.
- Mashele, N. M., Thompson, L. J. and Downs, C. T. (2021) Uses of vultures in traditional medicines in the Kruger to Canyons Biosphere Region, South Africa. *J. Raptor Res.* **55**: 328–339.
- McKean, S. (2004) Traditional use of vultures: Some perspectives in the vultures of southern Africa – Quo Vadis? Pp. 195–201 in *Proceedings of a workshop on vulture research and conservation in southern Africa*. Birds of Prey Working Group.
- McKean, S., Mander, M., Diederichs, N., Ntuli, L., Mavundla, K., Williams, V. and Wakelin, J. (2013) The impact of traditional use on vultures in South Africa. *Vulture News* **65**: 15–36.
- MECV (2007) *Programme d'Action Nationale d'Adaptation à la Variabilité et aux Changements Climatiques*. Ouagadougou, Burkina Faso: Ministère de l'Environnement et du Cadre de Vie, Secrétariat permanent du conseil national de l'environnement et du développement durable.
- Mineau, P., Fletcher, M. R., Glaser, L. C., Thomas, N. J., Brassard, C., Wilson, L. K., Elliott, J. E., Lyon, L. A., Henny, C. J. and Bollinger, T. (1999) Poisoning

- of raptors with organophosphorus and carbamate pesticides with emphasis on Canada, US and UK. *J. Raptor Res.* **33**: 1–37.
- Monadjem, A., Kane, A., Botha, A., Kelly, C. and Murn, C. (2018) Spatially explicit poisoning risk affects survival rates of an obligate scavenger. *Sci. Rep.* **8**: 4364.
- Mullié, W. C., Couzi, F.-X., Diop, M. S., Piot, B., Peters, T., Reynaud, P. A. and Thiollay, J.-M. (2017) The decline of an urban Hooded Vulture *Necrosyrtes monachus* population in Dakar, Senegal, over 50 years. *Ostrich* **88**: 131–138.
- Mundy, P., Butchart, D., Ledger, J. and Piper, S. (1992) *The vultures of Africa*. London, UK: Academic Press.
- Murn, C. and Botha, A. (2018) A clear and present danger: Impacts of poisoning on a vulture population and the effect of poison response activities. *Oryx* **52**: 552–558.
- Nabaloum (2012a) *Tribunal de grande instance de Ouagadougou: Ils tuent 71 vautours et écopot de 18 mois de prison*. LeFaso.net, 19 January. <http://lefaso.net/spip.php?article45939> [accessed 20 April 2018].
- Nabaloum (2012b) *Braconnage: 26 vautours tués pour être exportés au Nigeria*. LeFaso.net, 11 avril. <https://lefaso.net/spip.php?article47381> [accessed 23 July 2019].
- Nikolaus, G. (2011) The fetish culture in West Africa: an ancient tradition as a threat to endangered bird life? Pp. 145–155 in K.L. Schuchmann, ed. *Tropical vertebrates in a changing world*. Bonn, Germany: Zoologisches Forschungsmuseum Alexander Koenig.
- Nosazeogie, E., Tende, T. and Monadjem, A. (2018) Hooded Vultures *Necrosyrtes monachus* nearly extirpated from Edo State, Nigeria: A report on the avian scavenger community. *Ostrich* **89**: 265–273.
- Ogada D. L. (2014) The power of poison: Pesticide poisoning of Africa's wildlife. *Ann. N. Y. Acad. Sci.* **1322**: 1–20.
- Ogada, D. L. and Keesing, F. (2010) Decline of raptors over a three-year period in Laikipia, central Kenya. *J. Raptor Res.* **44**: 129–135.
- Ogada D. L. and Buij R. (2011) Large declines of the Hooded Vulture *Necrosyrtes monachus* across its African range. *Ostrich* **82**: 101–113.
- Ogada D. L., Keesing F. and Virani M. Z. (2012) Dropping dead: Causes and consequences of vulture population declines worldwide. *Ann. N. Y. Acad. Sci.* **1249**: 57–71.
- Ogada, D. L., Shaw, P., Beyers, R. L., Buij, R., Murn, C., Thiollay, J. M., Beale, C. M., Holdo, R. M., Pomeroy, D., Baker, N., Krüger, S. C., Botha, A., Virani, M. Z., Monadjem, A. and Sinclair, A. R. E. (2016a) Another continental vulture crisis: Africa's vultures collapsing toward extinction. *Conserv. Lett.* **9**: 89–97.
- Ogada D., Botha A. and Shaw P. (2016b) Ivory poachers and poison: drivers of Africa's declining vulture populations. *Oryx* **50**: 593–596.
- Pantović, U. and Andevski, J. (2018) *Review of the problem of poison use and vulture poisoning in the Balkan Peninsula*. The Netherlands: Vulture Conservation Foundation.
- Patton, M. (2002) *Qualitative research and evaluation methods*. 3rd edition. Thousand Oaks, CA: Sage.
- Pauli, J. N., Donadio, E. and Lambertucci, S.A. (2018) The corrupted carnivore: how humans are rearranging the return of the carnivore-scavenger relationship. *Ecology* **99**: 2122–2124.
- Pavez, E. F. and Estades, C. F. (2016) Causes of admission to a rehabilitation center for Andean Condors (*Vultur gryphus*) in Chile. *J. Raptor Res.* **50**: 23–32.
- Portier, B. (2007) Les oiseaux. In *Nazinga*. Les presses agronomiques de Gembloux.
- Richards, N., Ogada, D., Buij, R. and Botha, A. (2017) The killing fields: The use of pesticides and other contaminants to poison wildlife in Africa. *Encyclopedia of the Anthropocene* **5**: 161–167.
- Safford, R., Andevski, J., Botha, A., Bowden, C. G. R., Crockford, N., Garbett, R., et al., (2019) Vulture conservation: the case for urgent action. *Bird Conserv. Internatn.* **29**: 1–9.
- Saidu, Y. and Buij, R. (2013) Traditional medicine trade in vulture parts in northern Nigeria. *Vulture News* **65**: 4–14.
- Spradley, J. (1979) *The ethnographic interview*. Long Grove, IL: Waveland Press.
- Thiollay, J.-M. (2006a) Large bird declines with increasing human pressure in savanna woodlands (Burkina Faso). *Biodivers. Conserv.* **15**: 2085–2108.
- Thiollay J.-M. (2006b) The decline of raptors in West Africa: Long-term assessment and the role of protected areas. *Ibis* **148**: 240–254.
- Thiollay, J.-M. (2007a) Raptor declines in West Africa: comparisons between protected, buffer and cultivated areas. *Oryx* **41**: 322–329.
- Thiollay, J.-M. (2007b) Raptor population decline in West Africa. *Ostrich* **78**: 405–413.
- Virani, M. Z., Kendall, C., Njoroge, P. and Thomsett, S. (2011) Major declines in the abundance of vultures and other scavenging raptors in and around the Masai Mara ecosystem, Kenya. *Biol. Conserv.* **144**: 746–752.
- Weesie, P. D. M. and Belemsobgo, U. (1997) Les rapaces diurnes du ranch de gibier de Nazinga (Burkina Faso): liste commentée, analyse du peuplement et cadre biogéographique. *Alauda* **65**: 263–278.