

---

Department of Applied Mathematics  
Faculty of EEMCS



University of Twente  
*The Netherlands*

---

P.O. Box 217  
7500 AE Enschede  
The Netherlands

Phone: +31-53-4893400

Fax: +31-53-4893114

Email: [memo@math.utwente.nl](mailto:memo@math.utwente.nl)  
[www.math.utwente.nl/publications](http://www.math.utwente.nl/publications)

---

**Memorandum No. 1700**

**A continuation of spanning 2-connected  
subgraphs in truncated rectangular grid graphs**

A.N.M. SALMAN, H.J. BROERSMA AND

C.A. RODGER<sup>1</sup>

December, 2003

ISSN 0169-2690

---

<sup>1</sup>Department of Discrete and Statistical Sciences, Auburn University, Auburn, Alabama 36849, USA

**A CONTINUATION OF SPANNING 2-CONNECTED  
SUBGRAPHS IN TRUNCATED RECTANGULAR GRID  
GRAPHS**

A. N. M. Salman

Department of Applied Mathematics  
Faculty of Electrical Engineering, Mathematics and Computer Science  
University of Twente  
P.O. Box 217, 7500 AE Enschede, The Netherlands  
*a.n.m.salman@math.utwente.nl*

H. J. Broersma

Department of Applied Mathematics  
Faculty of Electrical Engineering, Mathematics and Computer Science  
University of Twente  
P.O. Box 217, 7500 AE Enschede, The Netherlands  
*h.j.broersma@math.utwente.nl*

C.A. Rodger

Department of Discrete and Statistical Sciences,  
Auburn University, Auburn, Alabama 36849, USA  
*rodgec1@mail.auburn.edu*

**Abstract**

A grid graph is a finite induced subgraph of the infinite 2-dimensional grid defined by  $Z \times Z$  and all edges between pairs of vertices from  $Z \times Z$  at Euclidean distance precisely 1. An  $m \times n$ -rectangular grid graph is induced by all vertices with coordinates 1 to  $m$  and 1 to  $n$ , respectively. A natural drawing of a (rectangular) grid graph  $G$  is obtained by drawing its vertices in  $\mathbb{R}^2$  according to their coordinates. We consider a subclass of the rectangular grid graphs obtained by deleting some vertices from the corners. Apart from the outer face, all (inner) faces of these graphs have area one (bounded by a 4-cycle) in a natural drawing of these graphs. We determine which of these graphs contain a Hamilton cycle, i.e. a cycle containing all vertices, and solve the problem of determining a spanning 2-connected subgraph with as few edges as possible for all these graphs.

**Keywords:** (rectangular) grid graph, Hamilton cycle, spanning 2-connected subgraph

**AMS Subject Classifications:** 05C40, 05C85

## 1 Introduction

The *infinite grid graph*  $G^\infty$  is defined by the set of vertices  $V = \{(x, y) \mid x \in \mathbb{Z}, y \in \mathbb{Z}\}$  and the set of edges  $E$  between all pairs of vertices from  $V$  at Euclidean distance precisely 1. For any integers  $m \geq 1$  and  $n \geq 1$ , the *rectangular grid graph*  $R(m, n)$  is the (finite) subgraph of  $G^\infty$  induced by  $V(m, n) = \{(x, y) \mid 1 \leq x \leq m, 1 \leq y \leq n, x \in \mathbb{Z}, y \in \mathbb{Z}\}$  (and just containing all edges from  $G^\infty$  between pairs of vertices from  $V(m, n)$ ). This graph  $R(m, n)$  is also known as the *product graph*  $P_m \times P_n$  of two disjoint paths  $P_m$  and  $P_n$ . A *grid graph* is a graph that is isomorphic to a subgraph of  $R(m, n)$  induced by a subset of  $V(m, n)$  for some integers  $m \geq 1$  and  $n \geq 1$ . It is clear that a grid graph  $G = (V, E)$  is a *planar graph*, i.e. it can be drawn in the plane  $\mathbb{R}^2$  in such a way that the edges only intersect at the vertices of the graph. In such a drawing, the regions of  $\mathbb{R}^2 \setminus (V \cup E)$  are called the *faces* of  $G$ . Exactly one of the faces is unbounded; this is called the *outer face*; the others are its *inner faces*. The *natural drawing* of a grid graph is just described by drawing its vertices in  $\mathbb{R}^2$  according to their coordinates. A *solid grid graph* is a grid graph all of whose inner faces have area one (are bounded by a cycle on four vertices) in a natural drawing. A grid graph that is not solid contains inner faces (in a natural drawing) that have area larger than one; these faces are called *holes*. A subgraph  $H$  of a graph  $G = (V, E)$  is called a *spanning subgraph* if  $V(H) = V$ . A connected graph is called *2-connected* if it remains connected if at most one vertex is removed. A *Hamilton cycle* in a graph  $G = (V, E)$  is a cycle containing every vertex of  $V$ , i.e. a spanning 2-connected subgraph in which every vertex has degree 2 (the number of edges is  $|V|$ ).

Itai, Papadimitriou and Szwarcfiter [2] have proved that deciding whether a given grid graph has a Hamilton cycle is an NP-complete problem. This implies that the problem of finding a spanning 2-connected subgraph with as few edges as possible is also NP-hard for grid graphs. It has been conjectured that the first problem remains NP-complete when it is restricted to solid grid graphs. However, Umans and Lenhart [9] recently proved that this problem is polynomially solvable for solid grid graphs, by presenting a complicated algorithm with time complexity  $O(|V|^4)$ . In a recent paper of Sheffield [8] the work of [2] has been extended to grid graphs with a small number of holes. For the second problem the complexity is not known when it is restricted to solid grid graphs. It remains an open problem –what the complexity of both problems is– when we restrict ourselves to grid graphs with a fixed number of holes.

Motivated by the above problems, we study the problem of the existence of a Hamilton cycle and the problem of determining a spanning 2-connected subgraph with as few edges as possible for a number of classes of grid graphs without holes, obtained from rectangular grid graphs by deleting vertices from the corners in a natural drawing. This is a continuation of the work started in [3], [4] and [6]. In the other papers, [5] and [7], we study the same problem for classes of grid graphs

with a few holes. For all graphs of the defined classes we are able to solve the second problem. All solutions are of the same type: first, we use the well-known *Grinberg-condition* and the properties of bipartite graphs to derive a lower bound for the number of edges in a spanning 2-connected subgraph. Secondly, we show by construction that this lower bound is in fact the optimum value.

## 2 Truncated rectangular grid graphs

We now introduce the classes of grid graphs which we call *truncated rectangular grid graphs*.

For  $m \geq 3$ ,  $n \geq 3$ ,  $0 \leq k \leq \min\{m-2, n-2\}$  and  $0 \leq l \leq \min\{m-2, n-2\}$  we define a *1-corner truncated rectangular grid graph*  $R(m, n)^{-1(k,l)}$  as the subgraph obtained from  $R(m, n)$  by deleting  $k \times l$  vertices from one corner in  $V(m, n)$  together with their incident edges in a natural drawing. For illustration, consider  $R(11, 13)^{-1(2,3)}$  in Figure 1(a).

For  $m \geq 6$ ,  $n \geq 6$ ,  $1 \leq k \leq \min\{\frac{m-4}{2}, \frac{n-4}{2}\}$  and  $1 \leq l \leq \min\{\frac{m-4}{2}, \frac{n-4}{2}\}$  we define a *2-corner truncated rectangular grid graph*  $R(m, n)^{-2(k,l)}$  as the subgraph obtained from  $R(m, n)$  by deleting  $k \times l$  vertices from two opposite corners in  $V(m, n)$  together with their incident edges in a natural drawing. For illustration, consider  $R(11, 13)^{-2(2,3)}$  in Figure 1(b).

For  $m \geq 6$ ,  $n \geq 6$ ,  $1 \leq k \leq \min\{\frac{m-4}{2}, \frac{n-4}{2}\}$  and  $1 \leq l \leq \min\{\frac{m-4}{2}, \frac{n-4}{2}\}$  we define a *3-corner truncated rectangular grid graph*  $R(m, n)^{-3(k,l)}$  as the subgraph obtained from  $R(m, n)$  by deleting  $k \times l$  vertices from three corners in  $V(m, n)$  together with their incident edges in a natural drawing. For illustration, consider  $R(11, 13)^{-3(2,3)}$  in Figure 1(c).

For  $m \geq 6$ ,  $n \geq 6$ ,  $1 \leq k \leq \min\{\frac{m-4}{2}, \frac{n-4}{2}\}$  and  $1 \leq l \leq \min\{\frac{m-4}{2}, \frac{n-4}{2}\}$  we define a *4-corner truncated rectangular grid graph*  $R(m, n)^{-4(k,l)}$  as the subgraph obtained from  $R(m, n)$  by deleting  $k \times l$  vertices from each corner in  $V(m, n)$  together with their incident edges in a natural drawing. For illustration, consider  $R(11, 13)^{-4(2,3)}$  in Figure 1(d).




Figure 1: Truncated rectangular grid graphs : (a)  $R(11, 13)^{-1(2,3)}$  (b)  $R(11, 13)^{-2(2,3)}$  (c)  $R(11, 13)^{-3(2,3)}$  (d)  $R(11, 13)^{-4(2,3)}$

Spanning 2-connected subgraphs with a minimum number of edges for the 1-

corner truncated rectangular grid graph  $R(m, n)^{-1(k, k)}$  and for the 4-corner truncated rectangular grid graph  $R(m, n)^{-4(k, k)}$  were studied in [4]. Subsequently, in [6] these results were generalized to the following theorem.

**Theorem 1** *Let  $R(m, n)^{-1(k, l)}$  and  $R(m, n)^{-4(k, l)}$  denote the 1-corner truncated rectangular grid graph and the 4-corner truncated rectangular grid graph as defined above, respectively. Then:*

(i)  $R(m, n)^{-1(k, l)}$  contains a spanning 2-connected subgraph with (at most)  $|V| + 1$  edges and is hamiltonian if and only if both  $m \cdot n$  and  $k \cdot l$  are even or both  $m \cdot n$  and  $k \cdot l$  are odd;

(ii)  $R(m, n)^{-4(k, l)}$  contains a spanning 2-connected subgraph with (at most)  $|V| + 3$  edges and is hamiltonian if and only if  $m \cdot n$  is even. The bound  $|V| + 3$  is best possible for any odd numbers  $m, n, k$  and  $l$ .

Our main result in this paper characterizes which of the other truncated rectangular grid graphs are hamiltonian and guarantees the existence of a spanning 2-connected subgraph with at most two edges more than their number of vertices. We postpone the proofs and constructions (figures) until the next section.

**Theorem 2** *Let  $R(m, n)^{-2(k, l)}$  and  $R(m, n)^{-3(k, l)}$  denote the 2-corner truncated rectangular grid graph and the 3-corner truncated rectangular grid graph as defined above, respectively. Then:*

(i)  $R(m, n)^{-2(k, l)}$  contains a spanning 2-connected subgraph with

- $|V|$  edges if  $m \cdot n$  is even and at least one of  $k$  and  $l$  is even if both  $m$  and  $n$  are even;
- $|V| + 2$  edges if  $m$  and  $n$  are even and  $k$  and  $l$  are odd;
- $|V| + 1$  edges in all other cases.

*These numbers of edges are all best possible;*

(ii)  $R(m, n)^{-3(k, l)}$  contains a spanning 2-connected subgraph with

- $|V|$  edges if both  $m \cdot n$  and  $k \cdot l$  are even;
- $|V| + 2$  edges if all of  $m, n, k$  and  $l$  are odd;
- $|V| + 1$  edges in all other cases.

*These numbers of edges are all best possible.*

### 3 Proofs and constructions

A useful necessary condition for hamiltonicity is the following result due to Grinberg [1].

**Lemma 3** *Suppose a planar graph  $G$  has a Hamilton cycle  $H$ . Let  $G$  be drawn in the plane, and let  $r_i$  denote the number of faces inside  $H$  bounded by  $i$  edges in this planar embedding. Let  $r'_i$  be the number of faces outside  $H$  bounded by  $i$  edges. Then*

the numbers  $r_i$  and  $r'_i$  satisfy the following equation.

$$\sum_i (i-2)(r_i - r'_i) = 0.$$

We use this lemma to show that  $R(m, n)^{-2(k, l)}$  contains no Hamilton cycle if  $m \cdot n$  is odd, and that  $R(m, n)^{-3(k, l)}$  contains no Hamilton cycle if  $m \cdot n$  and  $k \cdot l$  have a different parity.

**Corollary 4**  $R(m, n)^{-2(k, l)}$  contains no Hamilton cycle if  $m \cdot n$  is odd.

*Proof.* There is exactly one face with  $2(m + n - 2)$  edges and there are exactly  $(m - 1)(n - 1) - 2k \cdot l$  faces with four edges in the planar (natural) drawing of the 2-corner truncated rectangular grid graph  $R(m, n)^{-2(k, l)}$ . Let this graph be hamiltonian. Then by Lemma 3 we have

$$(2(m + n - 2) - 2)(-1) + (4 - 2)(r_4 - r'_4) = 0.$$

Hence

$$r_4 - r'_4 = m + n - 3. \quad (1)$$

It is easy to check that the number of faces with four edges is

$$r_4 + r'_4 = (m - 1)(n - 1) - 2k \cdot l. \quad (2)$$

From equation (1) and (2) we obtain

$$2r_4 = m \cdot n - 2k \cdot l - 2. \quad (3)$$

So,  $m \cdot n$  is even. □

**Corollary 5**  $R(m, n)^{-3(k, l)}$  contains no Hamilton cycle if  $m \cdot n$  and  $k \cdot l$  have a different parity.

*Proof.* There is exactly one face with  $2(m + n - 2)$  edges and there are exactly  $(m - 1)(n - 1) - 3k \cdot l$  faces with four edges in the planar (natural) drawing of the 3-corner truncated rectangular grid graph  $R(m, n)^{-3(k, l)}$ . Let this graph be hamiltonian. Then by Lemma 3 and using a similar method as in the proof of Corollary 4, we obtain

$$2r_4 = m \cdot n - 3k \cdot l - 2. \quad (4)$$

So, either  $m \cdot n$  and  $k \cdot l$  are even or  $m \cdot n$  and  $k \cdot l$  are odd. □

**Lemma 6**  $R(m, n)^{-2(k, l)}$  contains no spanning 2-connected subgraph with at most  $|V| + 1$  edges if both  $m$  and  $n$  are even and both  $k$  and  $l$  are odd.

*Proof.* First, consider a bipartition  $(S, T)$  of  $R(m, n)$  for some positive even integers  $m$  and  $n$ . Assume that one of the corner vertices is in  $S$ . Then one easily shows that the opposite corner vertex is also in  $S$ , whereas the two other corner vertices are in  $T$  and that  $|S| = |T|$ . This can be proved by induction on  $m$  and  $n$ , and removing the cycle of the outer face if  $m, n \geq 4$ .

Secondly, consider a bipartition  $(S, T)$  of  $R(k, l)$  for odd  $k$  and  $l$ . Assume that one of the corner vertices is in  $S$  (if  $m, n \geq 3$ ; otherwise consider an end vertex). Then we can show that all corner vertices (end vertices) are in  $S$ , and that  $|S| = |T| + 1$ . This can be proved by induction on  $m$  and  $n$ , and removing the cycle of the outer face if  $m, n \geq 3$ .

So if we remove the two opposite corner  $R(k, l)$ 's from  $R(m, n)$ , we reduce  $|S|$  by two more units than  $|T|$ , implying that  $R(m, n)^{-2(k, l)}$  has a bipartition  $(S', T')$  with  $|T'| = |S'| + 2$ . In any spanning 2-connected subgraph  $G$  of  $R(m, n)^{-2(k, l)}$  all vertices in  $T'$  have degree at least 2, hence  $|E(G)| \geq 2|T'| = |T'| + |S'| + 2 = |V(G)| + 2$ . This completes the proof of Lemma 6. □

**Lemma 7**  $R(m, n)^{-3(k, l)}$  contains no spanning 2-connected subgraph with at most  $|V| + 1$  edges if all of  $m, n, k$  and  $l$  are odd.

*Proof.* Consider a bipartition  $(S, T)$  of  $R(m, n)$  for odd  $m$  and  $n$ . Assume that one of the corner vertices is in  $S$ . By the same arguments as in the proof of Lemma 6, then all corner vertices are in  $S$ , and  $|S| = |T| + 1$ . The same holds for  $R(k, l)$  if  $k$  and  $l$  are odd. So if we remove the three corner  $R(k, l)$ 's from  $R(m, n)$ , we reduce  $|S|$  by three more units than  $|T|$ , implying that  $R(m, n)^{-3(k, l)}$  has a bipartition  $(S', T')$  with  $|T'| = |S'| + 2$ . In any spanning 2-connected subgraph  $G$  of  $R(m, n)^{-3(k, l)}$  all vertices in  $T'$  have degree at least 2, hence  $|E(G)| \geq 2|T'| = |T'| + |S'| + 2 = |V(G)| + 2$ . This completes the proof of Lemma 7. □

We complete the proof of Theorem 2 by showing, through construction, the existence of a Hamilton cycle or a spanning 2-connected subgraph with at most  $|V| + 2$  edges, in all cases where  $m = 10$  or  $11$ ,  $n = 12$  or  $13$ ,  $k = 2$  or  $3$  and  $l = 2$  or  $3$ . Meanwhile, for other values of  $m, n, k$  and  $l$ , it is not difficult to see from the patterns in the figures that follow how to extend the solutions.

A Hamilton cycle for  $R(11, 12)^{-2(3, 2)}$  is shown in Figure 2(a). The pattern in this figure can be used for finding a Hamilton cycle for the 2-corner truncated rectangular grid graph for either (any numbers  $m$  and  $k$ , and any even numbers  $n$  and  $l$ ) or (any numbers  $n$  and  $l$ , and any even numbers  $m$  and  $k$ ). In Figure 2(b) we show




Figure 2: A Hamilton cycle for : (a)  $R(11, 12)^{-2(3,2)}$  (b)  $R(11, 12)^{-2(2,3)}$  (c)  $R(11, 12)^{-2(3,3)}$

a Hamilton cycle for  $R(11, 12)^{-2(2,3)}$ . The pattern in Figure 2(b) can be used for finding a Hamilton cycle for the 2-corner truncated rectangular grid graph for either (any even numbers  $n$  and  $k$ , and any odd numbers  $m$  and  $l$ ) or (any even numbers  $m$  and  $l$ , and any odd numbers  $n$  and  $k$ ). In Figure 2(c) we show a Hamilton cycle for  $R(11, 12)^{-2(3,3)}$ . The pattern in Figure 2(c) can be used for finding a Hamilton cycle for the 2-corner truncated rectangular grid graph for either (any even number  $n$ , and any odd numbers  $m$ ,  $k$  and  $l$ ) or (any even number  $m$ , and any odd numbers  $n$ ,  $k$  and  $l$ ).




Figure 3: A spanning 2-connected subgraph for : (a)  $R(11, 13)^{-2(2,3)}$  with  $|V| + 1$  edges (b)  $R(11, 13)^{-2(2,2)}$  with  $|V| + 1$  edges (c)  $R(10, 12)^{-2(3,3)}$  with  $|V| + 2$  edges

A spanning 2-connected subgraph for  $R(11, 13)^{-2(2,3)}$  with  $|V| + 1$  edges is shown in Figure 3(a). The pattern in this figure can be used for finding such a spanning subgraph with  $|V| + 1$  edges for the 2-corner truncated rectangular grid graph for either (any number  $k$ , and any odd numbers  $m$ ,  $n$  and  $l$ ) or (any number  $l$ , and any odd numbers  $m$ ,  $n$  and  $k$ ). In Figure 3(b) we show a spanning 2-connected subgraph with  $|V| + 1$  edges for  $R(11, 13)^{-2(2,2)}$ . The pattern in Figure 3(b) can be used for finding a spanning 2-connected subgraph with  $|V| + 1$  edges for the 2-corner truncated rectangular grid graph for any even numbers  $k$  and  $l$  and any odd numbers  $m$  and  $n$ . In Figure 3(c) we show a spanning 2-connected subgraph with  $|V| + 2$  edges for  $R(10, 12)^{-2(3,3)}$ . The pattern in this figure can be used for finding a spanning 2-connected subgraph with  $|V| + 2$  edges for the 2-corner truncated rectangular grid graph for any even numbers  $m$ ,  $n$ , and any odd numbers  $k$  and  $l$ . This is the



optimum value for the minimum number of edges in such a spanning 2-connected subgraph.




Figure 4: A Hamilton cycle for : (a)  $R(11, 12)^{-3(3,2)}$  (b)  $R(11, 12)^{-3(2,3)}$

Hamilton cycles for  $R(11, 12)^{-3(3,2)}$  and  $R(11, 12)^{-3(2,3)}$  are shown in Figure 4. The pattern in Figure 4(a) can be used for finding a Hamilton cycle for the 3-corner truncated rectangular grid graph for either (any numbers  $m$  and  $k$ , and any even numbers  $n$  and  $l$ ) or (any numbers  $n$  and  $l$ , and any even numbers  $m$  and  $k$ ). The pattern in Figure 4(b) can be used for finding a Hamilton cycle for the 3-corner truncated rectangular grid graph for either (any even numbers  $n$  and  $k$ , and any odd numbers  $m$  and  $l$ ) or (any even numbers  $m$  and  $l$ , and any odd numbers  $n$  and  $k$ ).




Figure 5: A spanning 2-connected subgraph for : (a)  $R(10, 12)^{-3(3,3)}$  with  $|V| + 1$  edges (b)  $R(11, 12)^{-3(3,3)}$  with  $|V| + 1$  edges (c)  $R(11, 13)^{-3(2,2)}$  with  $|V| + 1$  edges (d)  $R(11, 13)^{-3(2,3)}$  with  $|V| + 1$  edges (e)  $R(11, 13)^{-3(3,3)}$  with  $|V| + 2$  edges

In Figure 5(a), Figure 5(b), Figure 5(c) and Figure 5(d) we show spanning 2-connected subgraphs with  $|V| + 1$  edges for 3-corner truncated rectangular grid graphs  $R(10, 12)^{-3(3,3)}$ ,  $R(11, 12)^{-3(3,3)}$ ,  $R(11, 13)^{-3(2,2)}$  and  $R(11, 13)^{-3(2,3)}$ , respectively. The pattern in Figure 5(a) can be used for finding such a spanning 2-connected subgraph for any even numbers  $m$  and  $n$ , and any odd numbers  $k$  and  $l$ . The pattern in Figure 5(b) can be used for finding such a spanning 2-connected subgraph for either (any even number  $n$ , and any odd numbers  $m$ ,  $k$  and  $l$ ) or (any even number  $m$ , and any odd numbers  $n$ ,  $k$  and  $l$ ). The pattern in Figure 5(c) can be used for finding such a spanning 2-connected subgraph for any even numbers  $k$  and  $l$ , and any odd numbers  $m$  and  $n$ . The pattern in Figure 5(d) can be used for

finding such a spanning 2-connected subgraph for either (any even number  $k$ , and any odd numbers  $m$ ,  $n$  and  $l$ ) or (any even number  $l$ , and any odd numbers  $m$ ,  $n$  and  $k$ ). Finally, in Figure 5(e) we show a spanning 2-connected subgraph with  $|V| + 2$  edges for  $R(11, 13)^{-3(3,3)}$ . The pattern in this last figure can be used for finding a spanning 2-connected subgraph with  $|V| + 2$  edges for the 3-corner truncated rectangular grid graph for any odd numbers  $m$ ,  $n$ ,  $k$  and  $l$ . This is the optimum value for the minimum number of edges in such a spanning 2-connected subgraph.

## References

- [1] E.J. Grinberg, Plane homogeneous graphs of degree three without hamiltonian circuits, *Latvian Math. Yearbook* **5** (1968), 51–58.
- [2] A. Itai, C.H. Papadimitriou, and J.L. Szwarcfiter, Hamilton paths in grid graphs, *SIAM Journal of Computing* **11** (4) (1982), 676–686.
- [3] A. N. M. SALMAN, E. T. BASKORO AND H. J. BROERSMA, A note concerning Hamilton cycles in some classes of grid graphs, *Proceedings ITB Sains dan Teknologi* **35A** (1) (2003), 65–70.
- [4] A.N.M. Salman, E.T. Baskoro and H.J. Broersma, Spanning 2-connected subgraphs in truncated rectangular grid graphs, *Journal of Indonesian Mathematical Society* **8** (4) (2002), 63–68.
- [5] A.N.M. Salman, E.T. Baskoro and H.J. Broersma, Spanning 2-connected subgraphs in alphabet graphs, special classes of grid graphs, *Accepted for Journal of Automata, Languages and Combinatorics* (2003).
- [6] A.N.M. Salman, E.T. Baskoro and H.J. Broersma, More on spanning 2-connected subgraphs in truncated rectangular grid graphs, *Accepted for Bulletin of the Institute of Combinatorics and Its Application* (2002).
- [7] A.N.M. Salman, H.J. Broersma and C.A. Rodger, More on spanning 2-connected subgraphs of alphabet graphs, special classes of grid graphs, *Preprint*, (2002).
- [8] S. Sheffield, Computing and sampling restricted vertex degree subgraphs and hamiltonian cycles, [www.arXiv.org/abs/math.CO.0008231](http://www.arXiv.org/abs/math.CO.0008231) (2001).
- [9] C. Umans and W. Lenhart, Hamiltonian cycles in solid grid graphs, *Proceedings of the 38th Annual Symposium on Foundations of Computer Science* (1997).