

AVERAGE RESISTANCE OF TOROIDAL GRAPHS*

WILBERT SAMUEL ROSSI[†], PAOLO FRASCA[†], AND FABIO FAGNANI[‡]

Abstract. The average effective resistance of a graph is a relevant performance index in many applications, including distributed estimation and control of network systems. In this paper, we study how the average resistance depends on the graph topology and specifically on the dimension of the graph. We concentrate on d -dimensional toroidal grids, and we exploit the connection between resistance and Laplacian eigenvalues. Our analysis provides tight estimates of the average resistance, which are key to studying its asymptotic behavior when the number of nodes grows to infinity. In dimension two, the average resistance diverges: in this case, we are able to capture its rate of growth when the sides of the grid grow at different rates. In higher dimensions, the average resistance is bounded uniformly in the number of nodes: in this case, we conjecture that its value is of order $1/d$ for large d . We prove this fact for hypercubes and when the side lengths go to infinity.

Key words. effective resistance, graph dimension, consensus, relative estimation, large graphs

AMS subject classifications. 93E10, 05C50, 94C15

DOI. 10.1137/130936117

1. Introduction. The effective resistance between nodes of a graph is a classical fundamental concept that naturally comes up when the graph is interpreted as an electrical network. For several decades, it has been known to play a key role in the theory of time-reversible Markov chains, because of its connections with escape probabilities and commute times [11, 10, 1, 20]. More generally, the notion of effective resistance has broad application in science: in chemistry, for instance, the total effective resistance (summed over all pairs of nodes) is known as the Kirchhoff index of the graph, where the graph of interest has the atoms as nodes and their bonds as edges. This classical index is linked to the properties of organic macromolecules [6] and to the vibrational energy of the atoms: the latter property has also been interpreted as a measure of vulnerability in complex networks [13].

Effective resistance in network systems. Recently, the average effective resistance of a graph has appeared as an important performance index in several network-oriented problems of control and estimation, where the nodes (or agents) collectively need to obtain estimates of given quantities with limited communication effort. One instance is the *consensus* problem, where a set of agents, each with a scalar value, has the goal of reaching a common state that is a weighted average of the initial values. This problem can be solved by a simple linear iterative algorithm, which has become very popular. The performance of this algorithm depends on the graph representing the communication between the agents, and the average effective resistance of this graph plays a key role [8, 14, 21]. Indeed, the average resistance determines both the convergence speed during the transient [17, section 3.4], [15] and the robustness against additive noise affecting the updates [28]: in the latter case, the

*Received by the editors September 9, 2013; accepted for publication (in revised form) June 15, 2015; published electronically August 18, 2015. This work has been partly supported by the Italian Ministry MIUR under grant PRIN-20087W5P2K.

[†]<http://www.siam.org/journals/sicon/53-4/93611.html>

[‡]Department of Applied Mathematics, University of Twente, 7500 AE Enschede, The Netherlands (w.s.rossi@utwente.nl, p.frasca@utwente.nl).

[‡]Dipartimento di Scienze Matematiche, Politecnico di Torino, corso Duca degli Abruzzi 24, 10129 Torino, Italy (fabio.fagnani@polito.it).

effective resistance of the graph is proportional to the mean deviation of the states from their average when time goes to infinity. Similar issues of robustness to disturbances for network systems, such as platooning of vehicles, have attracted much interest [2].

Another relevant problem is the *relative estimation* problem: each node is endowed with a value and these values have to be estimated by using noisy measurements of differences taken along the available edges. The expected error of the least-squares estimator is proportional to the average effective resistance of the graph [4]. This estimation problem arises in several applications, ranging from clock synchronization [12, 16] to self-localization of mobile robotic networks [5] and to statistical ranking from pairwise comparisons [19, 23]. Several distributed algorithms that solve the relative estimation problem have been recently studied [3, 25, 26, 9, 24].

In all the above situations, performance improves when the effective resistance is reduced. This observation motivates, for instance, the problem of allocating edge weights on the edges of given graph in order to minimize the average effective resistance [17]. Similarly, it motivates our interest in topologies ensuring small average resistance. More precisely, we consider families of graphs, and we ask whether the average resistance depends gracefully on the size.

Effective resistance and graph dimension. As we have argued, the average effective resistance of a graph is a relevant index in several problems. When one tries to understand the dependence of this index on the topology, it comes out that the notion of *dimension* of the graph plays an essential role. It is well known [3, 2] that in grid-like graphs of dimension d and size N (the cardinality of the set of vertices), the average effective resistance R_{ave} scales¹ in $N \rightarrow +\infty$ (and fixed d) as follows:

$$R_{\text{ave}} = \begin{cases} \Theta(N), & d = 1, \\ \Theta(\ln N), & d = 2, \\ \Theta(1), & d \geq 3. \end{cases}$$

Notwithstanding the history and the recent popularity of this problem, no estimate of the constants involved is available in the literature (except for the case $d = 1$). Especially significant is the lack of this information when $d \geq 3$, because it is not clear, in particular, what the behavior is of R_{ave} as a function of d and for $d \rightarrow +\infty$.

In this paper, we concentrate on regular grids constructed on d -dimensional tori as a benchmark example. Their interest is motivated by the ability to intuitively capture the notion of dimension and by their nice mathematical properties: recent applications in network systems include [18, 7, 8, 15, 2]. On such toroidal grids, we sharpen the above statements. First, in dimension $d = 2$, we compute the asymptotic proportionality constant and provide tight estimates that allow us to study the asymptotic behavior when the grid sides are unequal. Second, in toroidal grids with $d \geq 3$, we show that, when the side lengths tend to infinity, the average effective resistance is of order $1/d$. In fact, we conjecture that the order $1/d$ is valid for finite side lengths too.

Our analysis hinges on two facts: first, the average effective resistance can be computed using the eigenvalues of the Laplacian matrix associated to the graph; second, an explicit formula is available for the Laplacian eigenvalues of toroidal d -grids. Similar approaches have been taken elsewhere in the literature, namely, in [29]

¹Given two sequences $f, g : \mathbb{N} \rightarrow \mathbb{R}^+$, let $\ell^+ = \limsup_n f(n)/g(n)$ and $\ell^- = \liminf_n f(n)/g(n)$. We write that $f = O(g)$ when $\ell^+ < +\infty$; that $f = o(g)$ when $\ell^+ = 0$; that $f \sim g$ when $\ell^+ = \ell^- = 1$; and $f = \Theta(g)$ when $\ell^+, \ell^- \in (0, +\infty)$. Finally, we write $f = \Omega(g)$ when $g = O(f)$.

and in [2]. The paper [29] computes the effective resistances between pairs of nodes in d -dimensional grids by explicit formulas. Our work, instead, concerns estimates of average effective resistances in toroidal grids and their asymptotics for large N . The paper [2] also estimates the average resistance for large N : in comparison, the novelty of our work resides in more accurate estimates of the quantities involved, which are essential to capture the features of high-dimensional and irregular grids.

Paper structure. The rest of this paper is organized as follows. In section 2, we formally state our problem and present and discuss our main results. Their detailed derivation is provided in section 3, which also contains a mean-field approximation of the average resistance in dimension d . Finally, in section 4, we draw some conclusions about our work and future research.

2. Problem statement and main results. We consider an undirected graph $G = (V, E)$, where V is a finite set of vertices and E is a subset of unordered pairs of distinct elements of V called edges. We assume the graph to be connected and think of it as an electrical network with all edges having unit resistance. Given two distinct vertices $u, v \in V$, the effective resistance between u and v is defined as follows. Let there be a unit input current at node u and a unit output current at node v : using Ohm's and Kirchoff's laws, a potential W is then uniquely defined at every node (up to translation constants). We then define the effective resistance as $R_{\text{eff}}(u, v) := W_u - W_v$. Consequently, the average effective resistance of G is defined as

$$(2.1) \quad R_{\text{ave}}(G) := \frac{1}{2N^2} \sum_{u, v \in V} R_{\text{eff}}(u, v),$$

where $N = |V|$ is the size of the graph.

2.1. Toroidal d -dimensional grids. We now formally define the class of graphs we deal with. Consider the cyclic group \mathbb{Z}_M of integers modulo M and the product group $\mathbb{Z}_{M_1} \times \cdots \times \mathbb{Z}_{M_d}$. Let $e_j \in \mathbb{Z}_{M_1} \times \cdots \times \mathbb{Z}_{M_d}$ be the vector with all 0's except 1 in position j and define $S = \{\pm e_j \mid j = 1, \dots, d\}$. We define as the toroidal d -grid over $\mathbb{Z}_{M_1} \times \cdots \times \mathbb{Z}_{M_d}$ the graph $T_{M_1, \dots, M_d} = (\mathbb{Z}_{M_1} \times \cdots \times \mathbb{Z}_{M_d}, E_{M_1, \dots, M_d})$, where

$$E_{M_1, \dots, M_d} := \{\{(x_1, \dots, x_d), (y_1, \dots, y_d)\} \mid (x_1 - y_1, \dots, x_d - y_d) \in S\}.$$

In other words, we call toroidal d -grids those graphs where the vertexes are arranged on a Cartesian lattice in d dimensions, which has sides of length M_1, \dots, M_d and has edges between any vertex and its $2d$ nearest neighbors, with periodic boundary conditions. The total size of the graph is $N = M_1 \times \cdots \times M_d$. In the special case $M_1 = \cdots = M_d$, i.e., when all the M_i are equal to a specific M , we will use the notation T_{M^d} instead of $T_{M, \dots, M}$. In the special case when $M = 2$, we actually obtain degenerate grids on \mathbb{Z}_2^d , which are called hypercubes of dimension d and denoted by H_d : note that the size of H_d is $N = 2^d$ and the degree of each vertex is d .

2.2. Asymptotic results. We start by recalling the simple case $d = 1$, where the effective resistance can be directly computed. From the standard properties of series and parallel connections of resistors [20, pp. 119–120], one can see that $R_{\text{eff}}(v_0, v_0 + l) = \frac{l(M-l)}{M}$ and thus

$$(2.2) \quad R_{\text{ave}}(T_M) = \frac{1}{2M} \sum_{l=1}^{M-1} \frac{l(M-l)}{M} = \frac{M}{12} - \frac{1}{12M}.$$




FIG. 1. Left: R_{ave} in low-dimensional toroidal grids, as function of the size $N = M^d$, with the dashed lines representing the asymptotic trends $N/12$ and $\frac{1}{4\pi} \log N$. Right: R_{ave} in high-dimensional toroidal grids, as function of the dimension d , with the dashed line representing the trend $\frac{1}{2d}$.

This formula leads to the asymptotic relation

$$R_{\text{ave}}(T_M) \sim \frac{M}{12} \quad \text{for } M \rightarrow +\infty.$$

When $d \geq 2$, we prove in this paper that the following asymptotic relations hold.

THEOREM 2.1 (asymptotics). *Let T_{M^d} be the toroidal grid in $d \geq 2$ dimensions, with each side length being equal to M , and let $R_{\text{ave}}(T_{M^d})$ be its average effective resistance. Then,*

$$(2.3) \quad R_{\text{ave}}(T_{M^2}) \sim \frac{1}{2\pi} \ln M \quad \text{for } M \rightarrow +\infty$$

and

$$(2.4) \quad \lim_{M \rightarrow +\infty} R_{\text{ave}}(T_{M^d}) = \Theta\left(\frac{1}{d}\right) \quad \text{for } d \rightarrow +\infty.$$

The relations (2.3) and (2.4) follow immediately from the estimates provided below in Theorems 2.3 and 2.4. Furthermore, we conjecture that the statement (2.4) can be sharpened as follows.

CONJECTURE 2.2.

$$R_{\text{ave}}(T_{M^d}) = \Theta\left(\frac{1}{d}\right) \quad \text{for } d \rightarrow +\infty, M \text{ fixed.}$$

At the moment, we can only prove such a result in the degenerate case $M = 2$, corresponding to a hypercube, where

$$(2.5) \quad R_{\text{ave}}(H_d) \sim \frac{1}{d} \quad \text{as } d \rightarrow \infty.$$

Our results and conjecture are corroborated by numerical experiments, which are summarized in Figure 1. The left plot of Figure 1 shows the average effective resistances R_{ave} of four families of low-dimensional graphs as functions of the total size N of the graphs: $R_{\text{ave}}(T_M)$ and $R_{\text{ave}}(T_{M^2})$ follow the predicted linear (2.2) and

logarithmic (2.3) asymptotic trends, whereas $R_{\text{ave}}(T_{M^3})$ and $R_{\text{ave}}(T_{M^4})$ tend to a finite limit. The right plot of Figure 1 instead regards high-dimensional graphs and shows that R_{ave} decreases with d , when the side lengths M are kept fixed. If $d \geq 5$, then $R_{\text{ave}}(T_{M^d})$ for different M are roughly equal and inversely proportional to $2d$. This plot supports our conjecture that $R_{\text{ave}}(T_{M^d})$ is of order $1/d$, independent of M .

2.3. Estimates for finite toroidal grids. This subsection contains tight estimates of the average resistance in dimension d . These novel results are key to obtaining the asymptotic relations presented above. We begin with a pair of estimates in dimension two.

THEOREM 2.3 (torus T_{M_1, M_2}). *Let T_{M_1, M_2} be the toroidal grid in two dimensions with side lengths M_1 and M_2 , and let $R_{\text{ave}}(T_{M_1, M_2})$ be its average effective resistance. Suppose $4 \leq M_1 \leq M_2$. Then,*

$$\begin{aligned} R_{\text{ave}}(T_{M_1, M_2}) &\leq \frac{1}{2\pi} \log M_2 + \frac{1}{12} \frac{M_2}{M_1} + 1, \\ R_{\text{ave}}(T_{M_1, M_2}) &\geq \max \left\{ \frac{1}{12} \frac{M_2}{M_1} - \frac{1}{24}; \frac{1}{2\pi} \log M_1 - \frac{1}{12} \frac{M_2}{M_1} - \frac{1}{2} \right\}. \end{aligned}$$

In order to understand the consequences of Theorem 2.3, it is useful to fix specific relations between M_2 and M_1 and study the asymptotic behavior when the size $N = M_1 \times M_2$ of the graph tends to infinity. Preliminarily, we observe that in the lower bound of Theorem 2.3, the former expression dominates when M_1 and M_2 grow with different rates, while the latter dominates when M_1 and M_2 have the same rate of growth. We then consider the following three relations between M_1 and M_2 :

1. $M_1 = c$, $M_2 = N/c$. Then,

$$\frac{1}{12} \frac{N}{c^2} - \frac{1}{24} \leq R_{\text{ave}}(T_{c, N/c}) \leq \frac{1}{12} \frac{N}{c^2} + \frac{1}{2\pi} \log N + 1.$$

In this case, $R_{\text{ave}}(T_{c, N/c}) \sim \frac{N}{12c^2}$ as $N \rightarrow +\infty$: we may interpret this linear growth as reminiscent of the one-dimensional case.

2. $M_1 = \sqrt[3]{N}$, $M_2 = \sqrt[3]{N^{c-1}}$ with $c > 2$. Then,

$$\frac{1}{12} N^{\frac{c-2}{c}} - \frac{1}{24} \leq R_{\text{ave}}(T_{\sqrt[3]{N}, \sqrt[3]{N^{c-1}}}) \leq \frac{1}{12} N^{\frac{c-2}{c}} + \frac{1}{2\pi} \frac{c-1}{c} \log N + 1.$$

In this case, $R_{\text{ave}}(T_{\sqrt[3]{N}, \sqrt[3]{N^{c-1}}}) \sim N^{\frac{c-2}{c}}/12$ as $N \rightarrow +\infty$, which is sublinear and proportional to the ratio between M_2 and M_1 .

3. $M_1 = \sqrt{N/c}$, $M_2 = \sqrt{cN}$ with $c = \frac{M_2}{M_1}$. Then,

$$\frac{1}{4\pi} \log N - \frac{\log c}{4\pi} - \frac{c}{12} - \frac{1}{2} \leq R_{\text{ave}}(T_{\sqrt{N/c}, \sqrt{cN}}) \leq \frac{1}{4\pi} \log N + \frac{c}{12} + \frac{\log c}{4\pi} + 1.$$

In this case, $R_{\text{ave}}(T_{\sqrt{N/c}, \sqrt{cN}}) \sim \frac{1}{4\pi} \log N$ as $N \rightarrow +\infty$. That is, taking M_1 proportional to M_2 makes $R_{\text{ave}}(T_{M_1, M_2})$ grow logarithmically with N : this order of growth must be contrasted against the linear growth that characterizes one-dimensional graphs and against the two previous examples. In fact, this is the lowest asymptotic average effective resistance reachable by a bidimensional toroidal grid.

Next, we provide a pair of bounds valid when $d \geq 3$: for simplicity, we assume that the lengths along each of the d dimensions are all equal to M .

THEOREM 2.4 (torus T_{M^d}). *Let T_{M^d} be the toroidal grid in $d \geq 3$ dimensions, with each side length being equal to M , and let $R_{\text{ave}}(T_{M^d})$ be its average effective*

resistance. Provided $M \geq 4$, it holds that

$$\begin{aligned} R_{\text{ave}}(T_{M^d}) &\leq \frac{8}{d+1} \left(1 + \frac{1}{M}\right)^{d+1} + \frac{d}{4M^{d-2}} \left(\frac{1}{3} + \frac{(d-1)\log M}{\pi}\right), \\ R_{\text{ave}}(T_{M^d}) &\geq \frac{1}{4d}. \end{aligned}$$

Notice that if $d \geq 3$ is fixed and M diverges, then Theorem 2.4 yields $R_{\text{ave}}(T_{M^d}) = \Theta(1)$ as $M \rightarrow +\infty$. This fact is well known: the difficulty here lies in finding a tight upper bound, which can reveal the dependence on d and imply (2.4).

We conclude the presentation of our main results with the relevant estimates for the hypercube, corresponding to the case $M = 2$.

THEOREM 2.5 (hypercube). *Let H_d be a d -dimensional hypercube graph and $R_{\text{ave}}(H_d)$ be its average effective resistance. When $d \geq 2$, the following estimates hold:*

$$\frac{1}{2} \frac{1}{d+1} \leq R_{\text{ave}}(H_d) \leq \frac{2}{d+1}.$$

3. Resistance and eigenvalues. We have seen in the previous section that the average effective resistance of the one-dimensional ring graph can be computed from the effective resistance between any pair of nodes. Indeed, in that case, effective resistances can be directly computed using simple properties of electrical networks. However, this approach is not viable for d -dimensional tori with $d \geq 2$. Instead, we can rely on the fact that for any graph $R_{\text{ave}}(G)$ can be expressed in terms of its Laplacian eigenvalues. Given a graph G , the Laplacian of G , $L(G) \in \mathbb{R}^{V \times V}$ is the matrix defined by

$$L(G)_{uu} = |\{v \in V \mid \{u, v\} \in E\}|, \quad L(G)_{uv} = \begin{cases} -1 & \text{if } \{u, v\} \in E, \\ 0 & \text{otherwise,} \end{cases} \quad u \neq v.$$

It is well known that its eigenvalues can be ordered to satisfy $0 = \lambda_1 < \lambda_2 \leq \dots \leq \lambda_N$ and the following relation holds true [17, eq. (15)]:

$$(3.1) \quad R_{\text{ave}}(G) = \frac{1}{N} \sum_{i \geq 2} \frac{1}{\lambda_i}.$$

We are going to use (3.1) in order to prove our results.² Indeed, the eigenvalues of the Laplacian can be exactly computed for the toroidal grid T_{M_1, \dots, M_d} using a discrete Fourier transform [15]

$$(3.2) \quad \lambda_{\mathbf{h}} = \lambda_{h_1, \dots, h_d} = 2d - 2 \sum_{i=1}^d \cos \frac{2\pi h_i}{M_i}, \quad \mathbf{h} = (h_1, \dots, h_d) \in \mathbb{Z}_{M_1} \times \dots \times \mathbb{Z}_{M_d}.$$

This formula leads to the key expression

$$(3.3) \quad R_{\text{ave}}(T_{M_1, \dots, M_d}) = \frac{1}{M_1 \cdots M_d} \sum_{\mathbf{h} \neq \mathbf{0}} \frac{1}{2d - 2 \sum_{i=1}^d \cos \left(\frac{2\pi h_i}{M_i} \right)},$$

on which most of our derivations are based (excluding section 3.4).

²Note that using the Laplacian eigenvalues and eigenvectors, it is possible to compute the effective resistance between any pair of nodes [29, eq. (11)]: $R_{\text{eff}}(v, u) = \sum_{i \geq 2} \frac{1}{\lambda_i} |\psi_i(v) - \psi_i(u)|^2$, where $\psi_i(v)$ is the component v of the eigenvector associated to the eigenvalue λ_i of the Laplacian of G . Actually, from this formula and the definition of $R_{\text{ave}}(G)$, one easily deduces (3.1), which only requires the knowledge of the eigenvalues.

3.1. Bounds for the 2-torus T_{M_1, M_2} . We provide here the proof of Theorem 2.3. As explained before, we resort to the Laplacian eigenvalues, which for T_{M_1, M_2} read $\lambda_{i,j} = 4 - 2 \cos(2\pi i/M_1) - 2 \cos(2\pi j/M_2)$ with $i \in \{0, \dots, M_1 - 1\}$ and $j \in \{0, \dots, M_2 - 1\}$. Hence,

$$R_{\text{ave}}(T_{M_1, M_2}) = \frac{1}{M_1 M_2} \sum_{(i,j) \neq \mathbf{0}} \frac{1}{4 - 2 \cos(2\pi i/M_1) - 2 \cos(2\pi j/M_2)}.$$

In order to estimate this quantity, we are going to interpret certain partial sums as upper/lower Riemann sums of suitable integrals, similarly to what is done in [2]. However, it will be essential to single out some “one-dimensional” contributions to the overall sum. To this goal, we recall that

$$R_{\text{ave}}(T_M) = \frac{1}{M} \sum_{i \geq 1} \frac{1}{2 - 2 \cos(2\pi i/M)},$$

since the eigenvalues of T_M are $\lambda_i = 2 - 2 \cos(2\pi i/M)$ with $i \in \{0, \dots, M - 1\}$.

Proof of Theorem 2.3. In order to prove the upper bound, we rewrite $R_{\text{ave}}(T_{M_1, M_2})$ as

$$(3.4) \quad R_{\text{ave}}(T_{M_1, M_2}) = \frac{1}{M_2} R_{\text{ave}}(T_{M_1}) + \frac{1}{M_1} R_{\text{ave}}(T_{M_2}) + \mathring{R}_{\text{ave}}(T_{M_1, M_2}),$$

where

$$\mathring{R}_{\text{ave}}(T_{M_1, M_2}) = \frac{1}{M_1 M_2} \sum_{i \neq 0} \sum_{j \neq 0} \frac{1}{\lambda_{i,j}}.$$

The first two terms in (3.4) are easily bounded with the explicit formula (2.2):

$$(3.5) \quad \frac{1}{M_2} R_{\text{ave}}(T_{M_1}) + \frac{1}{M_1} R_{\text{ave}}(T_{M_2}) \leq \frac{M_1}{12M_2} + \frac{M_2}{12M_1}.$$

Concerning $\mathring{R}_{\text{ave}}(T_{M_1, M_2})$, by symmetry it holds that

$$\mathring{R}_{\text{ave}}(T_{M_1, M_2}) = \frac{1}{M_1 M_2} \sum_{i=1}^{M_1-1} \sum_{j=1}^{M_2-1} \frac{1}{\lambda_{i,j}} \leq \frac{4}{M_1 M_2} \sum_{i=1}^{\lfloor M_1/2 \rfloor} \sum_{j=1}^{\lfloor M_2/2 \rfloor} \frac{1}{\lambda_{i,j}}.$$

Consider the function

$$(3.6) \quad f(x, y) = \frac{1}{4 - 2 \cos(2\pi x) - 2 \cos(2\pi y)}$$

and notice that $\frac{1}{\lambda_{i,j}} = f(\frac{i}{M_1}, \frac{j}{M_2})$. For a fixed \bar{y} , f is decreasing for $x \in (0, 1/2]$, and vice versa for fixed \bar{x} , f is decreasing for $y \in (0, 1/2]$. It follows that, for each pair i, j with $1 \leq i \leq \lfloor M_1/2 \rfloor$ and $1 \leq j \leq \lfloor M_2/2 \rfloor$,

$$\frac{1}{M_1 M_2} \frac{1}{\lambda_{i,j}} \leq \int_{\frac{i-1}{M_2}}^{\frac{i}{M_2}} \int_{\frac{j-1}{M_1}}^{\frac{j}{M_1}} f(x, y) \, dx \, dy.$$

Define the region $D = [0, 1/2] \times [0, 1/2]$ and $D^* = D \setminus ([0, 1/M_1] \times [0, 1/M_2])$ as in




FIG. 2. The regions D , D^* , and C , which are useful in the proof of the upper bound of Theorem 2.3.

Figure 2 (left) to estimate

$$(3.7) \quad \begin{aligned} \mathring{R}_{\text{ave}}(T_{M_1, M_2}) &= \frac{4}{M_1 M_2} \sum_{i=1}^{\lfloor M_1/2 \rfloor} \sum_{j=1}^{\lfloor M_2/2 \rfloor} \frac{1}{\lambda_{i,j}} \\ &\leq \frac{4}{M_1 M_2} f\left(\frac{1}{M_1}, \frac{1}{M_2}\right) + 4 \iint_{D^*} f(x, y) dx dy. \end{aligned}$$

The term for $i = 1, j = 1$ is kept aside, because of the singularity in the origin. Next, instead of computing the integral in (3.7) in closed form, we observe that

$$\begin{aligned} f(x, y) &= \frac{1}{4 - 2 \cos(2\pi x) - 2 \cos(2\pi y)} \\ &\leq \frac{1}{(2\pi x)^2 + (2\pi y)^2 - \frac{(2\pi x)^4}{12} - \frac{(2\pi y)^4}{12}} \\ &\leq \frac{1}{(2\pi)^2(x^2 + y^2) - \frac{(2\pi)^4}{12}(x^2 + y^2)^2} = g(\sqrt{x^2 + y^2}), \end{aligned}$$

where we defined the function $g : (0, \frac{\sqrt{3}}{\pi}) \rightarrow \mathbb{R}^+$ as

$$(3.8) \quad g(r) = \frac{1}{4\pi^2 r^2 \left(1 - \frac{\pi^2}{3} r^2\right)}.$$

Unfortunately, g does not provide an useful upper bound because it has a singularity in $\frac{\sqrt{3}}{\pi}$. We instead use the continuous modification

$$\tilde{g}(\rho) = \begin{cases} \frac{1}{4\pi^2 \rho^2 \left(1 - \frac{\pi^2}{3} \rho^2\right)} & \text{if } 0 < \rho < \frac{1}{2}, \\ \frac{1}{\pi^2 \left(1 - \frac{\pi^2}{12}\right)} & \text{if } \rho \geq \frac{1}{2}, \end{cases}$$

which is decreasing in $(0, \frac{\sqrt{3}}{\sqrt{2}\pi})$ and such that $f(x, y) \leq \tilde{g}(\sqrt{x^2 + y^2})$ for all $(x, y) \in D$. We now use this bound to estimate the right-hand side of (3.7). Regarding the first term, using that $M_2 \geq M_1 \geq 4$, we obtain

$$(3.9) \quad \frac{4}{M_1 M_2} \tilde{g}\left(\sqrt{\frac{1}{M_2^2} + \frac{1}{M_1^2}}\right) \leq \frac{4}{M_1 M_2} \tilde{g}(1/M_1) \leq \frac{2}{\pi^2} \frac{M_1}{M_2}.$$

On the other hand, defining $C = \{(x, y) \in \mathbb{R}^2 : \frac{1}{M_2^2} \leq x^2 + y^2 \leq \frac{1}{4}\}$ as illustrated in Figure 2 (right), we can estimate the second term with polar coordinates:

$$\begin{aligned}
4 \iint_{D^*} f(x, y) dx dy &= 4 \iint_{D^*} \tilde{g}(\rho) \rho d\rho d\theta \\
&\leq 4 \iint_C \tilde{g}(\rho) \rho d\rho d\theta + 4 \iint_{D^* \setminus C} \tilde{g}(\rho) \rho d\rho d\theta \\
&\leq 4 \int_0^{\frac{\pi}{2}} \int_{\frac{1}{M_2}}^{\frac{1}{2}} \frac{1}{4\pi^2 \rho^2 (1 - \frac{\pi^2}{3} \rho^2)} \rho d\rho d\theta + \left(1 - \frac{\pi}{4}\right) \tilde{g}\left(\frac{1}{2}\right) \\
&\leq \frac{2}{\pi^2} \frac{M_1}{M_2} + \frac{1}{2\pi} \int_{\frac{1}{M_2}}^{1/2} \frac{1}{\rho - \frac{\pi^2}{3} \rho^3} d\rho + \frac{1}{6} \\
&= \frac{1}{2\pi} \left[\log \rho - \frac{1}{2} \log \left(1 - \frac{\pi}{3} \rho^2\right) \right]_{\frac{1}{M_2}}^{1/2} + \frac{1}{6} \\
&\leq \frac{1}{2\pi} \log M_2 - \frac{1}{4\pi} \log \left(1 - \frac{\pi}{12}\right) + \frac{1}{6} \\
(3.10) \quad &\leq \frac{1}{2\pi} \log M_2 + \frac{1}{5}.
\end{aligned}$$

Using bounds (3.9) and (3.10) in (3.7), we obtain

$$(3.11) \quad \dot{R}_{\text{ave}}(T_{M_1, M_2}) \leq \frac{1}{2\pi} \log M_2 + \frac{2}{\pi^2} \frac{M_1}{M_2} + \frac{1}{5}.$$

Now using (3.11) and (3.5) in (3.4), we finally get

$$R_{\text{ave}}(T_{M_1, M_2}) \leq \frac{1}{2\pi} \log M_2 + \frac{M_2}{12M_1} + \left(\frac{2}{\pi^2} + \frac{1}{12}\right) \frac{M_1}{M_2} + \frac{1}{5},$$

and the thesis follows since $\frac{M_1}{M_2} \leq 1$.

The first estimate of the lower bound can be proved easily: it is enough to neglect in the expression of $R_{\text{ave}}(T_{M_1, M_2})$ all terms that have $i > 0$ or $j > 0$. Then,

$$\begin{aligned}
R_{\text{ave}}(T_{M_1, M_2}) &\geq \frac{1}{M_2} R_{\text{ave}}(T_{M_1}) + \frac{1}{M_1} R_{\text{ave}}(T_{M_2}) \\
&= \frac{1}{M_2} \left(\frac{M_1}{12} - \frac{1}{12M_1} \right) + \frac{1}{M_1} \left(\frac{M_2}{12} - \frac{1}{12M_2} \right) \\
&\geq \frac{1}{12} \left(\frac{M_2}{M_1} + \frac{M_1}{M_2} \right) - \frac{1}{6M_2 M_1} \geq \frac{1}{12} \frac{M_2}{M_1} - \frac{1}{24}.
\end{aligned}$$

To prove the second estimate, we use an approach similar to that of the upper bound. Since a symmetric domain is convenient, we define the index sets

$$\begin{aligned}
\Gamma_+ &= \mathbb{Z}_{M_1} \times \mathbb{Z}_{M_2} \setminus \{(0, 0)\}, \\
\Gamma^+ &= \Gamma_+ \cup \{M_1\} \times \{1, 2, \dots, M_2 - 1\} \cup \{1, 2, \dots, M_1 - 1\} \times \{M_2\}
\end{aligned}$$

to write

$$(3.12) \quad R_{\text{ave}}(T_{M_1, M_2}) = \frac{1}{M_1 M_2} \sum_{\Gamma_+} \frac{1}{\lambda_{i,j}} = \bar{R}_{\text{ave}}(T_{M_1, M_2}) - \frac{1}{M_2} R_{\text{ave}}(T_{M_1}) - \frac{1}{M_1} R_{\text{ave}}(T_{M_2}),$$




FIG. 3. Left plot: Regions E and D^* . Right plot: In order to illustrate how the Riemann sum is built, dots on the corners of the grey rectangles indicate the interpolation points, whose values are assumed on each rectangle. The contributions of the dashed parts of the rectangles are disregarded in the integral, without compromising the validity of inequality (3.13).

where $\bar{R}_{\text{ave}}(T_{M_1, M_2}) = \frac{1}{M_1 M_2} \sum_{\Gamma^+} \frac{1}{\lambda_{i,j}}$. To estimate $\bar{R}_{\text{ave}}(T_{M_1, M_2})$, we consider the function $f(x, y)$ as defined in the proof of the upper bound and the domain E , defined (Figure 3) as

$$E = [0, 1] \times [0, 1] \setminus \left(\left(\left[0, \frac{1}{M_1} \right] \cup \left[1 - \frac{1}{M_1}, 1 \right] \right) \times \left(\left[0, \frac{1}{M_2} \right] \cup \left[1 - \frac{1}{M_2}, 1 \right] \right) \right),$$

and we notice that

$$(3.13) \quad \bar{R}_{\text{ave}}(T_{M_1, M_2}) \geq \iint_E f(x, y) dx dy = 4 \iint_{D^*} f(x, y) dx dy,$$

where the equality exploits the symmetry of f . Since $f(x, y) \geq (4\pi^2)^{-1}(x^2 + y^2)^{-1}$, we obtain

$$\begin{aligned} \bar{R}_{\text{ave}}(T_{M_1, M_2}) &\geq \frac{1}{\pi^2} \iint_{D^*} \frac{1}{x^2 + y^2} dx dy \\ &\geq \frac{1}{2\pi} \int_{\delta}^{1/2} \frac{1}{\rho^2} \rho d\rho = \frac{1}{2\pi} (\log(\delta^{-1}) - \log 2) \end{aligned}$$

with $\delta = \sqrt{\frac{1}{M_1^2} + \frac{1}{M_2^2}}$. If we observe that $\frac{1}{M_1^2} + \frac{1}{M_2^2} \leq \frac{2}{M_1^2}$, we get

$$(3.14) \quad \bar{R}_{\text{ave}}(T_{M_1, M_2}) \geq \frac{1}{2\pi} \log(M_1) - \frac{1}{4}.$$

Now using (3.14) inside (3.12) together with the exact calculation (2.2), we finally obtain

$$R_{\text{ave}}(T_{M_1, M_2}) \geq \frac{1}{2\pi} \log(M_1) - \frac{M_2}{12M_1} - \frac{M_1}{12M_2} - \frac{1}{4} \geq \frac{1}{2\pi} \log(M_1) - \frac{M_2}{12M_1} - \frac{1}{2}.$$

This inequality concludes the proof of the second estimate for the lower bound and hence the proof of the theorem. \square

3.2. Continuous approximation of $R_{\text{ave}}(T_{M^d})$. We consider here the quantity $\gamma(d)$, defined as

$$(3.15) \quad \gamma(d) := \int_{[0,1]^d} \frac{1}{2d - 2 \sum_{i=1}^d \cos(2\pi x_i)} d\mathbf{x},$$

and prove an upper and lower bound of order $1/d$. In the proof of Theorem 2.4, this quantity will play the role of a “continuous” approximation of $R_{\text{ave}}(T_{M^d})$.

LEMMA 3.1. *If $d \geq 3$, then*

$$\frac{1}{4d} \leq \gamma(d) \leq \frac{4}{d}.$$

Proof. The lower bound is trivial: the integrand is not smaller than $\frac{1}{4d}$ over all the domain. What follows is devoted to proving the upper bound. By symmetry,

$$\gamma(d) = 2^d \int_{[0,\frac{1}{2}]^d} \frac{1}{2d - 2 \sum_{i=1}^d \cos(2\pi x_i)} d\mathbf{x},$$

and then we define the following three subsets of $[0, \frac{1}{2}]^d$:

$$\begin{aligned} A &= \left\{ \mathbf{x} \in \left[0, \frac{1}{2}\right]^d \quad \text{s.t.} \quad \|\mathbf{x}\|_2 \leq \frac{1}{\pi} \right\}, \\ B &= \left\{ \mathbf{x} \in \left[0, \frac{1}{2}\right]^d \quad \text{s.t.} \quad \|\mathbf{x}\|_2 \geq \frac{1}{\pi} \quad \text{and} \quad x_i \leq \frac{1}{\pi} \quad \forall i \right\}, \\ C &= \left\{ \mathbf{x} \in \left[0, \frac{1}{2}\right]^d \quad \text{s.t.} \quad \exists \quad x_i \geq \frac{1}{\pi} \right\}, \end{aligned}$$

such that $A \cup B \cup C = [0, \frac{1}{2}]^d$. Correspondingly, we define

$$\begin{aligned} \mathcal{I}_d^A &= 2^d \int_A \frac{1}{2d - 2 \sum_{i=1}^d \cos(2\pi x_i)} d\mathbf{x}, \\ \mathcal{I}_d^B &= 2^d \int_B \frac{1}{2d - 2 \sum_{i=1}^d \cos(2\pi x_i)} d\mathbf{x}, \\ \mathcal{I}_d^C &= 2^d \int_C \frac{1}{2d - 2 \sum_{i=1}^d \cos(2\pi x_i)} d\mathbf{x}, \end{aligned}$$

so that $\gamma(d) = \mathcal{I}_d^A + \mathcal{I}_d^B + \mathcal{I}_d^C$.

We begin by a bound on \mathcal{I}_d^A . First, we work on the denominator of the integrand, using the inequality $1 - \cos x \geq \frac{x^2}{2} - \frac{x^4}{24}$ to show

$$\begin{aligned} 2 \sum_{i=1}^d (1 - \cos(2\pi x_i)) &\geq 4\pi^2 \sum_{i=1}^d x_i^2 - \frac{16\pi^4}{12} \sum_{i=1}^d x_i^4 \\ &\geq 4\pi^2 \left(\sum_{i=1}^d x_i^2 - \frac{\pi^2}{3} \sum_{i=1}^d \sum_{j=1}^d x_i^2 x_j^2 \right) \\ &= 4\pi^2 \left(1 - \frac{\pi^2}{3} \sum_{i=1}^d x_i^2 \right) \sum_{i=1}^d x_i^2. \end{aligned}$$

With the last expression, in polar coordinates we obtain

$$\begin{aligned}\mathcal{I}_d^A &\leq 2^d \int_A \frac{1}{4\pi^2 \left(\sum_{i=1}^d x_i^2 \right) \left(1 - \frac{\pi^2}{3} \sum_{i=1}^d x_i^2 \right)} d\mathbf{x} \\ &= \int_0^{\frac{1}{\pi}} \frac{2\pi^{\frac{d}{2}}}{\Gamma(\frac{d}{2})} \rho^{d-1} \frac{1}{4\pi^2 \rho^2 \left(1 - \frac{\pi^2}{3} \rho^2 \right)} d\rho \\ &= \frac{\pi^{\frac{d}{2}-2}}{2\Gamma(\frac{d}{2})} \int_0^{\frac{1}{\pi}} \frac{\rho^{d-3}}{1 - \frac{\pi^2}{3} \rho^2} d\rho.\end{aligned}$$

The change of variables involving the Gamma function has cleared the singularity in zero, and the new integrand is an increasing function. Then,

$$\mathcal{I}_d^A \leq \frac{\pi^{\frac{d}{2}-2}}{2\Gamma(\frac{d}{2})} \int_0^{\frac{1}{\pi}} \frac{\left(\frac{1}{\pi}\right)^{d-3}}{\left[1 - \frac{\pi^2}{3} \left(\frac{1}{\pi}\right)^2\right]} d\rho = \frac{3}{4\pi^{\frac{d}{2}} \Gamma(\frac{d}{2})}.$$

Since $x^{(1-\gamma)x-1} < \Gamma(x)$ if $x > 1$ (see [27]), where $\gamma \simeq 0.577$ is the Euler–Mascheroni constant, we have

$$(3.16) \quad \mathcal{I}_d^A \leq \frac{3d}{8\pi^{\frac{d}{2}} \left(\frac{d}{2}\right)^{(1-\gamma)\frac{d}{2}}}.$$

Next, we estimate \mathcal{I}_d^B . Recall definition (3.6) and notice that the function

$$f(\mathbf{x}) := \frac{1}{2d - 2 \sum_{i=1}^d \cos(2\pi x_i)}$$

is decreasing in every direction i , when $\mathbf{x} \in [0, \frac{1}{2}]$. Then, defining $g(\rho)$ as in (3.8), we have

$$(3.17) \quad \mathcal{I}_d^B \leq 2^d \mu(B) g\left(\frac{1}{\pi}\right) \leq \frac{3}{8} \left(\frac{2}{\pi}\right)^d,$$

where $\mu(B)$ denotes the measure of B , and $B \subset [0, \frac{1}{\pi}]^d$.

Finally, we consider \mathcal{I}_d^C . Let $\Omega = \{0, 1\}^d$ and for all $\omega \in \Omega$, define the set $C_\omega \subset C$ as $C_\omega = \{\mathbf{x} \in C \text{ s.t. } x_i \geq \frac{1}{\pi} \text{ iff } \omega_i = 1\}$. Clearly, $\bigcup_{\omega \neq \mathbf{0}} C_\omega = C$. Then,

$$\mathcal{I}_d^C = 2^d \sum_{\omega \neq \mathbf{0}} \int_{C_\omega} \frac{1}{2d - 2 \sum_{i=1}^d \cos(2\pi x_i)} d\mathbf{x}.$$

For a fixed $\omega \in \Omega$, we denote by l_ω the number of 1's in ω (that is, the so-called Hamming weight of ω), and we notice that

$$\mu(C_\omega) = \left(\frac{1}{\pi}\right)^{d-l_\omega} \left(\frac{1}{2} - \frac{1}{\pi}\right)^{l_\omega}.$$

Moreover, the function $f(\mathbf{x})$ is symmetric under permutations of the components of \mathbf{x} . Then,

$$f(\mathbf{x}) \leq f\left(\frac{1}{\pi}\omega\right) = \frac{1}{2(1 - \cos(2))} \frac{1}{l_\omega} \quad \text{if } \mathbf{x} \in C_\omega.$$

Since clearly there are $\binom{d}{l}$ elements in Ω with Hamming weight l , we can argue that

$$\begin{aligned} \mathcal{I}_d^C &\leq 2^d \sum_{l=1}^d \binom{d}{l} \frac{1}{2l(1-\cos(2))} \left(\frac{1}{\pi}\right)^{d-l} \left(\frac{1}{2} - \frac{1}{\pi}\right)^l \\ &= \frac{1}{2(1-\cos(2))} \sum_{l=1}^d \binom{d}{l} \left(\frac{2}{\pi}\right)^{d-l} \left(1 - \frac{2}{\pi}\right)^l \frac{1}{l} \\ &\leq \frac{1}{(1-\cos(2))(1-\frac{2}{\pi})} \frac{1}{d+1}, \end{aligned}$$

where the last inequality follows from standard manipulations on the binomials. This bound can be replaced by a simpler

$$(3.18) \quad \mathcal{I}_d^C \leq \frac{3}{d},$$

and we are able to conclude the proof by combining (3.16), (3.17), and (3.18) to get $\gamma(d) = \mathcal{I}_d^A + \mathcal{I}_d^B + \mathcal{I}_d^C \leq \frac{4}{d}$. \square

3.3. Bounds for the d -torus T_{M^d} . We proceed with the proof of Theorem 2.4, containing the bounds for $R_{\text{ave}}(T_{M^d})$ when $d \geq 3$. Notice that, when all the side lengths are equal to M , the general expression (3.3) becomes

$$(3.19) \quad R_{\text{ave}}(T_{M^d}) = \frac{1}{M^d} \sum_{\mathbf{h} \neq \mathbf{0}} \frac{1}{2d - 2 \sum_{i=1}^d \cos\left(\frac{2\pi h_i}{M}\right)}.$$

Proof of Theorem 2.4. The lower bound can be easily proved by observing that for all $\mathbf{h} \neq \mathbf{0}$, $\frac{1}{\lambda_{\mathbf{h}}} \geq \frac{1}{4d}$. Moreover, since $\frac{1}{\lambda_{(1,0,\dots,0)}} = \frac{1}{2-2\cos(\frac{2\pi}{M})} \geq \frac{1}{2d}$,

$$R_{\text{ave}}(T_{M^d}) \geq \frac{1}{M^d} \left[(M^d - 2) \frac{1}{4d} + \frac{2}{4d} \right] = \frac{1}{4d}.$$

In order to prove the upper bound, let us consider the terms in the sum (3.19) for which $\mathbf{h} \succ \mathbf{0}$, i.e., those for which all $h_i > 0$. Define

$$\mathring{R}_{\text{ave}}(T_{M^d}) = \frac{1}{M^d} \sum_{\mathbf{h} \succ \mathbf{0}} \frac{1}{2d - 2 \sum_{i=1}^d \cos\left(\frac{2\pi h_i}{M}\right)}$$

(where $\mathbf{h} \succ \mathbf{0}$ means that $h_i > 0$ for all i) and observe that

$$R_{\text{ave}}(T_{M^d}) = \sum_{m=1}^d \binom{d}{m} \frac{1}{M^{d-m}} \mathring{R}_{\text{ave}}(T_{M^m}).$$

It is crucial to observe that, with $\gamma(m)$ defined at (3.15),

$$\mathring{R}_{\text{ave}}(T_{M^m}) \leq \gamma(m)$$

for any $m \geq 1$, since we can see $\mathring{R}_{\text{ave}}(T_{M^m})$ as a lower Riemann sum of the integral. When $m \geq 3$, Lemma 3.1 gives

$$\mathring{R}_{\text{ave}}(T_{M^m}) \leq \frac{4}{m},$$

while for $m = 2$ we use the bound (3.11) on $\dot{R}_{\text{ave}}(T_{M_1, M_2})$ from the proof regarding T_{M_1, M_2} . For $m = 1$, notice that $\dot{R}_{\text{ave}}(T_M) = R_{\text{ave}}(T_M)$, and hence we can use (2.2). We thus obtain

$$\begin{aligned} R_{\text{ave}}(T_{M^d}) &\leq \binom{d}{1} \frac{1}{M^{d-1}} \frac{M}{12} + \binom{d}{2} \frac{1}{M^{d-2}} \left[\frac{1}{2\pi} \log M + 1 \right] + \sum_{m=3}^d \binom{d}{m} \frac{1}{M^{d-m}} \frac{4}{m} \\ &\leq \frac{4}{M^d} \sum_{m=1}^d \binom{d}{m} M^m \frac{1}{m} + \frac{d}{4M^{d-2}} \left[\frac{1}{3} + \frac{(d-1) \log M}{\pi} \right]. \end{aligned}$$

After noting that

$$\begin{aligned} \frac{4}{M^d} \sum_{m=1}^d \binom{d}{m} M^m \frac{1}{m} &\leq \frac{4}{M^d} \sum_{m=1}^d \binom{d}{m} M^m \frac{2}{m+1} \\ &\leq \frac{8}{M^{d+1}} \sum_{m=1}^d \binom{d+1}{m+1} \frac{M^{m+1}}{d+1} \\ &\leq \frac{8}{d+1} \frac{1}{M^{d+1}} \sum_{n=0}^{d+1} \binom{d+1}{n} M^n \\ &= \frac{8}{d+1} \left(1 + \frac{1}{M} \right)^{d+1}, \end{aligned}$$

the thesis follows immediately. \square

3.4. Analysis for the hypercube H_d . The eigenvalues³ of the hypercube H_d are $\lambda_m = 2m$ for $m \in \{0, \dots, d\}$, where the eigenvalue λ_m has multiplicity $\binom{d}{m} = \frac{d!}{m!(d-m)!}$. We thus obtain that

$$R_{\text{ave}}(H_d) = \frac{1}{2^d} \sum_{m=1}^d \frac{1}{2m} \binom{d}{m}.$$

Proof of Theorem 2.5. For the lower bound, we have

$$\begin{aligned} R_{\text{ave}}(H_d) &\geq \frac{1}{2^{d+1}} \sum_{m=1}^d \frac{1}{m+1} \binom{d}{m} \\ &= \frac{1}{2^{d+1}} \sum_{m=1}^d \frac{1}{d+1} \binom{d+1}{m+1}. \end{aligned}$$

By the change of variables $m' = m + 1$ and $d' = d + 1$, we compute $\sum_{m=1}^d \binom{d+1}{m+1} = 2^{d+1} - d - 2$ and conclude that

$$(3.20) \quad R_{\text{ave}}(H_d) \geq \left(1 - \frac{d+2}{2^{d+1}} \right) \frac{1}{d+1}.$$

³Note that these eigenvalues cannot be computed using (3.2) with $M = 2$ because H_d is a degenerate case of T_{2^d} .

For the corresponding upper bound, we have

$$\begin{aligned} R_{\text{ave}}(H_d) &\leq \frac{1}{2^{d+1}} \sum_{m=1}^d \frac{2}{m+1} \binom{d}{m} \\ &= \frac{1}{2^{d+1}} \sum_{m=1}^d \frac{2}{d+1} \binom{d+1}{m+1} \leq \frac{2}{d+1}. \quad \square \end{aligned}$$

Proof of (2.5). In order to prove the asymptotic trend (2.5), from the definition of $R_{\text{ave}}(H_d)$ and using Pascal's rule, we compute

$$\begin{aligned} R_{\text{ave}}(H_d) &= \frac{1}{2} R_{\text{ave}}(H_{d-1}) + \frac{1}{2^{d+1}} \frac{1}{d} \sum_{k=1}^d \binom{d}{k} \\ &= \frac{1}{2} R_{\text{ave}}(H_{d-1}) + \frac{1}{2d} \left(1 - \frac{1}{2^d} \right). \end{aligned}$$

We have thus shown that the sequence $R_{\text{ave}}(H_d)$ can be constructed recursively by the above formula and defining $R_{\text{ave}}(H_0) = 0$. This recursion implies that

$$\begin{aligned} R_{\text{ave}}(H_d) &= \sum_{i=1}^d \frac{1}{2^{d-i}} \frac{1}{2i} \left(1 - \frac{1}{2^i} \right) \\ &= \sum_{i=1}^d \frac{1}{2^{d+1}} \frac{2^i - 1}{i}. \end{aligned}$$

Consequently, $R_{\text{ave}}(H_d) \leq \frac{1}{2^{d+1}} \sum_{i=1}^d \frac{2^i}{i}$, and we claim that

$$(3.21) \quad \lim_{d \rightarrow +\infty} \frac{\frac{1}{2^{d+1}} \sum_{i=1}^d \frac{2^i}{i}}{\frac{1}{d}} = 1.$$

This fact can be shown true as follows. Let $a_d = \frac{1}{2^{d+1}} \sum_{i=1}^d \frac{2^i}{i}$. Then, it is immediate to verify that a_d satisfies the recursion

$$\begin{cases} a_0 = 0, \\ a_{d+1} = \frac{1}{2} \left(1 + \frac{1}{d} \right) a_d + \frac{1}{2} \quad \text{for } d \geq 0, \end{cases}$$

and—by induction—that if $d \geq 3$, then $a_d > 1$, and if $d \geq 5$, then $a_{d+1} < a_d$. Then, a_d must have a finite limit $\ell \geq 1$. Also, note that

$$a_{d+1} = \frac{1}{2} \left(1 + \frac{1}{d} \right) a_d + \frac{1}{2} \leq \frac{1}{2} a_d + \frac{4}{3} \frac{1}{d} + \frac{1}{2}.$$

By taking the limit on both sides of the inequality, we obtain that $\ell \leq 1$. Finally, the desired (2.5) follows by combining (3.20) and (3.21). \square

4. Conclusion. The average effective resistance of a graph is an important performance index in several problems of distributed control and estimation, where toroidal grid graphs are exemplary d -dimensional graphs. In these graphs, the asymptotical dependence of the average effective resistance on the network size is well known,

but limited information was available about the constants involved in such relations and about the dependence on the dimension d .

We have expressed the average effective resistance of a graph in terms of a sum of the inverse Laplacian eigenvalues and found new estimates of this quantity: these estimates are key to our refined asymptotic analysis. For bidimensional toroidal grids, we have identified the proportionality constant of the leading term and have studied the case when the grid sides have unequal lengths. In grids with $d \geq 3$ and equal side lengths, we conjectured that the average effective resistance is inversely proportional to the dimension d . This conjecture is supported by numerical evidences and by several partial results.

Our results have been derived for toroidal grids, but we believe that they provide more general insights about the role of graph dimension in network estimation problems. Indeed, scaling properties deduced on toroidal grid graphs can typically be extended, with due care, to less structured graphs: works in this direction include [3, 5, 21, 22]. We envisage that our results on high-dimensional graphs can undergo similar extensions and thus cover more realistic networks in engineering and social sciences.

Acknowledgment. The authors wish to thank F. Garin for fruitful conversations on the topics of this paper.

REFERENCES

- [1] D. ALDOUS AND J. FILL, *Reversible Markov Chains and Random Walks on Graphs*, <http://www.stat.berkeley.edu/~aldous/index.html>.
- [2] B. BAMIEH, M. R. JOVANOVIC, P. MITRA, AND S. PATTERSON, *Coherence in large-scale networks: Dimension-dependent limitations of local feedback*, IEEE Trans. Automat. Control, 57 (2012), pp. 2235–2249.
- [3] P. BAROOAH AND J. P. HESPAÑHA, *Estimation from relative measurements: Algorithms and scaling laws*, IEEE Control Syst. Mag., 27 (2007), pp. 57–74.
- [4] P. BAROOAH AND J. P. HESPAÑHA, *Estimation from relative measurements: Electrical analogy and large graphs*, IEEE Trans. Signal Process., 56 (2008), pp. 2181–2193.
- [5] P. BAROOAH AND J. P. HESPAÑHA, *Error scaling laws for linear optimal estimation from relative measurements*, IEEE Trans. Inform. Theory, 55 (2009), pp. 5661–5673.
- [6] D. BONCHEV, E. J. MARKEI, AND A. H. DEKMEZIAN, *Long chain branch polymer chain dimensions: Application of topology to the Zimm–Stockmayer model*, Polymer, 43 (2002), pp. 203–222.
- [7] R. CARLI, F. FAGNANI, A. SPERANZON, AND S. ZAMPIERI, *Communication constraints in the average consensus problem*, Automatica, 44 (2008), pp. 671–684.
- [8] R. CARLI, F. GARIN, AND S. ZAMPIERI, *Quadratic indices for the analysis of consensus algorithms*, in Proceedings of the Information Theory and Applications Workshop, San Diego, 2009, pp. 96–104.
- [9] A. CARRON, M. TODDESCATO, R. CARLI, AND L. SCHENATO, *An asynchronous consensus-based algorithm for estimation from noisy relative measurements*, IEEE Trans. Control Network Syst., 1 (2014), pp. 283–295.
- [10] A. K. CHANDRA, P. RAGHAVAN, W. L. RUZZO, AND R. SMOLENSKY, *The electrical resistance of a graph captures its commute and cover times*, in Proceedings of the ACM Symposium on Theory of Computing, 1989, pp. 574–586.
- [11] P. G. DOYLE AND J. L. SNELL, *Random Walks and Electric Networks*, Carus Math. Monogr. 22, Mathematical Association of America, Washington, DC, 1984.
- [12] J. ELSON, R. M. KARP, C. H. PAPADIMITRIOU, AND S. SHENKER, *Global synchronization in sensornets*, in Proceedings of the LATIN 2004: Theoretical Informatics, Springer, New York, 2004, pp. 609–624.
- [13] E. ESTRADA AND N. HATANO, *A vibrational approach to node centrality and vulnerability in complex networks*, Phys. A, 389 (2010), pp. 3648–3660.

- [14] F. GARIN AND L. SCHENATO, *A survey on distributed estimation and control applications using linear consensus algorithms*, in Networked Control Systems, A. Bemporad, M. Heemels, and M. Johansson, eds., Lecture Notes in Control and Inform. Sci. 406, Springer, New York, 2010, pp. 75–107.
- [15] F. GARIN AND S. ZAMPIERI, *Mean square performance of consensus-based distributed estimation over regular geometric graphs*, SIAM J. Control Optim., 50 (2012), pp. 306–333.
- [16] A. GIRIDHAR AND P. R. KUMAR, *Distributed clock synchronization over wireless networks: Algorithms and analysis*, in Proceedings of the IEEE Conference on Decision and Control, 2006, pp. 4915–4920.
- [17] A. GHOSH, S. BOYD, AND A. SABERI, *Minimizing effective resistance of a graph*, SIAM Rev., 50 (2008), pp. 37–66.
- [18] S. KOSE AND E. G. FRIEDMAN, *Fast algorithms for power grid analysis based on effective resistance*, in Proceedings of the IEEE Symposium on Circuits and Systems, Paris, 2010, pp. 3661–3664.
- [19] A. N. LANGVILLE AND C. D. MEYER, *Who's # 1?: The Science of Rating and Ranking*, Princeton University Press, Princeton, NJ, 2012.
- [20] D. A. LEVIN, Y. PERES, AND E. L. WILMER, *Markov Chains and Mixing Times*, AMS, Providence, RI, 2008.
- [21] E. LOVISARI, F. GARIN, AND S. ZAMPIERI, *Resistance-based performance analysis of the consensus algorithm over geometric graphs*, SIAM J. Control Optim., 51 (2013), pp. 3918–3945.
- [22] E. LOVISARI AND S. ZAMPIERI, *Performance metrics in the average consensus problem: A tutorial*, Ann. Rev. Control, 36 (2012), pp. 26–41.
- [23] B. OSTING, C. BRUNE, AND S. J. OSHER, *Optimal data collection for improved rankings expose well-connected graphs*, J. Mach. Learn. Res., 15 (2014), pp. 2981–3012.
- [24] C. RAVAZZI, P. FRASCA, R. TEMPO, AND H. ISHII, *Ergodic randomized algorithms and dynamics over networks*, IEEE Trans. Control Network Syst., 2 (2015), pp. 78–87.
- [25] W. S. ROSSI, P. FRASCA, AND F. FAGNANI, *Transient and limit performance of distributed relative localization*, in Proceedings of the IEEE Conference on Decision and Control, Maui, HI, 2012, pp. 2744–2748.
- [26] W. S. ROSSI, P. FRASCA, AND F. FAGNANI, *Limited benefit of cooperation in distributed relative localization*, in Proceedings of the IEEE Conference on Decision and Control, Florence, 2013, pp. 5427–5431.
- [27] A. S. SHABANI, *Notes on the upper and lower bounds of two inequalities for the gamma function*, Hacet. J. Math. Stat., 39 (2010), pp. 11–15.
- [28] L. XIAO, S. BOYD, AND S.-J. KIM, *Distributed average consensus with least-mean-square deviation*, J. Parallel Distrib. Comput., 67 (2007), pp. 33–46.
- [29] F. Y. WU, *Theory of resistor networks: The two-point resistance*, J. Phys. A, 37 (2004), pp. 6653–6673.