

ANNALES MATHÉMATIQUES



BLAISE PASCAL

D. VAMSHEE KRISHNA, B. VENKATESWARLU & T. RAMREDDY

**Coefficient inequality for transforms of parabolic starlike
and uniformly convex functions**

Volume 21, n° 2 (2014), p. 39-56.

http://ambp.cedram.org/item?id=AMBP_2014__21_2_39_0

© Annales mathématiques Blaise Pascal, 2014, tous droits réservés.

L'accès aux articles de la revue « Annales mathématiques Blaise Pascal » (<http://ambp.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://ambp.cedram.org/legal/>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

*Publication éditée par le laboratoire de mathématiques
de l'université Blaise-Pascal, UMR 6620 du CNRS
Clermont-Ferrand — France*

cedram

*Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>*

Coefficient inequality for transforms of parabolic starlike and uniformly convex functions

D. VAMSHEE KRISHNA
B. VENKATESWARLU
T. RAMREDDY

Abstract

The objective of this paper is to obtain sharp upper bound to the second Hankel functional associated with the k^{th} root transform $[f(z^k)]^{\frac{1}{k}}$ of normalized analytic function $f(z)$ belonging to parabolic starlike and uniformly convex functions, defined on the open unit disc in the complex plane, using Toeplitz determinants.

1. Introduction

Let A denote the class of all functions $f(z)$ of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (1.1)$$

in the open unit disc $E = \{z : |z| < 1\}$. Let S be the subclass of A consisting of univalent functions. Let the functions F and G be analytic in the unit disc E . Then F is said to be subordinate to G , written $F \prec G$, if there exists an analytic function $w(z)$ in the open unit disc E satisfying $w(0) = 1$ and $|w(z)| < 1, \forall z \in E$ called the Schwarz's function such that

$$F(z) = G(w(z)), \forall z \in E. \quad (1.2)$$

If $F \prec G$ and $G(z)$ is univalent in the open unit disc E , then the subordination is equivalent to $F(0) = G(0)$ and $\text{range } F(z) \subseteq \text{range } G(z)$. For a univalent function in the class A , it is well known that the n^{th} coefficient is bounded by n . The bounds for the coefficients give information about the geometric properties of these functions. For example, the bound for the second coefficient of normalized univalent function readily yields

Keywords: Analytic function, parabolic starlike and uniformly convex functions, upper bound, second Hankel functional, positive real function, Toeplitz determinants.

Math. classification: 30C45, 30C50.

the growth and distortion properties for univalent functions. The Hankel determinant of f for $q \geq 1$ and $n \geq 1$ was defined by Pommerenke [18] as

$$H_q(n) = \begin{vmatrix} a_n & a_{n+1} & \cdots & a_{n+q-1} \\ a_{n+1} & a_{n+2} & \cdots & a_{n+q} \\ \vdots & \vdots & \vdots & \vdots \\ a_{n+q-1} & a_{n+q} & \cdots & a_{n+2q-2} \end{vmatrix}, (a_1 = 1).$$

This determinant has been considered by many authors in the literature . For example, Noor [17] determined the rate of growth of $H_q(n)$ as $n \rightarrow \infty$ for the functions in S with bounded boundary. Ehrenborg [8] studied the Hankel determinant of exponential polynomials. The Hankel transform of an integer sequence and some of its properties were discussed by Layman in [13]. In 1966, Pommerenke [18] investigated the Hankel determinant of areally mean p -valent functions, also studied by Noonan and Thomas [16], univalent functions as well as starlike functions. In the recent years, several authors have investigated bounds for the Hankel determinant of functions belonging to various subclasses of univalent and multivalent functions [1, 12, 11]. In particular cases, $q = 2, n = 1, a_1 = 1$ and $q = 2, n = 2, a_1 = 1$, the Hankel determinant simplifies respectively to

$$H_2(1) = \begin{vmatrix} a_1 & a_2 \\ a_2 & a_3 \end{vmatrix} = a_3 - a_2^2,$$

$$H_2(2) = \begin{vmatrix} a_2 & a_3 \\ a_3 & a_4 \end{vmatrix} = a_2 a_4 - a_3^2.$$

We refer to $H_2(2)$ as the second Hankel determinant. It is fairly well known that for the univalent functions of the form given in (1.1) the sharp inequality $|H_2(1)| = |a_3 - a_2^2| \leq 1$ holds true [7]. For a family \mathcal{T} of functions in S , the more general problem of finding sharp estimates for the functional $|a_3 - \mu a_2^2|$ ($\mu \in \mathbb{R}$ or $\mu \in \mathbb{C}$) is popularly known as the Fekete-Szegő problem for \mathcal{T} . Ali [3] found sharp bounds for the first four coefficients and sharp estimate for the Fekete-Szegő functional $|\gamma_3 - t\gamma_2^2|$, where t is real for the inverse function of f defined as $f^{-1}(w) = w + \sum_{n=2}^{\infty} \gamma_n w^n$ when it belongs to the class of strongly starlike functions of order α ($0 < \alpha \leq 1$) denoted by $\widetilde{ST}(\alpha)$. R. M. Ali, S. K. Lee, V. Ravichandran and S. Supramaniam [5] obtained sharp bounds for the Fekete-Szegő functional denoted by $|b_{2k+1} - \mu b_{k+1}^2|$ associated with the k^{th} root transform $[f(z^k)]^{\frac{1}{k}}$ of the function given in (1.1), belonging to

certain subclasses of S . The k^{th} root transform for the function f given in (1.1) is defined as

$$F(z) := [f(z^k)]^{\frac{1}{k}} = z + \sum_{n=1}^{\infty} t_{kn+1} z^{kn+1} \quad (1.3)$$

Motivated by the results obtained by R. M. Ali, S. K. Lee, V. Ravichandran and S. Supramaniam [5], in the present paper, we obtain sharp upper bound to the functional $|t_{k+1}t_{3k+1} - t_{2k+1}^2|$, called the second Hankel determinant for the k^{th} root transform of the function f when it belongs to certain subclasses of S , defined as follows.

Definition 1.1. A function $f(z) \in A$ is said to be parabolic starlike function, if and only if

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| < \operatorname{Re} \left\{ \frac{zf'(z)}{f(z)} \right\}, \forall z \in E \quad (1.4)$$

The class of all parabolic starlike functions is introduced by Ronning [20] and is denoted by S_p . Geometrically, (see [4]) S_p is the class functions f , for which $\left\{ \frac{zf'(z)}{f(z)} \right\}$ takes its value in the interior of the parabola in the right half plane symmetric about the real axis with vertex at $(\frac{1}{2}, 0)$.

Definition 1.2. A function $f \in A$ is said to be in UCV , if and only if

$$\left| \frac{zf''(z)}{f'(z)} \right| < \operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\}, \forall z \in E. \quad (1.5)$$

Goodman [9] introduced the class UCV of uniformly convex functions consisting of convex functions $f \in A$ with the property that for every circular arc γ contained in the unit disc E with centre also in E , the image arc $f(\gamma)$ is a convex arc. Ma and Minda [15] and Ronning [20] independently developed a one-variable characterization for the functions in the class UCV . From the Definitions 1.1 and 1.2, we have the relation between UCV and S_p is given in terms of an Alexander type Theorem [2] by Ronning (see [4]) as follows.

$$f \in UCV \Leftrightarrow zf' \in S_p. \quad (1.6)$$

Further, Ali [4] obtained sharp bounds on the first four coefficients and Fekete-Szegö inequality for the functions in the class S_p . Ali and Singh [6] showed that the normalized Riemann mapping function $q(z)$ from E onto the domain $D = \{w = u + iv : v^2 < 4u\} = \{w : |w - 1| < 1 + \operatorname{Re}(w)\}$,

denotes the parabolic region in the right half plane of the complex plane given by

$$q(z) = \left[1 + \frac{4}{\pi^2} \left\{ \log \frac{1 + \sqrt{z}}{1 - \sqrt{z}} \right\}^2 \right] = \left[1 + \sum_{n=1}^{\infty} B_n z^n \right], \forall z \in E. \quad (1.7)$$

It can be observed that if $f(z) \in S_p$ then

$$\frac{zf'(z)}{f(z)} \prec q(z), \forall z \in E, \quad (1.8)$$

where $q(z)$ is given in (1.7).

Some preliminary lemmas required for proving our results are as follows:

2. Preliminary Results

Let \mathbb{P} denote the class of functions consisting of p , such that

$$p(z) = 1 + c_1 z + c_2 z^2 + c_3 z^3 + \dots = \left[1 + \sum_{n=1}^{\infty} c_n z^n \right], \quad (2.1)$$

which are regular in the open unit disc E and satisfy $\operatorname{Re}\{p(z)\} > 0$ for any $z \in E$. Here $p(z)$ is called the Caratheodory function [7].

Lemma 2.1. ([19, 21]) *If $p \in \mathbb{P}$, then $|c_k| \leq 2$, for each $k \geq 1$ and the inequality is sharp for the function $\left(\frac{1+z}{1-z}\right)$.*

Lemma 2.2. ([10]) *The power series for $p(z) = 1 + \sum_{n=1}^{\infty} c_n z^n$ given in (2.1) converges in the open unit disc E to a function in \mathbb{P} if and only if the Toeplitz determinants*

$$D_n = \begin{vmatrix} 2 & c_1 & c_2 & \cdots & c_n \\ c_{-1} & 2 & c_1 & \cdots & c_{n-1} \\ c_{-2} & c_{-1} & 2 & \cdots & c_{n-2} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ c_{-n} & c_{-n+1} & c_{-n+2} & \cdots & 2 \end{vmatrix}, n = 1, 2, 3, \dots$$

and $c_{-k} = \bar{c}_k$, are all non-negative. They are strictly positive except for $p(z) = \sum_{k=1}^m \rho_k P_0(e^{it_k z})$, $\rho_k > 0$, t_k real and $t_k \neq t_j$, for $k \neq j$, where $P_0(z) = \left(\frac{1+z}{1-z}\right)$; in this case $D_n > 0$ for $n < (m - 1)$ and $D_n = 0$ for $n \geq m$.

COEFFICIENT INEQUALITY FOR TRANSFORMS

This necessary and sufficient condition found in [10] is due to Carathéodory and Toeplitz. We may assume without restriction that $c_1 > 0$. On using Lemma 2.2, for $n = 2$, we have

$$D_2 = \begin{vmatrix} 2 & c_1 & c_2 \\ \bar{c}_1 & 2 & c_1 \\ \bar{c}_2 & \bar{c}_1 & 2 \end{vmatrix} = [8 + 2Re\{c_1^2 c_2\} - 2|c_2|^2 - 4|c_1|^2] \geq 0,$$

which is equivalent to

$$2c_2 = \{c_1^2 + x(4 - c_1^2)\}, \text{ for some } x, |x| \leq 1. \quad (2.2)$$

For $n = 3$,

$$D_3 = \begin{vmatrix} 2 & c_1 & c_2 & c_3 \\ \bar{c}_1 & 2 & c_1 & c_2 \\ \bar{c}_2 & \bar{c}_1 & 2 & c_1 \\ \bar{c}_3 & \bar{c}_2 & \bar{c}_1 & 2 \end{vmatrix} \geq 0$$

and is equivalent to

$$|(4c_3 - 4c_1 c_2 + c_1^3)(4 - c_1^2) + c_1(2c_2 - c_1^2)^2| \leq 2(4 - c_1^2)^2 - 2|(2c_2 - c_1^2)|^2. \quad (2.3)$$

From the relations (2.2) and (2.3), after simplifying, we get

$$4c_3 = \{c_1^3 + 2c_1(4 - c_1^2)x - c_1(4 - c_1^2)x^2 + 2(4 - c_1^2)(1 - |x|^2)z\} \\ \text{for some } z, \text{ with } |z| \leq 1. \quad (2.4)$$

To obtain our results, we refer to the classical method initiated by Libera and Zlotkiewicz [14] and used by several authors in the literature.

3. Main Results

Theorem 3.1. *If f given by (1.1) belongs to S_p and F is the k^{th} root transformation of f given by (1.3) then*

$$|t_{k+1}t_{3k+1} - t_{2k+1}^2| \leq \left[\frac{8}{k\pi^2} \right]^2$$

and the inequality is sharp.

Proof. For $f(z) = z + \sum_{n=2}^{\infty} a_n z^n \in S_p$, by virtue of Definition 1.1, we have

$$\left[\frac{zf'(z)}{f(z)} \right] \prec q(z), \quad \forall z \in E. \quad (3.1)$$

By the subordination principle, there exist a Schwarz's function $w(z)$ such that

$$\left[\frac{zf'(z)}{f(z)} \right] \prec [q\{w(z)\}], \forall z \in E. \quad (3.2)$$

Define a function $h(z)$ such that

$$h(z) = \left[\frac{zf'(z)}{f(z)} \right] = 1 + b_1z + b_2z^2 + b_3z^3 + \dots = \left[1 + \sum_{n=1}^{\infty} b_nz^n \right] \\ \Leftrightarrow [zf'(z)] = [f(z)h(z)]. \quad (3.3)$$

Using the series representations for $f(z)$, $f'(z)$ and $h(z)$ in (3.3), we have

$$z \left\{ 1 + \sum_{n=2}^{\infty} na_nz^{n-1} \right\} = \left\{ z + \sum_{n=2}^{\infty} a_nz^n \right\} \left\{ 1 + \sum_{n=1}^{\infty} b_nz^n \right\}. \quad (3.4)$$

Upon simplification, we obtain

$$1 + a_2z + 2a_3z^2 + 3a_4z^3 + \dots = 1 + b_1z + (b_1a_2 + b_2)z^2 + \\ (b_1a_3 + b_2a_2 + b_3)z^3 + \dots \quad (3.5)$$

Equating the coefficients of like powers of z , z^2 and z^3 respectively on both sides of (3.5), after simplifying, we get

$$a_2 = b_1; \quad a_3 = \frac{1}{2} (b_2 + b_1^2); \quad a_4 = \frac{1}{3} \left(b_3 + \frac{3}{2}b_1b_2 + \frac{b_1^3}{2} \right). \quad (3.6)$$

Since $q(z)$ is univalent in the open unit disc E and $h(z) \prec q(z)$, define a function

$$p(z) = \left[\frac{1+w(z)}{1-w(z)} \right] = \left[\frac{1+q^{-1}\{h(z)\}}{1-q^{-1}\{h(z)\}} \right] = 1 + c_1z + c_2z^2 + c_3z^3 + \dots, \quad (3.7)$$

where $p(z)$ is given in (2.1). Solving $w(z)$ in terms of $p(z)$ in the relation (3.7) and replacing $p(z)$ by its equivalent expression in series, we have

$$w(z) = \left[\frac{p(z) - 1}{p(z) + 1} \right] = \left[\frac{(1 + c_1z + c_2z^2 + c_3z^3 + \dots) - 1}{(1 + c_1z + c_2z^2 + c_3z^3 + \dots) + 1} \right].$$

Upon simplification, we obtain

$$w(z) = \frac{1}{2} \left\{ c_1z + \left(c_2 - \frac{c_1^2}{2} \right) z^2 + \left(c_3 - c_1c_2 + \frac{c_1^3}{4} \right) z^3 + \dots \right\}. \quad (3.8)$$

COEFFICIENT INEQUALITY FOR TRANSFORMS

Using the expansion of $\log(1+x) = \left\{x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots\right\}$ for $q(z)$ given in (1.7), after simplifying, we get

$$\left[\log\left(\frac{1+\sqrt{z}}{1-\sqrt{z}}\right)\right]^2 = \left\{4z + \frac{8}{3}z^2 + \frac{92}{45}z^3 + \frac{176}{105}z^4 + \dots\right\}. \quad (3.9)$$

From the relations (1.7) and (3.9), we obtain

$$\begin{aligned} q(z) &= \left\{1 + \frac{16}{\pi^2}z + \frac{32}{3\pi^2}z^2 + \frac{368}{45\pi^2}z^3 + \frac{704}{105\pi^2}z^4 \dots\right\} \\ &= [1 + B_1z + B_2z^2 + B_3z^3 + \dots] = \left[1 + \sum_{n=1}^{\infty} B_n z^n\right]. \end{aligned} \quad (3.10)$$

Equating the coefficients of like powers of z , z^2 and z^3 respectively, on both sides of (3.10), we get

$$\begin{aligned} B_1 &= \frac{16}{\pi^2}; \quad B_2 = \frac{32}{3\pi^2}; \quad B_3 = \frac{368}{45\pi^2} \dots, \\ B_n &= \frac{16}{n\pi^2} \sum_{k=0}^{n-1} \frac{1}{(2k+1)}, \quad n = 2, 3, 4, \dots \end{aligned} \quad (3.11)$$

From the relations (3.2) and (3.3), we have

$$h(z) = [q\{w(z)\}]. \quad (3.12)$$

In view of (3.12), using (3.8) in (3.10) along with the equivalent expression for $h(z)$ given in (3.3), upon simplification, (3.12) is equivalent to

$$\begin{aligned} [1 + b_1z + b_2z^2 + b_3z^3 + \dots] &= \left[1 + \frac{1}{2}B_1c_1z + \left\{\frac{1}{2}B_1\left(c_2 - \frac{c_1^2}{2}\right) + \frac{1}{4}B_2c_1^2\right\}z^2 + \right. \\ &\left. \left\{\frac{1}{2}B_1\left(c_3 - c_1c_2 + \frac{c_1^3}{4}\right) + \frac{1}{2}B_2c_1\left(c_2 - \frac{c_1^2}{2}\right) + \frac{1}{8}B_3c_1^3\right\}z^3 + \dots\right]. \end{aligned} \quad (3.13)$$

Equating the coefficients of like powers of z , z^2 and z^3 respectively, on both sides of (3.13), we have

$$\begin{aligned} b_1 &= \frac{1}{2}B_1c_1; \quad b_2 = \left\{\frac{1}{2}B_1\left(c_2 - \frac{c_1^2}{2}\right) + \frac{1}{4}B_2c_1^2\right\}; \\ b_3 &= \left\{\frac{1}{2}B_1\left(c_3 - c_1c_2 + \frac{c_1^3}{4}\right) + \frac{1}{2}B_2c_1\left(c_2 - \frac{c_1^2}{2}\right) + \frac{1}{8}B_3c_1^3\right\}. \end{aligned} \quad (3.14)$$

Simplifying the relations (3.11) and (3.14), we get

$$b_1 = \frac{8c_1}{\pi^2}; \quad b_2 = \frac{8}{\pi^2} \left(c_2 - \frac{c_1^2}{6} \right); \quad b_3 = \frac{8}{\pi^2} \left(c_3 - \frac{c_1 c_2}{3} + \frac{2}{45} c_1^3 \right). \quad (3.15)$$

From the relations (3.6) and (3.15), upon simplification, we obtain

$$\begin{aligned} a_2 &= \frac{8c_1}{\pi^2}; \quad a_3 = \frac{8}{2\pi^2} \left[c_2 - \left\{ \frac{1}{6} - \frac{8}{\pi^2} \right\} c_1^2 \right]; \\ a_4 &= \frac{8}{3\pi^2} \left[c_3 - \left\{ \frac{1}{3} - \frac{12}{\pi^2} \right\} c_1 c_2 + \left\{ \frac{2}{45} - \frac{2}{\pi^2} + \frac{32}{\pi^4} \right\} c_1^3 \right]. \end{aligned} \quad (3.16)$$

For a function f given by (1.1), a computation shows that

$$\begin{aligned} [f(z^k)]^{\frac{1}{k}} &= \left[z^k + \sum_{n=2}^{\infty} a_n z^{nk} \right]^{\frac{1}{k}} \\ &= \left[z + \frac{1}{k} a_2 z^{k+1} + \left\{ \frac{1}{k} a_3 + \frac{(1-k)}{2k^2} a_2^2 \right\} z^{2k+1} \right. \\ &\quad \left. + \left\{ \frac{1}{k} a_4 + \frac{(1-k)}{k^2} a_2 a_3 + \frac{(1-k)(1-2k)}{6k^3} a_2^3 \right\} z^{3k+1} + \dots \right]. \end{aligned} \quad (3.17)$$

From the equations (1.3) and (3.16) together with (3.17), after simplifying, we get

$$\begin{aligned} t_{k+1} &= \frac{8c_1}{k\pi^2}; \quad t_{2k+1} = \frac{8}{2k\pi^2} \left[c_2 - \left\{ \frac{1}{6} - \frac{8}{k\pi^2} \right\} c_1^2 \right]; \\ t_{3k+1} &= \frac{8}{3k\pi^2} \left[c_3 - \left\{ \frac{1}{3} - \frac{12}{k\pi^2} \right\} c_1 c_2 + \left\{ \frac{2}{45} - \frac{2}{k\pi^2} + \frac{32}{k^2\pi^4} \right\} c_1^3 \right]. \end{aligned} \quad (3.18)$$

Substituting the values of t_{k+1} , t_{2k+1} and t_{3k+1} from (3.18) in the second Hankel determinant $|t_{k+1}t_{3k+1} - t_{2k+1}^2|$ to the k^{th} transformation for the function $f \in S_p$, upon simplification, we obtain

$$|t_{k+1}t_{3k+1} - t_{2k+1}^2| = \frac{16}{9k^2\pi^4} \left| 12c_1c_3 - c_1^2c_2 - 9c_2^2 + \left\{ \frac{17}{60} - \frac{192}{k^2\pi^4} \right\} c_1^4 \right|, \quad (3.19)$$

which is equivalent to

$$|t_{k+1}t_{3k+1} - t_{2k+1}^2| = \frac{16}{9k^2\pi^4} |d_1c_1c_3 + d_2c_1^2c_2 + d_3c_2^2 + d_4c_1^4|, \quad (3.20)$$

$$\text{where } d_1 = 12; \quad d_2 = -1; \quad d_3 = -9; \quad d_4 = \left\{ \frac{17}{60} - \frac{192}{k^2\pi^4} \right\}. \quad (3.21)$$

COEFFICIENT INEQUALITY FOR TRANSFORMS

Substituting the values of c_2 and c_3 from (2.2) and (2.4) respectively from Lemma 2.2 on the right-hand side of (3.20), we have

$$\begin{aligned}
 & |d_1c_1c_3 + d_2c_1^2c_2 + d_3c_2^2 + d_4c_1^4| \\
 &= |d_1c_1 \times \frac{1}{4}\{c_1^3 + 2c_1(4 - c_1^2)x - c_1(4 - c_1^2)x^2 + 2(4 - c_1^2)(1 - |x|^2)z\} + \\
 &\quad d_2c_1^2 \times \frac{1}{2}\{c_1^2 + x(4 - c_1^2)\} + d_3 \times \frac{1}{4}\{c_1^2 + x(4 - c_1^2)\}^2 + d_4c_1^4|. \quad (3.22)
 \end{aligned}$$

Using the triangle inequality and the fact that $|z| < 1$, we get

$$\begin{aligned}
 4|d_1c_1c_3 + d_2c_1^2c_2 + d_3c_2^2 + d_4c_1^4| &\leq |(d_1 + 2d_2 + d_3 + 4d_4)c_1^4 + 2d_1c_1(4 - c_1^2) \\
 &\quad + 2(d_1 + d_2 + d_3)c_1^2(4 - c_1^2)|x| - \\
 &\quad \left\{ (d_1 + d_3)c_1^2 + 2d_1c_1 - 4d_3 \right\} (4 - c_1^2)|x|^2|. \quad (3.23)
 \end{aligned}$$

From the relation (3.21), we can now write

$$\begin{aligned}
 \{(d_1 + 2d_2 + d_3 + 4d_4) = \frac{32}{15k^2\pi^4} (k^2\pi^4 - 360); \quad d_1 = 12; \\
 (d_1 + d_2 + d_3) = 2\}. \quad (3.24)
 \end{aligned}$$

$$\left\{ (d_1 + d_3)c_1^2 + 2d_1c_1 - 4d_3 \right\} = 3(c_1 + 2)(c_1 + 6). \quad (3.25)$$

Since $c_1 \in [0, 2]$, using the result $(c_1 + a)(c_1 + b) \geq (c_1 - a)(c_1 - b)$, where $a, b \geq 0$ in the relation (3.25), we get

$$- \left\{ (d_1 + d_3)c_1^2 + 2d_1c_1 - 4d_3 \right\} \leq -3(c_1 - 2)(c_1 - 6). \quad (3.26)$$

Substituting the calculated values from (3.24) and (3.26) on the right-hand side of (3.23), we have

$$\begin{aligned}
 4|d_1c_1c_3 + d_2c_1^2c_2 + d_3c_2^2 + d_4c_1^4| &\leq \left| \frac{32}{15k^2\pi^4} (k^2\pi^4 - 360) c_1^4 + 24c_1(4 - c_1^2) \right. \\
 &\quad \left. + 4c_1^2(4 - c_1^2)|x| - 3(c_1 - 2)(c_1 - 6)(4 - c_1^2)|x|^2 \right|. \quad (3.27)
 \end{aligned}$$

Choosing $c_1 = c \in [0, 2]$, applying triangle inequality and replacing $|x|$ by μ on the right-hand side of (3.27), we get

$$\begin{aligned} 4|d_1c_1c_3 + d_2c_1^2c_2 + d_3c_2^2 + d_4c_1^4| &\leq \left[\frac{32}{15k^2\pi^4} \{360 - k^2\pi^4\} c^4 + 24c(4 - c^2) \right. \\ &\quad \left. + 4c^2(4 - c^2)\mu + 3(c - 2)(c - 6)(4 - c^2)\mu^2 \right] \\ &= F(c, \mu), \text{ for } 0 \leq \mu = |x| \leq 1, \end{aligned} \quad (3.28)$$

$$\begin{aligned} \text{where } F(c, \mu) &= \frac{32}{15k^2\pi^4} \{360 - k^2\pi^4\} c^4 + 24c(4 - c^2) + 4c^2(4 - c^2)\mu \\ &\quad + 3(c - 2)(c - 6)(4 - c^2)\mu^2. \end{aligned} \quad (3.29)$$

We next maximize the function $F(c, \mu)$ on the closed region $[0, 1] \times [0, 2]$. Differentiating $F(c, \mu)$ in (3.29) partially with respect to μ , we obtain

$$\frac{\partial F}{\partial \mu} = [4c^2 + 6(c - 2)(c - 6)\mu] \times (4 - c^2). \quad (3.30)$$

For $0 < \mu < 1$ and for fixed c with $0 < c < 2$, from (3.30), we observe that $\frac{\partial F}{\partial \mu} > 0$. Consequently, $F(c, \mu)$ becomes an increasing function of μ and hence it cannot have a maximum value at any point in the interior of the closed region $[0, 1] \times [0, 2]$. Moreover, for fixed $c \in [0, 2]$, we have

$$\max_{0 \leq \mu \leq 1} F(c, \mu) = F(c, 1) = G(c). \quad (3.31)$$

Therefore, replacing μ by 1 in (3.29), upon simplification, we obtain

$$G(c) = \frac{1}{15k^2\pi^4} \{11520 - 137k^2\pi^4\} c^4 - 8c^2 + 144, \quad (3.32)$$

$$G'(c) = \frac{4}{15k^2\pi^4} \{11520 - 137k^2\pi^4\} c^3 - 16c. \quad (3.33)$$

From (3.33), we observe that $G'(c) \leq 0$, for every $c \in [0, 2]$ and for all values of k . Therefore, $G(c)$ is a monotonically decreasing function of c in the interval $[0, 2]$ and hence its maximum value occurs at $c = 0$ only. From (3.32), we get

$$\max_{0 \leq c \leq 2} G(c) = G(0) = 144. \quad (3.34)$$

Simplifying the relations (3.28) and (3.34), we obtain

$$|d_1c_1c_3 + d_2c_1^2c_2 + d_3c_2^2 + d_4c_1^4| \leq 36. \quad (3.35)$$

COEFFICIENT INEQUALITY FOR TRANSFORMS

From the relations (3.20) and (3.35), after simplifying, we get

$$|t_{k+1}t_{3k+1} - t_{2k+1}^2| \leq \left[\frac{8}{k\pi^2} \right]^2. \quad (3.36)$$

By setting $c_1 = c = 0$ and selecting $x = -1$ in (2.2) and (2.4), we find that $c_2 = -2$ and $c_3 = 0$. Using these values in (3.35), we observe that equality is attained, which shows that our result is sharp. For these values, we derive that

$$p(z) = \frac{1 - z^2}{1 + z^2} = 1 - 2z^2 + 2z^4 - \dots \text{ and } w(z) = -z^2. \quad (3.37)$$

Therefore, in this case the extremal function is $\left[\frac{zf'(z)}{f(z)} \right] = \frac{1-z^2}{1+z^2}$. This completes the proof of our Theorem 3.1. \square

Theorem 3.2. *If f given by (1.1) belongs to UCV and F is the k^{th} root transformation of f given by (1.3) then*

$$|t_{k+1}t_{3k+1} - t_{2k+1}^2| \leq \left[\frac{8}{3k\pi^2} \right]^2$$

and the inequality is sharp.

Proof. Let $f(z) = z + \sum_{n=2}^{\infty} a_n z^n \in \text{UCV}$, from the Definition 1.2, we have

$$\left[1 + \frac{zf''(z)}{f'(z)} \right] \prec q(z), \quad \forall z \in E.$$

By the subordination principle, there exist a Schwarz's function $w(z)$ such that

$$\left[1 + \frac{zf''(z)}{f'(z)} \right] \prec [q\{w(z)\}], \forall z \in E. \quad (3.38)$$

Define a function $h(z)$ such that

$$\begin{aligned} h(z) &= \left[1 + \frac{zf''(z)}{f'(z)} \right] = \{1 + b_1 z + b_2 z^2 + b_3 z^3 + \dots\} = \left[1 + \sum_{n=1}^{\infty} b_n z^n \right] \\ &\Leftrightarrow [f'(z) + zf''(z)] = [f'(z)h(z)]. \end{aligned} \quad (3.39)$$

Replacing $f'(z)$, $f''(z)$ and $h(z)$ by their equivalent expressions in series in the expression (3.39), we have

$$\left[\left\{ 1 + \sum_{n=2}^{\infty} na_n z^{n-1} \right\} + z \left\{ \sum_{n=2}^{\infty} n(n-1)a_n z^{n-2} \right\} \right] = \left[\left\{ 1 + \sum_{n=2}^{\infty} na_n z^{n-1} \right\} \left\{ 1 + \sum_{n=1}^{\infty} b_n z^n \right\} \right].$$

Upon simplification, we obtain

$$1 + 2a_2 z + 6a_3 z^2 + 12a_4 z^3 + \dots = 1 + b_1 z + (2b_1 a_2 + b_2) z^2 + (3b_1 a_3 + 2b_2 a_2 + b_3) z^3 + \dots \quad (3.40)$$

Equating the coefficients of like powers of z , z^2 and z^3 respectively on both sides of (3.40), after simplifying, we get

$$a_2 = \frac{b_1}{2}; \quad a_3 = \frac{1}{6}(b_2 + b_1^2); \quad a_4 = \frac{1}{24}(2b_3 + 3b_1 b_2 + b_1^3). \quad (3.41)$$

Applying the same procedure as described in Theorem 3.1, we obtain

$$a_2 = \frac{4}{\pi^2} c_1; \quad a_3 = \frac{4}{3\pi^2} \left[c_2 - \left\{ \frac{1}{6} - \frac{8}{\pi^2} \right\} c_1^2 \right]; \\ a_4 = \frac{2}{3\pi^2} \left[c_3 - \left\{ \frac{1}{3} - \frac{12}{\pi^2} \right\} c_1 c_2 + \left\{ \frac{2}{45} - \frac{2}{\pi^2} + \frac{32}{\pi^4} \right\} c_1^3 \right]. \quad (3.42)$$

From the equations (1.3) and (3.17) together with (3.42), after simplifying, we get

$$t_{k+1} = \frac{4c_1}{k\pi^2}; \quad t_{2k+1} = \frac{4}{3k\pi^2} \left[c_2 - \left\{ \frac{1}{6} - \frac{2(k+3)}{k\pi^2} \right\} c_1^2 \right]; \\ t_{3k+1} = \frac{2}{3k\pi^2} \left[c_3 + \left\{ \frac{-1}{3} + \frac{4(k+2)}{k\pi^2} \right\} c_1 c_2 + \left\{ \frac{2}{45} - \frac{2(k+2)}{3k\pi^2} + \frac{16(k+1)}{k^2\pi^4} \right\} c_1^3 \right]. \quad (3.43)$$

Substituting the values of t_{k+1} , t_{2k+1} and t_{3k+1} from (3.43) in the second Hankel determinant $|t_{k+1}t_{3k+1} - t_{2k+1}^2|$ to the k^{th} transformation for the

COEFFICIENT INEQUALITY FOR TRANSFORMS

function $f \in UCV$, upon simplification, we obtain

$$|t_{k+1}t_{3k+1} - t_{2k+1}^2| = \frac{4}{405k^2\pi^8} \times \left| 270\pi^4 c_1 c_3 + 30\pi^2 \{-\pi^2 + 12\} c_1^2 c_2 - 180\pi^4 c_2^2 + \{7\pi^4 - 60\pi^2 - 720 \left(1 + \frac{3}{k^2}\right)\} c_1^4 \right|. \quad (3.44)$$

The above expression is equivalent to

$$|t_{k+1}t_{3k+1} - t_{2k+1}^2| = \frac{4}{405k^2\pi^8} |d_1 c_1 c_3 + d_2 c_1^2 c_2 + d_3 c_2^2 + d_4 c_1^4|, \quad (3.45)$$

where $d_1 = 270\pi^4$; $d_2 = 30\pi^2 \{-\pi^2 + 12\}$;

$$d_3 = -180\pi^4; \quad d_4 = \left\{ 7\pi^4 - 60\pi^2 - 720 \left(1 + \frac{3}{k^2}\right) \right\}. \quad (3.46)$$

Applying the same procedure as described in Theorem 3.1, we get

$$4|d_1 c_1 c_3 + d_2 c_1^2 c_2 + d_3 c_2^2 + d_4 c_1^4| \leq |(d_1 + 2d_2 + d_3 + 4d_4) c_1^4 + 2d_1 c_1 (4 - c_1^2) + 2(d_1 + d_2 + d_3) c_1^2 (4 - c_1^2)| x| - \left\{ (d_1 + d_3) c_1^2 + 2d_1 c_1 - 4d_3 \right\} (4 - c_1^2) |x|^2|. \quad (3.47)$$

Using the values of d_1, d_2, d_3 and d_4 from (3.46), upon simplification, we obtain

$$(d_1 + 2d_2 + d_3 + 4d_4) = \left\{ 58\pi^4 + 480\pi^2 - 2880 \left(1 + \frac{3}{k^2}\right) \right\};$$

$$d_1 = 270\pi^4; \quad (d_1 + d_2 + d_3) = \left\{ 60\pi^4 + 360\pi^2 \right\}. \quad (3.48)$$

$$\left\{ (d_1 + d_3) c_1^2 + 2d_1 c_1 - 4d_3 \right\} = 90\pi^4 (c_1 + 2)(c_1 + 4). \quad (3.49)$$

Since $c_1 \in [0, 2]$, using the result $(c_1 + a)(c_1 + b) \geq (c_1 - a)(c_1 - b)$, where $a, b \geq 0$ in the relation (3.49), we get

$$- \left\{ (d_1 + d_3) c_1^2 + 2d_1 c_1 - 4d_3 \right\} \leq -90\pi^4 (c_1 - 2)(c_1 - 4). \quad (3.50)$$

Substituting the calculated values from (3.48) and (3.50) on the right-hand side of (3.47), we obtain

$$4|d_1c_1c_3+d_2c_1^2c_2+d_3c_2^2+d_4c_1^4| \leq \left| \left\{ 58\pi^4 + 480\pi^2 - 2880 \left(1 + \frac{3}{k^2} \right) \right\} c_1^4 + 540\pi^4 c_1(4 - c_1^2) + 120\pi^2 (\pi^2 + 6) c_1^2(4 - c_1^2)|x| - 90\pi^4(c_1 - 2)(c_1 - 4)(4 - c_1^2)|x|^2 \right|. \quad (3.51)$$

Choosing $c_1 = c \in [0, 2]$, applying the triangle inequality and replacing $|x|$ by μ on the right-hand side of the above inequality, we have

$$4|d_1c_1c_3+d_2c_1^2c_2+d_3c_2^2+d_4c_1^4| \leq \left[\left\{ 2880 \left(1 + \frac{3}{k^2} \right) + 480\pi^2 - 58\pi^4 \right\} c^4 + 540\pi^4 c(4 - c^2) + 120\pi^2 (\pi^2 - 6) c^2(4 - c^2)\mu + 90\pi^4(c - 2)(c - 4)(4 - c^2)\mu^2 \right] = F(c, \mu), \text{ for } 0 \leq \mu = |x| \leq 1, \quad (3.52)$$

$$\text{where } F(c, \mu) = \left[\left\{ 2880 \left(1 + \frac{3}{k^2} \right) + 480\pi^2 - 58\pi^4 \right\} c^4 + 540\pi^4 c(4 - c^2) + 120\pi^2 (\pi^2 - 6) c^2(4 - c^2)\mu + 90\pi^4(c - 2)(c - 4)(4 - c^2)\mu^2 \right]. \quad (3.53)$$

Applying the same procedure as described in Theorem 3.1, we observe that $\frac{\partial F}{\partial \mu} > 0$, so that $F(c, \mu)$ is an increasing function of μ and hence its maximum value does not occur at any point in the interior of the closed region $[0, 1] \times [0, 2]$. Further, for fixed $c \in [0, 2]$, we have

$$\max_{0 \leq \mu \leq 1} F(c, \mu) = F(c, 1) = G(c). \quad (3.54)$$

Therefore, replacing μ by 1 in (3.53), upon simplification, we obtain

$$G(c) = \left\{ 2880 \left(1 + \frac{3}{k^2} \right) - 268\pi^4 + 1200\pi^2 \right\} c^4 - 120\pi^2 \left\{ 24 - \pi^2 \right\} c^2 + 2880\pi^4, \quad (3.55)$$

$$G'(c) = 4 \left\{ 2880 \left(1 + \frac{3}{k^2} \right) - 268\pi^4 + 1200\pi^2 \right\} c^3 - 240\pi^2 \left\{ 24 - \pi^2 \right\} c. \quad (3.56)$$

From (3.56), for fixed $c \in [0, 2]$ and for every k , we observe that $G'(c) \leq 0$, which shows that $G(c)$ is a monotonically decreasing function of c and hence it attains the maximum value at $c = 0$ only. From (3.55), we get

$$\max_{0 \leq c \leq 2} G(c) = G(0) = 2880\pi^4. \quad (3.57)$$

From (3.52) and (3.57), upon simplification, we obtain

$$|d_1c_1c_3 + d_2c_1^2c_2 + d_3c_2^2 + d_4c_1^4| \leq 720\pi^4. \quad (3.58)$$

Simplifying the relations (3.45) and (3.58), we get

$$|t_{k+1}t_{3k+1} - t_{2k+1}^2| \leq \left[\frac{8}{3k\pi^2} \right]^2. \quad (3.59)$$

If we set $c_1 = c = 0$ and take $x = 1$ in (2.2) and (2.4), we find that $c_2 = 2$ and $c_3 = 0$. Using these values in (3.58), we see that equality is attained, which shows that our result is sharp. For these values, we derive that

$$p(z) = \frac{1+z^2}{1-z^2} = 1 + 2z^2 + 2z^4 + \dots \text{ and } w(z) = z^2. \quad (3.60)$$

Therefore, the extremal function in this case is $\left[1 + \frac{zf''(z)}{f'(z)} \right] = \frac{1+z^2}{1-z^2}$. This completes the proof of our Theorem 3.2. \square

Remark 3.3. For the choice of $k = 1$, the result coincides with that of VamsheeKrishna and RamReddy [22].

Acknowledgement: The authors would like to express their sincere thanks to the esteemed Referee(s) for their careful readings, valuable suggestions and comments, which helped to improve the presentation of the paper.

References

- [1] A. ABUBAKER et M. DARUS – “Hankel determinant for a class of analytic functions involving a generalized linear differential operators”, *Int. J. Pure Appl. Math.* **69(4)** (2011), p. 429 – 435, MR 2847841 | Zbl 1220.30011.
- [2] J. W. ALEXNADER – “Functions which map the interior of the unit circle upon simple regions”, *Annal. of Math.* **(2)17** (1915), p. 12 –22, MR 1503516 | JFM 45.0672.02.

- [3] R. M. ALI – “Coefficients of the inverse of strongly starlike functions”, *Bull. Malays. Math. Sci. Soc.(second series)* **26(1)** (2003), p. 63 – 71, MR 2055766 (2005b:30011) | Zbl 1185.30010.
- [4] ———, “Starlikeness associated with parabolic regions”, *Int. J. Math. Math. Sci.* **4** (2005), p. 561–570, MR 2172395 (2006d:30011) | Zbl 1077.30011.
- [5] R. M. ALI, S. K. LEE, V. RAVICHANDRAN et S. SUPRAMANIAM – “The Fekete-szegő coefficient functional for transforms of analytic functions”, *Bull. Iran. Math. Soc.* **35(2)** (2009), p. 119–142, MR 2642930.
- [6] R. M. ALI et V. SINGH – “Coefficients of parabolic starlike functions of order ρ ”, World Sci. Publ. River Edge, New Jersey, 1995, 1995 MR 1415158 [97 h:30008]., p. 23 – 36.
- [7] P. L. DUREN – *Univalent functions*, 259, Grundlehren der Mathematischen Wissenschaften, New York, Springer-verlag XIV, 328, 1983, MR 0708494 | Zbl 0514.30001.
- [8] R. EHRENBORG – “The hankel determinant of exponential polynomials”, *Amer. Math. Monthly* **107(6)** (2000), p. 557–560, MR 1767065 (2001c:15009) | Zbl 0985.15006.
- [9] A. W. GOODMAN – “On uniformly convex functions”, *Ann. Polon. Math.* **56 (1)** (1991), p. 87 – 92, MR 1145573 (93a:30009) | Zbl 0744.30010.
- [10] U. GRENANDER et G. SZEGÖ – *Toeplitz forms and their applications*, Second edition. Chelsea Publishing Co., New York, 1984, MR 0890515 | Zbl 0611.47018.
- [11] A. JANTENG, S. A. HALIM et M. DARUS – “Coefficient inequality for a function whose derivative has a positive real part”, *J. Inequal. Pure Appl. Math.* **7(2)** (2006), p. 1–5, MR 2221331 | Zbl 1134.30310.
- [12] ———, “Hankel determinant for starlike and convex functions”, *Int. J. Math. Anal., (Ruse)* **4 (no. 13-16)** (2007), p. 619–625, MR 2370200 | Zbl 1137.30308.
- [13] J. W. LAYMAN – “The hankel transform and some of its properties”, *J. Integer Seq.* **4 (1)** (2001), p. 1–11, MR 1848942 | Zbl 0978.15022.

COEFFICIENT INEQUALITY FOR TRANSFORMS

- [14] R. J. LIBERA et E. J. ZLOTKIEWICZ – “Coefficient bounds for the inverse of a function with derivative in \mathbb{P} ”, *Proc. Amer. Math. Soc.* **87** (1983), p. 251–257, MR 0681830 | Zbl 0488.30010.
- [15] W. C. MA et D. MINDA – “Uniformly convex functions”, *Ann. Polon. Math.* **57(2)** (1992), p. 165 – 175, MR 1182182.
- [16] J. W. NOONAN et D. K. THOMAS – “On the second hankel determinant of areally mean p - valent functions”, *Trans. Amer. Math. Soc.* **223(2)** (1992), p. 337 – 346, MR 0422607 | Zbl 0346.30012.
- [17] K. I. NOOR – “Hankel determinant problem for the class of functions with bounded boundary rotation”, *Rev. Roum. Math. Pures Et Appl.* **28(8)** (1983), p. 731 – 739, MR 0725316 | Zbl 0524.30008.
- [18] C. POMMERENKE – “On the coefficients and Hankel determinants of univalent functions”, *J. London Math. Soc.* **41** (1966), p. 111–122, MR 0185105 | Zbl 0138.29801.
- [19] C. POMMERENKE – *Univalent functions*, Vandenhoeck and Ruprecht, Gottingen, 1975, MR 0507768 | Zbl 0298.30014.
- [20] F. RONNING – “A survey on uniformly convex and uniformly starlike functions”, *Ann. Univ. Mariae Curie - Sklodowska Sect. A.* **47** (1993), p. 123 – 134, MR 1344982 | Zbl 0879.30004.
- [21] B. SIMON – *Orthogonal polynomials on the unit circle, part 1. classical theory*, AMS Colloquium Publ. 54, Part 1, American Mathematical Society, Providence, RI, 2005, MR 2105088 | Zbl 1082.42020.
- [22] D. VAMSHEEKRISHNA et T. RAMREDDY – “Coefficient inequality for uniformly convex functions of order α ”, *J. Adv. Res. Pure Math.* **5(1)** (2013), p. 25–41, MR 3020966.

D. VAMSHEE KRISHNA
 Department of Mathematics
 GIT, GITAM University
 Visakhapatnam- 530 045, A.P., India.
 vamsheekrishna1972@gmail.com

B. VENKATESWARLU
 Department of Mathematics
 GIT, GITAM University
 Visakhapatnam- 530 045, A.P., India.
 bvlmaths@gmail.com

D. VAMSHEE KRISHNA, B. VENKATESWARLU & T. RAMREDDY

T. RAMREDDY
Department of Mathematics,
Kakatiya University,
Warangal- 506 009, A.P., India.
reddytr2@gmail.com