

Original Paper

Stimulation of Suicidal Erythrocyte Death by PRIMA-1

Caterina Faggio^a Kousi Alzoubi^b Salvatrice Calabrò^{a,b} Florian Lang^b^aDepartment of Biological and Environmental Sciences, University of Messina, Agata-Messina, Italy;^bDepartment of Physiology, University of Tübingen, Tübingen, Germany**Key Words**

Phosphatidylserine • PRIMA-1 • Calcium • Ceramide • Eryptosis

Abstract

Background: The anticarcinogenic drug PRIMA-1 (p53 reactivation and induction of massive apoptosis 1) induces suicidal death of tumor cells, an effect in large part attributed to the up-regulation of the proapoptotic transcription factor p53. Erythrocytes are lacking gene transcription but are nevertheless able to enter eryptosis, a suicidal erythrocyte death characterized by cell shrinkage and cell membrane scrambling with phosphatidylserine translocation to the erythrocyte surface. Stimulators of eryptosis include increase of cytosolic Ca²⁺-activity ([Ca²⁺]_i) and ceramide formation. The present study tested whether PRIMA-1 stimulates eryptosis. **Methods:** Phosphatidylserine exposure at the cell surface was estimated from annexin V binding, cell volume from forward scatter, [Ca²⁺]_i from Fluo3-fluorescence, ceramide abundance from binding of specific antibodies, and ROS formation from DCFDA fluorescence. **Results:** A 48 h exposure of human erythrocytes to PRIMA-1 (25 μM) significantly increased the percentage of annexin-V-binding cells without significantly influencing [Ca²⁺]_i or forward scatter. PRIMA-1 (100 μM) induced annexin-V-binding was not significantly blunted by removal of extracellular Ca²⁺ or by the caspase-3 inhibitor zVAD. PRIMA-1 (100 μM) further increased the ceramide abundance at the cell surface and ROS formation. **Conclusions:** PRIMA-1 stimulates phosphatidylserine translocation at the erythrocyte cell membrane, an effect at least partially due to up-regulation of ceramide abundance and ROS formation.

Copyright © 2015 S. Karger AG, Basel

C. Faggio and K. Alzoubi contributed equally and share first authorship.

Prof. Dr. Florian Lang

Physiologisches Institut der Universität Tübingen
Gmelinstr. 5, D-72076 Tübingen (Germany)
Tel. +49 7071 29 72194, Fax +49 7071 29 5618, E-Mail florian.lang@uni-tuebingen.de

Introduction

PRIMA-1 (p53 reactivation and induction of massive apoptosis 1 or APR-246), a widely used investigational drug for cancer therapy successfully tested in phase I/II clinical trials [1-3], triggers apoptosis of tumor cells and augments the tumor cell apoptosis following cytostatic treatment, radiation or hypoxia [1, 3-29]. The substance is at least partially effective by reactivating the proapoptotic transcription factor p53 [1, 3-5, 7, 8, 11-14, 16, 18-21, 23, 25-27, 30]. Moreover, PRIMA-1 up-regulates the related transcription factors p63 and p73 [2, 26]. PRIMA-1 may further stimulate autophagy [31]. In addition, PRIMA-1 derivatives have been shown to stimulate ceramide formation [32].

Erythrocytes lack nuclei and are unable to execute transcription factor dependent gene expression. Nevertheless, erythrocytes may enter suicidal cell death or eryptosis, which is characterized by cell shrinkage [33] and cell membrane scrambling with exposure of phosphatidylserine at the cell surface [34]. Signaling involved in the stimulation of eryptosis includes increase of cytosolic Ca^{2+} activity ($[Ca^{2+}]_i$) [34], formation of ceramide [35], oxidative stress [36], caspase activation [37-41], activation of casein kinase 1 α [42, 43], Janus-activated kinase JAK3 [44], protein kinase C [45], or p38 kinase [46], as well as inhibition or knockout of AMP activated kinase AMPK [47], cGMP-dependent protein kinase [38], PAK2 kinase [48], sorafenib sensitive kinases [49] and sunitinib sensitive kinases [50].

The present study tested, whether PRIMA-1 is able to stimulate eryptosis. To this end, phosphatidylserine surface abundance, cell volume, $[Ca^{2+}]_i$, ceramide abundance, and ROS formation were determined in human erythrocytes from healthy individuals prior to and following treatment with PRIMA-1.

Materials and Methods

Ethics statement

All experiments in this manuscript have been approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki. The study is approved by the ethics committee of the University of Tübingen (184/2003 V).

Erythrocytes, solutions and chemicals

Fresh Lithium-Heparin-anticoagulated blood samples were kindly provided by the blood bank of the University of Tübingen. The blood was centrifuged at 120 rcf for 20 min at 23°C and the platelets and leukocytes-containing supernatant was disposed. Erythrocytes were incubated *in vitro* at a hematocrit of 0.4% in Ringer solution containing (in mM) 125 NaCl, 5 KCl, 1 MgSO₄, 32 N-2-hydroxyethylpiperazine-N-2-ethanesulfonic acid (HEPES), 5 glucose, 1 CaCl₂; pH 7.4 at 37°C for 48 h. Where indicated, erythrocytes were exposed to PRIMA-1 (Sigma Aldrich, Schnelldorf, Germany) at the indicated concentrations, whereby 5 mg PRIMA-1 were solved in 270 μ l H₂O to yield a 100 mM stock solution.

Analysis of annexin-V-binding and forward scatter

After incubation under the respective experimental condition, 50 μ l cell suspension was washed in Ringer solution containing 5 mM CaCl₂ and then stained with Annexin-V-FITC (1:200 dilution; ImmunoTools, Friesoythe, Germany) in this solution at 37°C for 20 min under protection from light. In the following, the forward scatter (FSC) of the cells was determined, and annexin-V-FITC fluorescence intensity was measured with an excitation wavelength of 488 nm and an emission wavelength of 530 nm on a FACS Calibur (BD, Heidelberg, Germany).

Measurement of intracellular Ca²⁺

After incubation, erythrocytes were washed in Ringer solution and then loaded with Fluo-3/AM (Biotium, Hayward, USA) in Ringer solution containing 5 mM CaCl₂ and 5 μ M Fluo-3/AM. The cells were incubated at 37°C for 30 min and washed twice in Ringer solution containing 5 mM CaCl₂. The Fluo-3/AM-loaded erythrocytes were resuspended in 200 μ l Ringer. Then, Ca²⁺-dependent fluorescence intensity

was measured with an excitation wavelength of 488 nm and an emission wavelength of 530 nm on a FACS Calibur.

Determination of ceramide formation

To determine ceramide abundance, a monoclonal antibody-based assay was used. After incubation, cells were stained for 1 h at 37°C with 1 µg/ml anti-ceramide antibody (clone MID 15B4; Alexis, Grünberg, Germany) in phosphate-buffered saline (PBS) containing 0.1 % bovine serum albumin (BSA) at a dilution of 1:10. After two washing steps with PBS-BSA, cells were stained for 30 min with polyclonal fluorescein-isothiocyanate (FITC)-conjugated goat anti-mouse IgG and IgM specific antibody (Pharmingen, Hamburg, Germany) diluted 1:50 in PBS-BSA. Unbound secondary antibody was removed by repeated washing with PBS-BSA. Samples were then analyzed by flow cytometric analysis at an excitation wavelength of 488 nm and an emission wavelength of 530 nm.

Quantification of reactive oxidant species (ROS)

Oxidative stress was determined utilizing 2',7'-dichlorodihydrofluorescein diacetate (DCFDA). After incubation, a 50 µl suspension of erythrocytes was washed in Ringer solution and then stained with DCFDA (Sigma, Schnellendorf, Germany) in Ringer solution containing DCFDA at a final concentration of 10 µM. Erythrocytes were incubated at 37°C for 30 min in the dark and then washed three times in Ringer solution. The DCFDA-loaded erythrocytes were resuspended in 200 µl Ringer solution, and ROS-dependent fluorescence intensity was measured at an excitation wavelength of 488 nm and an emission wavelength of 530 nm on a FACS Calibur (BD).

Statistics

Data are expressed as arithmetic means ± SEM. As indicated in the figure legends, statistical analysis was made using ANOVA with Tukey's test as post-test and *t* test as appropriate. *n* denotes the number of different erythrocyte specimens studied. Since different erythrocyte specimens used in distinct experiments are differently susceptible to triggers of eryptosis, the same erythrocyte specimens have been used for control and experimental conditions.


Results

The present study explored whether PRIMA-1 was capable to induce eryptosis, the suicidal erythrocyte death. The decisive hallmark of eryptosis is phospholipid scrambling of the cell membrane with phosphatidylserine translocation to the cell surface.

In order to identify phosphatidylserine exposing erythrocytes, binding of labeled Annexin-V was determined utilizing flow cytometry. The measurements were performed following incubation of erythrocytes for 48 hours in Ringer solution without or with presence of PRIMA-1 (10 – 100 µM) prior to the measurements. As illustrated in Fig. 1, a 48 h exposure to PRIMA-1 enhanced the percentage of annexin-V-binding erythrocytes, an effect reaching statistical significance at 25 µM PRIMA-1. PRIMA-1 treatment thus resulted in erythrocyte cell membrane scrambling with subsequent translocation of phosphatidylserine to the cell surface. An extended dose response curve is provided in Fig.1C,D. Calculation of an EC50 from log[agonist] vs. normalized response (variable slope) utilizing Graphpad Prism software yielded a value of 3.5 mM. In a separate series, the percentage of Annexin-V binding erythrocytes was significantly ($p < 0.05$) lower when 100 µM PRIMA-1 was added for 24 hours and the erythrocytes were subsequently exposed for additional 24 hours without PRIMA-1 (3.0 ± 0.5 %, $n = 8$) than when PRIMA-1 was added for 48 hours (5.0 ± 0.7 %, $n = 8$).

Alterations of cell volume were evidenced by alterations of forward scatter, which was again determined in flow cytometry. Forward scatter was quantified after incubation of the erythrocytes for 48 h in Ringer solution without or with PRIMA-1 (10 – 100 µM). As illustrated in Fig. 2, incubation of human erythrocytes in Ringer solution with PRIMA-1 tended to slightly decrease forward scatter, an effect, however, not reaching statistical significance.

Fig. 1. Effect of PRIMA-1 on phosphatidylserine exposure. A. Original histogram of annexin-V-binding of erythrocytes (M1) following exposure for 48 h to Ringer solution without (grey area) and with (black line) presence of 100 μM PRIMA-1. B. Arithmetic means \pm SEM of erythrocyte annexin-V-binding (n = 20) following incubation for 48 h to Ringer solution with-



out (white bar) or with (black bars) presence of PRIMA-1 (10 - 100 μM). *($p < 0.05$), ***($p < 0.001$), indicates significant difference from the absence of PRIMA-1 (ANOVA). C,D. The percentage of erythrocytes binding annexin-V as a linear (C) and semilogarithmic (D) function of PRIMA-1 concentration (n=8).




Fig. 2. Effect of PRIMA-1 on erythrocyte forward scatter. A. Original histogram of forward scatter of erythrocytes following exposure for 48 h to Ringer solution without (grey area) and with (black line) presence of 100 μM PRIMA-1. B. Arithmetic means \pm SEM (n = 20) of the normalized erythrocyte forward scatter (FSC) following incubation for 48 h to Ringer solution without (white bar) or with (black bars) presence of PRIMA-1 (10 - 100 μM). No significant differences were observed (ANOVA).

Alterations of cytosolic Ca^{2+} activity ($[\text{Ca}^{2+}]_i$) were estimated utilizing Fluo3 fluorescence, which was again determined in flow cytometry. As shown in Fig. 3A,B, a 48 h exposure to PRIMA-1 (10 - 100 μM) did not significantly modify Fluo3 fluorescence, indicating that PRIMA-1 did not alter appreciably $[\text{Ca}^{2+}]_i$. Further experiments were performed to explore whether the PRIMA-1-induced cell membrane scrambling required entry of extracellular





Fig. 3. Effect of PRIMA-1 on erythrocyte Ca²⁺ activity and Ca²⁺ dependence of PRIMA-1-induced phosphatidylserine exposure. A. Original histogram of Fluo3 fluorescence in erythrocytes following exposure for 48 h to Ringer solution without (grey area) and with (black line) presence of 100 μM PRIMA-1. B. Arithmetic means ± SEM (n = 20) of the Fluo3 fluorescence (arbitrary units) in erythrocytes exposed for 48 h to Ringer solution without (white bar) or with (black bars) presence of PRIMA-1 (10 – 100 μM). No significant differences were observed (ANOVA). C. Original histograms of annexin-V-binding of erythrocytes following exposure for 48 h to Ringer solution in the presence (left) or absence (right) of calcium, without (grey area) and with (black line) presence of 100 μM PRIMA-1. D. Arithmetic means ± SEM (n = 8) of erythrocyte forward scatter after a 48 h treatment with Ringer solution without (white bars) or with (black bars) 100 μM PRIMA-1 in the presence (left bars, +Ca²⁺) and absence (right bars, -Ca²⁺) of calcium. *** (p < 0.001) indicates significant difference from the absence of PRIMA-1.

Fig. 4. PRIMA-1-induced annexin-V-binding in presence of caspase inhibitor zVAD. Arithmetic means ± SEM (n = 8) of the percentage of annexin-V-binding erythrocytes after a 48 h treatment with Ringer solution without (white bars) or with (black bars) 100 μM PRIMA-1 in the absence (left bars) and presence (right bars) of caspase inhibitor zVAD (10 μM). *** (p < 0.001) indicates significant difference from the absence of PRIMA-1 (ANOVA).



Ca²⁺. To this end, the erythrocytes were exposed for 48 h to 100 μM PRIMA-1 in the presence or nominal absence of extracellular Ca²⁺. As shown in Fig. 3C, removal of extracellular Ca²⁺ did not significantly affect the increase of annexin-V-binding following PRIMA-1 treatment.




Fig. 5. Effect of PRIMA-1 on ceramide formation. A. Original histogram of ceramide surface abundance of erythrocytes following exposure for 48 h to Ringer solution without (grey shadow) and with (black line) presence of 100 μM PRIMA-1. B. Arithmetic means \pm SEM (n = 8) of ceramide abundance after a 48 h incubation in Ringer solution without (white bar) or with (black bars) 1-200 μM PRIMA-1. * (p < 0.05), *** (p < 0.001) indicates significant difference from the absence of PRIMA-1 (ANOVA).




Fig. 6. Effect of PRIMA-1 on reactive oxygen species. A. Original histogram of 2',7'-dichlorodihydrofluorescein diacetate (DCFDA) fluorescence in erythrocytes following exposure for 48 hours to Ringer solution without (grey shadow) and with (black line) presence of 100 μM PRIMA-1. B. Arithmetic means \pm SEM (n = 8) of erythrocyte DCFDA fluorescence following incubation for 48 hours to Ringer solution without (white bar) or with (black bar) presence of PRIMA-1 (1 - 200 μM). ** (p < 0.01), *** (p < 0.001) indicates significant difference from the absence of PRIMA-1 (ANOVA).

Instead, PRIMA-1 significantly increased the percentage of annexin-V-binding erythrocytes even in the absence of extracellular Ca^{2+} . Thus, the effect of PRIMA-1 on phosphatidylserine translocation was mediated by a mechanism other than entry of extracellular Ca^{2+} .

A further series of experiments addressed the putative involvement of caspases. To this end, erythrocytes were exposed to 100 μM PRIMA-1 for 48 h either in the absence or presence of the pancaspase inhibitor zVAD (1 or 10 μM). As illustrated in Figure 4, zVAD did not significantly modify the effect of PRIMA-1 on annexin V binding.

As cell membrane scrambling could be triggered without requirement of increased $[\text{Ca}^{2+}]_i$ by ceramide, a further series of experiments explored, whether PRIMA-1-induced cell membrane scrambling was paralleled by formation of ceramide. To this end, the ceramide abundance at the erythrocyte surface was determined utilizing a specific anti-ceramide antibody. As shown in Fig. 5, a 48 h exposure of erythrocytes to PRIMA-1 increased the abundance of ceramide at the erythrocyte surface, an effect reaching statistical significance at 100 μM .

In order to test whether PRIMA-1 enhances oxidative stress, reactive oxygen species (ROS) were determined utilizing 2',7'-dichlorodihydrofluorescein diacetate (DCFDA). As illustrated in fig.6, a 48 h treatment with PRIMA-1 increased DCFDA fluorescence, an effect reaching statistical significance at 100 μM .

Discussion

The present study reveals that exposure of human erythrocytes to high PRIMA-1 concentrations is followed by stimulation of cell membrane scrambling with phosphatidylserine translocation and increase of ceramide abundance at the erythrocyte surface. Phosphatidylserine exposure at the cell surface is a hallmark of eryptosis, the suicidal death of erythrocytes [34].

The cell membrane scrambling following PRIMA-1 treatment was not paralleled by an increase of $[\text{Ca}^{2+}]_i$. Moreover, removal of extracellular Ca^{2+} did not significantly modify the stimulation of eryptosis by PRIMA-1. Instead, PRIMA-1 triggered phosphatidylserine exposure even in the absence of extracellular Ca^{2+} . Thus, PRIMA-1 was effective by mechanisms other than increase of $[\text{Ca}^{2+}]_i$. Moreover, the effect of PRIMA-1 was not significantly modified by the caspase inhibitor zVAD and was thus not dependent on caspase activation. Instead PRIMA-1 stimulated the formation of ceramide, an effect well known to stimulate eryptosis [34]. PRIMA-1 derivatives have most recently been shown to stimulate ceramide formation in tumor cells [32]. Moreover, PRIMA-1 induced oxidative stress, a well known stimulator of erythrocyte cell membrane scrambling [34].

PRIMA-1 tended to decrease cell volume, an effect, however, not reaching statistical significance. Thus, PRIMA-1 failed to significantly trigger the second hallmark of eryptosis. Moreover, PRIMA-1 apparently did not trigger membrane blebbing with formation of small particles, a further hall mark of eryptosis. This observation parallels the lack of PRIMA-1 effect on $[\text{Ca}^{2+}]_i$. Eryptotic cell shrinkage is usually caused by increase of $[\text{Ca}^{2+}]_i$ with subsequent activation of Ca^{2+} sensitive K^+ channels, K^+ exit, hyperpolarization of the cell membrane, Cl^- exit and thus cellular loss of KCl with osmotically obliged water [33]. The possibility must be kept in mind that PRIMA-1 triggers a programmed necrosis, a suicidal death distinct from eryptosis [51].

The PRIMA-1 concentration (25 μM) required for statistically significant stimulation of erythrocyte cell membrane scrambling was higher than those triggering apoptosis in tumor cells [12, 16]. PRIMA-1 tended to increase phosphatidylserine exposure at lower concentrations (10 μM), an effect, however, not reaching statistical significance. It must be kept in mind that erythrocytes may be sensitized to the effect of PRIMA-1 by other xenobiotics stimulating cell membrane scrambling [35, 49, 50, 52-82] or by diseases associated with enhanced cell membrane scrambling, such as sepsis, malaria, sickle cell disease, Wilson's disease, iron deficiency, malignancy, metabolic syndrome, diabetes, hepatic failure, renal insufficiency, hemolytic uremic syndrome, hyperphosphatemia and phosphate depletion [34, 83, 84].

The sensitization of erythrocytes for cell membrane scrambling by ceramide may, at least in theory, be relevant in malaria. The malaria pathogen *Plasmodium* triggers eryptosis by induction of oxidative stress, which activates several ion channels of the host cell membrane including Ca^{2+} -permeable erythrocyte cation channels [85, 86]. The Ca^{2+} entry through unselective cation channels triggers eryptosis with subsequent clearance of the infected erythrocytes from circulating blood [87]. Several genetic erythrocyte disorders including sickle-cell trait, beta-thalassemia-trait, homozygous Hb-C and homozygous G6PD-deficiency enhance the susceptibility of erythrocytes for eryptosis and thus confer some protection against a severe course of malaria [34, 88-90]. Moreover, the clinical course of malaria is favourably influenced by clinical conditions with accelerated eryptosis, such as iron deficiency [91]. Eryptosis triggering xenobiotics shown to favourably influence the clinical course of malaria include lead [92], chlorpromazine [93] or NO synthase inhibitors [94]. Whether or not PRIMA-1 may influence the clinical course of malaria remains to be tested.

On the other hand, eryptosis may lead to anemia due to phagocytosis and subsequent removal of phosphatidylserine exposing erythrocytes. Clinically overt anemia is observed as soon as the rate of eryptosis exceeds the formation of new erythrocytes [34]. The binding of phosphatidylserine exposing erythrocytes to endothelial CXCL16/SR-PSO may in addition lead to adherence of eryptotic erythrocytes to the vascular wall [95]. Phosphatidylserine exposing erythrocytes may further stimulate blood clotting and thrombosis [96-98]. As a result phosphatidylserine exposing erythrocytes may impair microcirculation [35, 96, 99-102].

In conclusion, PRIMA-1 stimulates erythrocyte cell membrane scrambling with phosphatidylserine translocation to the erythrocyte surface, an effect at least partially due to stimulation of ceramide formation.

Disclosure Statement

The authors declare that they have no conflict of interest.

Acknowledgements

The authors acknowledge the meticulous preparation of the manuscript by Tanja Loch. The study was supported by the Deutsche Forschungsgemeinschaft and the Open Access Publishing Fund of Tuebingen University.

References

- 1 Qiang W, Jin T, Yang Q, Liu W, Liu S, Ji M, He N, Chen C, Shi B, Hou P: PRIMA-1 selectively induces global DNA demethylation in p53 mutant-type thyroid cancer cells. *J Biomed Nanotechnol* 2014;10:1249-1258.
- 2 Shen J, van den Bogaard EH, Kouwenhoven EN, Bykov VJ, Rinne T, Zhang Q, Tjabringa GS, Gilissen C, van Heeringen SJ, Schalkwijk J, van Bokhoven H, Wiman KG, Zhou H: APR-246/PRIMA-1(MET) rescues epidermal differentiation in skin keratinocytes derived from EEC syndrome patients with p63 mutations. *Proc Natl Acad Sci U S A* 2013;110:2157-2162.
- 3 Wang T, Lee K, Rehman A, Daoud SS: PRIMA-1 induces apoptosis by inhibiting JNK signaling but promoting the activation of Bax. *Biochem Biophys Res Commun* 2007;352:203-212.
- 4 Aryee DN, Niedan S, Ban J, Schwentner R, Muehlbacher K, Kauer M, Kofler R, Kovar H: Variability in functional p53 reactivation by PRIMA-1(Met)/APR-246 in Ewing sarcoma. *Br J Cancer* 2013;109:2696-2704.
- 5 Bykov VJ, Issaeva N, Selivanova G, Wiman KG: Mutant p53-dependent growth suppression distinguishes PRIMA-1 from known anticancer drugs: a statistical analysis of information in the National Cancer Institute database. *Carcinogenesis* 2002;23:2011-2018.
- 6 Bykov VJ, Zache N, Stridh H, Westman J, Bergman J, Selivanova G, Wiman KG: PRIMA-1(MET) synergizes with cisplatin to induce tumor cell apoptosis. *Oncogene* 2005;24:3484-3491.
- 7 Bykov VJN, Issaeva B, Shilov A, Hultcrantz M, Pugacheva E, Chumakov P, Bergman J, Wiman KG, Selivanova G: Restoration of the tumor suppressor function to mutant p53 by a low molecular weight compound. *Nat Med* 1985;8:282-288.
- 8 Charlot JF, Nicolier M, Pretet JL, Mouglin C: Modulation of p53 transcriptional activity by PRIMA-1 and Pifithrin-alpha on staurosporine-induced apoptosis of wild-type and mutated p53 epithelial cells. *Apoptosis* 2006;11:813-827.
- 9 Cory AH, Chen J, Cory JG: Effects of PRIMA-1 on wild-type L1210 cells expressing mutant p53 and drug-resistant L1210 cells lacking expression of p53: necrosis vs. apoptosis. *Anticancer Res* 2006;26:1289-1295.
- 10 Duan W, Gao L, Wu X, Wang L, Nana-Sinkam SP, Otterson GA, Villalona-Calero MA: MicroRNA-34a is an important component of PRIMA-1-induced apoptotic network in human lung cancer cells. *Int J Cancer* 2010;127:313-320.

- 11 Izetti P, Hautefeuille A, Abujamra AL, de Farias CB, Giacomazzi J, Alemar B, Lenz G, Roesler R, Schwartsmann G, Osvaldt AB, Hainaut P, Ashton-Prolla P: PRIMA-1, a mutant p53 reactivator, induces apoptosis and enhances chemotherapeutic cytotoxicity in pancreatic cancer cell lines. *Invest New Drugs* 2014;10.1007/s10637-014-0090-9
- 12 Kobayashi N, Abedini M, Sakuragi N, Tsang BK: PRIMA-1 increases cisplatin sensitivity in chemoresistant ovarian cancer cells with p53 mutation: a requirement for Akt down-regulation. *J Ovarian Res* 2013;6:7.
- 13 Lambert JM, Gorzov P, Veprintsev DB, Soderqvist M, Segerback D, Bergman J, Fersht AR, Hainaut P, Wiman KG, Bykov VJ: PRIMA-1 reactivates mutant p53 by covalent binding to the core domain. *Cancer Cell* 2009;15:376-388.
- 14 Lee K, Wang T, Paszczynski AJ, Daoud SS: Expression proteomics to p53 mutation reactivation with PRIMA-1 in breast cancer cells. *Biochem Biophys Res Commun* 2006;349:1117-1124.
- 15 Li Y, Mao Y, Brandt-Rauf PW, Williams AC, Fine RL: Selective induction of apoptosis in mutant p53 premalignant and malignant cancer cells by PRIMA-1 through the c-Jun-NH2-kinase pathway. *Mol Cancer Ther* 2005;4:901-909.
- 16 Liang Y, Besch-Williford C, Hyder SM: PRIMA-1 inhibits growth of breast cancer cells by re-activating mutant p53 protein. *Int J Oncol* 2009;35:1015-1023.
- 17 Magrini R, Russo D, Ottaggio L, Fronza G, Inga A, Menichini P: PRIMA-1 synergizes with adriamycin to induce cell death in non-small cell lung cancer cells. *J Cell Biochem* 2008;104:2363-2373.
- 18 Messina RL, Sanfilippo M, Vella V, Pandini G, Vigneri P, Nicolosi ML, Giani F, Vigneri R, Frasca F: Reactivation of p53 mutants by prima-1 [corrected] in thyroid cancer cells. *Int J Cancer* 2012;130:2259-2270.
- 19 Nahi H, Lehmann S, Mollgard L, Bengtzen S, Selivanova G, Wiman KG, Paul C, Merup M: Effects of PRIMA-1 on chronic lymphocytic leukaemia cells with and without hemizygous p53 deletion. *Br J Haematol* 2004;127:285-291.
- 20 Nahi H, Merup M, Lehmann S, Bengtzen S, Mollgard L, Selivanova G, Wiman KG, Paul C: PRIMA-1 induces apoptosis in acute myeloid leukaemia cells with p53 gene deletion. *Br J Haematol* 2006;132:230-236.
- 21 Piantino CB, Reis ST, Viana NI, Silva IA, Morais DR, Antunes AA, Dip N, Srougi M, Leite KR: Prima-1 induces apoptosis in bladder cancer cell lines by activating p53. *Clinics (Sao Paulo)* 2013;68:297-303.
- 22 Rao CV, Patlolla JM, Qian L, Zhang Y, Brewer M, Mohammed A, Desai D, Amin S, Lightfoot S, Kopelovich L: Chemopreventive effects of the p53-modulating agents CP-31398 and Prima-1 in tobacco carcinogen-induced lung tumorigenesis in A/J mice. *Neoplasia* 2013;15:1018-1027.
- 23 Rehman A, Chahal MS, Tang X, Bruce JE, Pommier Y, Daoud SS: Proteomic identification of heat shock protein 90 as a candidate target for p53 mutation reactivation by PRIMA-1 in breast cancer cells. *Breast Cancer Res* 2005;7:R765-774.
- 24 Rieber M, Strasberg-Rieber M: Hypoxia, Mn-SOD and H(2)O(2) regulate p53 reactivation and PRIMA-1 toxicity irrespective of p53 status in human breast cancer cells. *Biochem Pharmacol* 2012;84:1563-1570.
- 25 Rokaeus N, Klein G, Wiman KG, Szekely L, Mattsson K: PRIMA-1(MET) induces nucleolar accumulation of mutant p53 and PML nuclear body-associated proteins. *Oncogene* 2007;26:982-992.
- 26 Rokaeus N, Shen J, Eckhardt I, Bykov VJ, Wiman KG, Wilhelm MT: PRIMA-1(MET)/APR-246 targets mutant forms of p53 family members p63 and p73. *Oncogene* 2010;29:6442-6451.
- 27 Russo D, Ottaggio L, Penna I, Foggetti G, Fronza G, Inga A, Menichini P: PRIMA-1 cytotoxicity correlates with nucleolar localization and degradation of mutant p53 in breast cancer cells. *Biochem Biophys Res Commun* 2010;402:345-350.
- 28 Shi H, Lambert JM, Hautefeuille A, Bykov VJ, Wiman KG, Hainaut P, Caron de Fromental C: In vitro and in vivo cytotoxic effects of PRIMA-1 on hepatocellular carcinoma cells expressing mutant p53ser249. *Carcinogenesis* 2008;29:1428-1434.
- 29 Supiot S, Zhao H, Wiman K, Hill RP, Bristow RG: PRIMA-1(met) radiosensitizes prostate cancer cells independent of their MTP53-status. *Radiother Oncol* 2008;86:407-411.
- 30 Cui B, Yang Q, Guan H, Shi B, Hou P, Ji M: PRIMA-1, a mutant p53 reactivator, restores the sensitivity of TP53 mutant-type thyroid cancer cells to the histone methylation inhibitor 3-Deazaneplanocin A. *J Clin Endocrinol Metab* 2014;99:E962-970.
- 31 Russo D, Ottaggio L, Foggetti G, Masini M, Masiello P, Fronza G, Menichini P: PRIMA-1 induces autophagy in cancer cells carrying mutant or wild type p53. *Biochim Biophys Acta* 2013;1833:1904-1913.
- 32 Soans E, Evans SC, Cipolla C, Fernandes E: Characterizing the sphingomyelinase pathway triggered by PRIMA-1 derivatives in lung cancer cells with differing p53 status. *Anticancer Res* 2014;34:3271-3283.

- 33 Lang PA, Kaiser S, Myssina S, Wieder T, Lang F, Huber SM: Role of Ca²⁺-activated K⁺ channels in human erythrocyte apoptosis. *Am J Physiol Cell Physiol* 2003;285:C1553-C1560.
- 34 Lang E, Qadri SM, Lang F: Killing me softly - suicidal erythrocyte death. *Int J Biochem Cell Biol* 2012;44:1236-1243.
- 35 Abed M, Towhid ST, Mia S, Pakladok T, Alesutan I, Borst O, Gawaz M, Gulbins E, Lang F: Sphingomyelinase-induced adhesion of eryptotic erythrocytes to endothelial cells. *Am J Physiol Cell Physiol* 2012;303:C991-999.
- 36 Tesoriere L, Attanzio A, Allegra M, Cilla A, Gentile C, Livrea MA: Oxysterol mixture in hypercholesterolemia-relevant proportion causes oxidative stress-dependent eryptosis. *Cell Physiol Biochem* 2014;34:1075-1089.
- 37 Bhavsar SK, Bobbala D, Xuan NT, Foller M, Lang F: Stimulation of suicidal erythrocyte death by alpha-lipoic acid. *Cell Physiol Biochem* 2010;26:859-868.
- 38 Foller M, Feil S, Ghoreschi K, Koka S, Gerling A, Thunemann M, Hofmann F, Schuler B, Vogel J, Pichler B, Kasinathan RS, Nicolay JP, Huber SM, Lang F, Feil R: Anemia and splenomegaly in cGKI-deficient mice. *Proc Natl Acad Sci USA* 2008;105:6771-6776.
- 39 Foller M, Mahmud H, Gu S, Wang K, Floride E, Kucherenko Y, Luik S, Laufer S, Lang F: Participation of leukotriene C(4) in the regulation of suicidal erythrocyte death. *J Physiol Pharmacol* 2009;60:135-143.
- 40 Lau IP, Chen H, Wang J, Ong HC, Leung KC, Ho HP, Kong SK: In vitro effect of CTAB- and PEG-coated gold nanorods on the induction of eryptosis/erythroptosis in human erythrocytes. *Nanotoxicology* 2012;6:847-856.
- 41 Maellaro E, Leoncini S, Moretti D, Del Bello B, Tanganelli I, De Felice C, Ciccoli L: Erythrocyte caspase-3 activation and oxidative imbalance in erythrocytes and in plasma of type 2 diabetic patients. *Acta Diabetol* 2013;50:489-495.
- 42 Kucherenko Y, Zelenak C, Eberhard M, Qadri SM, Lang F: Effect of casein kinase 1alpha activator pyrvinium pamoate on erythrocyte ion channels. *Cell Physiol Biochem* 2012;30:407-417.
- 43 Zelenak C, Eberhard M, Jilani K, Qadri SM, Macek B, Lang F: Protein kinase CK1alpha regulates erythrocyte survival. *Cell Physiol Biochem* 2012;29:171-180.
- 44 Bhavsar SK, Gu S, Bobbala D, Lang F: Janus kinase 3 is expressed in erythrocytes, phosphorylated upon energy depletion and involved in the regulation of suicidal erythrocyte death. *Cell Physiol Biochem* 2011;27:547-556.
- 45 Klari BA, Lang PA, Kempe DS, Niemoeller OM, Akel A, Sobiesiak M, Eisele K, Podolski M, Huber SM, Wieder T, Lang F: Protein kinase C mediates erythrocyte "programmed cell death" following glucose depletion. *Am J Physiol Cell Physiol* 2006;290:C244-C253.
- 46 Gatidis S, Zelenak C, Fajol A, Lang E, Jilani K, Michael D, Qadri SM, Lang F: p38 MAPK activation and function following osmotic shock of erythrocytes. *Cell Physiol Biochem* 2011;28:1279-1286.
- 47 Foller M, Sopjani M, Koka S, Gu S, Mahmud H, Wang K, Floride E, Schleicher E, Schulz E, Munzel T, Lang F: Regulation of erythrocyte survival by AMP-activated protein kinase. *FASEB J* 2009;23:1072-1080.
- 48 Zelenak C, Foller M, Velic A, Krug K, Qadri SM, Viollet B, Lang F, Macek B: Proteome analysis of erythrocytes lacking AMP-activated protein kinase reveals a role of PAK2 kinase in eryptosis. *J Proteome Res* 2011;10:1690-1697.
- 49 Lupescu A, Jilani K, Zelenak C, Zbidah M, Qadri SM, Lang F: Hexavalent chromium-induced erythrocyte membrane phospholipid asymmetry. *Biometals* 2012;25:309-318.
- 50 Shaik N, Lupescu A, Lang F: Sunitinib-sensitive suicidal erythrocyte death. *Cell Physiol Biochem* 2012;30:512-522.
- 51 LaRocca TJ, Stivison EA, Hod EA, Spitalnik SL, Cowan PJ, Randis TM, Ratner AJ: Human-specific bacterial pore-forming toxins induce programmed necrosis in erythrocytes. *MBio* 2014;5:e01251-01214.
- 52 Zhang R, Xiang Y, Ran Q, Deng X, Xiao Y, Xiang L, Li Z: Involvement of calcium, reactive oxygen species, and ATP in hexavalent chromium-induced damage in red blood cells. *Cell Physiol Biochem* 2014;34:1780-1791.
- 53 Arnold M, Bissinger R, Lang F: Mitoxantrone-induced suicidal erythrocyte death. *Cell Physiol Biochem* 2014;34:1756-1767.
- 54 Firat U, Kaya S, Cim A, Buyukbayram H, Gokalp O, Dal MS, Tamer MN: Increased caspase-3 immunoreactivity of erythrocytes in STZ diabetic rats. *Exp Diabetes Res* 2012;2012:316384.
- 55 Ganesan S, Chaurasiya ND, Sahu R, Walker LA, Tekwani BL: Understanding the mechanisms for metabolism-linked hemolytic toxicity of primaquine against glucose 6-phosphate dehydrogenase deficient

- human erythrocytes: evaluation of eryptotic pathway. *Toxicology* 2012;294:54-60.
- 56 Bissinger R, Lupescu A, Zelenak C, Jilani K, Lang F: Stimulation of eryptosis by cryptotanshinone. *Cell Physiol Biochem* 2014;34:432-442.
- 57 Jilani K, Qadri SM, Lang F: Geldanamycin-induced phosphatidylserine translocation in the erythrocyte membrane. *Cell Physiol Biochem* 2013;32:1600-1609.
- 58 Zbidah M, Lupescu A, Jilani K, Lang F: Stimulation of suicidal erythrocyte death by fumagillin. *Basic Clin Pharmacol Toxicol* 2013;112:346-351.
- 59 Lupescu A, Bissinger R, Warsi J, Jilani K, Lang F: Stimulation of erythrocyte cell membrane scrambling by gedunin. *Cell Physiol Biochem* 2014;33:1838-1848.
- 60 Jacobi J, Lang E, Bissinger R, Frauenfeld L, Modicano P, Faggio C, Abed M, Lang F: Stimulation of erythrocyte cell membrane scrambling by mitotane. *Cell Physiol Biochem*. 2014;33:1516-1526.
- 61 Lupescu A, Bissinger R, Herrmann T, Oswald G, Jilani K, Lang F: Induction of suicidal erythrocyte death by novobiocin. *Cell Physiol Biochem* 2014;33:670-680.
- 62 Polak-Jonkisz D, Purzyc L: Ca Influx versus Efflux during Eryptosis in Uremic Erythrocytes. *Blood Purif* 2012;34:209-210.
- 63 Qian EW, Ge DT, Kong SK: Salidroside protects human erythrocytes against hydrogen peroxide-induced apoptosis. *J Nat Prod* 2012;75:531-537.
- 64 Bissinger R, Modicano P, Frauenfeld L, Lang E, Jacobi J, Faggio C, Lang F: Estramustine-induced suicidal erythrocyte death. *Cell Physiol Biochem* 2013;32:1426-1436.
- 65 Vota DM, Maltaner RE, Wenker SD, Nesse AB, Vittori DC: Differential erythropoietin action upon cells induced to eryptosis by different agents. *Cell Biochem Biophys* 2013;65:145-157.
- 66 Weiss E, Cytlak UM, Rees DC, Osei A, Gibson JS: Deoxygenation-induced and Ca(2+) dependent phosphatidylserine externalisation in red blood cells from normal individuals and sickle cell patients. *Cell Calcium* 2012;51:51-56.
- 67 Zappulla D: Environmental stress, erythrocyte dysfunctions, inflammation, and the metabolic syndrome: adaptations to CO2 increases? *J Cardiometab Syndr* 2008;3:30-34.
- 68 Abed M, Herrmann T, Alzoubi K, Pakladok T, Lang F: Tannic acid induced suicidal erythrocyte death. *Cell Physiol Biochem* 2013;32:1106-1116.
- 69 Malik A, Bissinger R, Calabrò S, Faggio C, Jilani K, Lang F: Aristolochic Acid Induced Suicidal Erythrocyte Death. *Kidney Blood Press Res* 2014;39:408-419.
- 70 Abed M, Zoubi KA, Theurer M, Lang F: Effect of dermaseptin on erythrocytes. *Basic Clin Pharmacol Toxicol* 2013;113:347-352.
- 71 Ahmed MS, Langer H, Abed M, Voelkl J, Lang F: The uremic toxin acrolein promotes suicidal erythrocyte death. *Kidney Blood Press Res* 2013;37:158-167.
- 72 Ghashghaenia M, Cluitmans JC, Toulany M, Saki M, Koberle M, Lang E, Dreischer P, Biedermann T, Duszenko M, Lang F, Bosman GJ, Wieder T: Age Sensitivity of NFkappaB Abundance and Programmed Cell Death in Erythrocytes Induced by NFkappaB Inhibitors. *Cell Physiol Biochem* 2013;32:801-813.
- 73 Abed M, Feger M, Alzoubi K, Pakladok T, Frauenfeld L, Geiger C, Towhid ST, Lang F: Sensitization of erythrocytes to suicidal erythrocyte death following water deprivation. *Kidney Blood Press Res* 2013;37:567-578.
- 74 Alzoubi K, Honisch S, Abed M, Lang F: Triggering of Suicidal Erythrocyte Death by Penta-O-galloyl-beta-D-glucose. *Toxins (Basel)* 2014;6:54-65.
- 75 Jilani K, Lang F: Carmustine-induced phosphatidylserine translocation in the erythrocyte membrane. *Toxins (Basel)* 2013;5:703-716.
- 76 Jilani K, Enkel S, Bissinger R, Almilaji A, Abed M, Lang F: Fluoxetine induced suicidal erythrocyte death. *Toxins (Basel)* 2013;5:1230-1243.
- 77 Lupescu A, Bissinger R, Jilani K, Lang F: Triggering of suicidal erythrocyte death by celecoxib. *Toxins (Basel)* 2013;5:1543-1554.
- 78 Lupescu A, Jilani K, Zbidah M, Lang F: Patulin-induced suicidal erythrocyte death. *Cell Physiol Biochem* 2013;32:291-299.
- 79 Arnold M, Lang E, Modicano P, Bissinger R, Faggio C, Abed M, Lang F: Effect of nitazoxanide on erythrocytes. *Basic Clin Pharmacol Toxicol* 2014;114:421-426.
- 80 Oswald G, Alzoubi K, Abed M, Lang F: Stimulation of suicidal erythrocyte death by ribavirin. *Basic Clin Pharmacol Toxicol* 2014;114:311-317.

- 81 Bissinger R, Malik A, Jilani K, Lang F: Triggering of Erythrocyte Cell Membrane Scrambling by Salinomycin. *Basic Clin Pharmacol Toxicol* 2014;10.1111/bcpt.12250
- 82 Gao M, Lau PM, Kong SK: Mitochondrial toxin betulinic acid induces in vitro eryptosis in human red blood cells through membrane permeabilization. *Arch Toxicol* 2014;88:755-768.
- 83 Voelkl J, Alzoubi K, Mamar AK, Ahmed MS, Abed M, Lang F: Stimulation of suicidal erythrocyte death by increased extracellular phosphate concentrations. *Kidney Blood Press Res* 2013;38:42-51.
- 84 Lang E, Gatidis S, Freise NF, Bock H, Kubitz R, Laueremann C, Orth HM, Klindt C, Schuier M, Keitel V, Reich M, Liu G, Schmidt S, Xu HC, Qadri SM, Herebian D, Pandyra AA, Mayatepek E, Gulbins E, Lang F, Haussinger D, Lang KS, Foller M, Lang PA: Conjugated bilirubin triggers anemia by inducing erythrocyte death. *Hepatology* 2014;10.1002/hep.27338
- 85 Duranton C, Huber S, Tanneur V, Lang K, Brand V, Sandu C, Lang F: Electrophysiological properties of the Plasmodium falciparum-induced cation conductance of human erythrocytes. *Cell Physiol Biochem* 2003;13:189-198.
- 86 Kirk K: Membrane transport in the malaria-infected erythrocyte. *Physiol Rev* 2001;81:495-537.
- 87 Foller M, Bobbala D, Koka S, Huber SM, Gulbins E, Lang F: Suicide for survival--death of infected erythrocytes as a host mechanism to survive malaria. *Cell Physiol Biochem* 2009;24:133-140.
- 88 Ayi K, Giribaldi G, Skorokhod A, Schwarzer E, Prendergast PT, Arese P: 16alpha-bromoepiandrosterone, an antimalarial analogue of the hormone dehydroepiandrosterone, enhances phagocytosis of ring stage parasitized erythrocytes: a novel mechanism for antimalarial activity. *Antimicrob Agents Chemother* 2002;46:3180-3184.
- 89 Ayi K, Turrini F, Piga A, Arese P: Enhanced phagocytosis of ring-parasitized mutant erythrocytes: a common mechanism that may explain protection against falciparum malaria in sickle trait and beta-thalassemia trait. *Blood* 2004;104:3364-3371.
- 90 Cappadoro M, Giribaldi G, O'Brien E, Turrini F, Mannu F, Ulliers D, Simula G, Luzzatto L, Arese P: Early phagocytosis of glucose-6-phosphate dehydrogenase (G6PD)-deficient erythrocytes parasitized by Plasmodium falciparum may explain malaria protection in G6PD deficiency. *Blood* 1998;92:2527-2534.
- 91 Koka S, Foller M, Lamprecht G, Boini KM, Lang C, Huber SM, Lang F: Iron deficiency influences the course of malaria in Plasmodium berghei infected mice. *Biochem Biophys Res Commun* 2007;357:608-614.
- 92 Koka S, Huber SM, Boini KM, Lang C, Foller M, Lang F: Lead decreases parasitemia and enhances survival of Plasmodium berghei-infected mice. *Biochem Biophys Res Commun* 2007;363:484-489.
- 93 Koka S, Lang C, Boini KM, Bobbala D, Huber SM, Lang F: Influence of chlorpromazine on eryptosis, parasitemia and survival of Plasmodium berghei infected mice. *Cell Physiol Biochem* 2008;22:261-268.
- 94 Koka S, Lang C, Niemoeller OM, Boini KM, Nicolay JP, Huber SM, Lang F: Influence of NO synthase inhibitor L-NAME on parasitemia and survival of Plasmodium berghei infected mice. *Cell Physiol Biochem* 2008;21:481-488.
- 95 Borst O, Abed M, Alesutan I, Towhid ST, Qadri SM, Foller M, Gawaz M, Lang F: Dynamic adhesion of eryptotic erythrocytes to endothelial cells via CXCL16/SR-PSOX. *Am J Physiol Cell Physiol* 2012;302:C644-C651.
- 96 Andrews DA, Low PS: Role of red blood cells in thrombosis. *Curr Opin Hematol* 1999;6:76-82.
- 97 Chung SM, Bae ON, Lim KM, Noh JY, Lee MY, Jung YS, Chung JH: Lysophosphatidic acid induces thrombogenic activity through phosphatidylserine exposure and procoagulant microvesicle generation in human erythrocytes. *Arterioscler Thromb Vasc Biol* 2007;27:414-421.
- 98 Zwaal RF, Comfurius P, Bevers EM: Surface exposure of phosphatidylserine in pathological cells. *Cell Mol Life Sci* 2005;62:971-988.
- 99 Closse C, Dachary-Prigent J, Boisseau MR: Phosphatidylserine-related adhesion of human erythrocytes to vascular endothelium. *Br J Haematol* 1999;107:300-302.
- 100 Gallagher PG, Chang SH, Rettig MP, Neely JE, Hillery CA, Smith BD, Low PS: Altered erythrocyte endothelial adherence and membrane phospholipid asymmetry in hereditary hydrocytosis. *Blood* 2003;101:4625-4627.
- 101 Pandolfi A, Di Pietro N, Sirolli V, Giardinelli A, Di Silvestre S, Amoroso L, Di Tomo P, Capani F, Consoli A, Bonomini M: Mechanisms of uremic erythrocyte-induced adhesion of human monocytes to cultured endothelial cells. *J Cell Physiol* 2007;213:699-709.
- 102 Wood BL, Gibson DF, Tait JF: Increased erythrocyte phosphatidylserine exposure in sickle cell disease: flow-cytometric measurement and clinical associations. *Blood* 1996;88:1873-1880.