

Feeding behaviour and daily ration of the spotted flounder *Citharus linguatula* (Osteichthyes: Pleuronectiformes) in the central Tyrrhenian Sea

PAOLO CARPENTIERI, TIZIANA CANTARELLI, FRANCESCO COLLOCA,
ALESSANDRO CRISCOLI and GIANDOMENICO ARDIZZONE

Animal and Human Biology Department, University of Rome "La Sapienza", Viale dell'Università 32, 00185 Rome, Italy.
E-mail: paolo.carpentieri@uniroma1.it

SUMMARY: Spotted flounder *Citharus linguatula* (Linnaeus, 1758) is one of the most abundant species within demersal fish assemblages next to the continental shelf (up to 200 m depth) of the Mediterranean Sea. In spite of the ecological importance of this species in the central Tyrrhenian Sea, many aspects of its biology (e.g. feeding behaviour) are poorly described. In order to analyze how feeding habits are related to the ontogenetic changes in this species, 2636 specimens (between 9 and 26 cm of total length TL) were collected from November 2000 to January 2002. During ontogenetic development spotted flounder occupied different trophic levels: diet shifted from mysids, predominantly in the smaller spotted flounder (<11 cm TL), towards fish in the larger individuals (>19 cm TL). Before the transition to the ichthyophagous phase, a more generalist foraging behaviour including decapods (*Processa* sp., *Alpheus glaber*), molluscs (*Turitella* sp.) and benthic fish (*Lesueurigobius* spp., *Gobius niger*, *Callionymus* spp., *Arnoglossus laterna*) dominated the diet. Seasonal analysis of the diet showed an increase in the presence of mysids in summer and autumn, and of decapods in summer and spring. Teleosteans are present throughout the year without significant variations. Daily ration values, computed from the Eggers method, ranged from 1.57% to 2.83% wet body weight, showing a negative relationship between reproduction and daily consumption rate.

Keywords: spotted flounder, feeding, Mediterranean Sea, ontogenesis, Pleuronectiformes.

RESUMEN: COMPORTAMIENTO ALIMENTICIO Y RACIÓN DIARIA DE LA SOLLETA *CITHARUS LINGUATULA* (OSTEICHTHYES: PLEURONECTIFORMES) EN EL MAR TIRRENO CENTRAL. – La solleta, *Citharus linguatula*, es una de las especies más abundantes entre las comunidades de peces de la plataforma continental (hasta los 200 m de profundidad) del Mediterráneo. A pesar de su importancia ecológica en el mar Tirreno central, muchos aspectos de la biología de *C. linguatula* (e.g. alimentación) continúan siendo desconocidos. Con el objetivo de investigar cómo los hábitos alimenticios están relacionados con los cambios ontogénicos en esta especie, se analizaron 2636 individuos, entre 9 y 26 cm de longitud total (LT), que fueron muestreados entre noviembre de 2000 y enero de 2002. Durante el desarrollo ontogénico *C. linguatula* ocupa diferentes niveles tróficos: la alimentación varía desde los misidáceos, que son predominantes para las solletas más pequeñas (<11 cm LT), hasta los peces, que son la presa más importante para los individuos de mayor talla (>19 cm LT). Antes de llegar a la fase piscívora, el régimen alimenticio es más generalista, incluyendo crustáceos decápodos (*Processa* sp., *Alpheus glaber*), moluscos (*Turitella* sp.) y peces bentónicos (*Lesueurigobius* spp., *Gobius niger*, *Callionymus* spp., *Arnoglossus laterna*). Los análisis estacionales de la dieta muestran un aumento de la presencia de misidáceos en verano y otoño, y un aumento de decápodos en verano y primavera. Los teleosteos están presentes todo el año sin ninguna variación significativa. Los valores de la tasa diaria de alimentación, calculados con el método de Eggers, variaron entre 1.57% y 2.83% del peso húmedo individual, con una relación negativa entre la reproducción y el consumo diario.

Palabras clave: solleta, alimentación, Mediterráneo, desarrollo ontogénico, Pleuronectiformes.

INTRODUCTION

The spotted flounder *Citharus linguatula* (Linnaeus, 1758) is a benthic species distributed in the whole Mediterranean basin and along the Atlantic coasts of Portugal, Morocco and Angola (Nielsen, 1986). It generally inhabits sandy and muddy bottoms of the continental shelf up to 200 m depth. In the Mediterranean basin the species provides an important contribution to commercial landings: it is regularly found at fish markets and is an important by-catch accompanying species for the trawling fleet. Together with *Lepidorhombus boschii* and *Solea vulgaris*, it is the most important and abundant resource of the Mediterranean Sea in the group of flatfishes (Sartor *et al.*, 2002). In the Italian part of the Mediterranean it is exploited mainly by a fleet that uses a peculiar kind of trawl, called *rapido*, and it is caught as an associated species of the target common sole (*S. vulgaris*) and scallops (*Aequipecten opercularis*, *Chlamys glabra* and *Pecten jacobaeus*).

Most of the available information on spotted flounder comes from studies on distribution and abundance (Jardas, 1984; García-Rodríguez and Esteban, 2000; Sartor *et al.*, 2002), age, growth (Vassilopoulou and Papaconstantinou, 1994; Çakır *et al.*, 2005) and sexual cycle and reproduction (Sabatés, 1988). However, its feeding habits have been poorly investigated (Planas and Vives, 1956; Redon *et al.*, 1994). Despite the increasing interest in life-cycle and energetic demands (e.g. feeding ecology and daily ration) of the demersal communities, no information is available, at least for the Mediterranean, on the ontogenesis and daily ration of this species. As Wootton (1990) pointed out, the foraging behaviour of all animals in a community is the key feature of the community stability and structure.

The aim of this paper is (1) to provide information on the daily and seasonal feeding behaviour of *C. linguatula*; (2) to analyze its ontogenesis-related diet changes; (3) to produce estimates of its food consumption based on daily ration models.

MATERIAL AND METHODS

Study area

Spotted flounder sampling was carried out on the continental shelf of the Tyrrhenian Sea. This area extends down to 120-150 m depth, covering 15-30 km of distance from the shoreline (Fig. 1). The sea surface temperature ranges from 12-15°C in the winter season to 27-28°C in the summer; near the bottom (from a depth of 100 m down to 200 m) temperature is constant around 13°C.

Most of the fishing area is characterized by sandy-muddy bottoms (SFBC-VTC benthic assemblage, Pérès and Picard [1964]) and muddy bottoms (VTC) between 10 and 150 m depth.

Sampling

Size-stratified samples were obtained monthly during a ground fish survey on board a commercial *rapido* trawler. The *rapido* trawl used is made of an iron frame provided with skids and a toothed bar on its lower side; a wooden plank is fitted to the front of the iron frame to act as a spoiler and keep the gear in contact with the seabed. The net is 4.8 m long and consists of two portions: the first one connected to the *rapido* mouth has a stretched mesh size of 24 by 30 cm; the second one is the codend, with a mesh size of 39.9 cm. The




FIG. 1. – Study area. Points represent the trawl stations located on the continental shelf of the central western coasts of Italy.

lower portion of the net is protected by a reinforced rubber diamond-mesh matting (stretched mesh size 24*30 mm).

Two *rapido* trawls are usually towed at a speed of about 5.5-6 knots and each trawl-haul covered a distance of approximately 20 km. The fishing activity is carried out for about 14-16 hours from night (about 01:00) to dusk (about 19:00) without stopping. A total of about 6-8 hauls per fishing day are performed. To standardize the data, the hauls were pooled according to the time of the day into 2-hour time steps.

In order to emphasize the main diet changes in relation to the growth processes, spotted flounder were collected monthly from November 2000 to the end of January 2002. Seasons were defined as follows: samples carried out after 23 September were classified as autumn hauls; after 21 December as winter hauls; after 21 March as spring hauls and after 21 June as summer hauls. Samples were collected at least every 2 hours over a period of about 20 h to test for a diel cycle.

Samples of *C. linguatula* were frozen immediately after catch and stored at -20°C to prevent further digestion of prey. Later, fish were brought back to the laboratory where individual total body weight (BW, measured to the nearest 0.01 g), and total length (TL, rounded to the next half centimetre) were measured. Stomachs were removed and preserved in 70% ethanol solution, and prey were identified to the lowest possible taxonomic level, counted, and weighed (to the nearest 0.01 mg). When the state of digestion was more advanced, prey were sorted and grouped into unidentified Osteichthyes, Crustacea or Mollusca. The collected data of stomach contents were classified with respect to the time of the day, fish length and season.

Sex and maturity stages were analyzed macroscopically. The maturity scale for partial spawner fish was used to classify the stage of maturation (Holden and Raitt, 1974).

Data analysis

The size at which 50% of the fish were mature (TL_{50}) was calculated using the following logistic function (Sparre *et al.*, 1989):

$$P_{TL} = 1/(1 + e^{(S_1 - S_2 TL_{50})})$$

where P_{TL} is the percentage of mature individuals for length classes; S_1 and S_2 are constants and TL_{50} is given by S_1/S_2 .

Size-related dietary variations were investigated by dividing all the specimens into 2 cm length-classes, with the exception of the largest individuals (>21 cm TL) all included into a single length-class.

Hierarchical cluster analysis and non-metric multidimensional scaling (NMDS), based on Bray-Curtis similarity and on the IRI%, were used for classification and ordination of spotted-flounder size classes (Clarke and Warwick, 1994).

The contribution of each food item to the diet, for the different length classes and for the size dietary groups individuated by the NMDS and Cluster Analysis was evaluated using the index of relative importance (IRI, Pinkas *et al.*, 1971) as modified by Hacunda (1981). This index, expressed as:

$$IRI\% = IRI^{-\sum IRI} * 100$$

incorporates the percentage by number (N%), wet weight (W%) and frequency of occurrence (F%) (Hyslop, 1980).

The diel feeding periodicity was investigated by means of stomach fullness index, expressed as the percentage ratio of the weight of a fish's stomach contents to its total body weight ($FI\% = W_s/W_t * 100$) (Tudela and Palomera, 1995).

One-way ANOVA was used to test for differences in stomach fullness (FI%) between relative fish abundance (expressed as number of individuals per hour n^{h-1}) and day/night hauls, and for differences between seasons and size groups. The differences were considered significant at $P < 0.05$.

The estimate of daily ration, based on the ratio of the percent stomach weight to predator weight (%BW), was obtained through the model of Eggers (1977):

$$C_{(24E)} = F_{24} R_{24}$$

where F_{24} is the mean stomach fullness index over 24 h and R_{24} is the instantaneous gastric evacuation rate. This model assumes that the fullness indexes at the beginning and at the end of a 24 h period are equal (Boisclair and Marchand, 1993). When this assumption cannot be met and the weight of food in the stomach at the beginning and at the end of the 24 h cycles are significantly different, Eggers (1979) proposed a corrected model:

$$C_{(24EC)} = C_{(24E)} + (F_{t=24} - F_{t=0})$$

where $F_{t=24}$ and $F_{t=0}$ are the values of stomach fullness index at the end and the beginning of the sampling period.

In order to estimate daily ration the instantaneous coefficient of gastric evacuation R must be obtained. Since the instantaneous food evacuation rate of this species had never been evaluated before, the values used here were obtained from temperature-based models (R' - Durbin *et al.*, 1983; Worobec, 1984; Macpherson, 1985) and determined from our experimental data set following the R_{max} procedure (Héroux and Magnan, 1996).

The first model, the most widely accepted in the literature to describe food evacuation in most fish species, is probably more appropriate for describing the evacuation of small, relatively easily digested particles from the stomach (Worobec, 1984; Macpherson, 1985). This model assumes $R' = 0.0175T - 0.0442$ for

TABLE 1. – Number of fish samples and stomach contents collected during the four sampling periods. For each season the value of the stomach fullness (FI), the length range and the mean standard length of the collected samples are reported.

Season	n. of fish	n. of stomachs	FI	length range (cm)	mean length (cm)
Autumn	1080	589	0.55	9.5 - 25	14.4
Summer	559	376	0.67	10.5 - 26	15.4
Winter	505	256	0.51	10 - 24	14.2
Spring	492	272	0.55	9 - 23.5	15.0

fish feeding on polychaetes or small crustaceans, and $R' = 0.00408T - 0.00527$ for fish eating predators, where T is the temperature in Celsius degrees. Evacuation rate R' depends primarily on water temperature and on the prey food type, whereas it is not influenced by food size and by the frequency of feeding (Elliott and Persson, 1978; Durbin *et al.*, 1983; Macpherson, 1985). The constant temperature regime in the study area, around $13^{\circ}\text{C} \pm 0.5$ down to 50-200 m depth all year around (Ifremer, 1997), means that our estimates of daily ration were probably free of temperature influence.

The R_{max} procedure assumes the occurrence of a period of non-feeding between two consecutive sampling times. Once the period of non-feeding is identified, the stomach can be assumed to be exclusively egesting and R_{max} can be computed as the difference between the initial and final stomach contents, following an exponential model:

$$R_{max} = \ln FI_{(T+1)} - \ln FI_{(T)/T}$$

where $FI_{(T+1)}$ and $FI_{(T)}$ are the values of the stomach fullness index at the end and at the beginning of the sampling period over a time interval T . In this study the gastric evacuation rate R_{max} was calculated for two different fish size groups: specimens under 13 cm TL and specimens over 13 cm TL (the switch of 13 cm corresponds to the attainment of maturity size). Thereafter, daily ration values were calculated for the two groups mentioned and for the different seasons.

RESULTS

Diet in relation with length

A total of 2636 specimens with a mean standard length of 14.7 cm (range: 9-26 cm TL) and a mean weight of 26.2 g (range: 7.5-146 g) were collected for diet analysis (Table 1). Size at maturity (L_{50}) was estimated to be 13.3 cm ($R = 0.95$; 91.31 of the variance explained) (Fig. 2); individuals over 18 cm in length were almost all female (70%).

The total number of preys was about 1800 divided into 54 different food items (Table 2). Cluster and NMDS analysis (stress=0.01), based on the IRI%, identified three cluster groups below 65% similarity, separated along a length-size gradient (Fig. 3a). The first group (A) contained fish between 9 and 12.9 cm TL;




FIG. 2. – Size at maturity of *Citharus linguatula*. Points represent the percentage of mature individuals for each size class.




FIG. 3. – Dendrogram (a) and NMDS plot (b) based on the IRI% values, using group-average clustering from Bray-Curtis similarity on diet data.

the second group (B) contained fish between 13 and 18.9 cm TL; the third group (C) contained specimens longer than 19 cm TL (Fig. 3b). Mysids (*Leptomysis gracilis* IRI= 50.2%) and small teleosts (Gobiidae IRI= 36.3%) dominated the diet of group A. Seasonal dietary patterns of group A, shown as mysids, were important in the stomachs of the spotted flounder throughout the year (Fig. 4), with a significant increase ($P < 0.01$) in summer (IRI% = 55.2) and autumn (IRI% = 51.9). During both periods, mysids were represented by various species, *Gastrosaccus spinifer*, *Anchialina agilis* and *Leptomysis apiops*, though the greatest contribution of this group to the diet is due to the presence of *L. gracilis*. Teleosts were the most frequent prey item in winter (IRI% = 51.7) and spring (IRI% = 52.1), but were also an important prey item throughout the year.

TABLE 2. – Prey items reported as index of relative importance (IRI %) for the seven size classes. The three size groups (A, B and C) identified by the cluster analysis are indicated.

Size group	A		III	B		V	C	
	I	II		IV	VI		VII	
Length (cm)	9.0-10.9	11.0-12.9	13.0-14.9	15.0-16.9	17.0-18.9	19.0-20.9	>21.0	
Number of stomachs	38	452	1016	742	297	70	21	
Mollusca								
<i>Allotheutis</i> spp.				0.001				
<i>Hinia incrassata</i>			0.02	0.005	0.01			
<i>Parvicardium scabrum</i>			0.001					
<i>Pusillina lineolata</i>			0.001					
<i>Sepietta</i> spp.				0.01				
Sepiolidae n.i.			0.01					
<i>Turritella biplicata</i>			0.03	0.03	0.02			
<i>Turritella communis</i>			0.002		0.02			
Polichaeta								
Polichaeta n.i.			0.001	0.001		0.15		
Crustacea								
<i>Alpheus glaber</i>		0.04	13.63	22.96	27.09	22.74	16.29	
<i>Anapagurus petiti</i>				0.001				
<i>Anchialina agilis</i>	0.27	0.04	0.002	0.001				
<i>Anilocra</i> sp.		0.05	0.01	0.02	0.13	0.17		
<i>Apseudes latrellei</i>					0.01			
<i>Athanas nitiscens</i>			0.001					
Axiidae n.i.			0.001	0.001				
<i>Brachicarpus biungulatus</i>		0.005		0.001				
<i>Callionassa</i> spp.				0.004				
Crangonidae n.i.	1.10	0.38	0.16	0.13	0.01			
Crustacea n.i.	2.44	0.36	0.07	0.12	0.18			
Decapoda n.i.	0.51	3.36	1.62	1.03	0.90	5.07		
<i>Erytrops</i> spp.	0.29	0.16						
<i>Gastrosaccus lobatus</i>				0.001				
<i>Gastrosaccus normani</i>	0.01	0.005			0.01			
<i>Gastrosaccus spinifer</i>	0.27		0.002	0.01				
<i>Gastrosaccus</i> spp.	2.55	0.04	0.01	0.004	0.02			
<i>Goneplax rhomboides</i>		0.01						
Isopoda n.i.		0.04	0.01	0.001				
<i>Leptomysis apiops</i>	2.50		0.01	0.00				
<i>Leptomysis gracilis</i>	49.48	49.37	18.61	12.63	4.82	1.38	1.33	
<i>Leptomysis</i> spp.	15.50	0.03	0.06	0.06				
<i>Lophogaster typicus</i>		0.02	0.02	0.001				
Misidacea n.i.		0.71	0.42	0.18	0.04			
Natantia n.i.	0.69	1.49	0.39	0.58	0.16			
<i>Philocheras bispinosus</i>	1.18	1.34	0.44	0.18	0.25	0.16		
<i>Philocheras monachantus</i>		0.01	0.02	0.06	0.02			
<i>Philocheras</i> spp.	1.81	2.12	0.29	0.33	0.25	0.16		
<i>Processa caniculata</i>			0.02	0.01	0.62			
<i>Processa elegantula</i>			0.07	0.04	0.19			
<i>Processa</i> spp.		0.08	0.11	0.12	0.45	0.75		
<i>Squilla mantis</i>			0.001	0.01				
<i>Upogebia</i> spp.					0.01			
<i>Upogebia tipica</i>			0.012		0.01			
Teleostea								
<i>Arnoglossus laterna</i>		0.02	0.003	0.19	0.05	5.95		
<i>Callionimus</i> spp.		0.05	0.002	0.03				
<i>Cepola macrophthalmia</i>	0.75	0.31	0.13	0.001	0.12			
<i>Deltentosteus quadrimaculatus</i>	0.76	0.82	2.86	2.36	4.47	1.25	13.54	
<i>Engraulis encrasicolus</i>				0.02	0.80	1.21		
Gobiidae n.i.	12.92	27.54	31.79	28.87	16.46	17.99	13.26	
<i>Gobius niger</i>		0.28	0.80	1.75	0.76	10.11	6.40	
<i>Leuseriogobius</i> spp.	4.21	4.09	1.56	1.78	0.71		1.51	
<i>Leuseriogobius sueri</i>			10.34	6.10	4.16	14.54	3.67	
<i>Serranus hepatus</i>				0.003			0.04	
Teleostea n.i.	2.78	7.23	16.47	20.35	37.26	18.37	43.96	

Group B showed a more heterogeneous diet, still characterized by a high occurrence of mysids (IRI= 12.7%), but also with a considerable amount of fish (IRI= 59%) and decapods (IRI= 24.2%) such as *Alpheus glaber*, *Processa* spp. and *Philocheras* spp. In this size group (spotted flounder from 13 to 18.9 cm TL) the presence of mollusc gasteropods (*Turritella*

spp., *Hynia incrassata*) with lower percentages (IRI= 0.8%) was recorded. Dietary patterns (Fig. 4) showed significant differences among seasons (P<0.05), mainly linked to the high occurrence of decapods in spring (IRI%= 30) and mysids in summer (IRI%= 10.5).

The proportion of teleosts increased strongly in Group C, where they accounted for about 80% in




Fig. 4. – Seasonal variation in percentage abundance (IRI%) of the different prey items for the three spotted flounder size groups identified by the Cluster and NMDS analysis.




Fig. 5. – Proportion (%) in number of fish prey occurring in the diet of *Citharus linguatula* during its growth.

number of the spotted flounder diet (Fig. 5). Teleosts were present throughout the year without significant variations ($P > 0.05$). The main prey items were *Deltentosteus quadrimaculatus* (IRI= 3.8%),

Leuseriogobius suerii (IRI= 10.9%) and *Gobius niger* (IRI= 10.5%), and other Pleuronectiformes such as *Arnoglossus laterna* (IRI= 2.7%). *A. glaber* (IRI= 20%) was the main prey item among crustaceans decapods. Slight seasonal differences ($P < 0.05$) in this size group (spotted flounder >19 cm TL) were mainly linked to the high occurrence of decapods in summer (IRI%= 38.5).

Diel feeding periodicity and daily ration

The diel pattern indicated that the presence of spotted flounder was more or less evenly distributed, with no significant relationship ($P > 0.05$) with either daylight or night hauls. Stomach fullness increased significantly ($P < 0.01$) by 10:00 and continued to rise until 15:00 (Fig. 6). During the early morning, just before sunrise (from 04:00 to 08:00 h) and before sunset (from




Fig. 6. – Relationship between time and stomach fullness expressed as %FI (weight of a fish's stomach contents to its total body weight) in the different sampling periods. Hauls were pooled according to the time of the day into 2-hours time steps.

TABLE 3. – Daily ration estimates (expressed as percentage of fish body weight - %BW) obtained with the Eggers (1979) model, both with the direct estimation of the consumption rate R_{max} and with the instantaneous evacuation rate from bibliographic review R' . SD, standard deviation.

Daily ration	Summer	Spring	Winter	Autumn
$R' = 0.18$ (<13 cm)	2.19	2.21	2.83	2.23
SD \pm	0.23	0.19	0.41	0.4
n.stomach	131	110	109	207
$R' = 0.07$ (>13 cm)	1.63	1.95	2.16	1.71
SD \pm	0.19	0.14	0.15	0.09
n.stomach	245	162	147	382
$R_{max} = 0.082$ (< 13 cm)	1.83	2.01	2.27	1.85
SD \pm	0.23	0.31	0.36	0.21
n.stomach	131	110	109	207
$R_{max} = 0.064$ (> 13 cm)	1.57	1.93	2.12	1.68
SD \pm	0.19	0.18	0.23	0.12
n.stomach	245	162	147	382

17:00 to 19:00) the feeding intensity was observed to be significantly low ($P < 0.01$).

Mature spotted flounder (>13 cm TL) showed a marked daily ration peak in winter (2.16% BW $R' = 0.07$; 2.12% BW $R_{max} = 0.064$), while lower values, from 1.57 ($R_{max} = 0.064$) to 1.63% BW ($R' = 0.07$), were obtained in summer (Table 3). Immature specimens (<13 cm TL) also showed a marked daily ration peak in winter (2.83% BW $R' = 0.18$; 2.27% BW $R_{max} = 0.082$), while in the other seasons the values were quite constant: from 1.83% BW to 2.01% BW with $R_{max} = 0.082$, and from 2.19% BW to 2.23% BW with $R' = 0.18$.

DISCUSSION

In the diet of the spotted flounder 54 different prey items belonging to five major groups were identified. Teleosts, decapods and mysids were the main prey items in all seasons, while molluscs and polychaetes were secondary and complementary preys. In the western Mediterranean, Planas and Vives (1956) and de Juan *et al.* (2007) found the diet of *C. linguatula* to consist mainly of crustaceans and fishes. Belghyti *et al.* (1993) reported that mysids appeared to be the main trophic group in the diet of spotted flounder along the Atlantic coasts of Morocco, with teleosts considered as secondary preys, and crustaceans and cephalopods as occasional preys.

During their ontogenetic development spotted flounder occupied different trophic levels. Diet shifted from mysids—the main preys for smaller specimens (<10.9 cm TL)—to fish (*L. sueri*, *G. niger*, *D. quadrimaculatus*), which in the larger individuals (>19 cm TL) represented almost the entire diet. Specimens between 11 and 19 cm TL showed a more heterogeneous diet, still characterized by the occurrence of mysids, but also with a considerable amount of fish, decapods and other groups such as polychaetes and gasteropod molluscs. This progressive passage from small preys to larger ones is typical for many Pleuronectiformes (Deniel, 1975; Braber and De Groot, 1973; Lockwood, 1984). Such changes in diet related to size are also

connected to morphologic changes associated with the growth of the fish, the increase of the mouth dimensions and probably a greater disposition to move (Wootton, 1990). As reported in the northwestern Mediterranean by de Juan *et al.* (2007), *C. linguatula* specimens of different length essentially preyed upon the same species and the only detectable changes were differences in ingested abundance of these preys.

Seasonal analysis of the diet showed an increase in the presence of mysids in summer and autumn, and an increase in the presence of decapods in summer and spring. Teleosts were present throughout the year without significant variations. Seasonal differences probably reflect temporal variability in prey resources throughout the year. In the western Mediterranean mysid abundance reaches a maximum in late summer-autumn (Franqueville, 1971), which coincides with the period of highest consumption by juvenile spotted flounder. The same seasonal trend, linked to the greater frequency index and numerical composition of the mysid *L. gracilis* in autumn, is confirmed by the study carried out in the Gulf of Valencia by Redon *et al.* (1994). This result has been reported as the main factor influencing diet composition of generalist and opportunistic feeders, such as some Pleuronectiformes species in the Mediterranean (Cabral *et al.*, 2002). In our study, the feeding activity of *C. linguatula* showed a marked daily temporal pattern, with a major peak occurring during daylight and a decline just before sunset. This cycle was found in all samples taken and could be linked to the biology of the species: the spotted flounder normally lives close to the bottom, feeding during daylight on benthic invertebrates and resting during the night (Nielsen, 1986). These results agree with a series of studies carried out on other Pleuronectiformes in which diurnal predator activity was evidenced (De Groot, 1971).

Among other factors, our findings showed that the main shift in the diet might be related to the attainment of sexual maturity. The size-at-maturity (13.3 cm TL) was positively associated with a shift in the diet towards bigger preys (i.e. decapods and fish). Under the first maturity size, diet was basically dominated by mysids and small crustaceans. The shift to utilization of large preys might be induced by gonads maturation. As reported by Ross (1978), the change in feeding patterns during growth can be related to particular phases of fish biological cycles. Such an ontogenetic diet shift could indicate that inter- and intraspecific partitioning of energy resources in fish vary with fish size and probably reduce intraspecific competition between immature and mature specimens.

Given the scant information available on the biology and feeding activity of the species studied, it is difficult to compare differences in daily consumption with its habits and metabolic requirements. Comparing our daily ration results with previously published data at similar temperature ranges (13°C), we noticed that the daily ration values (from 1.57% BW to 2.83% BW)

computed for spotted flounder fall well within the range of estimates obtained by species displaying a similar diet (from 1.4%BW to 3.82%BW; Cartes *et al.*, 2004; Carpentieri *et al.*, 2006). In the Atlantic Ocean for species with diets based on small crustaceans (e.g. *Coelorhynchus fasciatus* and *Pseudopleuronectes americanus*), Macpherson (1985) obtained a daily ration ranging from 0.55 to 2.15%BW whereas Worobec (1984) obtained a daily ration ranging from 1.27 to 3.31%BW.

Our findings showed a negative relationship between spawning and daily ration. The spawning activity of spotted flounder took place from July to October (in summer around 75% of the specimens over 13 cm TL showed ripe gonads), corresponding to the lowest values of daily ration computed for the species (from 1.57%BW to 1.63%BW). As already observed in other species, the reduction of food intake during spawning may also have been due to less space available in the body cavity for food, or could have been associated with a change in hormone levels at spawning that might reduce individual foraging (Hoar *et al.*, 1983; Fordham and Trippel, 1999).

The literature suggests that the interaction between feeding and reproduction is both highly variable and complex. Redon *et al.* (1994) found that the high percentage of empty stomachs was recorded in August to September, and may be related to the reproductive period of the spotted flounder (late summer), during which there is a significant decrease in food intake. In the western Mediterranean, Planas and Vives (1956) determined the spawning season of *C. linguatula* to be summer, corresponding to the maximum vacuity index values recorded in the same period. A similar pattern of decreasing daily ration during the reproduction period was already identified for other species in the Mediterranean (e.g. the synchronous hermaphrodite *Serranus hepatus* - Carpentieri *et al.*, 2006). In the Atlantic area, there is strong evidence that Atlantic herring (*Clupea harengus*), four-spot flounder (*Hippoglossina oblonga*) and red hake (*Urophycis chuss*) exhibit a “rester” pattern of feeding after spawning during the resting phase (Link and Burnett, 2001).

On the other hand, *Mullus barbatus* (Ünlüoglu *et al.*, 2002; Carpentieri *et al.*, 2006), *Macroramphosus scolopax* (Carpentieri *et al.*, 2006) and two Argentinidae, *Argentina sphyraena* and *Glossanodon leioglossus*, exhibited an increase in the daily consumption during spawning (Cohen, 1990; Carpentieri *et al.*, 2006). Link and Burnett (2001) demonstrated that the highest levels of daily consumption recorded for the Atlantic mackerel (*Scomber scombrus*) and silver hake (*Merluccius bilinearis*) occurred at times of increased sexual activity for both species. As these fish developed gonads and reached maturity, the total stomach contents were significantly higher than during the spent and resting stages (Link and Burnett, 2001).

The determination of the daily ration of *C. linguatula* presented here, applying the Eggers model, is the

first study of its kind carried out for this species in the Mediterranean. The advantage of this model was related to the use of field data with unstressed individuals under a natural feeding regime. Data obtained provided a broad overview of the ecology of the complex scenario present on the Mediterranean platform. The results are generally comparable with the estimated values found for species with similar trophic habits in both the Mediterranean and Atlantic areas. It is plausible that differences in daily ration estimates could be attributed to various factors, such as abundance of prey types, feeding rhythm and digestion rates.

REFERENCES

- Belghyti, D., P. Aguesse and C. Gabrion. – 1993. Ethologie alimentaire de *Citharus linguatula* et *Dicologlossa cuneata* sur les côtes atlantiques du Maroc. *Vie Milieu*, 43(2-3): 95-108.
- Boisclair, D. and F. Marchand. – 1993. The guts to estimate fish daily ration. *Can. J. Fish. Aquat. Sci.*, 50: 1969-1975.
- Braber, L. and S.J. De Groot. – 1973. The food of five flatfish species (Pleuronectiformes) in the southern North Sea. *Neth. J. Sea Res.*, 6: 163-172.
- Cabral, H.N., M. Lopes and R. Loeper. – 2002. Trophic niche overlap between flatfish in a nursery area on the Portuguese coast. *Sci. Mar.*, 66(3): 293-300.
- Çakır, D.T., B. Bayhan, B. Hoşsucu, A. Ünlüoglu and S. Akalyın. – 2005. Some Parameters of the Population Biology of Spotted Flounder (*Citharus linguatula*, Linnaeus, 1758) in Edremit Bay (North Aegean Sea). *Turk. J. Vet. Anim. Sci.*, 29: 1013-1018.
- Carpentieri, P., F. Colloca, A. Belluscio, A. Criscoli and G.D. Ardizzone. – 2006. Diel feeding periodicity and daily ration of shelf break species. *J. Mar. Biol. Assoc. U.K.*, 86: 853-860.
- Cartes, J.E., J. Rey, D. Lloris and L. Gil de Sola. – 2004. Influence of environmental variables on the feeding and diet of European hake (*Merluccius merluccius*) on the Mediterranean Iberian coasts. *J. Mar. Biol. Assoc. U.K.*, 84: 831-835.
- Clarke, K.R. and R.M. Warwick. – 1994. *Change in marine communities: an approach to statistical analysis and interpretation*. Plymouth Marine Laboratory, Plymouth.
- Cohen, D.M. – 1990. Argentinidae. p. 235-238. In J.C. Quero, J.C. Hureau, C. Karrer, A. Post and L. Saldanha (eds.). Check-list of the fishes of the eastern tropical Atlantic (CLOFETA). JNICT, Lisbon; SEI, Paris; and UNESCO, Paris. (1): 235.
- De Groot, S.J. – 1971. On the interrelationship between morphology of the alimentary tract, food and feeding behaviour in flatfishes (Pisces: Pleuronectiformes). *Neth. J. Sea Res.*, 5: 121-196.
- de Juan, S., J.E. Cartes and M. Demestre. – 2007. Effects of commercial trawling activities in the diet of the flat fish *Citharus linguatula* (Osteichthyes: Pleuronectiformes) and the starfish *Astropecten irregularis* (Echinodermata: Asteroidea). *J. Exp. Mar. Biol. Ecol.*, (349): 152-169.
- Deniel, C. – 1975. Régimes alimentaires d'*Arnoglossus thori* Kyle et d'*Arnoglossus imperialis* Rafinesque (Teleosteens-Botidae) en Baie de Douarnenez. *Rev. Trav. Inst. Pêches Marit.*, 39: 105-116.
- Durbin, E.G., A.G. Durbin, R.W. Langton and R.E. Bowman. – 1983. Stomach contents of silver hake, *Merluccius bilinearis*, and Atlantic cod, *Gadus morhua*, and estimation of their daily rations. *Fish. Bull. NOAA*, 81: 437-454.
- Eggers, D.M. – 1977. Factors in interpreting data obtained by diel sampling of fish stomachs. *J. Fish. Res. Bd Can.*, 34: 290-294.
- Eggers, D.M. – 1979. Comment on some recent methods for estimating food consumption by fish. *J. Fish. Res. Bd Can.*, 36: 1019-20.
- Elliott, J.M. and L. Persson. – 1978. The estimation of daily rates of food consumption for fish. *J. Anim. Ecol.*, 47: 977-991.
- Fordham, S.E. and E.A. Trippel. – 1999. Feeding behaviour of cod (*Gadus morhua*) in relation to spawning. *J. Appl. Ichthyol.*, 15: 1-9.
- Franqueville, C. – 1971. Macroplankton profond (Invertébrés) de la Méditerranée nord-occidentale. *Tethys*, 3(1): 11-56.

- García-Rodríguez, M. and A. Esteban. – 2000. Contribution to Knowledge of *Citharus linguatula* (Linnaeus, 1758) (Osteichthyes: Heterosomata) in the Iberian Mediterranean. In: Demersal resources in the Mediterranean, *Actes de Colloques IFREMER*, 26: 131-140.
- Hacunda, J.S. – 1981. Trophic relationships among demersal fishes in a coastal area of the Gulf of Maine. *Fish. Bull.*, 79(4): 775-788.
- Heroux, D. and P. Magnan. – 1996. In situ determination of food daily ration in fish: review and field evaluation. *Environ. Biol. Fishes*, 46: 61-74.
- Hoar, W.S., D.J. Randall and E.M. Donaldson. – 1983. Fish physiology, Vol. 9. Reproduction, Part A. Endocrine tissues and hormones. Academic Press, New York.
- Holden, M.J. and D.F.S. Raitt. – 1974. Manual of fisheries science. Part 2: Methods of resources investigation and their application. *FAO Fish. Tech. Paper*, 115, Rev. 1: 1-214.
- Hyslop, E.J. – 1980. Stomach contents analysis: a review of methods and their application. *J. Fish Biol.*, 17: 411-429.
- Ifremer. – 1997. Mediterranean hydrological atlas (CD rom). IFREMER edition, Brest, France.
- Jardas, I. – 1984. *Citharus macrolepidotus* (Bloch, 1787) (Pisces Pleuronectiformes) nourishment and the length-weight relationship in the Adriatic Sea. *Ichthyologia*, 16(1-2): 1-4.
- Link, J.S. and J. Burnett. – 2001. The relationship between stomach contents and maturity state for major northwest Atlantic fishes: new paradigms? *J. Fish Biol.*, 59: 783-794.
- Lockwood, S.J. – 1984. The daily food intake of 0-group plaice (*Pleuronectes platessa* L.) under natural conditions: changes with size and season. *J. Cons.*, 41(2): 181-193.
- Macpherson, E. – 1985. Daily ration and feeding periodicity of some fishes off the coast of Namibia. *Mar. Ecol. Prog. Ser.*, 26: 253-260.
- Nielsen, J. – 1986. Citharidae. pp 1286. In: P.J.P. Whitehead, M.L. Bauchot, J.C. Hureau, J. Nielsen and E. Tortonese (eds.), *Fish of the North-eastern Atlantic and the Mediterranean*. Vol. II, UNESCO, Paris.
- Pérès, J.M. and J. Picard. – 1964. Nouveau manuel de bionomie benthique de la mer Méditerranée. *Rec. Trav. Stat. Mar. Endoume*, 31(47): 1-137.
- Pinkas, L., M.S. Oliphant and I.L.K. Iverson. – 1971. Food habits of albacore, bluefin tuna and bonito in California waters. *Fish. Bull. NOAA*, 152: 1-150.
- Planas, A. and F. Vives. – 1956. Contribución al estudio de la solleta (*Citharus linguatula* Gunth) del Mediterraneo occidental. *Invest. Pesq.*, 3: 107-131.
- Redon, M.J., M.S. Morte and A. Sanz-Brau. – 1994. Feeding habits of the spotted flounder *Citharus linguatula* off the eastern coast of Spain. *Mar. Biol.*, 120: 197-201.
- Ross, S.T. – 1978. Trophic ontogeny of the leopard sea-robin, *Priodontus scitulus* (Pisces: Triglidae). *Fish. Bull. NOAA*, 76(1): 225-234.
- Sabatés, A. – 1988. Larval development and spawning of *Citharus linguatula* (Linnaeus, 1758) in the Western Mediterranean. *J. Plank. Res.*, 10: 1131-1140.
- Sartor, P., M. Sbrana, N. Ungaro, C.A. Marano, C. Piccinetti and C. Piccinetti Manfrin. – 2002. Distribution and abundance of *Citharus linguatula*, *Lepidorhombus boschii*, and *Solea vulgaris* (Osteichthyes: Pleuronectiformes) in the Mediterranean Sea. *Sci. Mar.*, 66(Suppl. 2): 83-102.
- Sparre, P., E. Ursin and S.C. Venema. – 1989. Introduction to tropical fish stock assessment. Part I. Manual. *FAO Fish. Tech. Paper*, 306(1): 1-337.
- Tudela, S. and I. Palomera. – 1995. Diel feeding intensity and daily ration in the anchovy *Engraulis encrasicolus* in the northwest Mediterranean Sea during the spawning period. *Mar. Ecol. Prog. Ser.*, 129: 55-61.
- Ünlüoğlu, A., B. Cihangir, M. Kaya, H.L. Benli and T. Katagan. – 2002. Variations in the feeding intensity and diet composition of red mullet (*Mullus barbatus*) during 24-h period in the summertime in Hisarönü Bay. *J. Mar. Biol. Assoc. U.K.*, 82: 527-528.
- Vassilopoulou, V. and C. Papaconstantinou. – 1994. Age, growth and mortality of the spotted flounder (*Citharus linguatula* Linnaeus, 1758) in the Aegean Sea. *Sci. Mar.*, 58: 261-267.
- Wootton, R.J. – 1990. *Ecology of teleosts fishes*. Chapman and Hall, London.
- Worobec, M.N. – 1984. Field estimation of the daily ration of winter flounder, *Pseudopleuronectes americanus* (Walbaum) in a southern New England Pond. *J. Exp. Mar. Biol. Ecol.*, 77: 183-196.

Scient. ed.: I. Palomera.

Received April 24, 2009. Accepted January 21, 2010.

Published online June 29, 2010.