



## Appraisal of ingestion and digestibility in growing rabbits using near infrared reflectance spectroscopy (NIRS) of feeds and faeces

Giorgia Meineri, Pier Giorgio Peiretti & Giorgio Masoero

To cite this article: Giorgia Meineri, Pier Giorgio Peiretti & Giorgio Masoero (2009) Appraisal of ingestion and digestibility in growing rabbits using near infrared reflectance spectroscopy (NIRS) of feeds and faeces, Italian Journal of Animal Science, 8:1, 75-82, DOI: [10.4081/ijas.2009.75](https://doi.org/10.4081/ijas.2009.75)

To link to this article: <https://doi.org/10.4081/ijas.2009.75>



Copyright 2009 Taylor & Francis Group LLC



Published online: 01 Mar 2016.



Submit your article to this journal [↗](#)



Article views: 22



View related articles [↗](#)



# Appraisal of ingestion and digestibility in growing rabbits using near infrared reflectance spectroscopy (NIRS) of feeds and faeces

Giorgia Meineri<sup>1</sup>, Pier Giorgio Peiretti<sup>2</sup>, Giorgio Masoero<sup>3</sup>

<sup>1</sup>Dipartimento di Produzioni Animali, Epidemiologia ed Ecologia. Università di Torino, Italy

<sup>2</sup>Istituto di Scienze delle Produzioni Alimentari. Consiglio Nazionale delle Ricerche, Torino, Italy

<sup>3</sup>Consiglio per la Ricerca e la Sperimentazione in Agricoltura. Torino, Italy

*Corresponding author:* Dr. Giorgia Meineri. Dipartimento di Produzioni Animali, Epidemiologia ed Ecologia. Università di Torino. Via Leonardo da Vinci 44, 10095 Grugliasco (TO), Italy - Tel. +39 011 6709209 - Fax: +39 011 6709240 - Email: giorgia.meineri@unito.it

---

*Paper received June 30, 2008; accepted October 5, 2008*

---

## ABSTRACT

The aim of this work was to examine whether faecal profiling using NIRS could be profitable for promote the Best Available Techniques (BAT) in the rational feeding of rabbits. A set of 51 feed samples, taken from 12 experimental diets, and of 66 dried grouped faeces samples, belonging to four nutrition experiments, with 130 *ad libitum* registered feed intakes (CV=25%), were submitted to a UV-Vis-NIRS scan (350-2500 nm) in order to calibrate the chemical composition and nutritional parameters, the ingestion aptitude and digestibility. A chemometric system has made it possible to contemporary use the spectrum of the input diet concatenated together with the spectra of the related output pool of the dried faeces. The daily measured feed intake, in absolute or in relative terms as ingestion per unit of metabolic weight, obtained a good resolution for the spectra of the feeds ( $R^2_{cv}=0.80$  and  $0.75$ , respectively), for the faeces ( $0.81$  and  $0.80$ ) and for the joint evaluation of the concatenated spectra ( $0.87$  and  $0.81$ ). The intake was positively correlated to the mineral, insoluble ash, protein, gross energy, crude fiber and acid detergent fiber (ADF) content in the feeds, and negatively correlated to the N-free extract, lignocellulose and all the digestibility coefficients, except crude fiber. Very significant improvements, on average equal to  $0.20 R^2$  points, were also provided to the digestibility coefficients when using the concatenated method; in decreasing order: neutral detergent fiber ( $R^2_{cv}=0.00$ ,  $0.18$  and  $0.50$  for the feeds, faeces and concatenated, respectively), ADF ( $0.00$ ,  $0.45$  and  $0.62$ ), ether extract ( $0.53$ ,  $0.52$  and  $0.86$ ), crude protein ( $0.53$ ;  $0.53$  and  $0.75$ ), and gross energy ( $0.61$ ;  $0.74$  and  $0.83$ ). The results corroborate previous knowledge and show the possibility of using NIRS faecal profiling in rabbit nutrition, which together with the NIRS of the feeds, could contribute to nitrogen monitoring.

*Key words:* Rabbits, Digestibility, Intake, Faecal profiling, Concatenated NIR Spectra.

## RIASSUNTO

VALUTAZIONE DELL'INGESTIONE E DELLA DIGERIBILITÀ DI MANGIMI IN CONIGLI  
IN ACCRESCIMENTO TRAMITE SPETTROSCOPIA NIR DEL MANGIME E DELLE FECI

*Lo scopo del presente lavoro è rivolto all'uso della diagnostica faecale mirata alla promozione delle Migliori Tecniche Disponibili (MTD, direttiva 96/61/CE) nell'allevamento razionale del coniglio. Un gruppo di 51*

mangimi derivati da 12 diete sperimentali e di 66 pool di feci provenienti da quattro esperimenti di alimentazione, con 130 controlli individuali di ingestione volontaria ( $CV=25\%$ ), furono sottoposti a scansione UV-Vis-NIRS (350-2500 nm) al fine di correlare i parametri nutrizionali e di composizione chimica con l'attitudine all'ingestione e la digeribilità. Un sistema chemometrico concatenato ha reso possibile il contemporaneo impiego degli spettri della dieta e dei relativi pool di feci. L'ingestione giornaliera misurata, in assoluto come consumo volontario e in relativo come appetibilità per unità di peso metabolico, ha evidenziato una buona risoluzione per lo spettro degli alimenti ( $R^2_{cv}=0,80$  e  $0,75$ , rispettivamente), delle feci ( $0,81$  e  $0,80$ ) e degli spettri concatenati ( $0,87$  e  $0,81$ ). Il consumo volontario era correlato positivamente a componente minerale, ceneri acido insolubili, proteina grezza, energia grezza, fibra grezza e fibra resistente al detergente acido (ADF) nei mangimi, e correlato negativamente a estrattivi inazotati, lignina e a tutti i coefficienti di digeribilità, ad eccezione di quelli della fibra grezza. Elaborando spettri concatenati dei mangimi (input) e delle relative feci (output), miglioramenti consistenti, mediamente pari a 0,20 punti di  $R^2_{cv}$  assoluto, corrispondenti all'87% come relativo, sono stati conseguiti per i coefficienti di digeribilità di fibra NDF (+41% per il metodo concatenato vs la media di mangimi e feci), ADF (+39%), estratto etereo (+34%), proteina grezza (+22%), energia grezza (+15%), fibra grezza (+11%), sostanza secca (+7%), sostanza organica (+6%) ed estrattivi inazotati (+3%). Questi risultati confermano quelli di precedenti ricerche e dimostrano la possibilità d'impiego nello studio della nutrizione dei conigli della spettroscopia NIRS, che, applicata alle feci unitamente alla lettura dei mangimi, potrebbe contribuire al monitoraggio dell'escrezione azotata di questa specie.

Parole chiave: Conigli, Digeribilità, Ingestione, Profilo fecale, Spettri NIR combinati.

## Introduction

The past five decades have witnessed the emergence of rapid methods which have become useful tools in the field of experimental biology and chemistry. Near infrared spectroscopy (NIRS) has been used for several years for the commercial testing of feeds; it is a rapid and convenient alternative to traditional chemical methods which are time consuming, polluting and expensive. In EU research programmes on rabbit nutrition (FAIR3-1651), the ERAFE group has examined collective feed and faeces databases from 164 experimental groups using NIRS (Xiccato *et al.*, 1999, 2003) and concluded that the method is reliable in the prediction of the chemical constituents, digestibility and energy values of rabbit compound feeds for a wide variation in chemical composition and ingredients. In the present work, our approach has been directed to faecal profiling using NIRS. This method was originally developed for free-range ruminants, where feeding is mainly

based on natural pastures which can have highly variable chemical compositions. The NIRS-NUTBAL programme (Mc Dryden, 2003) rules on the surveillance of nutritional scarcity in the field. In the research field, NIRS has been extended to the estimation of feed intake in ruminants in an easier way than the alkanes technique (Garnsworthy and Unal, 2004). In ruminants, NIR faecal profiling capitalises on the connection of organic matter digestibility to the N contents of the forages ( $R^2=0.82$ ; Garnsworthy and Unal, 2004). Decruyenaere *et al.* (2008) studied two large databases with 1034 forages and faeces in sheep and confirmed the relevance of faecal profiling. They also discovered the opportuneness of a joint elaboration of multi-NIR spectra, constituted by the input spectra of the feed concatenated with the output spectra of the individual or pooled faeces. In this way, the maximum  $R^2$  was achieved by fitting the grass OM intake to OM digestibility. The aim of the present work has been to examine whether faecal profiling using NIRS could be profitable

to promote the Best Available Techniques (BAT) in the modern feeding of rabbits.

## Material and methods

The database was obtained from four experiments related to studies on the inclusion of chia (*Salvia hispanica* L.) seeds (Meineri and Peiretti, 2007), false flax (*Camelina sativa* L.) seeds (Peiretti *et al.*, 2007), golden flax (*Linum usitatissimum*) seeds (Peiretti and Meineri, 2008b), and microalgae (*Spirulina platensis*) (Peiretti and Meineri, 2008a) in rabbit diets. The young cross-bred rabbits, originating from New Zealand White females, were reared in individual cages. The collection of the faeces lasted for 4 days, after which the 130 rabbits weighed on average  $2812 \pm 421$  grams (CV=15%) and their individual daily intake, on a fed basis, was  $160 \pm 40$  grams  $d^{-1}$  (CV=25%). Daily samples of the single faeces of the rabbits were pooled by groups and then stored at  $-20$  °C. After thawing, the samples were dried in a ventilated oven at  $60$  °C till constant weight and then ground to pass a 1mm sieve. Digestibility was calculated through the acid insoluble ash indirect method (Vogtmann *et al.*, 1975). After pooling of the faeces, the elaborated set consisted of 51 feed samples belonging to 12 experimental diets and 66 dried faeces pooled samples.

The spectroscopy scan was conducted using a Model LSP 350-2500P LabSpec Pro portable spectrophotometer (ASD, Analytical Spectral Devices Inc., Boulder, CO) which was equipped to collect spectra from 350 to 2500 nm. An A122100 ASD Model high-intensity reflectance probe was used as an external light source (a 2900 K colour temperature quartz halogen light) to illuminate the objects of interest. This probe can be used to collect reflectance spectra on an area as large as 25 mm in diameter. The reflected light was collected through a 04-

14766 ASD Model 1-m long fiber optic jump cable that consisted of a bundle of forty-four 200- $\mu$ m fibers. The spectra were collected with the LabSpec Pro software "sample spectrum count" option set at 20, meaning that 20 spectra were collected and averaged per sample.

All the spectra were mathematically pre-treated as Standard Normal Variates with Detrend (SNVD), derived and then smoothed using the 1441 mode. Multivariate statistical evaluation was performed through the Modified Partial Least Squares (MPLS) method of the WinISI II software (Infrasoft International, Port Matilda, PA, USA) which was chosen to perform the chemometric, using a cross-validation system to assess the optimal number of latent variables, and allowing one passage for the elimination of any outliers ( $t > 2$ ;  $H > 10$ ).


The spectra were processed separately according to their origin, feed or faeces. Then the two spectra of the feed and the corresponding faeces, each consisting of 2151 digits, were joined in a single concatenated spectrum (Figure 1) and mathematically pre-treated and calibrated in the same way as the 1441 mode. The coefficient of determination in cross validation mode ( $R^2_{cv}$ ) was used for the assessment evaluation of the equations, while the Standard Errors of Validation were compared with the available reference papers for digestibility values.

## Results and discussion

In comparison to the EU database of Xiccató *et al.* (2003), the standard deviations of the feed compounds (Table 1) were higher for the fibrous components, neutral detergent fiber (NDF), acid detergent fiber (ADF) and acid detergent lignin (ADL) and, consequently, for the gross energy and the organic matter content.


The correlation coefficients of the feed in-

Figure 1. A concatenated UV-Vis-NIR spectrum of a feed (left) and its related faeces (right).



X: 350-2500 nm axis repeated on 4302 digits; Y=Log (1/R).

Figure 2. Correlation coefficients of the daily feed intake with the chemical composition of the feeds and feces and with the digestibility coefficients.



Note: abbreviation of variables are reported in table 1 and 2, first column.

take with the chemical parameters on the DM basis of the feeds and faeces are shown in Figure 2 together with the digestibility coefficients. Strong positive relationships appeared for the ash, acid insoluble ash, crude protein, gross energy, crude fiber and ADF in the feeds, while the intake was depressed by high contents of N-free extract and ADL. Unlike to ruminants, where NDF plays a central role on the intake, the NDF does not apparently regulate ingestion ( $r=0.05$ ) in rabbits. On the other hand, the ADF in the feeds appeared more active in increasing the intake of rabbits (0.68) and this correlation was also reflected by the ADF of the faeces (0.44). The chemical composition of the faeces also appeared to be linked to the intake, but only weakly. On the other hand, a corresponding lowering in the digestibility coefficients should be expected for an increased level of intake and this was true for all the components of the feed, with the exception of the crude fiber in the feeds (0.43) and in faeces (0.44). The higher intakes were related to lower levels of the digestible energy contents in the experimental diets. The measured *ad-libitum* feeding parameters, in absolute terms as intake, and in relative terms as ingestion per unit of metabolic weight, obtained a good resolution from the spectra of the feeds ( $R^2_{cv}=0.80$  and 0.75, respectively), from the faeces (0.81 and 0.80) and from the joint evaluation of the concatenated spectra (0.87 and 0.81) (Tables 1 and 2).

Garnsworthy and Unal (2004) concluded that using NIRS to predict alkane concentrations in bovine faeces did not lead to accurate estimates of DMI, but a direct prediction of DMI by NIRS instead gave estimates with a similar accuracy to that derived using the traditional alkane technique.

Very significant improvements, on average by some 0.20 absolute  $R^2_{cv}$  points, corresponding to 87% as relative, were also

provided for the digestibility coefficients estimated by the concatenated spectra. The following appeared to be advantaged: NDF (+41% for concatenated *vs* the average of feeds and faeces), ADF (+39%), ether extract (+34%), crude protein (+22%), gross energy (+15%), crude fiber (+11%), dry matter (+7%), organic matter (+6%) and the N-free extract (+3%). These results, in spite of our higher standard deviations, agree well with the findings reported in EU experiments conducted by Xiccato *et al.* (1999; 2003). The Standard Error of Validation are compared in Tables 1 and 2, which show some improvement in the NIRS estimation of apparent digestibility coefficients for crude protein (SEV 1.3% *vs* 2.6%) and gross energy (1.5 *vs* 1.9 MJ  $kg^{-1}$  DM). Caution must be taken in extrapolating the NIRS estimates of the intake and ingestion parameters because, in spite of the 130 individual values of the intakes, with a 25% coefficient of variation, there were only 12 spectra in the experimental rations in the database, a rather limited number. This ingestion aspect, which is fundamental in ruminant nutrition, has been poorly investigated by rabbit scientists especially from the point of view of rapid NIRS utilization, considering the fact that no feeds tables exist that report information about the ingestion of simple feeds or forages.

## Conclusions

The results confirm the ability of the new chemometric method which uses the concatenated spectra of feeds and faeces in the analyses of experimental trials. They also corroborate results by previous researches and show the possibility of using NIRS faecal profiling in rabbit nutrition, which joined together with NIRS of the feeds, could optimize nitrogen monitoring. In the BAT challenge, further samples of dried and ground

Table 1. Calibration and cross-validation of feed composition and apparent digestibility to the UV-Vis-NIR Spectra of the feed or the concatenated feed-and-faeces spectra.

Composition	#	Mean	SD	Feeds				Joint		SEV X1c1	SEV X1c2	
				CV%	RSQ	SECV	R <sup>2</sup> cv	RSQ	SECV			R <sup>2</sup> cv
DM Intake	45	0.1612	0.0303	19%	0.99	0.01	0.80	0.99	0.0091	0.87	-	-
Ingest	49	0.0722	0.0086	12%	0.98	0.00	0.75	0.95	0.0030	0.81	-	-
DM	49	91.11	0.85	1%	0.67	0.56	0.59	-	-	-	0.48	-
OM	48	92.45	1.30	1%	0.97	0.56	0.81	-	-	-	0.86	-
ASH	48	7.55	1.30	17%	0.97	0.56	0.81	-	-	-	-	-
GE	49	18.01	1.56	9%	0.99	0.64	0.83	-	-	-	0.25	0.26
CP	47	17.35	1.44	8%	0.99	0.53	0.86	-	-	-	0.56	0.77
EE	47	6.31	1.48	23%	0.95	0.68	0.78	-	-	-	0.42	-
NFE	47	54.25	4.81	9%	0.98	2.08	0.81	-	-	-	-	-
CF	49	14.81	2.13	14%	0.97	1.07	0.74	-	-	-	1.60	-
NDF	48	24.65	5.56	23%	0.96	1.87	0.89	-	-	-	3.20	-
ADF	47	19.89	5.70	29%	0.95	2.69	0.78	-	-	-	1.40	-
ADL	48	3.97	0.96	24%	0.89	0.66	0.53	-	-	-	1.10	-
AIA	45	1.04	0.23	23%	0.97	0.09	0.86	-	-	-	-	-
Apparent Digestibility (%):												
_DM	46	67.09	4.09	6%	0.64	2.62	0.60	0.92	1.74	0.76	1.90	1.75
_OM	46	68.19	4.19	6%	0.69	2.50	0.65	0.94	1.68	0.79	-	-
_GE	48	66.65	3.67	6%	0.66	2.31	0.61	0.94	1.49	0.83	1.90	1.99
_CP	46	68.99	2.90	4%	0.82	2.02	0.53	0.92	1.30	0.75	2.60	-
_CF	46	27.98	7.38	26%	0.91	4.61	0.61	0.97	3.06	0.82	-	-
_EE	49	88.48	3.22	4%	0.90	2.20	0.53	0.98	1.22	0.86	-	-
_NFE	43	76.81	3.67	5%	0.96	1.43	0.86	0.95	1.25	0.87	-	-
_ADF	46	25.29	4.19	17%	0.22	4.24	0.00	0.90	3.49	0.62	-	-
_NDF	44	28.58	3.80	13%	0.14	3.89	0.00	0.77	2.85	0.50	-	-

#Number of valid spectra after removing of outliers; X1c1: Xiccato *et al.*, 1999; X1c2: Xiccato *et al.*, 2003.

Table 2. Calibration and cross-validation of feed composition and apparent digestibility to the UV-Vis-NIR Spectra of the feces or the concatenated feed-and-faeces spectra.

Composition	#	Mean	SD	CV%	RSQ	SECV	R <sup>2</sup> cv	Joint		SEV
								RSQ	SECV	
DM Intake	58	0.1669	0.0248	15%	0.94	0.01	0.81	0.99	0.0091	0.87
Ingest	58	0.0733	0.0072	10%	0.97	0.00	0.80	0.95	0.0030	0.81
DM	60	42.11	10.69	25%	0.97	3.67	0.88	-	-	-
OM	64	88.81	2.28	3%	0.99	0.66	0.92	-	-	-
ASH	64	11.19	2.28	20%	0.99	0.66	0.92	-	-	-
GE	59	17.90	0.47	3%	0.85	0.22	0.77	-	-	-
PG	59	16.29	1.60	10%	0.96	0.51	0.90	-	-	-
EE	59	1.99	0.36	18%	0.95	0.23	0.58	-	-	-
NFE	56	38.42	2.38	6%	0.96	1.05	0.80	-	-	-
CF	57	31.85	1.73	5%	0.97	0.72	0.83	-	-	-
NDF	55	50.02	10.82	22%	1.00	1.89	0.97	-	-	-
ADF	58	47.77	10.15	21%	0.99	1.97	0.96	-	-	-
ADL	58	10.37	1.82	18%	0.93	0.98	0.71	-	-	-
AIA	57	3.09	0.46	15%	0.95	0.19	0.83	-	-	-
Apparent Digestibility (%):										
_DM	59	67.08	3.38	5%	0.93	1.57	0.79	0.92	1.74	0.76
_OM	59	68.34	3.49	5%	0.94	1.55	0.80	0.94	1.68	0.79
_GE	59	67.60	3.05	5%	0.83	1.58	0.74	0.94	1.49	0.83
_CP	64	68.66	2.81	4%	0.70	1.92	0.53	0.92	1.30	0.75
_CF	56	30.19	7.03	23%	0.96	2.99	0.82	0.97	3.06	0.82
_EE	59	89.11	3.16	4%	0.73	2.20	0.52	0.98	1.22	0.86
_NFE	58	76.15	3.51	5%	0.95	1.50	0.82	0.95	1.25	0.87
_ADF	56	26.17	3.75	14%	0.78	2.78	0.45	0.90	3.49	0.62
_NDF	58	27.90	3.45	12%	0.52	3.14	0.18	0.77	2.85	0.50

Xic2: Xiccato et al., 2003.



rabbit faeces examined via NIRS, could substantially improve the estimation of the apparent digestibility of nitrogen in feeds ob-

tained only through the NIRS of feeds, with mutual advantages for the feed industry and herdsman.

## REFERENCES

- Decruyenaere, V., Lecomte, Ph., Demarquilly, C., Aufrere, J., Dardenne, P., Stilmant, D., Buldgen, A., 2008. Evaluation of green forage intake and digestibility in ruminants using near infrared reflectance spectroscopy (NIRS): Developing a global calibration. *Anim. Feed Sci. Tech.* 148:138-156.
- Garnsworthy, P.C., Unal, Y., 2004. Estimation of dry-matter intake and digestibility in group-fed dairy cows using near infrared reflectance spectroscopy. *Anim. Sci.* 79:327-334.
- Mc Dryden, L.G., 2003. Near Infrared Reflectance Spectroscopy: Applications in Deer Nutrition. A report for the Rural Industries Research and Development Corporation. RIRDC Publ., Kingston, Australia.
- Meineri, G., Peiretti, P.G., 2007. Apparent digestibility of mixed feed with increasing levels of chia (*Salvia hispanica* L.) seeds in rabbit diets. *Ital. J. Anim. Sci.* 6 (Suppl. 1) :778-780.
- Peiretti, P.G., Meineri, G., 2008a. Effects of diets with increasing levels of *Spirulina platensis* on the performance and apparent digestibility in growing rabbits. *Livest. Sci.* 118:173-177.
- Peiretti, P.G., Meineri, G., 2008b. Effects of whole golden flax seed supplementation on performance and feed digestibility in rabbits. *J. Anim. Vet. Adv.* 7:56-60.
- Peiretti, P.G., Mussa, P.P., Meineri, G., Perona, G., 2007. Apparent digestibility of mixed feed with increasing levels of false flax (*Camelina sativa* L.) seeds in rabbit diets. *J. Food Agric. Environ.* 5:85-88.
- Vogtmann, H., Frirter, P., Prabuck, A.L., 1975. A new method of determining metabolizability of energy and digestibility of fatty acids in broiler diets. *Brit. Poultry Sci.* 16:531-534.
- Xiccato, G., Trocino, A., Carazzolo, A., Meurens, M., Maertens, L., Caratano, R., 1999. Nutritive evaluation and ingredient prediction of compound feeds for rabbits by near-infrared reflectance spectroscopy (NIRS). *Anim. Feed Sci. Tech.* 77:201-212.
- Xiccato, G., Trocino, A., De Boever, J.L., Maertens, L., Carabano, R., Pascual, J.J., Perez, J.M., Gidenne, T., Falcao-E-Cunha, L., 2003. Prediction of chemical composition, nutritive value and ingredient composition of European compound feeds for rabbits by near infrared reflectance spectroscopy (NIRS). *Anim. Feed Sci. Tech.* 104:153-168.