

# The implication of ROS production on triflumuron-induced oxidative stress and genotoxicity in human colon carcinoma (HCT-116) cells

Toxicology Research and Application



Volume 4: 1–10

© The Author(s) 2020

DOI: 10.1177/2397847320931792

journals.sagepub.com/home/tor



Rim Timoumi<sup>1,2</sup>, Ines Amara<sup>1,2</sup>, Intidhar Ben Salem<sup>1,3</sup> ,  
Matia Franca Buratti<sup>4</sup>, Emanuella Testai<sup>4</sup>,  
and Salwa Abid-Essefi<sup>1</sup> 

## Abstract

The aim of this study is to evaluate the cytotoxic and the genotoxic effects of triflumuron (TFM) on human colon carcinoma cells (HCT-116). Indeed, TFM is used to protect vegetables, fruits, and domestic animals against a large spectrum of parasites causing animal and human disorders. However, studies revealing its toxicity and its mode of action in mammalian systems remain very limited. We monitored our work with the cytotoxicity assay starting with the cell viability test, the ROS generation, the malondialdehyde (MDA) production, the DNA fragmentation, and the measurement of some antioxidant enzymes activities such as catalase, superoxide dismutase, and the glutathione S-transferase. Also, we measured the mitochondrial transmembrane potential. We showed that TFM induced a dose-dependent cell death. This decrease in cell viability was accompanied by a significant reduction in the mitochondrial membrane potential. We also have shown that TFM induced oxidative stress as revealed by the generation of reactive oxygen species, the increase of the MDA levels, and the activation of the antioxidant enzymes. Moreover, our results indicated that TFM induced DNA damage in HCT-116 cells as monitored by the comet assay. We demonstrate, for the first time, the cytotoxic and the genotoxic potentials of TFM on human cultured cells.

## Keywords

Triflumuron, cytotoxicity, genotoxicity, oxidative stress

Date received: 18 April 2020; accepted: 10 May 2020

## Introduction

Excessive use of pesticides in the agricultural sector can cause chemical contamination leading to the loss of plant and animal lives.<sup>1</sup> Indeed, pesticides can pose problems for human health since pesticide residues are frequently detected in different compartments of the environment (surface water, groundwater, soil, etc.) and in products intended for human and animal food.<sup>2,3</sup>

Numerous epidemiological studies suggest that pesticides may be involved in the induction of cancer, neurodegenerative diseases, fertility, and reproductive disorders.<sup>3–7</sup> That is why it is necessary to look for new products with

<sup>1</sup>Laboratory for Research on Biologically Compatible Compounds (LRSBC), Faculty of Dental Medicine, University of Monastir, Monastir, Tunisia

<sup>2</sup>Higher Institute of Biotechnology of Monastir, University of Monastir, Monastir, Tunisia

<sup>3</sup>Faculty of Medicine of Sousse, University of Sousse, Tunisia

<sup>4</sup>Istituto Superiore di Sanità, Environment and Health Department, Roma, Italy

### Corresponding author:

Rim Timoumi, Laboratory for Research on Biologically Compatible Compounds (LRSBC), Faculty of Dental Medicine, Rue Avicenne, 5019 Monastir, Tunisia.

Email: timoumirim@yahoo.fr



higher selectivity for the target organisms and lower toxicity for human and for the environment.

Insect growth regulators (IGR) are among these new compounds. They act on the growth and development of insects. In fact, IGR were introduced following the appearance of insect's resistance to pyrethroids and organophosphate insecticides.<sup>8</sup> Among the IGR, the class of benzoylureas (disrupting molt) interferes with the insect's exoskeleton causing then its death.<sup>9</sup>

Benzoylurea insecticides are widely used for crop protection, especially fruits and vegetables and could be classified such as insecticides of fourth generation having high selectivity, low acute toxicity for mammals, and high biological activity.<sup>10</sup> In this context, we focused our interest in this study on triflumuron (TFM), a benzoylurea insecticide that is used worldwide. TFM shows interesting properties such as high selectivity, high biological activity, and low toxicity for animals.<sup>11</sup> TFM is used to protect crops (apples, tomatoes, trees, cotton, and potato) and domestic animals against a broad spectrum of parasites.<sup>12</sup>

Regarding the molecular level, TFM acts by inhibiting the synthesis of chitin which is a linear polymer composed of *N*-acetylglucosamines.<sup>13</sup> Thus, by blocking the transport of *N*-acetylglucosamine through the epithelial membrane, TFM could act as a general stressor making the insect more susceptible to diseases. By consequence, this facilitates the entry of pathogenic fungi into the insect by the weakening of their cuticle which accelerates their death.<sup>14</sup>

Particularly, TFM was used to control *Tuta absoluta*, a tomato leaf miner. Tomato is the first horticultural crop in Tunisia with a production area of 25,000 ha and a total harvest of 1.1 million metric tons.<sup>15</sup> To manage *Tuta absoluta*, a mixture of insecticides with different modes of action is used, such as spinosad, indoxacarb, pyrethroids, and TFM because organophosphate (OPTs) and carbamates are no more effective due to insect resistance. Furthermore, TFM is used also against locusts in the Sahel zone.<sup>16</sup>

Like all other pesticides, TFM is known for its toxic effects to mammals and especially to aquatic organisms and invertebrates. For example, in rats, dogs, and rabbits the administration of these compounds causes spasms, skin, and respiratory irritation resulting in sneezing and in several eye damages.<sup>17,18</sup> Repeated administration of TFM is also known to cause hemolytic anemia<sup>19</sup> and reproductive toxicity.<sup>20</sup>

Also, benzoylurea insecticides are known by their ovidical activity. Indeed, the treatment of insects of the genus *Aedes albopictus* with TFM inhibits the occlusion of eggs. Also, an abnormal morphology of the egg shell has developed.<sup>20</sup> In addition, TFM acts on the offspring of insects, especially at the time of pupal formation. Like most IGRs, it causes malformations of pupae from treated females<sup>21</sup> and causes death of the embryo.<sup>22</sup> It causes high rates of abortion in infected insects.<sup>23</sup> Insects from the pupae of treated females die hatching or a few days later.<sup>21</sup>

Moreover, oral administration of TFM to male Wistar rats for 28 days resulted in elevated hemoglobin accompanied by an increase in the number of reticulocytes.<sup>19</sup>

In human, the ingestion of an unspecified BPU caused toxic methemoglobinemia. The hospitalized patient manifested cyanosis, low oxygen saturation, and chocolate-brown colored blood, the methemoglobin concentration was 59.3%. Following treatment with ascorbic acid, there was evidence of hemolysis.<sup>24</sup>

Moreover, those symptoms were confirmed in humans. Recently, Ning et al.<sup>25</sup> reported that in Hep G2 cells, TFM was able to promote the metastasis of human liver cancer cells and that hypoxiainducible factor-1 $\alpha$  is potentially responsible for these changes.

Thus, we aimed to evaluate the cytotoxicity and the genotoxicity induced by this insecticide on human colon carcinoma cells (HCT-116), being one of the target cells exposed to this pesticide following the ingestion of contaminated products.

## Materials and methods

### Chemicals

TFM was supplied by Sigma-Aldrich (St Louis, Missouri, USA). 3-[4,5-Dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT), cell culture medium (RPMI 1640), fetal calf serum, phosphate-buffered saline (PBS), trypsin-EDTA, penicillin and streptomycin mixture, and L-glutamine (200 mM) were from GIBCO-BCL (UK). 2,7-Dichlorofluorescein diacetate (DCFH-DA) was supplied by Molecular Probes (Cergy-Pontoise, France). Low melting point agarose and normal melting point agarose were purchased from Sigma. All other chemicals used were of analytical grade.

### Cell culture and treatment

HCT-116 were cultured in RPMI, supplemented with 10% fetal bovin serum (FBS), 1% L-glutamine (200 mM), 1% mixture of penicillin (100 IU/mL) and streptomycin (100 lg/mL), at 37°C with 5% CO<sub>2</sub>.

### Cell toxicity assay (MTT assay)

The MTT assay (a tetrazolium salt reduction assay) provides sensitive measurements of the normal metabolic status of cells, particularly that of the mitochondrion, where measurements reflect early cellular redox changes.<sup>26</sup> HCT-116 cells ( $2.5 \times 10^4$  cells/well in 96-well plates) were incubated at 37°C for 24 h with different concentrations of TFM (100, 200, 400, 600, 800 and 1000  $\mu$ M). A negative control containing untreated cells was also evaluated. After treatment, the plates were incubated with the MTT solution (final concentration of 0.5 mg/mL) for 3 h. The dark-blue formazan crystals formed in intact cells were dissolved with DMSO, and the absorbance at 570 nm was measured with a

spectrophotometer microplate reader (Bioteck, Elx 800, France). The results were expressed as the percentage of MTT reduction relative to the absorbance's measured from negative control cells. All assays were performed in triplicate.

### *FDA/BET marking*

The fluorescent probe fluorescein diacetate (FDA) and the ethidium bromide (BET) were used to assess cell mortality. FDA/BET labeling allowed the distinction between live and dead cells. Indeed FDA and BET are two chloroforms that enter into the cells except that for the FDA it will be converted and cleaved only by esterases of living cells. To do this, cells were seeded in six-well plates ( $7.5 \times 10^6$  cell/well) and treated with the different concentrations of TFM (100, 200, 400, 600, 800 and 1000  $\mu\text{M}$ ). After 24 h exposure, cells were trypsinized and centrifuged (5 min at 3000 r/min). The pellet was resuspended in 100  $\mu\text{L}$  of the medium. Thus, 10  $\mu\text{L}$  FDA (10  $\mu\text{g}/\text{mL}$ ) and 10  $\mu\text{L}$  BET (100  $\mu\text{g}/\text{mL}$ ) were added to the mixture. Then the contents are spread on a slide and cells were visualized using a fluorescence microscope, and on 100 cells, we count the number of living and dead cells.

### *ROS determination and oxidative stress status*

Reactive oxygen species (ROS) are markers of oxidative status. Indeed, in case of cellular stress, it has an increased production of these ROS that interact with DNA, lipids, and proteins. The intracellular amounts of ROS were measured by a fluorometric assay with DCFH-DA used extensively to monitor oxidation in biological systems as a well-established compound to detect and quantify intracellular compounds produced such as superoxide radical, hydroxyl radical, and hydrogen peroxide ( $\text{H}_2\text{O}_2$ ).<sup>27,28</sup> The conversion of the non-fluorescent (DCFH-DA) to the highly fluorescent 2,7-dichlorofluorescein (DCF) product ( $\lambda_{\text{max}} = 522 \text{ nm}$ ) happens in many steps. The fluorescent probe, after diffusing in the cell membrane, is hydrolyzed by intracellular esterases to nonfluorescent dichlorofluorescein (DCFH), which is trapped inside the cells then oxidized to fluorescent DCF through the action of peroxides in the presence of ROS.<sup>29</sup>

HCT-116 cells were seeded on 24-well culture plates (Polylabo, Strasbourg, France) at  $10^5$  cells/well for 24 h of incubation. After cells were incubated with different concentrations of TFM, for 24 h at  $37^\circ\text{C}$ .  $\text{H}_2\text{O}_2$  (20  $\mu\text{M}$ ) was used as a positive control. After incubation, cells were treated with 20- $\mu\text{M}$  DCFH-DA. Intracellular production of ROS was measured after 30-min incubation at  $37^\circ\text{C}$  by fluorometric detection of DCF oxidation on a fluorimeter (Biotek FL 800 $\times$ ) with an excitation wavelength of 485 nm and emission wavelength of 522 nm. The DCF fluorescence intensity is proportional to the amount of ROS formed intracellularly.

### *Lipid peroxidation measurement*

The measurement of lipid peroxidation was determined according to Ohkawa et al.<sup>30</sup> by the measurement of the

level of malondialdehyde (MDA) which is an ultimate fragment of the degradation of the polyunsaturated fatty acids of the lipidic membrane.

To do this, cells were plated in six-well plates at  $7.5 \times 10^5$  cells/well and treated with the different concentrations of TFM for 24 h of incubation. A positive control using  $\text{H}_2\text{O}_2$  (20  $\mu\text{M}$ ) has been realized. Then, cells were collected in coded test tubes. Samples were mixed with 0.1 mL of 1.15% potassium chloride, 0.2 mL of 8.1% sodium-dodecylSulfate (SDS), 1.5 mL of 20% acetic acid adjusted to pH 3.5, and 1.5 mL of 0.8% thiobarbituric acid. The obtained mixture was vortexed and heated at  $95^\circ\text{C}$  for 2 h. After cooling to room temperature, a volume of 5 mL of mixture of *n*-butanol and pyridine (15:1, v:v) was added to each sample, and the mixture was shaken vigorously. A centrifugation (4000 r/min for 10 min) allowed isolating the supernatant fraction. Finally, the absorbance was measured at 546 nm. The concentration of MDA was determined according to a standard curve.

### *Protein extraction*

To obtain cell lysates, HCT-116 cells were seeded in six-well plates ( $10^6$  cells/well) and then treated with TFM for 24 h. Thus, cells were recovered in PBS, centrifuged and collected in cold lysis buffer solution for 30 min. A centrifugation makes it possible to obtain the protein extracts. Protein concentrations were determined in cell lysates using Protein Bio-Rad (France) assay.<sup>31</sup>

### *Measurement of SOD activity*

Superoxide dismutases (SODs) are ubiquitous metalloenzymes that catalyze the disproportionation of superoxide ions into  $\text{H}_2\text{O}_2$  and molecular oxygen. The measurement of SOD activity was carried out according to the method described by Markland and Marklund,<sup>32</sup> by following the autoxidation and illumination of pyrogallol at 440 nm for 3 min. One unit of SOD activity was calculated as the amount of protein that caused 50% pyrogallol autoxidation inhibition. The SOD activity is expressed as U/mg protein.

### *Measurement of CAT*

Catalases (CATs) are enzymes that intervene in the defense of the cell against oxidative stress by eliminating oxygen species ( $\text{H}_2\text{O}_2$ ). To measure the activity of this enzyme, 780  $\mu\text{L}$  of phosphate-buffered solution, 20  $\mu\text{L}$  of each sample, and 200  $\mu\text{L}$  of  $\text{H}_2\text{O}_2$  (the substrate of the enzyme) were placed in a quartz cuvette. Then, the optical density was measured at 240 nm for an interval of time of 1 min.<sup>33</sup> The activity of CAT was calculated using the molar extinction coefficient ( $0.04 \text{ mM}^{-1} \text{ cm}^{-1}$ ). The results were expressed as mmol/min/mg protein.

### *Measurement of GST*

The activity of this enzyme was determined by the formation of a chromophore, the 1-glutathione-2,4-dinitrobenzene

from 1-Cl-2,4 dinitrobenzene (CDNB), which was followed spectrophotometrically at 340 nm. To do this, in a cuve, a mixture of 830  $\mu\text{L}$  of phosphate buffer, 50  $\mu\text{L}$  of CDNB, 100 mL of glutathione reduced (glutathion [GSH]), and 20  $\mu\text{L}$  of each sample were realized. The absorbance was determined at 340 nm for an interval of 1 min.<sup>34</sup>

### MMP assay

The uptake of the cationic fluorescent dye rhodamine 123 has been investigated to estimate the mitochondrial membrane potential (MMP).<sup>35</sup> In a typical experiment, cells seeded in 96-well culture plates were treated for 24 h with TFM (100, 200, 400, 600, 800 and 1000  $\mu\text{M}$ ) then rinsed with PBS and 100  $\mu\text{L}$  of rhodamine 123 (1  $\mu\text{M}$ ) in PBS was added on the plates. Cells were incubated (37°C, 5%  $\text{CO}_2$ ) for 15 min. Next, the PBS solution containing un-absorbed rhodamine 123 was removed and replaced by fresh PBS. The uptake rhodamine 123 was measured by fluorimetric detection with an excitation wavelength of 488 nm and an emitting wavelength of 525 nm. The results were expressed as the percentage of uptaken rhodamine fluorescence relative to the fluorescence measured from negative control cells.

### DNA damage assessed by the comet assay

To measure DNA damage, caused by a toxic agent, in individual mammalian cells, the comet test was considered. For this, six-well plates were used to seed HCT-116 cells at  $7.5 \times 10^5$  cells/well. After 24 h of incubation, cells were cultured in the presence of TFM (100, 200, 400, 600, 800 and 1000  $\mu\text{M}$ ) and  $\text{H}_2\text{O}_2$  (20  $\mu\text{M}$ ) which served as a positive control. Cells already recovered in PBS were mixed with low melting agarose, and the mixture was then spread on a microscope slides covered with normal agarose. Thus, the slides were put in an alkaline lysis buffer solution for 1 h at 4°C. Then, electrophoresis step (30 min, at 25 V, 300 mA) was done before neutralizing slides using a Tris buffer solution for 15 min. After drying the slides, a staining with ethidium bromide (20  $\mu\text{g}/\text{mL}$ ) allowed to quantify the DNA damage using a fluorescence microscope. The experiment was done in triplicate. The damage is represented by an increase of DNA fragments that have migrated out of the cell nucleus during electrophoresis and formed an image of a “comet” tail. A total of 100 comets on each slide were visually scored according to the intensity of fluorescence in the tail and classified by one of five classes as described by Collins et al.<sup>36</sup> The total score was evaluated according to the following equation:

$$\begin{aligned} \text{Total score} = & (\% \text{ of cells in class } 0 \times 0) \\ & + (\% \text{ of cells in class } 1 \times 1) \\ & + (\% \text{ of cells in class } 2 \times 2) \\ & + (\% \text{ of cells in class } 3 \times 3) \\ & + (\% \text{ of cells in class } 4 \times 4) \end{aligned}$$

### Statistical analysis

Each experiment was done three times separately and data were expressed as the mean  $\pm$  standard deviation of the means. One-way ANOVA was used to assess differences among the groups followed by Dunnett’s post hoc test. When two groups were compared, differences were assessed by Student’s *t* test. Differences were considered significant at  $p < 0.05$ .

## Results

### TFM induces cell death in HCT-116

After treating HCT-116 cells with the increasing concentrations of TFM (100, 200, 400, 600, 800 and 1000  $\mu\text{M}$ ) for 24 h, the MTT test which is a cell viability test has been carried out. We showed that TFM significantly decreased cell viability ( $p < 0.05$ ) with a value of IC 50 around 400  $\mu\text{M}$  (Figure 1(a)).

Using another cell viability test, FDA/BET labeling, our results were presented in Figures 1(b) and 2(c). Thus, after cells exposure to different TFM concentrations (200, 400 and 800  $\mu\text{M}$ ), cell morphology was changed, visualized with an optical microscope (Figure 1(b)). After mixing cells with the fluorochromes, living cells appeared in green, while dead ones appeared in orange. Finally, cells were visualized using a fluorescence microscope (Figure 1(c)). In both cases, cells were photographed using a digital camera (original magnification 200 $\times$ ).

### Measurement of ROS production


The concentration range already used was tested to verify the proportion of the oxidative stress induced by TFM in the HCT-116 cells. Thus, the generation of ROS was measured via the production of fluorescent DCF. Our results showed that TFM is able to increase the level of ROS in the cellular model used in a dose-dependent manner (Figure 2).

### Induction of lipid peroxidation


After exposure of HCT-116 cells to different concentrations of TFM for 24 h, the MDA assay was performed. Our results showed that TFM is able to induce lipid peroxidation. Indeed, the level of MDA increased from  $0.057 \pm 0.0069$   $\mu\text{mol}$  MDA/mg of protein in the untreated cells to  $0.198 \pm 0.0023$   $\mu\text{mol}$  MDA/mg of proteins in cells treated with the highest concentration of TFM (Figure 3).

### Effect of TFM on antioxidant enzymes activities


To evaluate the effect of TFM on the activities of antioxidant enzymes, we measured the activity of CAT, SOD, and glutathione S-transferase (GST). Our results showed that TFM is able to increase the activities of these enzymes in HCT-116 cells, in a dose-dependent manner. Indeed, the




**Figure 1.** Cytotoxic effect of triflumuron on HCT-116 cells after 24-h treatment. Cells were treated with different concentrations. Cell viability was determined using the MTT assay and expressed as percentages of viability (a). Data are expressed as the mean  $\pm$  SD of three independent experiments. Values are significantly different ( $p < 0.05$ ) from control. Cell morphology changes were visualized using optical (b) and fluorescent (c) microscope. Then cell were photographed using a digital camera (original magnification  $\times 200$ ). MTT: 3-[4,5-Dimethylthiazol-2-yl]-, 2,5- diphenyltetrazolium bromide; SD: standard deviation; HCT-116: human colon carcinoma.



**Figure 2.** Levels of relative fluorescent DCF production after exposure of HCT-116 cells to different triflumuron concentrations for 24 h. H<sub>2</sub>O<sub>2</sub> (20 µM) was used as a positive control. Data are expressed as the mean  $\pm$  SD of three independent experiments. Values are significantly different ( $p < 0.05$ ) from control. \* $p < 0.05$ , statistically significant compared to the control group. \*\* $p < 0.01$ , statistically significant compared to the control group. \*\*\* $p < 0.001$ , statistically significant compared to the control group. DCF: 2,7-dichlorofluorescein; SD: standard deviation; H<sub>2</sub>O<sub>2</sub>: hydrogen peroxide; HCT-116: human colon carcinoma.




**Figure 3.** Induction of lipid peroxidation in HCT-116 cells, after 24 h of incubation with triflumuron measured by the production of MDA. H<sub>2</sub>O<sub>2</sub> (20 µM) was used as a positive control. Data are expressed as the mean  $\pm$  SD of three independent experiments. Values are significantly different ( $p < 0.05$ ) from control. \* $p < 0.05$ , statistically significant compared to the control group. \*\* $p < 0.01$ , statistically significant compared to the control group. \*\*\* $p < 0.001$ , statistically significant compared to the control group. MDA: malondialdehyde; SD: standard deviation; H<sub>2</sub>O<sub>2</sub>: hydrogen peroxide; HCT-116: human colon carcinoma.



**Figure 4.** Effects of triflumuron on antioxidant enzyme activities such as catalase (a), superoxide dismutase (b) and glutathion S-transferase (c), after incubation of HCT-116 cells with the tested concentrations of TFM for 24 h. Data are expressed as the mean  $\pm$  SD of three independent experiments. Values are significantly different ( $p < 0.05$ ) from control. \* $p < 0.05$ , statistically significant compared to the control group. \*\* $p < 0.01$ , statistically significant compared to the control group. \*\*\* $p < 0.001$ , statistically significant compared to the control group. HCT-116: human colon carcinoma; SD: standard deviation; TFM: triflumuron.

activities of these enzymes passed from  $113.80 \pm 4.28$  (USOD/mg of proteins),  $64.26 \pm 3.58$  (mmol/min/mg of proteins), and  $0.271 \pm 0.0036$  ( $\mu\text{mol}/\text{min}/\text{mg}$  of proteins) in the untreated cells to  $303.10 \pm 6.03$  (USOD/mg of proteins),  $201.64 \pm 3.23$  (mmol/min/mg of proteins), and  $1.48 \pm 0.18$  ( $\mu\text{mol}/\text{min}/\text{mg}$  of proteins) in cells treated with TFM at 1000  $\mu\text{M}$ , respectively, for SOD, CAT, and GST (Figure 4(a), (b) and (c)).



**Figure 5.** Triflumuron induces a loss of mitochondrial transmembrane potential on HCT-116. Cells were treated with TFM at the indicated concentrations for 24 h. Data are expressed as the mean  $\pm$  SD of three independent experiments. Values are significantly different ( $p < 0.05$ ) from control. \* $p < 0.05$ , statistically significant compared to the control group. \*\* $p < 0.01$ , statistically significant compared to the control group. \*\*\* $p < 0.001$ , statistically significant compared to the control group. HCT-116: human colon carcinoma; TFM: triflumuron; SD: standard deviation.

### Measurement of mitochondrial transmembrane potential ( $\Delta\Psi_m$ )


The measurement of mitochondrial transmembrane potential ( $\Delta\Psi_m$ ) after cell exposure to TFM was carried out using rhodamine 123. Indeed,  $\Delta\Psi_m$  is proportional to uptaken rhodamine 123.<sup>37</sup> Results of this test are indicated in Figure 5, where it has been found that the tested concentrations have significantly decreased this potential, indicating then the depolarization of mitochondria.

### TFM induces DNA damage

DNA damages were analyzed using the alkaline comet assay. After treatment of HCT-116 cells with different concentrations of TFM for 24 h, we noticed a marked increase in the DNA damages. These damages are classified into five classes from 0 to 5 (Figure 6(a)). Indeed, as shown in Figure 6(b), the total score passed from  $53 \pm 8.48$  in the control cells to  $264 \pm 12.72$  when cells were treated with TFM at 1000  $\mu\text{M}$ .  $\text{H}_2\text{O}_2$  (20  $\mu\text{M}$ )-treated cells (positive control) induced  $298.4 \pm 14.24$  of the total score of DNA damage. Moreover, after 24 h of TFM cell exposure to 200, 400, 600 and 1000  $\mu\text{M}$ . These damages are classified into five classes from 0 to 4 (Figure 6(b)).

## Discussion

Pesticides are molecules used to protect humans and their environment against attacks caused by harmful vectors.<sup>10</sup> Due to their increased use in many areas such as agriculture, industry, and traditional medicine, pesticides cause adverse effects on human health and threaten ecosystems.<sup>38</sup> In this context, several studies have revealed the role of pesticides in the occurrence of numerous pathologies, namely cancer, neurodegenerative, and cardiac disorders.<sup>4,39,40</sup> In this article, we investigated the mechanisms



**Figure 6.** (a) Induction of DNA damages in HCT-116 cells, following treatment with triflururon at 100, 200, 400, 600, 800, and 1000  $\mu\text{M}$ .  $\text{H}_2\text{O}_2$  (20  $\mu\text{M}$ ) was used as a positive control. DNA stand breaks were detected by the standard comet assay. Data are expressed as the mean  $\pm$  SD of three independent experiments. Values are significantly different ( $p < 0.05$ ) from control. \* $p < 0.05$ , statistically significant compared to the control group. \*\* $p < 0.01$ , statistically significant compared to the control group. \*\*\* $p < 0.001$ , statistically significant compared to the control group. (b) Different classes of DNA damages quantified by the comets test and visualized using fluorescent microscope after a 24-h cell exposure to different TFM concentrations. Then cell were photographed using a digital camera (original magnification  $\times 200$ ). HCT-116: human colon carcinoma; TFM: triflururon; SD: standard deviation.

by which TFM, a benzoylurea insecticide belonging to the IGRs, was used.<sup>18</sup> TFM inhibits the synthesis of chitin, the major constituent of the insect cuticle causing consequently its death.<sup>35</sup> However, studies highlighting the toxicity mechanism of TFM remain very limited.

The aim of the present study is to evaluate the mechanism of action of TFM-induced cytotoxicity in cultured HCT-116 cells which are intestinal human cells that represent the first barrier to meet for any toxic substance.

Our results indicated that exposure of HCT-116 cells to increased concentrations of TFM induces a dose-dependent cell death as revealed by MTT test. Moreover, cells labeling with FDA/BET showed that by increasing TFM

concentrations, cell morphology was changed and cell mortality was increased, which is in agreement with the results founded using the MTT assay (data not shown).

In fact, the tested concentrations represent 0.035, 0.071, 0.14, 0.21, 0.28, and 0.35 mg/mL, respectively, for 100, 200, 400, 600, 800, and 1000  $\mu\text{M}$ , of the environmental doses detected in TFM residues. TFM was detected in crops with values of 0.04 mg/kg and 0.03 mg/kg respectively in pear and wool.<sup>41,42</sup> Moreover, TFM was detected in tomatoes with value ranged from 0.5 mg/kg to 2.1 mg/kg.<sup>43</sup>

These results support the toxicity of benzoylurea pesticides. Indeed, another study demonstrated that Lufenuron, belonging to this family, causes toxic effects on aquatic

species where hemorrhages were observed in eyes, fins, and lids of fish exposed to this pesticide. Moreover, the histological analyses revealed that the gills structure have changed.<sup>44</sup>

Besides, we found that TFM-induced cell death was associated with a significant decrease in the MMP after 24 h of treatment, which suggest the involvement of mitochondrial pathway in cell apoptosis caused by TFM since the mitochondrial membrane depolarization is able to cause the release of cytochrome *c* followed by caspases activation and therefore the activation of the intrinsic apoptotic.<sup>45,46</sup>

Moreover, oxidative stress has been proposed as a mechanism linking to pesticides exposure, increasing the risk for the development of diseases such as cancer and neurodegenerative diseases. In addition to the increase in the production of free radicals, exposure to pesticides can also affect antioxidant capacity and defense mechanisms, as well as increase the lipid peroxidation.<sup>47,48</sup> In agreement, we observed, in HCT-116 cells, that TFM induces the production of ROS and stimulates the activities of antioxidant enzymes such as CAT, SOD, and GST. Moreover, the level of MDA, one of the major oxidation products of peroxidized polyunsaturated fatty acids, was greatly increased following cell treatment with TFM. These results confirm the ability of benzoylphenyl urea pesticides to induce oxidative stress in mammalian systems. In this context, diflufenuron (DFB) was described as an inductor of lipid peroxidation in Caco-2 cells. The same study showed that DFB was able to increase CAT, SOD, and Gpx activities.<sup>49,50</sup> DFB was also found cytotoxic toward Balb/c 3T3 cells leading to cell transforming activities.<sup>49</sup> In addition, DFB was found to be cytotoxic by decreasing cell viability in CHOK1 cells<sup>49</sup> and Hep G2 cells<sup>51</sup> using the MTT assay and the neutral red test. Besides, Lufenuron was also able to increase the levels of MDA generation and to disrupt the antioxidant parameters in the liver of exposed mice.<sup>52</sup> Our findings are in agreement with other research studies which showed that chlorfluazuron, a benzoylureas insecticide, induced cytotoxicity revealed via the MTT test in renal (HEK 293) and hepatic (Hep G2) human cells.<sup>53</sup> In addition, our previous study showed that TFM induced toxicity on a murine model after a short exposure.<sup>54</sup> In this study, we demonstrated that TFM is able to increase the levels of both carbonyl proteins and MDA as well as it altered the enzyme activities of antioxidant systems.<sup>54</sup> Similarly, the genotoxic effect of TFM was demonstrated in the bone marrow cells of male Balb/C mice.<sup>55</sup> Also, cytotoxicity and genotoxicity of TFM on hepatic and renal human cells were well clarified.<sup>56</sup>

Pesticides can be at the origin of genotoxic alterations. If a xenobiotic reacts with nuclear DNA, it may cause mutagenic and carcinogenic effects to the exposed organisms. Interestingly, our study clearly indicates that TFM induces significant DNA damage in HCT-116 cells. Therefore, the TFM-induced cell death in intestinal cells in this study can

be due to the increase in oxidative stress accompanied by mitochondrial dysfunction and DNA fragmentation.

In conclusion, TFM is considered as a potential tool for the control of disease vectors in public health and agriculture. However, its uses should be controlled in order to avoid the health problems that may arise from unconscious human uses.

### Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.



### Ethical approval

This article does not contain any studies on human participants or animals performed by any of the authors.

### Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study was supported by “Le Ministère Tunisien de l’Enseignement Supérieur et de la Recherche Scientifique.”

### ORCID iDs

Intidhar Ben Salem  <https://orcid.org/0000-0002-2033-4476>  
Salwa Abid-Essefi  <https://orcid.org/0000-0002-4965-5855>

### References

- Ralph S, Petras M, Pandrangi R, et al. Alkaline single-cell gel (Comet) assay and genotoxicity monitoring using two species of tadpoles. *Environ Mol Mutagen* 1996; **28**: 112–120.
- Cerejeira MJ, Viana P, Batista S, et al. Pesticides in Portuguese surface and ground waters. *Water Res* 2003; **37**: 1055–1063.
- Yuan Y, Chen C, Zheng C, et al. Residue of consumers in Zhejiang Province. *China Food control* 2014; **36**: 63–68.
- Akoto O, Oppong-Otoo J and Osei-Fosu P. Carcinogenic and non-carcinogenic risk of organochlorine pesticide residues in processed cereal-based complementary foods for infants and young children in Ghana. *Chemosphere* 2015; **132**: 193–199.
- Silva AR, Cardoso DN, Cruz A, et al. Ecotoxicity and genotoxicity of a binary combination of triclosan and carbendazim to *Daphnia magna*. *Ecotoxicol Environ Saf* 2015; **115**: 279–290.
- Yoshida M, Inoue K and Takahashi M. Predictive modes of action of pesticides in uterine adenocarcinoma development in rats. *J Toxicol Pathol* 2015; **28**: 207–216.
- Zhang X, Wu M, Yao H, et al. Pesticide poisoning and neurobehavioral function among farm workers in Jiangsu, People’s Republic of China. *Cortex* 2016; **74**: 396–404.
- Belinato TA, Martins AJ, Lima JBP, et al. Effect of trifluthiuron, a chitin synthesis inhibitor, on *Aedes aegypti*, *Aedes albopictus* and *Culex quinquefasciatus* under laboratory conditions. *Parasit Vectors* 2013; **6**: 683.
- Tunaz H and Uygun N. Insect growth regulators for insect pest control. *Turk J Agric For* 2004; **28**: 377–387.



10. Tomsej T and Hajgllov J. Determination of benzoylurea insecticides in apples by high-performance liquid chromatography. *J Chromatogr A* 1995; **704**: 513–517.
11. British Crop Protection Council and the Royal Society of Chemistry, The Pesticide Manual, 10th ed., British Crop Protection Council and the Royal Society of Chemistry, London, UK, 1997.
12. Lowden S, Gray S and Dawson K. Treatment of natural infestations of the biting louse (*Werneckiellaequi*) on horses using triflumuron, a benzoylurea derivative insect growth regulator. *Vet Parasitol* 2007; **148**: 295–300.
13. Merzendorfer H and Zimoch L. Chitin metabolism in insects: structure, function and regulation of chitin synthases and chitinases. *J Exp Biol* 2011; **206**: 4393–4412.
14. Irigaray JFSC, Marco-Mancebon V and Perez-Moreno I. The entomopathogenic fungus *Beauveria bassiana* and its compatibility with triflumuron: effects on the twospotted spider mite *Tetranychusurticae*. *Biol Control* 2003; **26**: 168–173.
15. Direction Generale de la Production Agricole (DGPA). Tunisian Ministry of Agriculture. Annual Report, 2009.
16. Pan Africa. Briefing on locust no 3. 2012. <http://www.pan-afrique.org/fr/Rapports/Briefing/Briefing%20locusts.pdf>.
17. Waller PJ and Lacey E. The effect of triflumuron (SIR8514) on the free-living stages of sheep nematodes. *Vet Parasitol* 1986; **2**: 119–126.
18. European Food Safety Authority, *EFSA J* 2011; **9**: 1941.
19. Tasheva M and Hristeva V. Comparative study on the effects of five benzoylphenylurea insecticides on haematological parameters in rats. *J Appl Toxicol* 1993; **13**: 67–68.
20. Suman DS, Yi W, Anwar L, et al. Ovicidal activity of three insect growth regulators against *Aedes* and *Culex* mosquitoes. *Acta Tropica* 2013; **128**: 103–109.
21. Ouédraogo A. Etude de l'efficacité et de la rémanence de la deltaméthrine et du triflumuron imprégné sur tissu (pour la lutte contre les glossines). Thèse de Doctorat en pharmacie (Diplôme d'état). Faculté de Médecine, de Pharmacie et d'Odonto stomatologie. Université du Mali, Bamako, 1998, p. 105.
22. Itard J. Les glossines ou mouches tsé-tsé. Etude et synthèse de l'IEMVT, 15. Département du Centre de Coopération Internationale en Recherche Agronomique pour le Développement, Paris, France, 1986, p. 155 .
23. Langley PA. Evaluation of the chitin synthesis inhibitor triflumuron for controlling the tsetse *Glossina morsitans morsitans* (Diptera: Glossinidae). *Bul Entomolog Res* 1995; **85**: 495–500.
24. D'sa SR, Victor P, Jagannati M, et al. Severe methemoglobinemia due to ingestion of toxicants. *Clin Toxicol* 2014; **52** (8): 897–900.
25. Ning X, Wang Y, Yan W, et al. Chitin synthesis inhibitors promote liver cancer cell metastasis via interfering with hypoxia-inducible factor 1 $\alpha$ . *Chemosphere* 2018; **206**: 231–237. DOI: 10.1016/j.chemosphere.
26. Mosman T. Rapid colorimetric assay for cellular growth and survival. Application to proliferation and cytotoxicity assays. *J Immunol Methods* 1983; **65**: 55–63.
27. Cathcart R, Schwiers E and Ames BN. Detection of picomole levels hydroperoxides using a fluorescent dichlorofluorescein assay. *Anal Biochem* 1983; **134**: 111–116.
28. Chen T and Wong YS. Selenocystine induces caspase-independent apoptosis in MCF-7 human breast carcinoma cells with involvement of p53 phosphorylation and reactive oxygen species generation. *Int J Biochem Cell Biol* 2009; **41**: 666–676.
29. Le Bel CP, Ischiropoulos H and Bondy SC. Evaluation of the probe 2 dichlorofluorescein as an indicator of reactive oxygen species formation and oxidative stress. *Chem Res Toxicol* 1992; **5**: 227–231.
30. Ohkawa H, Ohishi N and Yagi K. Assay of lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal Biochem* 1979; **95**: 351–358.
31. Wang W, Xiong W, Wan JL, et al. The decrease of PAMAM dendrimer-induced cytotoxicity by PEGylation via attenuation of oxidative stress. *Nanotechnology* 2009; **20**: 103–105.
32. Markland S and Marklund G. Involvement of the superoxide anion radical in the autoxidation of pyrogallol and a convenient assay for superoxide dismutase. *Eur J Biochem* 1974; **47**: 469–474.
33. Aebi H. Catalase in vitro. *Method Enzymol* 1984; **105**: 121–126.
34. Elia A, Dörr A, Prearo M, et al. Detoxification enzymes offreshwater crayfish *Procambarusclarkii* fed a diet enriched in selenium: Preliminary results. *Mar Fresh Behav Physiol* 2007; **40**: 195–199.
35. Cao J, Liu Y, Jia L, et al. Curcumin induces apoptosis through mitochondrial hyperpolarization and mtDNA damage in human hepatoma G2 cells. *Free Radic Biol Med* 2007; **43**: 968–975.
36. Collins AR, Dusinska M, Gedik CM., et al. Oxidative damage to DNA: Do we have a reliable biomarker? *Environ Heal Persp* 1996; **104**: 465–469.
37. Farnesi LC, Brito JM, Linss JG, et al. Physiological and morphological aspects of *Aedesegypti* developing larvae: effects of the chitin synthesis inhibitor novaluron. *PLoS One* 2012; **7**(1): e30363.
38. Juricek L and Coumoul X. Diet, pesticides and neurological diseases. *Noteb Nutrit Dietet* 2015; **9**: 74–80.
39. Kland MJ. Teratogenicity of pesticides and other environmental pollutants. *Stud Env Sci* 1988; **3**: 315–463.
40. Wilson TG and Cryan JR. Lufenuron, a chitin-synthesis inhibitor, interrupts development of *Drosophila melanogaster*. *J. Exp Zool* 1997; **278**: 37–44.
41. Aplada-Sarlis GE, Miliadis NG and Tsiropoulos. Dissipation of teflubenzuron and triflumuron residues in field-sprayed and cold-stored pears. *J Agric Food Chem* 1999; **47**(7): 2926–2929.
42. Fan Y, Huang S, Yu X, et al. Determination of diflubenzuron and triflumuron residues in greasy wool by accelerated solvent extraction technique and high performance liquid chromatography. *Se Pu* 2009; **27**(4): 463–466.
43. Martínez-Galeraa M, López-Lópezb T, Gil-Garcíaa MD, et al. Determination of benzoylureas in tomato by high-

- performance liquid chromatography using continuous on-line post-elution photoirradiation with fluorescence detection. *J Chromatogr A* 2001; **918**: 79–85.
44. Rafaela Leão Soares P, Lucas Corrêa de Andrade A, Pinheiro Santos T, et al. Acute and chronic toxicity of the benzoylurea pesticide, lufenuron, in fish, *Colossoma macropomum*. *Chemosphere* 2016; **161**: 412–421.
45. Raza H, John A and Benedict S. Acetylsalicylic acid-induced oxidative stress, cell cycle arrest, apoptosis and mitochondrial dysfunction in human hepatom HepG2 cells. *Eur J Pharmacol* 2011; **668**: 15–24.
46. Susin SA, Zamzami N and Kroemer G. Mitochondria as regulators of apoptosis: doubt no more. *Biochim Biophys Acta* 1998; **1366**: 151–165.
47. Abdollahi M and Karami-Mohajeri S. A comprehensive review on experimental and clinical findings in intermediate syndrome caused by organophosphate poisoning. *Toxicol Appl Pharmacol* 2012; **258**: 309–314.
48. Astiz M, Arnal N, de Alaniz MJT, et al. Occupational exposure characterization in professional sprayers: clinical utility of oxidative stress biomarkers. *Environ Toxicol Pharmacol* 2011; **32**: 249–258.
49. Bayoumi AE, Pérez-Pertejo Y, Zidan HZ, et al. Cytotoxic effects of two antimolting insecticides in mammalian CHO-K1 cells. *Ecotoxicol Environ Saf* 2003; **55**: 19–23.
50. Ilboudo S, Edwin F and Virginie R. In vitro impact of five pesticides alone or in combination on human intestinal cell line Caco-2. *Toxicol Rep* 2014; **1**: 474–489.
51. Delescluse C, Ledirac N, de Sousa G, et al. Cytotoxic effects and induction of cytochromes P450 1A1/2 by insecticides in hepatic or epidermal cells: binding capability to the Ah receptor. *Toxicol Lett* 1998; 96–97: **33–39**.
52. Perocco P, Colacci A and Grilli S. In vitro cytotoxic and cell transforming activities exerted by the pesticides cyanazine, dithianon, diflubenzuron, procymidone, and vinclozolin on BALB/c 3T3 cells. *Environ Mol Mutagen* 1993; **21**: 81–86.
53. Deivanayagam C, Asokan S and Rajasekar S. The study of lipid peroxidation, liver enzymes and antioxidant status in lufenuron treated mice, *mus musculus* species. *Int J Chem Tech Res* 2014; **6**: 5321–5328.
54. Timoumi R, Amara I, Neffati F, et al. Acute triflumuron exposure induces oxidative stress responses in liver and kidney of Balb/C mice. *Environ Sci Poll Res* 2019; **26**: 3723–3730.
55. Timoumi R, Amara I, Ayed Y, et al. Triflumuron induces genotoxicity in both mice bone marrow cells and human colon cancer cell line. *Toxicol Mech Methods* 2020; **21**: 1–33.
56. Timoumi R, Amara I, Ben Salem I, et al. Triflumuron induces cytotoxic effects on hepatic and renal human cell lines. *J Biochem Mol Toxicol* 2020; **30**: e22504.