

## **Histamine food poisonings: a systematic review and meta-analysis**

Fabio M. Colombo, Patrizia Cattaneo, Enrica Confalonieri, Cristian Bernardi

Dipartimento di Scienze Veterinarie per la Salute, la Produzione Animale e la Sicurezza Alimentare, Università degli Studi di Milano, Milano, Italy.

Corresponding author: Cristian Bernardi, Università Degli Studi di Milano, Dipartimento di Scienze Veterinarie per la Salute, la Produzione Animale e la Sicurezza Alimentare, Milano, Italy. via Grasselli, 7 20137 Milano, cristian.bernardi@unimi.it, Phone:00390250318506, Fax: 00390250318501

### **Summary**

**Objective:** Aim of this study was to assess the mean of histamine concentration in food poisoning.

**Design:** Systematic review and meta-analysis of reports published between 1959-2013.

**Study selection:** main criteria for inclusion of studies were: all report types that present outbreaks of “histamine poisoning” or “scombroid syndrome” from food, including histamine content and type of food. Health status of people involved must be non pathological.

**Results:** Fifty-five (55) reports were included, these studies reported 103 incidents. All pooled analyses were based on random effect model; histamine mean concentration in poisoning samples was 1107.21 mg/kg with a prediction 95% interval of 422.62 - 2900.78 mg/kg; heterogeneity index (I<sup>2</sup>) was 100 % (P<0.0001); prediction interval was 24.12 – 50822.78 mg/kg. Fish involved in histamine poisoning was mainly tuna or *Istiophoridae* species. No clues of association between concomitant conditions (female sex, alcohol consumption, previous

medication and consumption of histamine releasing food) and histamine poisoning, were highlighted.

## **Conclusions:**

This is the first systematic review and meta-analysis that analyzes all the available data on histamine poisoning outbreaks evaluating the histamine concentration in food involved. Histamine mean concentration in poisoning samples was fairly high. Our study suffers from some limitations, which are intrinsic of the studies included, for instance the lack of a complete anamnesis of each poisoning episode.

**Protocol registration:** methods were specified in advance and have been published as a protocol in PROSPERO database (18/07/2012 -CRD42012002566).

## **Keywords**

Scombroid syndrome, Scombrotxin fish poisoning, Foodborne illness, Outbreak, Histamine intoxication.

## INTRODUCTION

Scombroid syndrome/histamine poisoning occurs worldwide and it is considered one of - if not the most common form of toxicity caused by fish consumption (Dalgaard, Emborg, et al., 2008). The number of cases is increasing, in spite of the improved knowledge on seafood safety; this is due to a change in the way in which seafood, and mainly tuna, is eaten, that is as steaks or hamburger (Becker, Southwick, et al., 2001), or as canned tuna recipes (sandwiches, salads, pizza) (Cattaneo, Stella, 2001; Mclauchlin, Little, et al., 2006). Less is known about foods other than seafood and it is of utmost importance to assess the impact of all food types on this syndrome to implement specific prevention measures.

Periodically reviews on this item have been published (Lehane, Olley, 2000; Hungerford, 2010), although containing a lot of data they are not systematic reviews. Systematic review has not yet been performed on histamine poisoning. To assess histamine level of food associated to histamine poisonings, in the light of objective criteria, could lead to reliable information useful to control this hazard.

The general aim of this review is to perform the first systematic review about histamine food poisoning and meta-analysis of histamine content in food involved in these outbreaks.

## METHODS

According to the Cochrane Collaboration ([www.cochrane.org](http://www.cochrane.org)) guidelines, the methods of the analysis and inclusion criteria were specified in advance and documented in a protocol that has been published in the International prospective register of systematic reviews (PROSPERO

WEB site: <http://www.crd.york.ac.uk/PROSPERO>), on 18/07/2012 with registration number CRD42012002566.

## **Criteria for considering studies for this review**

### Types of studies

All report types of histamine food poisoning from food were considered for inclusion in the review. Reports of histamine poisonings from non-food sources (such as experimental studies with histamine administration) were not considered for inclusion. Only reports with histamine concentrations determined by chemical and ELISA methods were included. If the report was an experimental comparative one (e.g. experimental group versus control group) only data of group where occurred foodborne histamine intoxication were considered.

Eligible studies included any histamine poisoning outbreaks or single episodes that reported a measure of the histamine content and the type of the food involved in histamine poisoning.

The spatial interval for considering studies was set as worldwide. The time interval was set from 1959 through 2013, because in 1959 there was the first application of a specific and accurate quantitative method, the fluorimetric assay of histamine in tissues (Shore, Burkhalter, et al., 1959). Reports (abstract and full text) written in English, Italian, French, German, Portuguese and Spanish were considered; considering a full text in other languages was decided case-by case by the potential relevance for this review of its English abstract.

### **Population**

Only clinically healthy subjects were included; food allergic patients and other very sensitive people (due to serious illness or anomalous physical or psychic conditions), preschooler (< 6

years old) and very old (> 80 years old) people were excluded. If in a study nothing was reported about health status of people involved in histamine poisoning the health status was recorded as “unknown”.

## **Types of outcome measures**

### **Primary outcomes**

Number of histamine poisoning samples and histamine concentration in poisoning sample.

### **Secondary outcomes**

Concomitant conditions relevant to histamine poisoning were considered as listed in Maintz and Novak (2007): female sex, previous medication, food description (fish species, food recipe), consumption of alcohol during the meal; consumption of food recipe with suggested histamine-releasing capacities.

### **Search methods for identification of studies**

Search strategies were optimized to detect all reports of histamine poisonings from foods that met inclusion criteria. A main form of search strategy was designed and modified to meet settings of databases consulted. We systematically identified all potentially relevant reports through the main electronic databases (table A); additional search was conducted by analyzing references of the selected articles.

Characteristics of consulted databases, specific search strategy and number of reports obtained, searched database are shown in table A-1. Unpublished and ongoing studies were also considered and detected if existing. The main search strategy is presented in figure n.1, search terms included the following key word: "histamine", "scombroid syndrome", "histamine

poisoning", "food", "seafood", "meat products", "fish", "cheese", "beer", "wine", "biogenic amines". To improve the effectiveness of keywords in the search strategy a preliminary thesaurus study was performed. When multiple reports for a single study were present, it was used the most complete and updated version.

The literature search was conducted by two investigators (EC, FC) by aid of an information expert and by consulting with CB and PC. Two authors (CB and FC) independently selected potentially eligible studies for inclusion. The decision to include articles was made on the basis of the study title, then of the study abstract and finally of the full text; disagreements between reviewers were resolved by consensus; if no agreement was reached, a third author (PC) decided.

A data extraction sheet was developed and pilot-tested on a randomly-selected subgroup of included studies, data sheet was refined accordingly. One author (CB) extracted data from extraction sheet; data extracted were checked by a second author (FC). Disagreements were resolved by discussion between the two review authors; if no agreement was reached, a third author decided (PC).

A unique identifier of report was included in the characteristics recorded.

All quantitative measures of histamine content and measures of their variability; method of analysis used to determine food histamine content (if no method was mentioned the value was set to "unknown"); foods involved in histamine poisoning; primary and secondary outcome values; country or other identifier of geographic locations; people health category, i.e. if participants belonged to an excluded category and which was this category (if participants did not belong to above categories the status of "normal" was recorded); presence of "heterogeneous

food" (referring to more food types being associated to a single histamine mean value); other report characteristics useful to improve quality of information.

## **Assessment of risk of bias in individual studies**

Two reviewers (FC and CB) assessed the quality independently and any disagreements were resolved by discussion between the two review authors; if no agreement was reached, a third author decided (PC). Quality of included studies was considered a surrogate of risk of bias, so a quality score, of reports included in review based on additional relevant details other than inclusion criteria, was calculated. For each of the following 7 items a score of 1 was given if a value was present, 0 for absent value. The scores were then summed to give the final quality score (Murphy, Pfeiffer, et al., 2009). Variability estimate of histamine concentration, source of medical diagnosis (e.g. hospital m.d., family m.d.) or reasons given to present data as "histamine poisoning /scombroid syndrome", age, sex, health status, source of food involved in poisoning (restaurant, supermarket...), declaration of histamine content measurement method, number of patients involved in histamine poisoning; otherwise any element that could arise suspect of bias was recorded.

## **Summary measures**

Concomitant conditions ("risk factors") relevant to histamine poisoning" outcome were summarized as a contingency table of the declared risk factors versus the number of their occurrences. The "number of histamine poisonings" outcome was summarized as the overall sum of histamine poisoning samples.

The summary measure of histamine concentration in sample was set to “log- mean”; this term is defined as the value of the estimate of the mean of the logarithms of the raw data. If this log-mean value was not be given in reports it was calculated with documented methods to yield a log- mean and its standard error (Quan, Zhang, 2003; Higgins, White, et al., 2008).

## Unit of the analysis

The unit of the analysis was the "histamine poisoning sample". This unit is defined as one “histamine poisoning” that occurred to one group of people (for ‘group’ is meant one or more people) that ate one sample of food (for ‘sample of food’ is meant one or more foods that were involved in one poisoning).

## Histamine poisoning sample concept

One ”histamine poisoning sample” (as defined above) led to one observation for each of the three outcomes considered; the observation formats were: a count of one (1) case in “assessment of valid histamine poisoning cases outcome”, one histamine concentration log- mean in “histamine content” outcome and one list of values (i.e. the names of relevant concomitant factors) in “relevant concomitant factors” outcome. The number of patients involved in histamine poisoning sample was recorded. It was decided that all unexpected situations related to unit of analysis were assessed and managed and the management method recorded.

## **Methods to deal with missing data**

Missing variability data in poisoning samples (when a mean is given for more than one food specimens being involved in a single poisoning sample) was derived with documented statistical method that were recorded.



If a single poisoning sample (unit of analysis) was associated to more than one food type ("heterogeneous food category") and histamine values of single foods were given but not the mean, it was planned that histamine content value had to be recorded as the log- mean of the values and variability estimate had to be calculated, the single values being recorded. If any of single values were missing, it was planned that the mean and variability estimated had to be calculated and the presence of missing values recorded. Again, it was planned that if all, but one, values were missing histamine content had to be considered as a single value, this situation being recorded; moreover, all unexpected situations related to missing data had to be assessed and managed, possibly with documented methods that had to be recorded.

## **Synthesis of results\_methods**

Punctual estimates and their 95% confidence intervals were calculated across all selected studies on

statistical units according to the methods described above. Calculations were performed using the "metagen" procedure of "meta" package of R software (Schwarzer, 2010). As this meta-analysis was expected to yield a high degree of variability, the random effect model, described by DerSimonian and Laird (1986), was selected over the fixed effect model, because it incorporates within and between study variability. The chosen level of significance for statistical tests was  $P < 0.05$ . Heterogeneity, i.e. variability among records, was assessed by the I-squared (I<sup>2</sup>) statistic (Higgins et al., 2003). Ninety-five per cent (95%) prediction intervals were calculated by means of "metafor" R package (Viechtbauer, 2010).

## **Assessment of risk of bias across the studies**

In general, due to the nature of this systematic review, no selective reporting bias was assessed; it was planned that, if there were clues of selective reporting, authors of reports had to be contacted asking them about other results or outcomes not reported and that, if this issue was not resolved, to decide, with reasons, to exclude such reports. Decision had to be kept independently by CB and FC; if disagreement occurred PC had to keep final decision. Whatever the decision, the bias clues detected had to be recorded.

About management of reporting biases, being this concept difficult to apply due to the nature of this review, it was decided to discuss the publication bias issue according to data scenarios encountered during the review development.

## **Additional analyses**

Subgroup analysis about country or other identifier of geographic locations of histamine poisoning samples.

Subgroup analysis about groups: 1) fresh seafood, 2) frozen seafood, 3) canned seafood, 4) fermented seafood, 5) seafood other than 1,2,3,4; 6) cheese and dairy; 7) other foods.

Sensitivity analysis conducted by quality score or quality categories of the reports.

Sensitivity analysis on histamine concentration outcome conducting meta-analysis separately on two groups: one containing reports where variability was not derived (variability data value given in report) and one where variability was derived (variability data value not given in report, variability data inferred from other data).

## RESULTS

### **Study selection**

Searches yielded a total of 9390 references, after review and excluding duplicate reports 708 references were identified as potentially relevant. Of these, 556 records were included on the basis of title and abstract. We excluded 256 reports because they did not meet the adopted criteria and the full text of 300 reports was evaluated for report eligibility.

After excluding 248 full-text reports (corresponding to 285 poisoning samples), 52 reports (corresponding to 103 poisoning samples), listed in table 1, were included in overall quantitative synthesis for outcomes “number of poisoning samples” and “concomitant conditions”. Fourteen reports among them, corresponding to 15 poisoning samples, were selected for quantitative synthesis of outcome “histamine concentration in poisoning samples”. Selection process is summarized in figure 2.

### **Characteristics of included studies**

Below are summarized the characteristics of the 52 articles included; details are shown in table 1. The overall analysis comprised a total number of 1171 people involved in 103 episodes of histamine intoxication, ranged from 1 to 347 (person/poisoning sample). In these outbreaks the sources of food were reported in 50 episodes (missing = 53). On the known 50 sources, 17 were related to institutional or company food services, 20 to restaurants and only 9 (plus 4 unsure) linked to private home. Among the 103 poisoning samples, 101 were fish and seafood and only two were cheese. The raw data for each outbreaks of histamine intoxication are presented in the table 1. The meta-analysis of data from the 52 selected articles is summarized in Forest plot

(figure 3); the mean histamine concentration in studied episodes is 1107.21 mg/kg with a confidence interval of 422.62 – 2900.78 mg/kg. Heterogeneity index ( $I^2$ ) was 100 % ( $P < 0.0001$ ), log-prediction interval was 3.18 – 10.84, equivalent to 24.12- 50822.78 mg/kg. Secondary outcomes that are the concomitant conditions relevant to histamine poisoning were not evaluated, because in the most of included articles they are missing.

## **Risk of bias as quality score of individual reports**

Quality items values and the overall quality score are presented in table 2 for each included report.

## **Risk of bias across reports**

No elements pointing to selective reporting bias were detected. Publication bias was not assessed.

## **Additional Analyses**

Due to the nature of results about “concomitant conditions” outcome, this was not considered for additional analyses.

## **Subgroup analyses**

Number of poisoning samples and histamine concentration outcomes by geographic locations were not analyzed because of too many different locations.

Number of poisoning samples and histamine concentration outcomes by food categories was analyzed (respectively table 3 and figure 4).

## Sensitivity analyses

The following analyses were made: number of poisoning samples and histamine concentration outcomes by quality score categories; histamine concentration outcome by groups where variability was derived and where was not. Their results are respectively shown in table 4 and figures 5, 6.

## DISCUSSION

About the fish species associated with poisoning samples, it is worth noting that, n. 59 (out of 101) belonged to species associated with a high amount of histidine, according to EU legislation, that establishes a legal limit of histamine for "Particularly fish species of the families: *Scombridae*, *Clupeidae*, *Engraulidae*, *Coriphaenidae*, *Pomatomidae*, *Scomberosocidae*."(Communities, 2007), because these species are more likely to contain high histamine levels, as during spoilage some bacteria produce decarboxylase enzymes and convert histidine to histamine. Other 21 belonged to fish species without a legal limit in EU, and for a good 21 poisoning samples the species was unknown.

The data obtained by our review about canned tuna refute certain views that see this product as a main cause of poisoning. Among the 101 poisoning fish samples, only 22 consisted in canned products, mainly canned tuna (table 3) and all 22 poisoning samples were related to events happened before 1985, but two (Tsai, Kung, et al., 2005; Valentini, Levre, et al., 1991).

At present, canned tuna, and other canned fish belonging to species associated with the risk of histamine, have very low levels of histamine; this fact is likely due to the quality of canning process that is improving over the years due to widespread application of HACCP principles,

from the caught fish on the vessel to the processed product (Cattaneo, 2011; Guillier, Thebault, et al., 2011).

Other three episodes regarded canned tuna as ingredient (tuna salad and tuna sandwiches) (Stell, 1997; Predy, Honish, et al., 2003; Jantschitsch, Kinaciyani, et al., 2011). In all three, tuna cans had been opened hours or even a week before the preparation or the consumption, with likely post-processing contamination and consequent histamine production.

Fresh or frozen fish, diversely prepared and cooked, and fish products differently processed (not canned) were cause of poisoning in 79 episodes. The species or the family mainly reported were (number, % of 79): tuna (26, 32.9%); scombridae other than tuna (7, 8.8%); mahi mahi (3, 3.8%); species of the family *Istiophoridae* (total 8, 10.1%) such as *Makaira* spp (5), *Tetrapturus* spp (2), sailfish (1); swordfish (2); others species (12, 15.2%).

Among the "others", *Seriola lalandi* (n.3), *Chanos chanos* (n.1), *Arripis trutta* (n.4) were reported, fish species not considered in EU legislation, while having very high concentrations of histidine. Three other outbreaks (Eckstein, Serna, et al., 1999; Feldman, Werner, et al., 2005; Sinn, 2006) were attributed to *Lepidocybium flavobrunneum*, species whose meat has a very high content of wax ester that could cause gastrointestinal effects, but also has histidine levels as high as many Scombridae.

As to *Istiophoridae* and *Xiphidae* families, suborder Xiphiodei, in other countries they are associated with the risk of histamine because known to have very high free histidine levels or to be associated with SFP (Scombrotxin Fish Poisoning) (F.A.O., 2014). Interestingly the family *Istiophoridae* (Billfish) is placed in the Scombroidei suborder by Nelson (2006). Both Billfish and scombrids have common characteristics that could explain the frequency of episodes of

histamine intoxication caused by billfish. The complete list of fish species produced by our review can help to control imports and medical history of cases of suspected poisoning, as well as to cope with the problems arising from changes in international market trends of fishery products.

The source of poisoning (places where the poisoning samples were eaten) was not reported in 53 episodes (out of 103). The main reported sources were restaurants (20 cases, plus 3 unsure) and institutional foodservice, company or community canteens and cafeterias (17 cases), where the number of people involved is in terms of dozens or hundreds. The outbreaks occurred at home were 9 (plus 4 unsure); probably this kind of poisoning, involving a small number of persons for single episode, is little reported in literature and could indicate a reporting bias (under-reporting). Regarding the result of meta-analysis, the meta-mean of histamine concentration that summarizes the 14 reports (Foo, 1977; Molinari, Montagnoli, et al., 1989; Wu, Yang, et al., 1997; Su, Chou, et al., 2000; Becker, Southwick, et al., 2001; Hall, 2003; Emborg, Laursen, et al., 2005; Feldman, Werner, et al., 2005; Mclauchlin, Little, et al., 2006; Tsai, Hsieh, et al., 2007; Chen, Huang, et al., 2010; Chen, Lee, et al., 2011; Lee, Huang, et al., 2012) used for the statistical calculation (figure 3) is about 1000 ppm, a very high value if compared with what assumed by FDA (F.D.A., 2014) indicating, in most cases, histamine levels in illness-causing fish of about 200 or 500 ppm. On the other side, our result is in agreement with McLaughlin et al. (Mclauchlin, Little, et al., 2006) who wrote that ingestion of fish containing histamine at levels around 1000 ppm can result in illness. Shalaby (Shalaby, 1996) emphasized that poisoning does occur at histamine concentrations lower than 100 mg/100g and levels of histamine in fish of 5-20 mg/100g (50-200 ppm) are possibly toxic. This could be congruent with the lower limit of

overall predictive interval of histamine concentration from meta-analysis (24.12 ppm) although, due to the highest heterogeneity amount estimated, this value is questionable. Either way, EU maximum limit (Communities, 2007) seems to be proper to protect the consumer, also respect to the meeting report of FAO/WHO (F.A.O., 2014), where an oral NOAEL (No Observed Adverse Effect Level) of 50 mg was identified, from which was derived a histamine limit of 200 mg/kg, considering a service size of 250 g.

Due to the highest (100%) level of heterogeneity estimated for the overall meta-analysis the limits both for the meta-mean confidence interval and the predictive interval are questionable. More reliable are the values for subgroups where moderate amount of heterogeneity was estimated.

Subgroup analysis of histamine concentration outcome by food categories did not show significant difference between subgroups due to the overlapping of confidence interval. Moreover, the food category “fermented” (4) is missing, while categories “fresh” (1) and “canned” (3) consist of only one record and food category “other seafood” (5) is highly heterogeneous.

Sensitivity analysis of histamine concentration outcome by quality categories did not show separation of the values of quality categories (overlapping of confidence interval) but this cannot lead to declare absent the quality category effect, due to remarkable difference between the means of the categories, high degree of heterogeneity of each category and finally presence of single-record categories.

About sensitivity analysis of histamine concentration by derived or not variability, also this is poorly interpretable, due to high heterogeneity amount in each group and very unbalanced



sample size of the two group (2 vs 13). Moreover, the overlapping of confidence intervals is scarcely meaningful because its very large value in “variability derived” category.

Single-specimen poisoning samples were excluded from histamine concentration meta-analysis in order to not confound within - and between specimen variability.

## CONCLUSIONS

The main goal of our Systematic Review was to remove noise as more as possible from information about values of histamine in foods involved into poisoning; this goal has been reached by producing objective estimates.

To attribute precisely the responsibility of the poisoning event, increasing knowledge, allowing the food business operators to improve their practice or processing, as well as guaranteeing the customer also legally, it is fundamental to approach this topic with pragmatism. We hope that these estimates could be a valid reference for operators and consumers.

The estimate of the mean was found to be fairly high, its precision was unfortunately impaired by a lot of variability (heterogeneity).

Too few suitable data are presently available to conduct a reliable analysis on homogeneous subsets of food.

It is recommended that histamine poisoning episodes are recorded and published including the values of all important variables pointed out in this review, moreover, the variability within poisoning sample should be stated analyzing at least twice the histamine content for each sample.

About the conditions concomitant to the poisonings, the role of several health conditions, drugs and meal composition on the proceeding of an event of histamine (scombroid) poisoning has been underlined several times (Sattler, Hesterberg, et al., 1985; Taylor, 1986; Maintz, Novak,

2007; Hungerford, 2010). Alcoholic beverages can increase the seriousness of the episodes enhancing the absorption of histamine contained in the meal, but even if the importance of alcohol is reported in a previous review (Lehane, Olley, 2000) and other reports (Geiger, 1955; Zee, Simard, et al., 1981; Zimatkin, Anichtchik, 1999; Maintz, Novak, 2007), our results point out lack of this information, so it is recommended to physicians to include such items in the anamneses of the poisoning cases.

## SOURCES OF SUPPORT

### **Internal sources**

Università degli Studi di Milano (Italy): EC work in this systematic review was rewarded by a salary.

### **External sources**

Generale Conserve company, Italy: sponsored research about histamine poisoning in tunafish.

## DECLARATIONS OF INTEREST

This systematic review is a phase of a research commissioned from a food industry.

## ACKNOWLEDGEMENTS

Authors thank Generale Conserve for funding this systematic review.

Authors thank Angela Moccia for her contribute as information expert

# ACCEPTED MANUSCRIPT

## CONTRIBUTIONS OF AUTHORS

CB building protocol, study selection, data extraction and writing review ; EC bibliographic search; FC building protocol, study selection, data extraction and analysis and writing review; PC: building protocol, supervising of all phases and writing review.

## REFERENCES

- Anonymous (1985). Food poisoning and salmonella surveillance in England and Wales: 1983 *British medical journal (Clinical research ed.)*, **291**: 394-396.
- Anonymous (1988). Scombroid fish poisoning--New Mexico, 1987 *MMWR. Morbidity and mortality weekly report*, **37**.
- Anonymous (2000). Scombroid fish poisoning -- Pennsylvania, 1998 *MMWR: Morbidity & Mortality Weekly Report*, **49**: 398-400.
- Anonymous. (2007). Scombroid fish poisoning associated with tuna steaks--Louisiana and Tennessee, 2006. In *MMWR. Morbidity and mortality weekly report*, vol. 56 (pp. 817-819). United States.
- Becker, K., Southwick, K., Reardon, J., Berg, R., Maccormack, J. (2001). Histamine poisoning associated with eating tuna burgers *JAMA*, **285**(10): 1327-1330.
- Bedry, R., Gabinski, C., Paty, M. (2000). Diagnosis of scombroid poisoning by measurement of plasma histamine *N Engl J Med*, **342**: 520-521.
- Bremer, P. J., Fletcher, G. C., Osborne, C. (2003). Scombrototoxin in seafood. In, (pp. 1-9): New Zealand Institute for Crop & Food Research Limited.
- Cattaneo, P. (2011). Sindrome sgombroide -intossicazione da istamina *Food in*, **2**: 5-55.
- Cattaneo, P., Stella, S. (2001). Un episodio di intossicazione da istamina per consumo di pizza al tonno. *Arch. Vet. Ital.*, **52**(5/6): 13.
- Chen, H. C., Huang, Y. R., Hsu, H. H., Lin, C. S., Chen, W. C., Lin, C. M., Tsai, Y. H. (2010). Determination of histamine and biogenic amines in fish cubes (*Tetrapturus angustirostris*) implicated in a food-borne poisoning *Food Control*, **21**: 13-18.

- Chen, H. C., Kung, H. F., Chen, W. C., Lin, W. F., Hwang, D. F., Lee, Y. C., Tsai, Y. H. (2008). Determination of histamine and histamine-forming bacteria in tuna dumpling implicated in a food-borne poisoning *Food Chemistry*, **106**: 612-618.
- Chen, H. C., Lee, Y. C., Hwang, D. F., Chiou, T. K., Tsai, Y. H. (2011). Determination of histamine in mahi-mahi fillets (*Coryphaena hippurus*) implicated in a foodborne poisoning *Journal of Food Safety*, **31**: 320-325.
- Chen, H. C., Lee, Y. C., Lin, C. M., Hwang, D. F., Tsai, Y. H. (2010). Determination of histamine and bacterial isolation in marlin fillets (*Makaira nigricans*) implicated in a foodborne poisoning *Journal of Food Safety*, **30**: 699-710.
- Chianea, D., Drouillard, I., Puyhardy, J. M., Corbe, H., Adam, F., Bietrix, P., Cariou, M., Helies, C., Boutin, J. P. (1998). Collective histamine poisoning confirmed by a rapid high pressure liquid chromatography mass spectrometry technique *Ann Biol Clin*, **56**: 578-579.
- Communities, C. O. T. E. (2007). Commission Regulation (EC) No 1441/2007 of 5 December 2007 amending Regulation (EC) No 2073/2005 on microbiological criteria for foodstuffs. In *Official Journal of the European Union*, vol. L322 (pp. 12 - 13).
- D'aloia, A., Vizzardi, E., Pina, P. D., Bugatti, S., Magro, F. D., Raddino, R., Curnis, A., Cas, L. D. (2011). A scombroid poisoning causing a life-threatening acute pulmonary edema and coronary syndrome in a young healthy patient *Cardiovasc. Toxicol.*, **11**: 280-283.
- Dalgaard, P., Emborg, J., Kjølby, A., Sørensen, N. D., Ballin, N. Z., Dalgaard, P., Emborg, J., Kjølby, A., Sørensen, N. D., Ballin, N. Z. (2008). Histamine and biogenic amines: formation and importance in seafood. In T. Børresen, T. Børresen (Eds.), *Improving seafood products for the consumer*, (pp. 292-324). Cambridge: British Welding Research Association.

- Demoncheaux, J. P., Michel, R., Mazenot, C., Duflos, G., Iacini, C., De Laval, F., Saware, E. M., Renard, J. C. (2012). A large outbreak of scombroid fish poisoning associated with eating yellowfin tuna (*Thunnus albacares*) at a military mass catering in Dakar, Senegal. In *Epidemiol Infect*, vol. 140 (pp. 1008-1012).
- Dersimonian, R., Laird, N. (1986). Meta-analysis in clinical trials *Control. Clin. Trials* **7**: 177-188.
- Doeglas, H. M., Huisman, J., Nater, J. P. (1967). Histamine intoxication after cheese *Lancet*, **2**: 1361-1362.
- Eckstein, M., Serna, M., Delacruz, P., Mallon, W. K. (1999). Out-of-hospital and emergency department management of epidemic scombroid poisoning *Acad Emerg Med*, **6**: 916-920.
- Emborg, J., Dalgaard, P. (2006). Formation of histamine and biogenic amines in cold-smoked tuna: an investigation of psychrotolerant bacteria from samples implicated in cases of histamine fish poisoning *Journal of Food Protection*, **69**: 897-906.
- Emborg, J., Laursen, B. G., Dalgaard, P. (2005). Significant histamine formation in tuna (*Thunnus albacares*) at 2(degrees)C - Effect of vacuum- and modified atmosphere-packaging on psychrotolerant bacteria *International journal of food microbiology*, **101**: 263-279.
- F.A.O. (2014). Joint FAO/WHO Expert Meeting on the Public Health Risks of Histamine and other Biogenic Amines from Fish and Fishery Products. In).
- F.D.A. (2014). Scombrototoxin (Histamine) Formation. In *Fish and Fishery Products Hazards and Controls Guidance* 4th Edition ed.).

- Feldman, K. A., Werner, S. B., Cronan, S., Hernandez, M., Horvath, A. R., Lea, C. S., Au, A. M., Vugia, D. J. (2005). A large outbreak of scombroid fish poisoning associated with eating escolar fish (*Lepidocybium flavobrunneum*) *Epidemiol Infect*, **133**: 29-33.
- Fernández, M. A., González, J. M. B., Rozas, S. F. (2001). Escombroid intoxicación por consumo de bonito *Emergencias*.
- Foo, L. Y. (1975a). The content of histamine and fish food poisoning *N Z Med J*, **82**: 381-383.
- Foo, L. Y. (1975b). Scombroid-type poisoning induced by the ingestion of smoked kahawai *N Z Med J*, **81**: 476-477.
- Foo, L. Y. (1977). Scombroid poisoning: Recapitulation on the role of histamine *N Z Med J*, **85**: 425-427.
- Geiger, E. (1955). Role of histamine in poisoning with spoiled fish *Science*, **121**: 865-866.
- Gellert, G. A., Ralls, J., Brown, C., Huston, J., Merryman, R. (1992). Scombroid fish poisoning. Underreporting and prevention among noncommercial recreational fishers *Western Journal of Medicine British Medical Journal*, **157**: 645-647.
- Guillier, L., Thebault, A., Gauchard, F., Pommepeuy, M., Guignard, A., Malle, P. (2011). A risk-based sampling plan for monitoring of histamine in fish products *Journal of Food Protection*, **74**: 302-310.
- Guly, H. R., Grant, I. C. (2006). Case of the month: lesson of the week: don't forget scombroid *Emergency Medicine Journal*, **23**: 955-956.
- Hall, M. (2003). Something fishy: six patients with an unusual cause of food poisoning! *Emerg. Med.*, **15**: 293-295.

- Higgins, J. P. T., Thompson, S. G., Deeks, J. J., Altman, D. G. (2003). Measuring inconsistency in metaanalyses *Brit. Med. J.* , **327**: 557-560.
- Higgins, J. P. T., White, I. R., Anzures-Cabrera, J. (2008). Meta-analysis of skewed data: Combining results reported on log-transformed or raw scales. *Stat Med*, **27**: 6072–6092.
- Hobbs, G. (1983). Food poisoning and fish *The Journal of the Royal Society for the Promotion of Health*, **103**: 144-149.
- Hungerford, J. M. (2010). Scombroid poisoning: a review. In *Toxicon*, vol. 56 (pp. 231-243). England: Published by Elsevier Ltd.
- Hwang, D. F., Chang, S. H., Shiua, C. Y., Tuu-Jyi, C. (1997). High-performance liquid chromatographic determination of biogenic amines in fish implicated in food poisoning *J Chrom B*, **693**: 23-29.
- Iguchi, S., Sugita, M., Nomura, T., Sekii, H., Ookubo, H., Yamaguchi, N. (2008). A local outbreak of scombroid fish poisoning in Japan *Clin. Toxicol.*, **46**: 621-621.
- Jantschitsch, C., Kinaciyar, T., Manafi, M., Safer, M., Tanew, A. (2011). Severe scombroid fish poisoning: An underrecognized dermatologic emergency *J. Am. Acad. Dermatol.*, **65**: 246-247.
- Jiang, D. D. S., Liu, C. M., Wang, S. C., Lin, C. W., Chen, H. R. (2009). Histamine-Induced Allergic Outbreak in among Junior/Senior High School Students-A Case-Control Study *Epidemiology Bulletin*, **25**(5): 316-331.
- Kanki, M., Yoda, T., Ishibashi, M., Tsukamoto, T. (2004). Photobacterium phosphoreum caused a histamine fish poisoning incident *International journal of food microbiology*, **92**: 79-87.
- Kelso, J. M., Lin, F. L. (2009). Skin testing for scombroid poisoning *Annals of Allergy, Asthma & Immunology*, **103**: 447-447.



- Kim, R. (1979). Flushing syndrome due to mahimahi (scombroid fish) poisoning *Arch. Dermatol.*, **115**: 963-965.
- Kow-Tong, C., Malison, M. D. (1987). Outbreak of scombroid fish poisoning, Taiwan *Am J Public Health*, **77**: 1335-1336.
- Leask, A., Yankos, P., Ferson, M. J. (2004). Fish, so foul! Foodborne illness caused by combined fish histamine and wax ester poisoning *Communicable diseases intelligence*, **28**: 83-85.
- Lee, Y. C., Huang, T. C., Lin, C. S., Lin, C. M., Tsai, Y. H. (2012). Determination of Histamine and Histamine-forming Bacteria in Striped Marlin Fillets (*Tetrapturus audax*) Implicated in a Food-borne Poisoning *Toxicon*, **60**: 161-162.
- Lehane, L., Olley, J. (2000). Histamine fish poisoning revisited *International journal of food microbiology*, **58**: 1-37.
- Maintz, L., Novak, N. (2007). Histamine and histamine intolerance *Am J Clin Nutr*, **85**(5): 1185-1196.
- Mclauchlin, J., Little, C. L., Grant, K. A., Mithani, V. (2006). Scombrototoxic fish poisoning *J. Public Health*, **28**(1): 61-62.
- Molinari, G., Montagnoli, G., Pellegrini, G., Caroli, G. (1989). [Hygiene and health importance of histamine as an unhealthy factor in several food products] *Annali di igiene : medicina preventiva e di comunità*, **1**: 637-646.
- Muller, G. J., Lamprecht, J. H., Barnes, J. M., De Villiers, R. V. P., Honeth, B. R., Hoffman, B. A. (1992). Scombroid poisoning. Case series of 10 incidents involving 22 patients *S Afr Med J*, **81**: 427-430.

- Murphy, G., Pfeiffer, R., Camargo, M. C., Rabkin, C. S. (2009). Meta-analysis shows that prevalence of Epstein–Barr virus-positive gastric cancer differs based on sex and anatomic location *Gastroenterology*, **137**(3): 824-833.
- Nalinee Hongchumpon Oupparong, T. P. J., Et Al. (2010). Scombrototoxin Food Poisoning Outbreak among Frozen Seafood Factory Workers Samut Prakan Province, Thailand, July 2007 *Outbreak, Surveillance and Investigation Reports*, **2**(1): 5.
- Nelson, J. S. (2006). *Fishes of the World* (4th edition ed.): Hoboken, NJ: John Wiley & So.
- Ohnuma, S., Higa, M., Hamanaka, S., Matsushima, K., Yamamuro, W. (2001). An outbreak of allergy-like food poisoning *Intern Med*, **40**: 833-835.
- Predy, G., Honish, L., Hohn, W., Jones, S. (2003). Was it something she ate? Case report and discussion of scombroid poisoning *CMAJ: Canadian Medical Association Journal*, **168**: 587-588.
- Quan, H., Zhang, J. (2003). Estimate of standard deviation for a log-transformed variable using arithmetic means and standard deviations *Stat Med*, **22**: 2723–2736.
- Sanchez-Guerrero, I. M., Vidal, J. B., Escudero, A. I. (1997). Scombroid fish poisoning: A potentially life-threatening allergic-like reaction *J Allergy Clin Immunol*, **100**: 433-434.
- Sanders, W. E. (1987). Intoxications from the seas: ciguatera, scombroid, and paralytic shellfish poisoning *Infect Dis Clin North Am*, **1**: 665-676.
- Sattler, J., Hesterberg, R., Lorenz, W., Schmidt, U., Crombach, M., Stahlknecht, C. D. (1985). Inhibition of human and canine diamine oxidase by drugs used in an intensive care unit: relevance for clinical side effects? *AGENTS ACTIONS*, **16**: 91-94.

Schulze, K., Reusse, V., Tillack, J. (1979). Histamine food poisoning after consumption of sardines in oil *Archiv fuer Lebensmittelhygiene*, **30**: 56-59.

Schwarzer G. (2010). Meta-Analysis with R. R package version 1.6-0.

Shalaby, A. R. (1996). Significance of biogenic amines to food safety and human health *Food Research International*, **29**: 675-690.

Shore, P. A., Burkhalter, A., Cohn, V. H. (1959). A method for the fluorometric assay of histamine in tissues *J Pharmacol Exp Ther*, **127**(3): 182-186.

Sinn, G. (2006). Histamine poisoning and gastroenteritis breakout after participation in a school lunch with fish burgers *Gesundheitswesen*, **68**: 216-217.

Stell, I. M. (1997). Trouble with tuna: Two cases of scombrototoxin poisoning *J Accid Emerg Med*, **14**: 110-111.

Su, S. C., Chou, S. S., Chang, P. C., Hwang, D. F. (2000). Determination of biogenic amines in fish implicated in food poisoning by micellar electrokinetic capillary chromatography *J. Chromatogr. B Biomed. Sci. Appl.*, **749**: 163-169.

Taylor, S. L. (1986). Histamine food poisoning: Toxicology and clinical aspects *Crit Rev Toxicol*, **17**: 91-128.

Taylor, S. L., Keefe, T. J., Windham, E. S., Howell, J. F. (1982). Outbreak of histamine poisoning associated with consumption of Swiss cheese *Journal of Food Protection*, **45**: 455-457.

Tsai, Y. H., Hsieh, H. S., Chen, H. C., Cheng, S. H., Chai, T. J., Hwang, D. F. (2007). Histamine level and species identification of billfish meats implicated in two food-borne poisonings *Food Chemistry*, **104**: 1366-1371.

- Tsai, Y. H., Kung, H. F., Chen, H. C., Chang, S. C., Hsu, H. H., Wei, C. I. (2007). Determination of histamine and histamine-forming bacteria in dried milkfish (*Chanos chanos*) implicated in a food-borne poisoning *Food Chemistry*, **105**: 1289-1296.
- Tsai, Y. H., Kung, H. F., Lee, T. M., Chen, H. C., Chou, S. S., Wei, C. I., Hwang, D. F. (2005). Determination of histamine in canned mackerel implicated in a food borne poisoning *Food Control*, **16**: 579-585.
- Valentini, P., Levre, E., Molinari, G., Brunetti, M., Domenici, D., Caroli, G. (1991). Histamine and histamine-producing enterobacteria in canned tuna fish responsible for the scombrototoxic syndrome *Igiene Moderna*, **95**: 154-164.
- Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software*, **36**(3): 1-48.
- Wu, M. L., Yang, C. C., Yang, G. Y., Cer, J., Deng, J. F. (1997). Scombroid fish poisoning: An overlooked marine food poisoning *Vet. Hum. Toxicol.*, **39**: 236-241.
- Zee, J. A., Simard, R. E., Vaillancourt, R., Boudreau, A. (1981). Effect of *Lactobacillus brevis*, *Saccharomyces uvarum* and grist composition on amine formation in beers *Canadian Institute Food Science and Technoogy*, **14**(4): 321-325.
- Zimatkin, S. M., Anichtchik, O. V. (1999). Alcohol-histamine interactions *Alcohol Alcohol.*, **34**: 141-147.

Table A: Characteristics of searched database.

Database category	Datab	Data	Database-	Fin
(some database belong to more than one category, in this case a semicolon (“;”) is used to separate multiple categories)	ase	base	specific search	al
	name	time	strategy	nu
		interv	(sometimes	mbe
		al	three contiguous	r of
			underscores	ite
			“___” are used	ms
			to separate	retri
			search steps or	eve
			multiple	d
			searches – this	fro
			symbol doesn’t	m
			belong to search	data
			strategy syntax;	bas
			moreover	e
			sometimes, due	
			to syntax	
			complexity,	
			only the main	
			structure of	
			search is	

			reported )	
Dissertations and theses database	DAR T EUR OPE	-	Keywords = histamine poisoning Keywords = histamine intoxication Keywords = "scombroid syndrome"	1
	ProQu est Disser tation & These s Datab ase	-	histamine AND (poisoning OR intoxication) OR "scombroid syndrome"	0
Full text journals available electronically	BioM ed Centr	1997- 2013	((histamine AND (poisoning OR	0

	al		intoxication)) OR scombroid and syndrome and (Exact and phrase) and in and All and fields and (full and text)) (All words) in All fields (full text) from 1997 to 2013 ____ 13 (All words) in All fields (full text) from 1997 to 2013 ____ (histamine AND (poisoning OR intoxication)) OR scombroid syndrome (Exact phrase)	
--	----	--	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--

			<p>in All fields (full text) (All words) in All fields (full text) ___ histamine AND (poisoning OR intoxication) (All words) in All fields (full text) ___ poisoning OR intoxication (All words) in All fields (full text) ___ scombroid syndrome (Exact phrase) in All fields (full text) ___ intoxication (All words) in All</p>	
--	--	--	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--



			fields (full text) ____ poisoning (All words) in All fields (full text) ____ histamine (All words) in All fields (full text) from 1997 to 2013	
	High Wire Press	01/01 /1959 - 31/12 /2013	(histamine and poisoning) or (histamine and intoxication) or "scombroid syndrome"	15
	PubM ed Centr al (PMC )	01/01 /1959 - 31/12 /2013	(((((("Histamine/ poisoning"[Mes h] OR "Histamine/toxi city"[Mesh]) AND	51

			<p>(Humans[Mesh]      ))) OR      ("scombrototoxin"      [Supplementary      Concept] OR      "saurine"[Suppl      ementary      Concept])))) OR      (((("Biogenic      Amines/agonists      "[Mesh] OR      "Biogenic      Amines/poisoni      ng"[Mesh] OR      "Biogenic      Amines/toxicity      "[Mesh])) AND      ("Histamine"[M      esh]) AND      (Humans[Mesh]      )) AND      (Humans[Mesh]</p>	
--	--	--	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--

			)	
General bibliographic database	CENT RAL	1959- 2012	#1 MeSH descriptor: [Biogenic Amines] explode all trees ____ #2 #1 from 1959 to 2012, in Trials ____ #3 #1 or #2 from 1959 to 2012, in Trials ____ #4"scombroid fish poisoning" or "scombroid food poisoning" or "scombroid poisoning" or "scombroid syndrome" or "scombroid type	1

			poisoning" or "scombroide intoxicatie" or "scombrototoxic" or "scombrototoxic fish" or "scombrototoxic fish poisoning" or "scombrototoxic poisoning" or "scombrotoxico sis" or "scombrotoxin" or "scombrotoxin poisoning" or "scombrotoxins " from 1959 to 2012, in Trials —	
--	--	--	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--

			<p>#5“histamine poisoning”or "histamine intoxication" from 1959 to 2012, in Trials ___ #6 #3 or #4 or #5 from 1959 to 2012, in Trials ___ #7MeSH descriptor: [Foodborne Diseases] explode all trees ___ #8 "food poisoning" from 1959 to 2012, in Trials ___ #9 #7 or #8 from 1959 to 2012, in Trials ___</p>	
--	--	--	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--

		<p>#10#6 and #9  from 1959 to  2012, in Trials  ___ #11 MeSH  descriptor:  [Food] explode  all trees ___  #12food ___  #13 #11 or #12  from 1959 to  2012, in Trials  ___ #14 MeSH  descriptor:  [Fishes] explode  all trees ___ #15  fish ___ #16  #14 or #15 from  1959 to 2012, in  Trials ___ #17  #13 or #16 ___  #18 #10 and  #17</p>
--	--	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

	Cinah	01/01	S1 (MH	24
	1	/1959	"Histamine/PO"	
		-	) ___ S2	
		31/12	(MH	
		/2013	"Histamine")	
			___ S3	
			(MH	
			"Toxicology")	
			OR (MH	
			"Toxins+") ___	
			S4	
			scombrotxin*	
			___ S5	
			biogenic* ___	
			S6 S1 OR	
			S4 ___ S7	
			(S6) OR (S2	
			AND S3) ___	
			S8 (S5)	
			AND (S2) ___	
			S9 S7 ___	
			S10 (S7) OR	

			(S8) ___ S11 (MH "Poisoning+") OR TI POISON* ___ S12 (S2 AND S11) ___ S13 S10 OR S12	
	FSTA (Food Scien ce and Techn ology Abstr acts)	1959 - Curre nt	1. scombroid poisoning/ ___ 2. histamine/ ___ 3. poisoning/ ___ 4. FOOD POISONING/ ___ 5. exp toxicity/ ___ 6. (POISON* or TOXIC*).hw,ti. ___ 7. HISTAMINE.ti.	82



			___ 8. 6 and 7 ___ 9. 3 or 4 or 5 ___ 10. 2 and 9 ___ 11. 1 or 8 or 10 ___ 12. saurine.hw,ti. ___ 13. 11 or 12	
	SciFin der	1959- 2012	Main structure of search: #1 “biogenic amines”.to ___ #2 “scombroid syndrome” ___ #3 “histamine poisoning” ___ #4 #1 or #2 or #3 ___ #5 “food poisoning” ___ #6 #4 and #5 ___ #7 food ___ #8 fish ___ #9 #7 or #8 ___	0

			#10 #6 and #9	
	Scirus	1950-2013	("food poisoning" and ("biogenic amines" or "scombroid syndrome" or "scombroid fish poisoning" or "scombroid food poisoning" or "scombroid poisoning" or "scombroid syndrome" or "scombroid type poisoning" or "scombroid intoxication" or "scombrototoxic" or "scombrototoxic fish" or	87

			"scombrototoxic fish poisoning" or "scombrotoxic poisoning" or scombrototoxicosis or scombrotoxin or "scombrotoxin poisoning" or scombrotoxins or "scombroid toxin" or "histamine poisoning" or "histamine intoxication")) and (food or fish)	
	Scopus	>1958 & =201	(TITLE(scombroid* OR scrombotox*))	83

		3	OR (INDEXTERM S(scombrotax* OR scombroid*)) OR (TITLE(scombr otox* OR scombroid*)) OR (TITLE(histami ne) AND TITLE(poison* OR toxic*))	
General bibliographic database; Conference abstracts or proceedings	Pubmed	01/01 /1959 - 2013/ 12/31	(((("Histamine/ poisoning"[Mes h] OR "Histamine/toxi city"[Mesh]) AND (Humans[Mesh] ))) OR	102

			("scombrototoxin" [Supplementary Concept] OR "saurine"[Suppl ementary Concept])) OR (((("Biogenic Amines/agonists "[Mesh] OR "Biogenic Amines/poisoni ng"[Mesh] OR "Biogenic Amines/toxicity "[Mesh])) AND ("Histamine"[M esh]) AND (Humans[Mesh] )) AND (Humans[Mesh] )	
	Web	1985-	ti=histamine	54

	of Scien ce	01-01 - 2012- 11-05	<p>____</p> <p>Databases=SCI-</p> <p>EXPANDED,</p> <p>SSCI, A&amp;HCI,</p> <p>CPCI-S, CPCI-</p> <p>SSH</p> <p>Timespan=All</p> <p>Years ____</p> <p>Lemmatization=</p> <p>On ____</p> <p>ti=(toxic* or</p> <p>poison*) ____</p> <p>Databases=SCI-</p> <p>EXPANDED,</p> <p>SSCI, A&amp;HCI,</p> <p>CPCI-S, CPCI-</p> <p>SSH</p> <p>Timespan=All</p> <p>Years ____</p> <p>Lemmatization=</p> <p>On ____ #2</p> <p>AND #1 ____</p>	
--	-------------------	------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--

			<p>Databases=SCI- EXPANDED, SSCI, A&amp;HCI, CPCI-S, CPCI- SSH</p> <p>Timespan=All Years ____</p> <p>Lemmatization= On ____</p> <p>ti=(biogen* and amin*) ____</p> <p>Databases=SCI- EXPANDED, SSCI, A&amp;HCI, CPCI-S, CPCI- SSH</p> <p>Timespan=All Years ____</p> <p>Lemmatization= On ____ #4 AND #2 ____</p> <p>Databases=SCI-</p>	
--	--	--	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--

			EXPANDED, SSCI, A&HCI, CPCI-S, CPCI- SSH Timespan=All Years ____ Lemmatization= On ____ ti=((scombr* and (poison* or toxic* or syndro*)) or scombrottox* or saurine) ____ Databases=SCI- EXPANDED, SSCI, A&HCI, CPCI-S, CPCI- SSH Timespan=All Years ____ Lemmatization=	
--	--	--	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--



			<p>On ___ #6 OR #3 ___ Databases=SCI- EXPANDED, SSCI, A&amp;HCI, CPCI-S, CPCI- SSH Timespan=All Years ___ Lemmatization= On ___ #7 OR #5 ___ Databases=SCI- EXPANDED, SSCI, A&amp;HCI, CPCI-S, CPCI- SSH Timespan=All Years ___ Lemmatization= On</p>	
General bibliographic database ; Other reviews,	Emba	<196	histamine:ti	151

guidelines and reference lists as sources of studies	se	6- 2013	AND (poison*:ti OR toxic*:ti) OR scombroid:ti OR scombrottox*:ti AND [embase]/lim	
General bibliographic database ; Subject-specific bibliographic database	Googl e schola r	1959- 2013	The following 4 searches were performed and their results combined A[1959-2000] = food "biogenic amines" OR "scombroid syndrome" OR "histamine poisoning" "food poisoning" ____	180

			___ A[2001-2012] = food "biogenic amines" OR "scombroid syndrome" OR "histamine poisoning" "food poisoning" ___ ___ B [1959-2000] = (fish "biogenic amines" OR "scombroid syndrome" OR "histamine poisoning" "food poisoning") ___ B [2001-2012] = (fish	
--	--	--	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--

			"biogenic amines" OR "scombroid syndrome" OR "histamine poisoning" "food poisoning")	
Grey literature database		-	The following 2 searches were performed and their results merged: histamine and (poisoning or intoxication) "scombroid syndrome"	0
	EAG LE (Open Grey)	1959- 2013	((histamine AND (poisoning OR intoxication) )	0

			OR “scombroid syndrome”	
	The National Technical Information Service (NTIS)	1964-2013	The following 2 searches were performed and their results merged: histamine and (poisoning or intoxication) “scombroid syndrome”	3
Other reviews, guidelines and reference lists as sources of studies	Medline (Ovid)	1959-2013	exp histamine/ _____ histamine.mp. _____ 1 or 2 _____ exp poisoning/ _____ poisoning.mp.	0

			<p>___ ___ 4 or 5</p> <p>___</p> <p>intoxication.mp.</p> <p>___ ___</p> <p>intoxication.mp.</p> <p>___ ___ 7 or 8</p> <p>___ 6 or 9 ___ 3</p> <p>and 10 ___</p> <p>'scombroid</p> <p>syndrome'.mp.</p> <p>___ 'scombroid</p> <p>syndrome'.mp.</p> <p>___ 12 or 13</p> <p>___ ___ 11 or</p> <p>14 ___ - ___</p> <p>___ ___ ___</p> <p>16 limit 15 to</p> <p>"review articles"</p> <p>___ 17</p> <p>'systematic</p> <p>review'.mp. ___</p> <p>18 16 and 17</p>	
--	--	--	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--

			<p>___ - ___ 19 18</p> <p>___ 20 limit 19</p> <p>to yr="1959 -</p> <p>Current"</p>	
	The cochr ane librar y	1959- 2013	<p>#1 MeSH descriptor: [Histamine] explode all trees and with qualifiers: [Poisoning - PO] ___ #2 MeSH descriptor: [Histamine] explode all trees and with qualifiers: [Toxicity - TO] ___ #3 scombrotax*:ti (Word</p>	0

			<p>variations have been searched) ___ #4 saurine:ti,ab,kw (Word variations have been searched) ___ #5 MeSH descriptor: [Biogenic Amines] explode all trees and with qualifiers: [Poisoning - PO] ___ #6 MeSH descriptor: [Biogenic Amines] explode all trees and with</p>	
--	--	--	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--



			<p>qualifiers:          [Toxicity - TO]          ___ #7 MeSH</p> <p>descriptor:          [Histamine]</p> <p>explode all trees          ___ #8 (((#5          or #6) and #7)          or #3 or #2 or          #1) ___ #9</p> <p>scombroid:ti,ab,          kw (Word          variations have          been searched)</p>	
	Africa n Index Medic us	-	. histamine and poisoning . histamine and intoxication . "scombroid syndrome"	0
	FAO	1959- 2013	((histamine AND	0

		(poisoning OR intoxication)) OR “scombroid syndrome”	
Index Medicus for the South-East Asia Region (IMS EAR)	-	. histamine and poisoning ____ . histamine and intoxication ____ .”scombroid syndrome”	2
IndMED	-	. histamine and poisoning ____ . histamine and intoxication ____ .”scombroid syndrome”	0
Korea	-	. histamine and	0

	Med		poisoning . histamine and intoxication . "scombroid syndrome"	
	LILA CS	1959- 2013	Sintassi LILACS ____ MH biogenic amines and (PD 1959 or PD 196\$ or PD 197\$ or PD 198\$ or PD 199\$ or PD 200\$ or PD 2010 or PD 2011 or PD 2012 or PD 2013) ____ [scombroid syndrome or scombroid fish	0

			poisoning or scombroid food poisoning or scombroid poisoning or scombroid syndrome or scombroid type poisoning or scombroid intoxicatio or scombrototoxic or scombrototoxic fish or scombrototoxic fish poisoning or scombrototoxic poisoning or scombrototoxicosi s or scombrototoxin or	
--	--	--	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--

			scombrototoxin poisoning or scombrototoxins or scombrototoxin ] ____ ____ and (PD 1959 or PD 196\$ or PD 197\$ or PD 198\$ or PD 199\$ or PD 200\$ or PD 2010 or PD 2011 or PD 2012 or PD 2013) ____ TI histamine and TI poisoning and (PD 1959 or PD 196\$ or PD 197\$ or PD 198\$ or PD	
--	--	--	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--

			199\$ or PD	
			200\$ or PD	
			2010 or PD	
			2011 or PD	
			2012 or PD	
			2013) ___ #1	
			___ MH food	
			poisoning or (TI	
			food and TI	
			poisoning) and	
			(PD 1959 or PD	
			196\$ or PD	
			197\$ or PD	
			198\$ or PD	
			199\$ or PD	
			200\$ or PD	
			2010 or PD	
			2011 or PD	
			2012 or PD	
			2013) ___ #1	
			and # 5 ___ -	
			___ - ___ - ___	

			-	
	Pantel eimon	-	1 ____ Keywords: histamine ____ 2 ____ Keywords: poisoning ____ 3 ____ Keywords: intoxication ____ 4 ____ Keywords: "scombroid syndrome" ____ 5 ____ Keywords: scombroid syndrome ____ 6 ____ Комбінація: 1 1 AND 2 ____ 7 ____ Комбінація: 1 1 AND 3 ____ 8	0

			— Комбінація: 4 4 OR 5 OR 6 OR 7	
Regional/National bibliographic database	Western Pacific Regional Index Medicus (WPRIM)	1959-2013	4#3 or #2 or #1 ___ 3 All:"scombroid syndrome" - Limits:1959-2013 ___ 2 All:histamine and All:intoxication -Limits:1959-2013 ___ 1 All:histamine and All:poisoning - Limits:1959-2013	0
Subject-specific bibliographic database	Bibliomap	-	1 Freetext: histamine ___ 3	0



	EPPI- Centr e databa se of health promo tion resear ch		Freetext: poisoning ___ 5 Freetext: intoxication ___ 10 Freetext: "scombroid syndrome" ___ 11 3 OR 5 ___ 12 1 AND 11 ___ 13 10 OR 12	
	Datab ase of Prom oting Healt h Effect iveness s Revie ws	-	1 Freetext: histamine ___ 3 Freetext: poisoning ___ 5 Freetext: intoxication ___ 7 Freetext: "scombroid syndrome" ___ 8 1 AND 5 ___ 9 1 AND 3 ___	0

	(DoP HER)		10 7 OR 8 OR 9	
	Global Health	-	(histamine AND poisoning) AND "scombroid syndrome" or (histamine AND intoxication) AND "scombroid syndrome"	0
	Intute	?-july 2011	. histamine and poisoning or. histamine and intoxication or ."scombroid syndrome"	0
	POPL INE (repro ductiv	1959- 2013	histamine AND poisoning or histamine AND intoxication or	0

	e health ) (free on the intern et)		“scombroid syndrome” <limit_to_years _1959-2013>	
	Turni ng Resea rch into Practi ce (TRIP ) databa se	1959- 2013	#4 ___ (#1 or #2 or #3) ___ #3 ___ ("scombroid syndrome") from:1959 to:2013 ___ #2 ___ (histamine poisoning) from:1959 to:2013 ___ #1 ___ (histamine intoxication) from:1959	16

			to:2013	
--	--	--	---------	--

Table 1: Characteristics of studies included in meta-analysis. N for number of people involved in histamine outbreak, MD for missing data. Type of food (food category): (1) fresh, (2) frozen, (3) canned, (4) fermented, (5) other seafood, (6) cheese, (7) other foods. Values of sample mean or sample standard deviation that were calculated have been rounded to two decimals.

Poisoning sample id / Reference	N	Locati on	Alco hol cons umpt ion	Pre viou s med icati on	Fe m ale se x	Health status	Heter ogene ous food	Hist ami ne- rele asin g cap aciti es foo d	Sour ce of food (food catego ry)	Type of food (food catego ry)	Sa m ple siz e	Sa mpl e mea n (mg /kg)	Sa mp le sta nda rd dev iati on (m g/k g)
Becker_2001_ ET_aai_1 <sup>2</sup>	1 1	North Caroli na	MD	MD	M D	unknown	no	yes	resta uran t	frozen tuna burger s (2)	2	299 5	353 .55
Becker_2001_	2	North	MD	MD	M	MD	yes	yes	resta	tuna	3	231	182

ET_aai_3 <sup>2</sup>		Caroli na			D				uran t	pieces in salad (5)		.83	.11
Bedry_2000_ EN_mi1_1 <sup>12</sup>	9	Borde aux, France	MD	MD	no	unknown	no	yes	MD	cooke d tuna (5)	M D	MD	M D
Bremer_2003 _ET_cca_1 <sup>13</sup>	5	Greate r Auckl and region (New Zealan d)	MD	MD	M D	unknown	MD	yes	MD	Hot smoke d kahaw ai ( <i>Arrip is trutta</i> ) (5)	2	190 - 985	M D
Bremer_2003 _ET_cca_2 <sup>13</sup>	2	Greate r Auckl and region (New	MD	MD	M D	unknown	MD	yes	MD	Hot smoke d kahaw ai ( <i>Arrip</i>	2	170 0- 210 0	M D

		Zealand)								<i>is trutta</i> (5)			
Bremer_2003 _ET_cca_3 <sup>13</sup>	1 6	Greater Auckland and region (New Zealand)	MD	MD	MD	unknown	MD	yes	MD	Hot smoke d kahawai ( <i>Arripis trutta</i> ) (5)	9	200 - 394 0	M D
Chen_2008_E T_hhh_1 <sup>14</sup>	7	Chiayi Prefecture, southern Taiwan	MD	MD	MD	unknown	yes	yes	MD	tuna dampplings (5)	3	160 8	59
Chen_2010_C B_zxc_1 <sup>15</sup>	3 4 7	Kaohsiung City	MD	MD	MD	unknown	no	yes	school cant	fried fish cubes	7	811 4.4	19. 94

		(Taiwan)							een	<i>(Tetra pturus angus tirostr is)</i> (5)			
Chen_2010_E T_fff_1 <sup>16</sup>	7	Kaohsiung City (Taiwan)	MD	MD	M D	unknown	no	yes	restaurant	marlin fillets (5)	2	456 .5	30. 41
Chen_2011_E T_eee_1 <sup>17</sup>	5 3	Kaohsiung City (Taiwan)	MD	MD	M D	unknown	no	yes	restaurant	mahi mahi fillets (5)	2	245	186 .68
Chianèa_1998 _ET_nnn_1 <sup>18</sup>	M D	West Africa	MD	MD	M D	unknown	no	yes	sell on Africa coast	smoked sword fish (5)	9	203 0- 475 0	M D
D'Aloia_2011	1	Brescia	MD	MD	ye	healthy	no	yes	MD	grilled	1	>10	M



_EN_5jb_1 <sup>19</sup>		a (Italy)			s					tuna (5)		000 00	D
Demoncheaux _2012_EN_99 w_1 <sup>20</sup>	7 1	Dakar, Seneg al	MD	MD	no	mainly healthy	no	yes	milit ary cate ring facil ity	yello wfin tuna (1)	2	490 0	145
Doeglas_1967 _ET_vvv_1 <sup>21</sup>	1	Rotter dam	MD	MD	no	unknown	no	no	MD	brand of very old Goud a chees e (6)	1	850	M D
Eckstein_199 9_EN_8f6_1 <sup>22</sup>	3 0	Los Angel es	MD	MD	M D	unknown	no	yes	cate ring	escola r <i>(Lepid ocybi um flavob</i>	5	111 - 767 0	M D

										<i>runne um)</i> (2)			
Emborg_2005 _CB_kju_1 <sup>23</sup>	8	Denm ark	MD	MD	M D	unknown	no	yes	cant een	chilli- marin ated steaks of Yello wfin tuna (5)	2	810 0	141 4.2 1
Emborg_2006 _CB_kjs_1 <sup>24</sup>	2	Denm ark	MD	MD	M D	unknown	no	yes	hom e	Cold- Smok ed Tuna (5)	2	454 8	123
Emborg_2006 _CB_kjs_2 <sup>24</sup>	1	Denm ark	MD	MD	M D	unknown	no	yes	priv ate part y	Cold- Smok ed Tuna (5)	2	197 2	4

Emborg_2006 _CB_kjs_3 <sup>24</sup>	1 0	Denm ark	MD	MD	M D	unknown	no	yes	buff et at a cant een	Cold- Smok ed Tuna (5)	2	914	8
Feldman_200 5_EN_555_1 <sup>2</sup> 5	4 2	Marin Count y, Califo rnia, USA	MD	MD	no	unknown	no	yes	retre at cent re	grilled escola r fish ( <i>Lepid ocybi um flavob runne um</i> ) (2)	4	282 5	741 .06
Fernandez_20 01_CB_3jd_3 26	3	Valde cilla (Spain )	MD	MD	no	unknown	no	yes	hom e	<i>Scom berida e</i> (1)	1	525 0	M D
Foo_1975_C B_h67_1 <sup>27</sup>		Napler (New Zealan	MD	MD	M D	unknown	no	yes	MD	kingfi sh (5)	1	758 0	M D

		d)											
Foo_1975_C B_jso_1 <sup>28</sup>		Wellington( New Zealand)	MD	MD	M D	unknown	no	yes	MD	kahawai (5)	1	800 0	M D
Foo_1977_C B_8sk_1 <sup>29</sup>		Petone (New Zealand)	MD	MD	M D	unknown	no	yes	MD	canne d skipjack (3)	12	567 .08	388 .08
Gellert_1992_82c_1 <sup>30</sup>	8	Santa Ana, California	MD	MD	M D	unknown	no	yes	fish caught by a recreational eater onal fisher	smoked yellowfin and skipjack tuna (5)	6	696 0	M D
Guly_2006_E N79j_1 <sup>31</sup>	5	Plymouth, UK	MD	MD	M D	unknown	no	yes	hotel buffet	fresh tuna (1)	1	>35 0	M D

									et				
Hall_2003_C B_h8s_1 <sup>32</sup>	6	Canberra	MD	no	no	unknown ;hypertension	no	yes	restaurants	yellowfin tuna (5)	2	480	14. 14
Hobbs_1982_ FC_55z_14 <sup>33</sup>	4	UK	MD	MD	M D	unknown	no	yes	MD	Canne d Tuna (3)	M D	105 0	M D
Hobbs_1982_ FC_55z_15 <sup>33</sup>	2	Malaysia	MD	MD	M D	unknown	no	yes	MD	Canne d Tuna (3)	M D	290 0	M D
Hobbs_1982_ FC_55z_16 <sup>33</sup>	2	UK	MD	MD	M D	unknown	no	yes	MD	Canne d Tuna (3)	M D	28	M D
Hobbs_1982_ FC_55z_18 <sup>33</sup>	1	UK	MD	MD	M D	unknown	no	yes	MD	Canne d Tuna (3)	M D	35	M D
Hobbs_1982_	2	UK	MD	MD	M	unknown	no	yes	MD	Canne	M	720	M

FC_55z_20 <sup>33</sup>					D					d Sardin es (3)	D		D
Hobbs_1982_ FC_55z_21 <sup>33</sup>	1	UK	MD	MD	M D	unknown	no	yes	MD	Canne d Anch ovies (3)	M D	680	M D
Hobbs_1982_ FC_55z_22 <sup>33</sup>	3	UK	MD	MD	M D	unknown	no	yes	MD	Canne d Sardin es (3)	M D	300 0	M D
Hobbs_1982_ FC_55z_23 <sup>33</sup>	1	UK	MD	MD	M D	unknown	no	yes	MD	Canne d Tuna (3)	M D	20	M D
Hobbs_1982_ FC_55z_24 <sup>33</sup>	1	UK	MD	MD	M D	unknown	no	yes	MD	Canne d Tuna (3)	M D	16	M D
Hobbs_1982_ FC_55z_25 <sup>33</sup>	1	UK	MD	MD	M D	unknown	no	yes	MD	Canne d	M D	260	M D

										Tuna (3)			
Hobbs_1982_ FC_55z_26 <sup>33</sup>	2	UK	MD	MD	M D	unknown	no	yes	MD	Canne d Tuna (3)	M D	640 0	M D
Hobbs_1982_ FC_55z_4 <sup>33</sup>	1	UK	MD	MD	M D	unknown	no	yes	MD	Canne d Mack erel (3)	M D	12. 5	M D
Hobbs_1982_ FC_55z_5 <sup>33</sup>	1	UK	MD	MD	M D	unknown	no	yes	MD	Canne d Pilcha rd (3)	M D	17	M D
Hobbs_1982_ FC_55z_7 <sup>33</sup>	1	UK	MD	MD	M D	unknown	no	yes	MD	Canne d Tuna (3)	M D	>10 000	M D
Hwang_1997 _CB_tre_1 <sup>34</sup>	3	MD	MD	MD	ye s	unknown	no	yes	eati ng hous	fried marlin fillet	M D	MD	M D

									e	(5)			
Iguchi_2008_ CB_qwe_135 <sup>3</sup> 5	3	Tokio	MD	MD	no	unknown	no	yes	resta uran t	sun- dried scomb roid fish (5)	1	160 0- 520 0	M D
Jantschitsch_2 011_EN_80u_ 1 <sup>36</sup>	2	Vienn a	MD	MD	no	unknown	yes	yes	resta uran t	tuna salad (3)	1	184 1	M D
Jiang_2009_C B_mbc_1 <sup>37</sup>	7 1	Kaohs iung City (Taiw an)	MD	MD	no	unknown	no	yes	scho ol	fried sailfis h fillet (5)	1	377 .4	M D
Kanki_2004_ EC_d8e_1 <sup>38</sup>	1	Osaka, Japan	no	no	no	unknown	no	yes	prob ably hom e	sardin e dried (5)	1	300 0	M D
Kelso_2009_ EC_80e_1 <sup>39</sup>	1	San Diego, Califo	MD	MD	ye s	unknown	no	yes	resta uran t	grilled tuna sandw	1	240 0	M D



		rnia								ich (5)			
Kim_1979_E C_71t_6 <sup>40</sup>	1	Honol ulu	MD	MD	ye s	unknown	no	yes	nurs ing hom e	mahi mahi (2)	1	630	M D
Kow- Tong_1987_E C_k&r_1 <sup>41</sup>	4 1	Taiwa n	MD	MD	M D	unknown	no	yes	emp loye e cafe teria	fried fish (white - tipped macke rel) (1)	1	100	M D
Leask_2004_ EC_kkk_1 <sup>42</sup>	9	South- easter n Sydne y	MD	MD	no	unknown	no	yes	mob ile cant een	fish curry (5)	1	200 9	M D
Lee_2012_EN _ii8_1 <sup>43</sup>	6 7	Kaohs iung City	no	no	no	unkown	no	yes	MD	fried fish ( <i>Tetra</i>	5	348	392 .07

		(Taiwan)								<i>pturus audax</i> ) (5)			
McLauchlin_2006_EN_u_13	4	London	MD	MD	MD	unknown	no	yes	restaurant	fresh tuna steaks (1)	2	3930	876.81
McLauchlin_2006_EN_u_23	16	London	MD	MD	MD	unknown	no	yes	restaurant	fresh tuna steaks (1)	1	5950	MD
McLauchlin_2006_EN_u_33	2	London	MD	MD	MD	unknown	no	yes	Braserie	fresh tuna steaks (1)	1	1700	MD
Missing_name_1988_FC_97y_1 <sup>44</sup>	2	Albuquerque	yes	MD	no	unknown	no	yes	MD	mahi mahi (2)	1	200	MD
Missing_name_2000_FC_y45_1 <sup>45</sup>	4	Pennsylvania	MD	MD	MD	unknown	no	yes	restaurant	tuna-spinach salad	1	>50	MD

										(5)			
Molinari_1989_EC_po1_1 <sup>4</sup> 6	MD	MD	MD	M D	unknown	no	yes	MD	smoked mackerel (5)	2	118	162	5 .63
Muller_1992_EC_3eq_1 <sup>47</sup>	Stellenbosch	MD	MD	M D	unknown	no	yes	restaurant home	Grilled or fried Cape yellowtail (Serio lalandii) (5)	1	250	M D	
Muller_1992_EC_3eq_2 <sup>47</sup>	Stellenbosch	MD	MD	M D	unknown	no	yes	restaurant home	Grilled or fried Cape yellowtail ( <i>Serio</i>	1	720	M D	

										<i>la</i> <i>laland</i> <i>ii) (5)</i>			
Muller_1992_ EC_3eq_3 <sup>47</sup>		Stelle nbosc h	MD	MD	M D	unknown	no	yes	resta uran t or hom e	Grille d or fried Cape yello wtail (Serio la laland ii) ( 5 )	1	162 5	M D
Nalinee Hongchumpo n Ouppapong_2 010_EC_knv_ 1 <sup>48</sup>	2 8	Samut Praka n Provin ce, Thaila nd	MD	MD	M D	unknown	no	yes	fact ory kitc hen and cafe teria	Fried ferme nted tuna (4)	1	446 .2	M D
Ohnuma_200	8	Yokoh	MD	MD	no	unknown	no	yes	resta	yello	1	310	M

1_EC_40e_1 <sup>4</sup> 9		ama							uran t	wfin tuna ( <i>Thun nus albac ares</i> ) sautè (1)		0	D
Predy_2003_ EN_88e_1 <sup>50</sup>	1	Edmo nton (Cana da)	MD	MD	ye s	unknown	yes	yes	coff ee shop	tuna fish salad (3)	1	350	M D
Sanchez- Guerrero_199 7_EN_8U_5 <sup>51</sup>	1 0	Lorca Murci a, Spain	MD	MD	M D	unknown	no	yes	groc er's shop	tuna ( <i>Thun nus thynn us</i> ) (5)	1	200 00	M D
Sanders_1987 _PC_qrs_1 <sup>52</sup>	2 3	New Jersey, U.S.A.	MD	MD	ye s	unknown	no	yes	mon aster y	yello wfin tuna broile	1	370 0	M D

										d and boiled (2)			
Schulze_1979 _PC_sss_1 <sup>53</sup>	1	Breme rhafen (Brem en harbo ur)	no	no	no	unknown	no	yes	MD	sardin es skin and viscer a free, in oil (3)	1	>50 0	M D
Sinn_2006_P C_nnn_1 <sup>54</sup>	2 0	Berlin er Stadt erzirk Charlo ttenbu rg- Wilme rsdorf (Berli n)	no	no	no	unknown	yes	yes	scho ol cant een	Butter fish- salmo n- potato patties (5)	1	565	M D

Stell_1997_E N_efb_1 <sup>55</sup>	1	London	MD	MD	no	unknown	no	yes	wine bar	cooked fresh tuna (1)	1	>20 00	M D
Stell_1997_E N_efb_2 <sup>55</sup>	7	London	MD	MD	no	unknown	yes	yes	MD	tuna mayo nnaise sandw iches (5)	1	>25 00	M D
Su_2000_PC_ aae_1 <sup>56</sup>		Pingtung County, southern Taiwan	MD	MD	MD	unknown	no	yes	public primary school	Marlin ( <i>M. Mazarina</i> ) fried fillets (5)	12	550 .78	26. 56
Taylor_1982_ PC_tta_1 <sup>57</sup>	6	Portsmouth, NH,	MD	MD	MD	unknown	no	no	aircraft carrier	swiss cheese (6)	3	187 0	M D

		USA							er cant een				
Tsai_2005_P C_bcd_1 <sup>58</sup>	3	Taipei Prefec ture, northe rn Taiwa n	MD	MD	M D	unknown	no	yes	groc ery	Canne d Mack erel (3)	3	153 9	98
Tsai_2007_P C_cde_1 <sup>59</sup>	3	Taina n Prefec ture, southe rn Taiwa n	MD	MD	M D	unknown	no	yes	groc ery	Dried milkfi sh ( <i>Chan os chano s</i> ) (5)	3	616	28
Tsai_2007_P C_efg_1 <sup>60</sup>	5 9	Pingtu ng, southe rn	MD	MD	M D	unknown	no	yes	resta uran t	Fried Billfis h meats	1	257 3	M D



		Taiwan								( <i>Makaira nigricans</i> ) (5)			
Tsai_2007_PC_efg_2 <sup>60</sup>	43	Taichung, central Taiwan	MD	MD	MD	unknown	no	yes	restaurant	Frozen Billfish meats ( <i>Xiphias gladius</i> ) (2)	5	2022.6	413.04
Valentini_1991_PC_ccc_1 <sup>61</sup>	3	Alta Val di Cecina (Italy)	MD	MD	MD	unknown	no	yes	canteen shop	canned tuna (3)	3	970	MD - 1050
Wu_1977_PC_aar_1 <sup>62</sup>	4	North eastern Taiwan	MD	MD	yes	unknown	no	yes	fast food stor	fried fish ( <i>Maka</i>	1	841.3	MD

		n							e	<i>ira</i> <i>spp.</i> ) (5)			
Wu_1977_PC _aar_2 <sup>62</sup>	4 8	North ern Taiwa n	MD	MD	M D	unknown	no	yes	emp loye e cafe teria	fried fish ( <i>Euth</i> <i>ynnus</i> <i>spp.</i> ) (5)	2	195 2	108 4.7 0
missing_name _1985_EN_3g 6_1 <sup>63</sup>		Engla nd and Wales	MD	MD	M D	unknown	no	yes	MD	fish (5)	1	<20 0	M D
missing_name _1985_EN_3g 6_10 <sup>63</sup>		Engla nd and Wales	MD	MD	M D	unknown	no	yes	MD	fish (5)	1	<20 0	M D
missing_name _1985_EN_3g 6_11 <sup>63</sup>		Engla nd and Wales	MD	MD	M D	unknown	no	yes	MD	fish (5)	1	<20 0	M D
missing_name _1985_EN_3g 6_12 <sup>63</sup>		Engla nd and Wales	MD	MD	M D	unknown	no	yes	MD	fish (5)	1	<20 0	M D
missing_name		Engla	MD	MD	M	unknown	no	yes	MD	fish	1	<20	M

_1985_EN_3g 6_13 <sup>63</sup>	nd and Wales			D					(5)		0	D
missing_name _1985_EN_3g 6_14 <sup>63</sup>	Engla nd and Wales	MD	MD	M D	unknown	no	yes	MD	fish (5)	1	<20 0	M D
missing_name _1985_EN_3g 6_15 <sup>63</sup>	Engla nd and Wales	MD	MD	M D	unknown	no	yes	MD	fish (5)	1	<20 0	M D
missing_name _1985_EN_3g 6_16 <sup>63</sup>	Engla nd and Wales	MD	MD	M D	unknown	no	yes	MD	fish (5)	1	<20 0	M D
missing_name _1985_EN_3g 6_17 <sup>63</sup>	Engla nd and Wales	MD	MD	M D	unknown	no	yes	MD	fish (5)	1	<20 0	M D
missing_name _1985_EN_3g 6_18 <sup>63</sup>	Engla nd and Wales	MD	MD	M D	unknown	no	yes	MD	fish (5)	1	<20 0	M D
missing_name _1985_EN_3g 6_19 <sup>63</sup>	Engla nd and Wales	MD	MD	M D	unknown	no	yes	MD	fish (5)	1	<20 0	M D
missing_name _1985_EN_3g	Engla nd and	MD	MD	M D	unknown	no	yes	MD	fish (5)	1	<20 0	M D

6_2 <sup>63</sup>	Wales												
missing_name _1985_EN_3g 6_20 <sup>63</sup>	England and Wales	MD	MD	M D	unknown	no	yes	MD	fish (5)	1	<20 0	M D	
missing_name _1985_EN_3g 6_21 <sup>63</sup>	England and Wales	MD	MD	M D	unknown	no	yes	MD	portug uese canned sardines (3)	1	800	M D	
missing_name _1985_EN_3g 6_22 <sup>63</sup>	England and Wales	MD	MD	M D	unknown	no	yes	MD	canned tuna from Peru (3)	1	>10 00	M D	
missing_name _1985_EN_3g 6_23 <sup>63</sup>	England and Wales	MD	MD	M D	unknown	no	yes	MD	canned tuna from Peru (5)	1	>10 00	M D	
missing_name _1985_EN_3g	England and	MD	MD	M D	unknown	no	yes	MD	smoked	1	>10 00	M D	

6_24 <sup>63</sup>	Wales								macke rel (5)			
missing_name _1985_EN_3g 6_3 <sup>63</sup>	Engla nd and Wales	MD	MD	M D	unknown	no	yes	MD	fish (5)	1	<20 0	M D
missing_name _1985_EN_3g 6_4 <sup>63</sup>	Engla nd and Wales	MD	MD	M D	unknown	no	yes	MD	fish (5)	1	<20 0	M D
missing_name _1985_EN_3g 6_5 <sup>63</sup>	Engla nd and Wales	MD	MD	M D	unknown	no	yes	MD	fish (5)	1	<20 0	M D
missing_name _1985_EN_3g 6_6 <sup>63</sup>	Engla nd and Wales	MD	MD	M D	unknown	no	yes	MD	fish (5)	1	<20 0	M D
missing_name _1985_EN_3g 6_7 <sup>63</sup>	Engla nd and Wales	MD	MD	M D	unknown	no	yes	MD	fish (5)	1	<20 0	M D
missing_name _1985_EN_3g 6_8 <sup>63</sup>	Engla nd and Wales	MD	MD	M D	unknown	no	yes	MD	fish (5)	1	<20 0	M D
missing_name _1985_EN_3g	Engla nd and	MD	MD	M D	unknown	no	yes	MD	fish (5)	1	<20 0	M D

6_9 <sup>63</sup>		Wales											
missing_name _2007_CB_dh i_1 <sup>64</sup>	6	Louisi ana	MD	MD	M D	unknown	no	yes	com pan y cafe teria	tuna steaks (5)	1	<20 0	M D

Table 2: Quality items values and overall quality score

Poisoning sample id / reference	Health status	Age	Sex	Source of medical diagnosis	Source of food	Histamine measurement method	Variability estimate	Quality score
Becker_2001_ET_aai_1 <sup>2</sup>	0	1	0	0	1	0	0	3
Becker_2001_ET_aai_3 <sup>2</sup>	0	1	0	0	1	0	0	3
Bedry_2000_EN_mi1_1 <sup>12</sup>	0	1	1	0	0	1	0	4
Bremer_2003_ET_cca_1 <sup>13</sup>	0	0	0	0	0	0	0	1
Bremer_2003_ET_cca_2 <sup>13</sup>	0	0	0	0	0	0	0	1
Bremer_2003_ET_cca_3 <sup>13</sup>	0	0	0	0	0	0	0	1
Chen_2008_ET_hhh_1 <sup>14</sup>	0	0	0	0	0	1	1	3
Chen_2010_CB_zxc_1 <sup>15</sup>	0	0	0	0	1	1	1	3
Chen_2010_ET_fff_1 <sup>16</sup>	0	0	0	0	1	1	0	3
Chen_2011_ET_eee_1 <sup>17</sup>	0	0	0	0	1	1	0	3
Chianèa_1998_ET_nnn_1 <sup>18</sup>	0	0	0	0	1	1	0	2
D'Aloia_2011_EN_5jb_1 <sup>19</sup>	1	1	1	0	0	0	0	4
Demoncheaux_2012_EN_99w _1 <sup>20</sup>	1	1	1	0	1	1	1	7
Doeglas_1967_ET_vvv_1 <sup>21</sup>	0	1	1	1	0	0	0	4

Eckstein_1999_EN_8f6_1 <sup>22</sup>	0	0	0	0	1	1	0	3
Emborg_2005_CB_kju_1 <sup>23</sup>	0	0	0	0	1	1	0	2
Emborg_2006_CB_kjs_1 <sup>24</sup>	0	0	0	0	1	1	1	3
Emborg_2006_CB_kjs_2 <sup>24</sup>	0	0	0	0	1	1	1	3
Emborg_2006_CB_kjs_3 <sup>24</sup>	0	0	0	0	1	1	1	3
Feldman_2005_EN_555_1 <sup>25</sup>	0	1	0	1	1	1	0	5
Fernandez_2001_CB_3jd_3 <sup>26</sup>	0	1	1	1	1	1	0	5
Foo_1975_CB_h67_1 <sup>27</sup>	0	0	0	0	0	1	0	1
Foo_1975_CB_jso_1 <sup>28</sup>	0	0	0	0	0	0	0	0
Foo_1977_CB_8sk_1 <sup>29</sup>	0	0	0	0	0	1	0	1
Gellert_1992_82c_1 <sup>30</sup>	0	1	0	0	1	0	0	3
Guly_2006_EN79j_1 <sup>31</sup>	0	0	0	1	1	0	0	3
Hall_2003_CB_h8s_1 <sup>32</sup>	0	1	1	1	1	0	0	4
Hobbs_1982_FC_55z_14 <sup>33</sup>	0	0	0	0	0	1	0	1
Hobbs_1982_FC_55z_15 <sup>33</sup>	0	0	0	0	0	1	0	1
Hobbs_1982_FC_55z_16 <sup>33</sup>	0	0	0	0	0	1	0	1
Hobbs_1982_FC_55z_18 <sup>33</sup>	0	0	0	0	0	1	0	1
Hobbs_1982_FC_55z_20 <sup>33</sup>	0	0	0	0	0	1	0	1
Hobbs_1982_FC_55z_21 <sup>33</sup>	0	0	0	0	0	1	0	1
Hobbs_1982_FC_55z_22 <sup>33</sup>	0	0	0	0	0	1	0	1
Hobbs_1982_FC_55z_23 <sup>33</sup>	0	0	0	0	0	1	0	1



Hobbs_1982_FC_55z_24 <sup>33</sup>	0	0	0	0	0	1	0	1
Hobbs_1982_FC_55z_25 <sup>33</sup>	0	0	0	0	0	1	0	1
Hobbs_1982_FC_55z_26 <sup>33</sup>	0	0	0	0	0	1	0	1
Hobbs_1982_FC_55z_4 <sup>33</sup>	0	0	0	0	0	1	0	1
Hobbs_1982_FC_55z_5 <sup>33</sup>	0	0	0	0	0	1	0	1
Hobbs_1982_FC_55z_7 <sup>33</sup>	0	0	0	0	0	1	0	1
Hwang_1997_CB_tre_1 <sup>34</sup>	0	0	1	0	1	1	1	4
Iguchi_2008_CB_qwe_135 <sup>35</sup>	0	0	0	1	1	0	0	2
Jantschitsch_2011_EN_80u_1 <sup>3</sup> 6	0	1	1	1	1	0	0	5
Jiang_2009_CB_mbc_1 <sup>37</sup>	0	0	1	0	1	0	0	2
Kanki_2004_EC_d8e_1 <sup>38</sup>	0	0	0	0	1	1	0	3
Kelso_2009_EC_80e_1 <sup>39</sup>	0	1	1	0	1	0	0	4
Kim_1979_EC_71t_6 <sup>40</sup>	0	1	1	0	1	0	0	4
Kow-Tong_1987_EC_k8r_1 <sup>41</sup>	0	0	0	0	1	1	0	3
Leask_2004_EC_kkk_1 <sup>42</sup>	0	1	1	0	1	0	0	4
Lee_2012_EN_ii8_1 <sup>43</sup>	0	0	0	0	0	0	0	1
McLauchlin_2006_EN_u_1 <sup>3</sup>	0	0	0	0	1	0	0	2
McLauchlin_2006_EN_u_2 <sup>3</sup>	0	0	0	0	1	0	0	2
McLauchlin_2006_EN_u_3 <sup>3</sup>	0	0	0	0	1	0	0	2
Missing_name_1988_FC_97y	0	0	1	1	0	0	0	2

_1 <sup>44</sup>								
Missing_name_2000_FC_y45 _1 <sup>45</sup>	0	0	0	1	1	1	0	3
Molinari_1989_EC_po1_1 <sup>46</sup>	0	0	0	0	0	1	0	1
Muller_1992_EC_3eq_1 <sup>47</sup>	0	0	0	1	1	1	0	3
Muller_1992_EC_3eq_2 <sup>47</sup>	0	0	0	1	1	1	0	3
Muller_1992_EC_3eq_3 <sup>47</sup>	0	0	0	1	1	1	0	3
Nalinee Hongchumpon Ouppapong_2010_EC_knv_1 <sup>4</sup> 8	0	1	1	0	1	0	0	4
Ohnuma_2001_EC_40e_1 <sup>49</sup>	1	1	1	1	1	1	0	7
Predy_2003_EN_88e_1 <sup>50</sup>	1	1	1	1	1	0	0	6
Sanchez- Guerrero_1997_EN_8U_5 <sup>51</sup>	0	0	0	0	1	0	0	2
Sanders_1987_PC_qrs_1 <sup>52</sup>	0	0	1	0	1	0	0	3
Schulze_1979_PC_sss_1 <sup>53</sup>	0	0	0	0	0	1	0	2
Sinn_2006_PC_nnn_1 <sup>54</sup>	0	0	0	0	1	0	0	2
Stell_1997_EN_efb_1 <sup>55</sup>	0	1	1	0	1	0	0	4
Stell_1997_EN_efb_2 <sup>55</sup>	0	0	0	0	0	0	0	1
Su_2000_PC_aae_1 <sup>56</sup>	0	0	0	0	1	1	1	4
Taylor_1982_PC_tta_1 <sup>57</sup>	0	0	0	0	1	1	0	3
Tsai_2005_PC_bcd_1 <sup>58</sup>	0	0	0	0	1	1	1	4

Tsai_2007_PC_cde_1 <sup>59</sup>	0	0	0	0	1	1	1	4
Tsai_2007_PC_efg_1 <sup>60</sup>	0	0	0	0	1	1	0	3
Tsai_2007_PC_efg_2 <sup>60</sup>	0	0	0	0	1	1	0	3
Valentini_1991_PC_ccc_1 <sup>61</sup>	0	0	0	0	1	1	0	3
Wu_1977_PC_aar_1 <sup>62</sup>	0	0	1	1	1	1	0	5
Wu_1977_PC_aar_2 <sup>62</sup>	0	0	0	0	1	1	0	3
missing_name_1985_EN_3g6 _1 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _10 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _11 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _12 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _13 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _14 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _15 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _16 <sup>63</sup>	0	0	0	0	0	0	0	0

missing_name_1985_EN_3g6 _17 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _18 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _19 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _2 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _20 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _21 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _22 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _23 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _24 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _3 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _4 <sup>63</sup>	0	0	0	0	0	0	0	0

missing_name_1985_EN_3g6 _5 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _6 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _7 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _8 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_1985_EN_3g6 _9 <sup>63</sup>	0	0	0	0	0	0	0	0
missing_name_2007_CB_dhi_ 1 <sup>64</sup>	0	0	0	0	1	0	0	1

Table n. 3: Sub-group analysis: number of poisoning samples by food category.

Food category id	Food category	Number of poisoning samples	Percent
1	fresh	9	8.74
2	frozen	7	6.80
3	canned	22	21.36
4	fermented	1	0.97
5	other seafood	62	60.19
6	cheese	2	1.94
7	other foods	0	0.00
Totals	all categories	103	100.00

Table n.4 Sensitivity analysis: number of poisoning samples by quality score.

Quality score	Number of poisoning samples	Percent
0	25	24.27
1	23	22.33
2	11	10.68
3	24	23.30
4	13	12.62
5	4	3.88
6	1	0.97
7	2	1.94
Totals	103	100.00

Excerpt from the systematic review protocol

*"The search strategies will be focused to detect all reports of histamine poisonings from foods that meet inclusion criteria, so will be optimized accordingly to this purpose. A main form of search strategy will be designed and could be modified to meet settings of databases consulted. The strategy adopted and database on which was performed will be recorded. Main key-words will be: "histamine", "scombroid syndrome", "histamine poisoning", food, seafood, "meat products", fish, cheese, beer, wine, "biogenic amines". To improve the effectiveness of key-words in the search strategy a preliminary thesaurus study will be performed.*

*Replicate reports of the same data will be detected and only one report will be considered for inclusion in this review; reasons for the choice will be given and recorded. "*

Main form of search strategyForeword

- if not specified, the mentioned keywords are meant to be searched both as "free search" keywords and "topic search" keywords:
- Key-words are "case-sensitive" written (lower case and upper case must be maintained when performing search)
- Logical operators and symbols are written in **bold** character

Legend

Suffix **.to** means: keyword will be searched only as "topic search" keyword

Suffix **.fr** means: keyword will be searched only as "free search"

<Keywords inside round brackets> means: phrase search

**x/x** (i.e. letter, slash, letter) means: degenerate letter, an appropriate "jolly character" has to be assigned to it.

Search Id	Key-words syntax	Notes
#1	"biogenic amines". <b>to</b>	-
#2	"scombroid syndrome"	-
#3	"histamine poisoning"	-
#4	<b>#1 or #2 or #3</b>	-
#5	"food poisoning"	-
#6	<b>#4 and #5</b>	-
#7	food	-
#8	fish	-
#9	<b>#7 or #8</b>	-
#10	<b>#6 and #9</b>	-

Figure 1



Figure 2. Selection of reports for systematic review




Figure 2.




Figure 3.




Figure 4.




Figure 5.




Figure 6.