

## What is changing in indications and treatment of hepatic hemangiomas. A review

Adriana Toro,\* Ahmed-Emad Mahfouz,<sup>†</sup> Annalisa Ardiri,<sup>‡</sup> Michele Malaguarnera,<sup>§</sup> Giulia Malaguarnera,<sup>||</sup> Francesco Loria,<sup>¶</sup> Gaetano Bertino,<sup>‡</sup> Isidoro Di Carlo<sup>\*\*</sup>,<sup>†</sup>

\* Department of Surgery, Taormina Hospital, Messina, Italy. <sup>†</sup> Department of Radiology, Hamad General Hospital, Doha Qatar.

<sup>‡</sup> Hepatology Unit, Department of Medical and Pediatric Science, University of Catania, Italy.

<sup>§</sup> International PhD Program in Neuropharmacology, University of Catania, Italy.

<sup>||</sup> University of Catania; Research Center "The Great Senescence", Cannizzaro Hospital, Italy.

<sup>¶</sup> Department of Radiology PO Palmi, ASP 5 RC, Italy.

<sup>\*\*</sup> Department of Surgical Sciences, Organ Transplantation and Advanced Technologies, University of Catania, Catania, Italy.

<sup>††</sup> Department of Surgery, Hamad Medical Hospital, Doha, Qatar.

### ABSTRACT

Hepatic cavernous hemangioma accounts for 73% of all benign liver tumors with a frequency of 0.4-7.3% at autopsy and is the second most common tumor seen in the liver after metastases. Patients affected by hemangioma usually have their tumor diagnosed by ultrasound abdominal examination for a not well defined pain, but pain persist after treatment of the hemangioma. The causes of pain can be various gastrointestinal pathologies including cholelithiasis and peptic ulcer disease. The malignant transformation is practically inexistent. Different imaging modalities are used to diagnosis liver hemangioma including ultrasonography, computed tomography (CT), magnetic resonance (MR) imaging, and less frequently scintigraphy, positron-emission tomography combined with CT (PET/CT) and angiography. Imaging-guided biopsy of hemangioma is usually not resorted to except in extremely atypical cases. The right indications for surgery remain rupture, intratumoral bleeding, Kasabach-Merritt syndrome and organ or vessels compression (gastric outlet obstruction, Budd-Chiari syndrome, etc.) represents the valid indication for surgery and at the same time they are all complications of the tumor itself. The size of the tumor do not represent a valid indication for treatment. Liver hemangiomas, when indication exist, have to be treated firstly by surgery (hepatic resection or enucleation, open, laproscopic or robotic), but in the recent years other therapies like liver transplantation, radiofrequency ablation, radiotherapy, trans-arterial embolization, and chemotherapy have been applied.

**Key words.** Hepatic surgery. Hemangiomas of the liver. Radiofrequency ablation. Benign tumors of the liver. Hepatic transplantation.

### INTRODUCTION

Hepatic cavernous hemangioma accounts for 73% of all benign liver tumors<sup>1</sup> with a frequency of 0.4-7.3%<sup>2</sup> at autopsy and is the second most common tumor seen in the liver after metastases.<sup>3</sup> These lesions consist of large vascular spaces lined by a single layer of endothelial cells and several authors agree that this vascular tumour is a benign,

congenital haematoma or tissue malformation that grows slowly from birth. The pathogenesis of this tumor is not still completely understood, but abnormal vasculogenesis and angiogenesis have been speculated to be involved.<sup>4</sup> The cavernous hemangioma endothelial cell differ from sinusoidal endothelial cell both derived from the human liver. Particularly the first cells exhibiting more activated angiogenesis capacity and forming abnormal capillary-like structures in vitro. The grown pattern of the cavernous hemangioma is still not understood. Ectasia rather than hypertrophy or hyperplasia usually is recorded.

Some hemangiomas have the receptor for the estrogens and they grown during the puberty, pregnancy (following ovarian stimulation therapy with clomiphene citrate and human chorionic gonadotrophin), or oral contapeptive use, androgen

Correspondence and reprint request: Isidoro Di Carlo, MD, PhD, FACS.  
Department of Surgical Sciences, Organ Transplantation, and Advanced Technologies, University of Catania, Cannizzaro Hospital, Via Messina 829, 95126 Catania, Italy  
Tel.: +39 095 7264863. Fax: +39 095 7263020.  
E-mail: idicarlo@unicat.it

*Manuscript received: May 12, 2014.*

*Manuscript accepted: May 21, 2014.*

or/and steroid administration.<sup>5</sup> These tumors are most commonly found in women, with a female:male ratio of up to 5:1, emphasizing the importance of excess of female sex hormones in these tumours.<sup>6,7</sup> In fact enlargement and rupture of these tumors have been reported in pregnancy and in menopausal women submitted to ormonal treatment.<sup>8</sup>

The size increase have been suggested from many studies, mainly case report.<sup>9-20</sup> The biggest study reported in literature compare 94 female patients that receiving hormonal treatment comparing with a group of patient untreated. In the treated group the increase in size was 22.7% compared with untreated patients that was only 9.7%.<sup>19</sup> But natural hystory is not completely defined; Moser et al. reported a large family of Italian origin in which three female patients in three successive generations had large symptomatic hepatic haemangiomas without any risk factor.<sup>21</sup>

In relation to their dimension the hemangiomas are defined as normal or giant. Different mesures are present in the literature and the majority of Authors call giant the hemangioma more than 4 cm.<sup>22-26</sup>



**Figure 1.** Large hemangioma of the left lobe of the liver that compresses the stomach.

Some Authors define the hemangiomas giant after 5 cm<sup>22,27</sup> and very small number of Authors define giant hemangiomas more than 10 cm.<sup>28</sup> The definition of giant hemangioma should be changed to indicate a minimum size of 10 cm. It is easier to accept that a 10 cm hemangioma can cause symptoms than 4 cm tumor. For example, a 10 cm tumor located in the left lobe of the liver compresses the stomach and premature satiety after a meal is a more credible and acceptable indication for surgery (Figure 1). On the opposite it is very difficult to imagine a giant hemangioma a tumor of 4 cm, a size less than a liver segment, that can be responsible of any symptoms in any location of the liver.<sup>29</sup>

### UPDATE IN DIAGNOSTIC METHODS


The importance of hemangioma, from the point of view of medical imaging comes from its relatively high incidence in comparison to other focal liver lesions, its occurrence as an incidental finding, and the need to differentiate it from other more serious focal liver lesions. The latter consideration is particularly important in two types of patients: patients with primary malignant neoplasm and patients with liver cirrhosis. In the patients with primary liver neoplasm, it is important to differentiate hemangioma from liver metastasis. In the patients with cirrhosis, it is important to differentiate it from hepatocellular carcinoma.

Different imaging modalities have been used in diagnosis of liver hemangioma including ultrasonography, computed tomography (CT), magnetic resonance (MR) imaging, and less frequently scintigraphy and positron-emission tomography combined with CT (PET/CT) and angiography. Imaging-guided biopsy of hemangioma is usually not resorted to except in extremely atypical cases. Specificity and sensitivity of all these procedures are reported in table 1.

**Table 1.** Sensitivity and specificity of the diagnostic methods.

Diagnostic method	Sensitivity (%)	Specificity (%)
Ultrasonography	96.9	60.3
Computed tomography	98.3	55.0
Magnetic resonance imaging	100	85.7
Tc-99m RBC blood pool scintigraphy	75	100
Angiography	na	na
PET/TC	na	na

na: not available.



**Figure 2.** Ultrasonography of the liver shows hemangioma with typical well-definition, homogenous hyperechogenicity, and close relation to a hepatic vein.

### Ultrasonography

On ultrasonography, hemangioma appears well defined, occasionally lobulated, homogeneously hyperechoic with posterior acoustic enhancement, and frequently located close to a portal or a hepatic vein (Figure 2). In most of the cases, hemangiomas show no change of size or morphological features on follow-up ultrasonography. Most of hemangiomas have no mass effect on the liver surface or adjacent hepatic vessels. However, large hemangiomas may bulge beyond the hepatic contour and exert mass effect on adjacent structures. Occasionally, hemangioma has a slightly hypoechoic center in comparison to its peripheral rim.

Acoustic radiation force impulse (ARFI) elastography is a new adjunct to ultrasonographic examination. Some preliminary studies have shown that stiffness was higher in malignant liver lesions compared to hemangiomas and focal nodular hyperplasia with some overlap,<sup>30</sup> while no significant difference was found between hemangioma and solid lesions on other studies.<sup>31</sup>

Pulse-Inversion Harmonic contrast-enhanced ultrasonography has also been tried for differentiation of hemangiomas and solid hepatic lesions such as hepatocellular carcinoma and metastasis. Most



**Figure 3.** CT of liver hemangioma shows the typical peripheral nodular enhancement of the hemangioma.

hemangiomas show peripheral globular or rim-like enhancement with progressive centripetal fill-in with no fast washout, while hepatocellular carcinoma and metastasis may show arterial enhancement followed by relatively fast washout.<sup>32,33</sup>

Color-Doppler Ultrasonography shows no flow within most of the hemangiomas. In some hemangiomas, however, there may be an intralesional arterio-portal shunt. In these cases, color Doppler ultrasonography may show flow inside the hemangioma and retrograde flow in the small portal branches around the hemangioma.<sup>34</sup>

### Computed tomography (CT)

On CT, hemangiomas show peripheral nodular enhancement, which progresses from the periphery of the lesion to its center until the whole lesion shows homogeneous prolonged enhancement, matching the enhancement of veins (Figure 3). Occasionally the hemangioma shows early enhancement on the arterial-phase CT followed by early filling-in of the lesion. This hyperdynamic behavior of some of the hemangiomas is attributed to intralesional arterioportal shunts. Small hemangiomas with high flow may show complete fill-in on the early phase of the contrast-enhanced CT.<sup>35</sup>

### Magnetic resonance (MR) imaging

On MR imaging, hemangioma appears well defined, homogeneously hyperintense on T2-weighted

images, homogeneously hypointense on T1-weighted images, with peripheral nodular enhancement that progresses to complete and prolonged fill-in on the dynamic gadolinium-enhanced images.<sup>36</sup> Hyperintensity of the hemangioma on T2-weighted images has been one of the important features differentiating it from solid neoplastic liver lesions. However, some of these lesions may show also similar hyperintensity on T2-weighted images. To differentiate the hyperintensity of hemangioma on T2-weighted images from these solid liver neoplasms, a useful method is to increase the TE of the T2-weighted images in steps. With such an increase in TE, the signal intensity of solid liver lesions would decrease, while that of hemangioma would remain high.<sup>37</sup> Occasionally hemangioma may show central nonenhancing component even on the delayed images of the dynamic gadolinium-enhanced images. This may be attributed to thrombosis, hyalinization, scarring, or cystic changes at the center of the hemangioma.<sup>38</sup> Hemangioma may appear similar to simple hepatic cyst on T2-weighted images as both lesions will be extremely well defined and extremely hyperintense on T2-weighted images. However, cysts tend to have lower signal intensity on T1-weighted images, lower signal intensity on proton-density images, no restriction of diffusion on the diffusion-weighted images, and no enhancement on the dynamic gadolinium-enhanced images. On the diffusion-weighted images, hemangioma tends to have less diffusion restriction than metastases and more diffusion restriction than cysts. Recently, tensor diffusion imaging has also been performed for focal liver lesions and has shown that fractional anisotropy (FA) of hemangioma is higher than that of cysts and lower but very close to that of metastases.<sup>39</sup>

#### Other imaging modalities

Tc-99m RBC blood pool scintigraphy shows high uptake of the radiotracer in hepatic hemangioma.<sup>40</sup> Angiography is seldom used for diagnosis of hemangioma. However, angiographic appearance of hemangioma is recognized from old publications performed before the era of modern sectional imaging, from angiographic studies performed for malignant liver neoplasm with incidentally co-existing hemangioma, and from angiography performed for embolization of giant hemangiomas. On angiography, hemangioma has a characteristic snowy-tree or cotton-wool appearance.<sup>41</sup> PET/CT shows no uptake of FDG by hemangioma of the liver, and may be

used to differentiate atypical hemangioma from metastasis in patients with primary malignant neoplasm.<sup>42</sup> Hemangioma may show uptake and accumulation of lipiodol for a long period after transcatheter arterial chemoembolization of hepatocellular carcinoma with incidentally co-existing hemangioma.<sup>43</sup>


### SPECIAL SITUATIONS OF LIVER HEMANGIOMAS ON IMAGING

#### Hemangioma in the fatty liver

Hemangioma may appear hypoechoic on ultrasonography if it occurs in fatty liver.<sup>44</sup> In this case, the diagnosis of hemangioma can be made by contrast-enhanced CT if the lesion is large enough or MR imaging if the lesion is small. Both CT and MR imaging can show the characteristic imaging features of the hemangioma and the typical appearance of the fatty liver. CT shows the fatty liver as low attenuation of the liver parenchyma on un-enhanced CT. In-phase and opposed-phase T1-weighted MR images are particularly useful in diagnosis of fatty liver.

#### Giant hemangioma

Giant hemangiomas, larger than 10 cm in diameter are clinically important because they may cause symptoms and may be complicated by hemorrhage, rupture, or intralesional hemolysis. Giant hemangiomas usually have central nonenhancing components and may have central hemorrhagic changes (Figure 4). Surgery or interventional radiological



**Figure 4.** Giant hemangioma of the left lobe of the liver. T2-weighted MR image shows hypointense hemorrhagic changes within the hyperintense hemangioma.

procedures are indicated for the treatment of giant hemangiomas. Ruptured hemangioma needs urgent surgical intervention.<sup>28,45</sup>

#### **Non-enhancing component within the hemangioma**

Persistent non-enhancing component within the hemangioma may be due to thrombosis, sclerosis, scarring, hyalinization, or cystic clefts within the hemangioma. Diagnosis of hemangioma remains to be easy if the hemangioma is large enough to show the characteristic features at the periphery of the lesion. If, however, the lesion is small with non-enhancing central component, it may be mistaken for hypovascular metastasis. In this case, diagnosis can be made by imaging follow-up if no primary malignant neoplasm is present and by PET/CT if there is a primary malignant neoplasm.

#### **High-flow hemangioma**

Hemangioma may have intra-lesional arterio-portal or arterio-venous shunts. This type of hemangioma may show total enhancement on the early phases of contrast-enhanced CT or dynamic gadolinium-enhanced images. The early enhancement of this hemangioma may mimic that of focal nodular hyperplasia. Diagnosis of hemangioma can be made in this case by the prolonged enhancement and the high signal intensity of the lesion on T2-weighted images. On color Doppler imaging high-flow hemangioma may show flow signal within the lesion and retrograde flow in the surrounding venous structures.<sup>34</sup> Hyperperfusion of the liver parenchyma around the high-flow hemangioma may be demonstrated on the arterial phase of contrast-enhanced CT and gadolinium-enhanced MR imaging.

#### **Very small hemangioma**

Hemangioma, less than 5 mm in diameter may be difficult to characterize on imaging because it is too small to show the typical peripheral nodular enhancement on contrast-enhanced CT or dynamic gadolinium-enhanced MRI. Such hemangiomas usually show total enhancement from the start. However, the rest of the imaging features, particularly the high signal intensity on T2-weighted images and the prolonged enhancement without significant washout would still be useful in characterization of hemangiomas.

#### **Hemangioma in infants and children**

Hemangioma in infants and children are more liable to cause hemolysis and consumption coagulopathy (Kasabach-Merritt syndrome) than hemangiomas in adults. On imaging follow-up, hemangioma in infants and children may show rapid resolution.<sup>46</sup>

#### **INDICATIONS TO TREATMENT**

Patients affected by hemangioma usually have their tumor diagnosed by ultrasound abdominal examination for a not well defined pain. A study reported in the literature 58% of the patients had abdominal pain at baseline, and in 50% of cases this pain was the reason for referral which led to the diagnosis of liver hemangioma.<sup>47</sup> However, in only 12.6% of the cases could the pain be attributed to the hemangioma; in the other patients, other gastro intestinal diseases, especially inflammatory bowel diseases and peptic ulcer disease, were also present.<sup>47</sup> In many studies reported in the literature pain persist after treatment of the hemangioma. In a study by Farges and colleagues,<sup>48</sup> pain disappeared in 54% of patients after treatment of associated disorders, and in 4 out of 11 patients, pain persisted even after tumor resection. In this series, pain also diminished in many patients even in those with hepatic hemangiomas, pain is thought to be present as a result of infarct and necrosis of the tumor<sup>49</sup> or the result of the tumor pressing on the liver capsule or adjacent organs.<sup>50</sup> The latter is especially important in large tumors and left-lobe lesions.<sup>48,51</sup> It is important to refer that many patients having pain with small lesion continue to have pain also if the tumor is not growing in size. Many of this symptoms are caused from other diseases especially inflammatory bowel disease at early stage. Furthermore there is no correlation in the majority of the study between the tumor size increase and the pain. But despite what, the abdominal pain remain the indication for surgical treatment, not supported by clinical evidence in a percentage between 48 and 86%.<sup>25,48,52</sup> The causes of pain should be critically analyzed in case of hepatic hemangioma because more than 50% of the patients with abdominal pain and liver hemangioma have been found to have various gastrointestinal pathologies including cholelithiasis and peptic ulcer disease.<sup>52</sup> Discomfort and anxiety have been reported as indication for surgery in case of hemangioma.<sup>53</sup> These two symptoms should never been considered as indication for surgery especially if tumor is small in size and cannot responsible for

**Table 2.** Results of the treatments.

Treatment	Mortality	Morbidity	Recurrence
Surgical Procedure	0-3%	10-27%	0%
Radiofrequency ablation	0%	33%	7,3%
Monoclonal antibody	na	na	na
Radiotherapy	na	na	na
TAE	0%	54%	0%
Chemotherapy	na	na	na
Liver transplantation	na	na	na

na: not available.

any problem.<sup>29</sup> The malignant transformation is practically inexistent and the patients have to be reassured or addressed to the psychiatrist. Patients operated for hemangioma for stress and anxiety due to the awareness of having a benign tumor, continued to have the same symptoms, even after surgery.<sup>51</sup> These patients operated with a wrong indications do not improve their quality of life and they are exposed at all the risks of the surgical operation.

The right indications for surgery remain strictly related to the tumor complications. In fact rupture, intratumoral bleeding, Kasabach-Merritt syndrome and organ or vessels compression (gastric outlet obstruction, Budd-Chiari syndrome, etc.) represents the valid indication for surgery and at the same time they are all complications of the tumor itself. The size of the tumor do not represent a valid indication for treatment.

The Kasabach Merritt syndrome (also known as hemangioma thrombocytopenia syndrome)<sup>54</sup> is a rare but life-threatening disease and represents a valid indication for immediate surgery in patients with hepatic hemangioma. This syndrome has been described in children in association with cutaneous hemangioma. The main characteristics of Kasabach-Merritt syndrome are enlarging hemangioma associated with thrombocytopenia and microangiopathic hemolytic anemia with acute and massive or chronic and low-grade consumptive coagulopathy. Clotting and fibrinolysis within the hemangioma represent the first event of the Kasabach-Merritt syndrome. The patients show in the laboratory test thrombocytopenia, anemia, hypofibrinogenemia but over all elevated fibrin degraded products (FDP). In patients with enlarged hemangioma with suspicion of Kasabach-Merritt syndrome can be monitored every 3 months in case of elevation of 10% of the previous basic value of the FDP this result can represent an indication to surgery.

Another issue concerning the indication to a treatment for hemangioma is still represented by the

location of the hemangiomas. In fact some tumors can be located close to the major vascular structures and in this way during their enlargement they can compress the major veins causing symptoms like in the case of Budd-Chiari syndrome. This is the reason why some authors have suggested an early prophylactic treatment even in case of asymptomatic patients in case of tumor located close to the confluence of hepatic veins that growing can provoke these problems.<sup>55</sup>

Traditionally, surgery had been advocated for most hemangiomas due to a concern for possible rupture.<sup>22</sup> However, over the last 25 years this paradigm has been challenged due to the relatively limited number of cases of rupture reported in the literature.<sup>56-58</sup> To date, less than 50 cases of spontaneous rupture have been reported,<sup>59</sup> while only 5 cases of traumatic rupture are known.<sup>57,60-63</sup> Hemangioma rupture have never been reported in parachuting, this confirms that the presence of hemangioma is not a risk for the spontaneous or traumatic rupture. All the methods to treat an hemangioma are reported in the next paragraph and the cumulative results for each treatment are reported in table 2.

## TREATMENT

### Surgical treatment

Liver hemangiomas, if indication exist, have to be treated firstly by surgery (hepatic resection or enucleation, open, laproscopic or robotic), but in the recent years other therapies like liver transplantation, radiofrequency ablation, radiotherapy, trans-arterial embolization, and chemotherapy have been applied.

First hemangioma have been treated by hepatic resection in 1898 by Hermann Pfannestil.<sup>64</sup> Since then the hepatic resection have been the most practiced therapy for this tumor. In 1988 almost a century after the first hepatic resection for hemangioma,

the enucleation have been reported in the literature.<sup>65</sup> Since this second technique have been used more and more. Also if do not exist prospective randomized study comparing the two technique the majority of reports present in the literature are in favour of enucleation. The enucleation is safe, minimize the blood loss, and have a less rate of complications in relation to the hepatic resection, decreasing the risk of biliary fistula and preserving the maximum amount of normal liver parenchyma.<sup>25,66-69</sup> However, the study populations in these reports were small, and the impact of the hemangioma location was not particularly analyzed. The enucleation is a useful technique that can be performed if the hemangioma has an external plane on the liver surface. The space between the liver and the hemangioma is an avascular plane consisting of liver tissue compressed by the growing hemangioma. Only a few blood vessels traverse the capsule, and these are easily controlled. The tumor can be shucked without blood loss by opening this layer between the capsule of the hemangioma and the liver. The amount of normal functioning liver parenchyma removed is minimal. If the hemangioma is deeply located in the liver parenchyma, or do not present a free surface from the glisson capsule, or occupe all entire lobe in this case the hepatic resection remain the method of choice.<sup>70</sup> The resection remain the choice also in case of multiple hemangiomas of the liver. The largest series concerning enucleation is reported in the literature and is related to 172 patients. In this report also the patients with deep hemangioma located in the liver have been treated by enucleation but with massive blood transfusions and their related risk increased and the procedure that became time consuming.<sup>70</sup>

Neonates with an abdominal mass and unidentified congestive heart failure are usually the patients and the symptoms of the diffuse hepatic hemangiomas, a rare condition characterized by diffuse replacement of liver parenchyma with hemangiomas lesions.<sup>71</sup> But definition of hemangiomas include also hemangiomas of the skin and involvement of at least two visceral organs.<sup>72</sup> Hepatic isolated hemangiomas of the liver has been reported as a very rare disease.<sup>73</sup> Although several cases of long-term adult survival of diffuse neonatal hemangiomas have been reported,<sup>74,75</sup> the etiology and natural history of diffuse hepatic hemangiomas remains unclear. The histological characteristics of hemangiomas includes the presence of large vascular channels in both the normal-appearing hepatic parenchyma and the cavernous tumor region.<sup>76</sup>

Recognition of the association of hepatic hemangiomas in patients with giant cavernous hemangiomas (GCHs) is important because management of hemangioma may depend on the presence and extent of hemangiomas. In case of large hamangioma surrounded by hemangiomas, the enucleation cannot be possible. In fact in this case the sheath of compressed liver tissue, clearly defining the border between the cavernous tissue and normal liver parenchyma, cannot be identified due to adjacent extensive hemangiomas. The enucleation can be avoided because can cause of massive hemorrhage, and hepatic resection is a preferable technique.

Surgery of large hemangiomas of the liver both by resection or by enucleation can cause torrential intraoperative hemorrhage, difficult to control with the high mortality risk for the patients<sup>48,49,77</sup> this is the reason why a vascular control before to start the procedure can be used to render safe the procedure.

Special attention is requested for hemangiomas located in the caudate lobe. Small hemangiomas may be enucleated or resected with no difficulty.<sup>78</sup> On the opposite resection of giant hemangiomas is more complicated, and isolation and excision of the lobe may be hazardous. The major risk in this second situation is the difficulty of the control of the profuse hemorrhage from the undersurface of the liver that can be very difficult to manage. In this case the ligation of the arterial branches to the caudate lobe along the border of the hepatoduodenal ligament, can shrink in size the hemangioma and becoming softer it is more is the manipulation and the dissection.<sup>79</sup> This technique is safer is supported by the vena cava and portal vessels encircled and ready to be clamped.

The Louisville consensus statement suggests laparoscopic liver resection as an option for lesions in the left lateral and inferior segments of the right lobe.<sup>80</sup> This technique has been described for liver hemangiomas. This technique pose some technical issues that concerning the location of the tumor, the absence of tactile feedback, the ideal technique for parenchymal transection that is not yet standardize and finally in case of large tumor the extraction can be difficult requiring a large incision. Despite the limitations and disadvantages of laparoscopic liver resection, which include a significant learning curve, bleeding that is more difficult to control laparoscopically, inadequate assessment of the liver for additional lesions, and increased risk for gas embolism, the increase in laparoscopic hepatic resections for benign tumors have recorded in the last

years based on the fact that these easy procedure are chosen by novice laparoscopic teams because of decreased initial difficulties.<sup>81</sup> The role of laparoscopic liver resection for liver tumors is unclear at present<sup>82-85</sup> but it can use in experienced groups with the right indications and good results. No case of laparoscopic enucleation has not yet reported.

The robotic procedure have until now very few reports.<sup>86,87</sup> It is a technique that like for the pilot for the airplane it will probably replace in the next 50 years the majority of the work of the surgeon. With the possibility to schedule the transection procedure by a computer the role of the surgeon will become just to supervise the procedure. But this is the future, for the present robotic surgery is not standardize for the resection of the hepatic hemangioma and the few experience do not permit to choice this technique as a first step and it should be performed in specialized center skilled both in hepatic and robotic surgery.

### Radiofrequency ablation

Radiofrequency ablation (RFA), used both percutaneously or laparoscopically, is a minimally invasive, safe, and effective treatment for primary and metastatic liver neoplasms.<sup>7,88</sup> Recently, percutaneous RFA therapy was also performed successfully for patients with hemangioma of the liver.<sup>89-92</sup> The mechanism of the RFA destruction of the hemangiomas is not completely understood. The effect of shrinkage of the tumor under the RFA is due to the action of the localized thermal injuries on the flat endothelial cells which constitute the wall of the widely dilated nonanastomotic vascular spaces. The temperature exceeding 90 °C damaging the endothelial layer lining the vascular structure may promote thrombosis.<sup>91</sup>

Both techniques percutaneous or laparoscopic can be used to treat the hemangioma with radiofrequency. Using the percutaneous technique tumor close to the gallbladder or other viscera cannot be treated, and at minimum 1 cm of distance need to avoid serious complications. Fullthickness burns of the stomach, small bowel, and colon have been observed when the edge of the thermal lesion was less than 1 cm from the surface of the liver in a pig liver model.<sup>26</sup>

On the opposite the laparoscopic approach due to the pneumoperitoneum elevates the diaphragm and the organ are displaced. This increase the operative space with facilitation of needle placement and at the same time avoiding damage of the closest viscera. Furthermore the laparoscopic approach permit

the use of intraoperative ultrasonography. This increase the possibility to determine real-time RF electrode placement and evaluate the efficacy of ablation. The success of the hemangioma ablation using radiofrequency depends also from the powerful of the used device and the length of the tines; both have to be maximized especially in largest hemangioma ( more than 10 cm).<sup>93</sup>

Advantages and disadvantages of the RF ablation depends from the benignity and hypervascularity of this tumor. The main 3 advantages of this techniques include at the first place the benignity of the tumor that does not require a safe margin of the liver parenchima surrounding the tumor. Second advantages is due to the content of the tumor. Infact as the hemangioma is constitute by blood positioned in cavities the effect of radiofrequency can lead to an obvious collapse of tumor tissue around the ablation zone. Third advantage of the hemangioma is strictly related to the benignity of the residual tumor, if it remains after the first treatment, does not need to be treated as soon as possible because neither develop rapid tumor progression nor metastasize to distant sites.<sup>94</sup>

The main disadvantage of the radiofrequency technique is the almost certain hemolysis due to the blood supply of this tumor. More is large the tumor and more is the risk and the gravity of this complication. Hemolysis can lead to various degrees of hemoglobinuria, hemolytic jaundice, anemia, or even renal damage accordingly, depending on its different severity.<sup>90</sup>

Furthermore other complications have been reported as esophageal perforation due to the closeness of the left lobe and ARDS.<sup>95</sup>

### Monoclonal antibody

Vascular endothelial growth factor (VEGF) is recognized as a regulator of blood vessel growth and it is postulated that also the hemangioma can grow under the action of the VEGF [96]. In the early 1970s, Folkman, *et al.*<sup>97</sup> identified a tumor-angiogenesis factor that was mitogenic to tumor capillary endothelial cells and suggested that blocking this factor might arrest tumor growth. The use of specific antibodies directed against VEGF abolishes the vascular endothelial growth-promoting activity *in vitro*.<sup>98</sup> Bevacizumab is a recombinant humanized monoclonal antibody (93% human and 7% murine) directed against VEGF which is used for the treatment of metastatic colorectal cancer in combination with 5-fluorouracil (5-FU)-based regimen.<sup>96</sup>



Bevacizumab have been reported as a medical treatment inducing a biological and clinical response in a patient of symptomatic hereditary hemorrhagic telangiectasia with liver involvement complicated by high-output cardiac failure, portal hypertension and cholestasis. There was a marked diminution of hepatic vascularity, a two-fold reduction in liver volume and normalization of cardiac output, thus alleviating the need for a liver transplantation. Therapy was well tolerated. The authors suggested that biological agents targeting angiogenic growth factors represent a novel form of therapy for these patients.<sup>99</sup> If the use of bevacizumab will be confirmed can be used not only for the hemangiomas of the liver but also for the hemangiomas in other parts of the body as well.

### Radiotherapy

Radiation therapy provides partial reduction in hemangioma size and relief of symptoms. Usually the dose of 30 Gy is given in 15 fractions over 3 weeks. With this method a complete clinical response have been reported over a period of 8-14 months.<sup>100</sup> This treatment can be effective with minimum morbidity if other treatment are nor advisable, but complications including radiation hepatitis, veno-occlusive disease, and hepatoma have to be taken in due account.<sup>101</sup> Although the biological mechanisms underlying the treatment efficacy of RT for haemangioma remain undetermined,<sup>102</sup> damage to vascular endothelial cells and smooth muscle cells is generally assumed to play a key role in the radiation effects, leading to vascular thrombosis, necrosis and fibrosis. Three stages can be described in this process.<sup>1</sup> Acute stage: interstitial oedema of the vascular endothelium develops;<sup>7</sup> transient subacute stage: organising changes occur in the central vein, which may progress to partial obliteration;<sup>103</sup> chronic stage: sclerosis or thrombosis of hepatic arterioles and portal tissues takes place against the healing process.

However, radiation therapy is rarely recommended as a first-line therapy for the liver hemangiomas owing to the concern of treatment-related liver toxicity and the long-term potential for secondary malignancies.<sup>101</sup> This treatment is reserved for massive hepatic haemangiomas associated with intractable congestive heart failure or hyperconsumptive coagulopathy in paediatric patients,<sup>101,104,105</sup> that cannot be treated with other therapies.

### TAE

Successful use of trans arterial embolization (TAE) before surgical resection of ruptured hepatic hemangioma was first reported by Yamamoto, *et al.* in 1991.<sup>106</sup> Since than this procedure have been also reported as a pre treatment before the elective surgical procedure, but results of this procedure are controversial<sup>107</sup> because of the fear of causing ischemia, intracavitary bleeding or infection.<sup>107</sup> The idea was to decrease the size of the hemangiomas, especially the tumors larger than 20 cm, by blocking its arterial supply and consequent shrinkage of the tumor, which facilitated subsequent mobilization of the liver, and consequently decreased intraoperative hemorrhage.<sup>108-110</sup> Recently, 98 patients affected by hemangiomas have been treated using trans arterial embolization using pingyangmycin-lipiodol emulsion that have proved to be a useful procedure as the only therapy of hemangiomas of the liver.<sup>111</sup> However, TAE may result in severe complications including ectopic embolizations and destructive biliary damage.<sup>112</sup>

Radiation therapy and TAE are generally considered palliative.

### Chemotherapy

The chemotherapy until now have not been extensively reported. Recently a very interesting case report reported by Hascimoto M and co-workers<sup>113</sup> have open new perspective concerning this kind of therapy. A 35 years old women affected by ovarian tumor with 6 hemangiomas of the liver have been submitted, before surgery, to five cycles of bleomycin/etoposide/cisplatin (BEP) therapy [cisplatin (20 mg/m<sup>2</sup>), etoposide (100 mg/m<sup>2</sup>) and bleomycin (5 mg/m<sup>2</sup>) on days 1-5. After this treatment also the hemangiomas reduced in size. At the intervention one of this tumor have been resected to be sure was not a metastasis. But the resected tumour was diagnosed as a typical haemangioma.

Although it is not clear why the haemangiomas decreased in size after systemic chemotherapy, some speculations can be proposed. First that development of hepatic haemangioma was associated with oestrogen so as the oophorectomy suppressed ovarian function, the absence of stimulation decreases also the haemangioma. Second, the decrease in the size of the haemangioma may have been in part due to the chemotherapy itself, especially to the direct devascularizing effect of bleomycin.<sup>111,114</sup> Third, a combination of the devascularizing effect of bleomycin

and suppression of ovarian function due to chemotherapy may have contributed to the reduction in size.

### Liver transplantation

Liver transplantation have been reported for giant hemangioma of the liver and life-threatening coagulopathy secondary to Kasabach Merritt syndrome<sup>115</sup> of for other indications.<sup>116</sup> It is an extreme life threatening treatment with very few indications.

### CONCLUSION

The hepatic cavernous hemangioma is the most common liver benign tumor. Liver hemangiomas, when indication exist, have to be treated firstly by surgery (hepatic resection or enucleation, open, laproscopic or robotic), but in the recent years other therapies like liver transplantation, radiofrequency ablation, radiotherapy, trans-arterial embolization, and chemotherapy have been applied. New prospective randomized studies are needed for evaluate which of these techniques is the best when the surgical procedure is not possible.

### REFERENCES

- Belli L, De Carlis L, Beati C, Rondinara G, Sansalone V, Brambilla G. Surgical treatment of symptomatic giant hemangiomas of the liver. *Surg Gynecol Obstet* 1992; 174: 474-8.
- Brouwers MA, Peeters PM, de Jong KP, Haagsma EB, Klompaker IJ, Bijleveld CM, Zwaveling JH, Slooff MJ. Surgical treatment of giant haemangioma of the liver. *Br J Surg* 1997; 84: 314-6.
- Mortele KJ, Ros PR. Benign liver neoplasms. *Clin Liver Dis* 2002; 6: 119-45.
- Giannitrapani L, Soresi M, La Spada E, Cervello M, D'Alessandro N, Montalto G. Sex hormones and risk of liver tumor. *Ann N Y Acad Sci* 2006; 1089: 228-36.
- Spitzer D, Krainz R, Graf AH, Menzel C, Staudach A. Pregnancy after ovarian stimulation and intrauterine insemination in a woman with cavernous macrohemangioma of the liver. A case report. *J Reprod Med* 1997; 42: 809-12.
- Reddy KR, Kligerman S, Levi J, Livingstone A, Molina E, Franceschi D, Badalamenti S, et al. Benign and solid tumors of the liver: relationship to sex, age, size of tumors, and outcome. *Am Surg* 2001; 67: 173-8.
- Gandolfi L, Leo P, Solmi L, Vitelli E, Verros G, Colecchia A. Natural history of hepatic haemangiomas: clinical and ultrasound study. *Gut* 1991; 32: 677-80.
- Ozakyol A, Kebapci M. Enhanced growth of hepatic hemangiomas in two adults after postmenopausal estrogen replacement therapy. *Tohoku J Exp Med* 2006; 210: 257-61.
- Sewell JH, Weiss K. Spontaneous rupture of hemangioma of the liver. A review of the literature and presentation of illustrative case. *Arch Surg* 1961; 83: 729-33.
- Morley JE, Myers JB, Sack FS, Kalk F, Epstein EE, Lannon J. Enlargement of cavernous haemangioma associated with exogenous administration of oestrogens. *S Afr Med J* 1974; 48: 695-7.
- Schwartz SI, Husser WC. Cavernous hemangioma of the liver. A single institution report of 16 resections. *Ann Surg* 1987; 205: 456-65.
- Graham E, Cohen AW, Soulen M, Faye R. Symptomatic liver hemangioma with intra-tumor hemorrhage treated by angiography and embolization during pregnancy. *Obstet Gynecol* 1993; 81: 813-6.
- Saegusa T, Ito K, Oba N, Matsuda M, Kojima K, Tohyama K, Matsumoto M, et al. Enlargement of multiple cavernous hemangioma of the liver in association with pregnancy. *Intern Med* 1995; 34: 207-11.
- Chui AK, Vass J, McCaughan GW, Sheil AG. Giant cavernous haemangioma: a rare indication for liver transplantation. *Aust N Z J Surg* 1996; 66: 122-4.
- Marques R, Taborda F, Jorge CS, Areias J, Rodrigues AM. Successful outcome in a pregnancy complicated by large hepatic hemangioma. *Acta Obstet Gynecol Scand* 1997; 76: 606-7.
- Conter RL, Longmire WP Jr. Recurrent hepatic hemangiomas. Possible association with estrogen therapy. *Ann Surg* 1988; 207: 115-9.
- Krasuski P, Poniecka A, Gal E, Wali A. Intrapartum spontaneous rupture of liver hemangioma. *J Matern Fetal Med* 2001; 10: 290-2.
- Cobey FC, Salem RR. A review of liver masses in pregnancy and a proposed algorithm for their diagnosis and management. *Am J Surg* 2004; 187: 181-91.
- Glinkova V, Shevah O, Boaz M, Levine A, Shirin H. Hepatic haemangiomas: possible association with female sex hormones. *Gut* 2004; 53: 1352-5.
- van Malenstein H, Maleux G, Monbaliu D, Verslype C, Komuta M, Roskams T, Laleman W, et al. Giant liver hemangioma: the role of female sex hormones and treatment. *Eur J Gastroenterol Hepatol* 2011; 23: 438-43.
- Moser C, Hany A, Spiegel R. Familial giant hemangiomas of the liver. Study of a family and review of the literature. *Praxis* 1998; 87: 461-8.
- Adam YG, Huvos AG, Fortner JG. Giant hemangiomas of the liver. *Ann Surg* 1970; 172: 239-45.
- Berliner L, el Ferzli G, Gianvito L, Worth MH Jr, Lowry J, Redmond P, Silich R. Giant cavernous hemangioma of the liver complicated by abscess and thrombosis. *Am J Gastroenterol* 1983; 78: 835-40.
- Kawarada Y, Mizumoto R. Surgical treatment of giant hemangioma of the liver. *Am J Surg* 1984; 148: 287-91.
- Yoon SS, Charny CK, Fong Y, Jarnagin WR, Schwartz LH, Blumgart LH, DeMatteo RP. Diagnosis, management, and outcomes of 115 patients with hepatic hemangioma. *J Am Coll Surg* 2003; 197: 392-402.
- Schnelldorfer T, Ware AL, Smoot R, Schleck CD, Harmsen WS, Nagorney DM. Management of giant hemangioma of the liver: resection versus observation. *J Am Coll Surg* 2010; 211: 724-30.
- Grieco MB, Miscall BG. Giant hemangiomas of the liver. *Surg Gynecol Obstet* 1978; 147: 783-7.
- van Tilborg AA, Nielsen K, Scheffer HJ, van den Tol P, van Waesberghe JH, Sietes C, Meijerink MR. Bipolar radiofrequency ablation for symptomatic giant (> 10 cm) hepatic cavernous haemangiomas: initial clinical experience. *Clin Radiol* 2013; 68: e9-e14.
- Di Carlo I, Toro A. Discomfort and Anxiety Should Never Be Considered Surgical Indications for Hemangioma of the Liver. *World J Surg* 2013 in press.
- Park H, Park JY, Kim do Y, Ahn SH, Chon CY, Han KH, Kim SU. Characterization of focal liver masses using acoustic

- radiation force impulse elastography. *World J Gastroenterol* 2013; 19: 219-26.
31. Frulio N, Laumonier H, Carteret T, Laurent C, Maire F, Balabaud C, Bioulac-Sage P, et al. Evaluation of liver tumors using acoustic radiation force impulse elastography and correlation with histologic data. *J Ultrasound Med* 2013; 32: 121-30.
  32. Streba CT, Ionescu M, Gheonea DI, Sandulescu L, Ciurea T, Saftoiu A, Vere CC, et al. Contrast-enhanced ultrasonography parameters in neural network diagnosis of liver tumors. *World J Gastroenterol* 2012; 18: 4427-34.
  33. Kim TK, Choi BI, Han JK, Hong HS, Park SH, Moon SK. Hepatic tumors: contrast agent enhancement patterns with pulse-inversion harmonic US. *Radiology* 2000; 216: 411-17.
  34. Lim KJ, Kim KW, Jeong WK, Kim SY, Jang YJ, Yang S, Lee JJ. Colour Doppler sonography of hepatic haemangiomas with arterioportal shunts. *Br J Radiol* 2012; 85: 142-6.
  35. Caseiro-Alves F, Brito J, Araujo AE, Belo-Soares P, Rodrigues H, Cipriano A, Sousa D, et al. Liver hemangioma: common and uncommon findings and how to improve the differential diagnosis. *Eur Radiol* 2007; 17: 1544-54.
  36. Fowler KJ, Brown JJ, Narra VR. Magnetic resonance imaging of focal liver lesion: approach to imaging diagnosis. *Hepatology* 2011; 54: 2227-37.
  37. Chan YL, Lee SF, Yu SC, Lai P, Ching AS. Hepatic malignant tumor versus cavernous hemangioma: differentiation on multiple breath-hold turbo spin echo MRI sequences with different T2 weighting and T2-relaxation time measurements on a single slice multi-echo sequence. *Clin Radiol* 2002; 57: 250-7.
  38. Vilgrain V, Boulos L, Vullierme MP, Denys A, Terris B, Menu Y. Imaging of atypical hemangioma of the liver with pathological correlation. *Radiographics* 2000; 20: 379-97.
  39. Erturk SM, Ichikawa T, Kaya E, Yapici O, Ozel A, Mahmutoglu AS, Basak M. Diffusion tensor imaging of cysts, hemangiomas, and metastases of the liver. *Acta Radiol* 2013; in press.
  40. Borse R, Mahapatra GN, Meht R, Plumber S, Dhuri S, Ali S. Scintigraphic finding of a silent hepatic haemangioma. *J Assoc Physicians India* 2010; 58: 637-40.
  41. Prasanna PM, Fredericks SE, Winn SS, Christman RA. Best cases from the AFIP: giant cavernous hemangioma. *Radiographics* 2010; 30: 1139-44.
  42. Imperiale A, Greget M, Chabrier G, Keomany J, Rust E, Detour J, Pessaux P, et al. Solitary hepatic metastasis from medullary thyroid carcinoma mimicking atypical hemangioma: insights from multimodality diagnostic approach by MRI, F-18 FDG and F-18 FDOPA PET/CT. *Clin Nucl Med* 2010; 35: 434-7.
  43. Chen RC, Lii JM, Chen WT, Tu HY, Chiang LC. Transcatheter arterial chemoembolization in patients with hepatocellular carcinoma and coexisting hepatic cavernous hemangioma. *Eur Radiol* 2006; 16: 1346-50.
  44. Marsh JI, Gibney RG, Li DKB. Hepatic hemangioma in the presence of fatty infiltration: an atypical sonographic appearance. *Gastrointest Radiol* 1989; 14: 262-4.
  45. Doklestiæ K, Stefanoviæ B, Karamarkovik A, Bumbasireviæ V, Stefanoviæ B, Gregoriæ P, Radenkoviæ D, et al. Spontaneous rupture of giant liver hemangioma: case report. *Srp Arh Celok Lek* 2013; 141: 95-9.
  46. Roebuck D, Sebire N, Lehmann E, Barnacle A. Rapidly involuting congenital haemangioma (RICH) of the liver. *Pediatr Radiol* 2012; 42: 308-14.
  47. Etemadi A, Golozar A, Ghassabian A, Zarei M, Hashemi Taheri AP, Dawsey SM, Malekzadeh R. Cavernous hemangioma of the liver: factors affecting disease progression in general hepatology practice. *Eur J Gastroenterol Hepatol* 2011; 23: 354-8.
  48. Farges O, Daradkeh S, Bismuth H. Cavernous hemangiomas of the liver: are there any indications for resection? *World J Surg* 1995; 19: 19-24.
  49. Ozden I, Emre A, Alper A, Tunaci M, Acarli K, Bilge O, Tebant Y, et al. Long-term results of surgery for liver hemangiomas. *Arch Surg* 2000; 135: 978-81.
  50. Buell JF, Tranchart H, Cannon R, Dagher I. Management of benign hepatic tumors. *Surg Clin North Am* 2010; 90: 719-35.
  51. Di Carlo I, Sofia M, Toro A. Does the psychological request of the patient justify the surgery for hepatic hemangioma? *Hepatogastroenterology* 2005; 52: 657-61.
  52. Herman P, Costa ML, Machado MA, Pugliese V, D'Albuquerque LA, Machado MC, Gama-Rodrigues JJ, et al. Management of hepatic hemangiomas: a 14-year experience. *J Gastrointest Surg* 2005; 9: 853-9.
  53. Yedibela S, Alibek S, Müller V, Aydin U, Langheinrich M, Lohmüller C, Hohenberger W, et al. Management of hemangioma of the liver: surgical therapy or observation? *World J Surg* 2013; 37: 1303-12.
  54. Kasabach HH, Merritt KK. Capillary hemangioma with extensive purpura: report of a case. *Am J Dis Child* 1940; 59: 1063.
  55. Choi J, Lee YJ, Hwang DW, Chon SH, Nagpal A, Park KM. Surgical treatment of giant hepatic hemangiomas: technical point of view. *Am Surg* 2011; 77: 48-54.
  56. Iqbal N, Saleem A. Hepatic hemangioma: a review. *Tex Med* 1997; 93: 48-50.
  57. Taitelbaum G, Hinchey EJ, Herba MJ, Lough J. Giant hemangioma of the liver. *Can J Surg* 1982; 25: 652-4.
  58. Trastek VF, van Heerden JA, Sheedy PF 2nd, Adson MA. Cavernous hemangiomas of the liver: resect or observe? *Am J Surg* 1983; 145: 49-53.
  59. Gilon D, Slater PE, Benbassat J. Can decision analysis help in the management of giant hemangioma of the liver? *J Clin Gastroenterol* 1991; 13: 255-8.
  60. Hotokezaka M, Kojima M, Nakamura K, Hidaka H, Nakano Y, Tsuneyoshi M, Jimi M. Traumatic rupture of hepatic hemangioma. *J Clin Gastroenterol* 1996; 23: 69-71.
  61. Kanazawa S, Douke T, Gotoh A, Nishi A, Yasui K, Kajiwara Y. [A case of giant cavernous hemangioma with hemoperitoneum due to blunt abdominal trauma; CT findings]. *Rinsho Hoshasen* 1990; 35: 979-82.
  62. Kocakusak A, Sunar H, Akinci M, Gulen M, Arikan S. Rupture of an incidental giant liver hemangioma caused by blunt abdominal trauma. *Ulus Travma Derg* 2002; 8: 176-8.
  63. Stayman JW Jr, Polsky HS, Blaum L. Case report. Ruptured cavernous hemangioma of the liver. *Pa Med* 1976; 79: 62-3.
  64. Nichols FC 3rd, van Heerden JA, Weiland LH. Benign liver tumors. *Surg Clin North Am* 1989; 69: 297-314.
  65. Alper A, Ariogul O, Emre A, Uras A, Okten A. Treatment of liver hemangiomas by enucleation. *Arch Surg* 1988; 123: 660-1.
  66. Hamaloglu E, Altun H, Ozdemir A, Ozenc A. Giant liver hemangioma: therapy by enucleation or liver resection. *World J Surg* 2005; 29: 890-3.
  67. Lerner SM, Hiatt JR, Salamandra J, Chen PW, Farmer DG, Ghobrial RM, Busuttill RW. Giant cavernous liver hemangiomas: effect of operative approach on outcome. *Arch Surg* 2004; 139: 818-21.
  68. Singh RK, Kapoor S, Sahni P, Chattopadhyay TK. Giant haemangioma of the liver: is enucleation better than resection? *Ann R Coll Surg Engl* 2007; 89: 490-3.

69. Blumgart LH. Liver resection for benign disease and for liver and biliary tumors. In: Blumgart LH, Fong Y (eds.). *Surgery of the liver and biliary tract*. 3rd Ed. Philadelphia: W.B. Saunders; 2000, p. 1639-713.
70. Fu XH, Lai EC, Yao XP, Chu KJ, Cheng SQ, Shen F, Wu MC, Lau WY. Enucleation of liver hemangiomas: is there a difference in surgical outcomes for centrally or peripherally located lesions? *Am J Surg* 2009; 198: 184-7.
71. Moon WS, Yu HC, Lee JM, Kang MJ. Diffuse hepatic hemangiomas in an adult. *J Korean Med Sci* 2000; 15: 471-4.
72. Lopriore E, Markhorst DG. Diffuse neonatal haemangiomas: new views on diagnostic criteria and prognosis. *Acta Paediatr* 1999; 88: 93-7.
73. Kim EH, Park SY, Ihn YK, Hwang SS. Diffuse hepatic hemangiomas without extrahepatic involvement in an adult patient. *Korean J Radiol* 2008; 9: 559-62.
74. Latifi HR, Siegel MJ. Diffuse neonatal hemangiomas: CT findings in an adult. *J Comput Assist Tomogr* 1992; 16: 971-3.
75. Ohnishi S, Miyagishima T, Nakagawa M, Kamata T, Kishimoto A, Choi GH, Kudo M, et al. Diffuse neonatal hemangiomas without cutaneous lesions in an adult a case report. *Angiology* 2002; 53: 235-7.
76. Lehmann FS, Beglinger C, Schnabel K, Terracciano L. Progressive development of diffuse liver hemangiomas. *J Hepatol* 1999; 30: 951-4.
77. Hanazaki K, Kajikawa S, Matsushita A, Monma T, Koide N, Nimura Y, Yazawa K, et al. Hepatic resection of giant cavernous hemangioma of the liver. *J Clin Gastroenterol* 1999; 29: 257-60.
78. Huang ZQ. *Huang Zhiqiang hepatic surgery operation*. 1st Ed. Beijing: People's military medical press; 2005, p. 275-8.
79. Xu LN, Huang ZQ. Resection of hepatic caudate lobe hemangioma: experience with 11 patients. *Hepatobiliary Pancreat Dis Int* 2010; 9: 487-91.
80. Buell JF, Cherqui D, Geller DA, O'Rourke N, Iannitti D, Dagher I, Koffron AJ, et al. The international position on laparoscopic liver surgery: The Louisville Statement, 2008. *Ann Surg* 2009; 250: 825-30.
81. Toro A, Gagner M, Di Carlo I. Has laparoscopy increased surgical indications for benign tumors of the liver? *Langenbecks Arch Surg* 2013; 398: 195-210.
82. Nguyen KT, Gamblin TC, Geller DA. World review of laparoscopic liver resection-2,804 patients. *Ann Surg* 2009; 250: 831-41.
83. Pulvirenti E, Toro A, Di Carlo I. An update on indications for treatment of solid hepatic neoplasms in noncirrhotic liver. *Future Oncol* 2010; 6: 1243-50.
84. Reich H, McGlynn F, DeCaprio J, Budin R. Laparoscopic excision of benign liver lesions. *Obstet Gynecol* 1991; 78: 956-8.
85. Polignano FM, Quyn AJ, de Figueiredo RS, Henderson NA, Kulli C, Tait IS. Laparoscopic versus open liver segmentectomy: prospective, case-matched, intention-to-treat analysis of clinical outcomes and cost effectiveness. *Surg Endosc* 2008; 22: 2564-70.
86. Giulianotti PC, Addeo P, Bianco FM. Robotic right hepatectomy for giant hemangioma in a Jehovah's Witness. *J Hepatobiliary Pancreat Sci* 2011; 18: 112-8.
87. Giulianotti PC, Coratti A, Sbrana F, Addeo P, Bianco FM, Buchs NC, Annechiario M, et al. Robotic liver surgery: results for 70 resections. *Surgery* 2011; 149: 29-39.
88. Pietrabissa A, Giulianotti P, Campatelli A, Di Candio G, Farina F, Signori S, Mosca F. Management and follow-up of 78 giant haemangiomas of the liver. *Br J Surg* 1996; 83: 915-8.
89. Sharpe EE 3rd, Dodd GD 3rd. Percutaneous radiofrequency ablation of symptomatic giant hepatic cavernous hemangiomas: report of two cases and review of literature. *J Vasc Interv Radiol* 2012; 23: 971-5.
90. Park SY, Tak WY, Jung MK, Jeon SW, Cho CM, Kweon YO, Kim KC. Symptomatic-enlarging hepatic hemangiomas are effectively treated by percutaneous ultrasonography-guided radiofrequency ablation. *J Hepatol* 2011; 54: 559-65.
91. Tak WY, Park SY, Jeon SW, Cho CM, Kweon YO, Kim SK, Choi YH, et al. Ultrasonography-guided percutaneous radiofrequency ablation for treatment of a huge symptomatic hepatic cavernous hemangioma. *J Clin Gastroenterol* 2006; 40: 167-70.
92. Cui Y, Zhou LY, Dong MK, Wang P, Ji M, Li XO, Chen CW, et al. Ultrasonography guided percutaneous radiofrequency ablation for hepatic cavernous hemangioma. *World J Gastroenterol* 2003; 9: 2132-4.
93. Ke S, Ding X, Gao J, Gao K, Qian X, Cao B, Li M, et al. Solitary huge hepatocellular carcinomas 10 cm or larger may be completely ablated by repeated radiofrequency ablation combined with chemoembolization: Initial experience with 9 patients. *Mal Med Rep* 2012; 5: 832-6.
94. Gao J, Ke S, Ding XM, Zhou YM, Qian XJ, Sun WB. Radiofrequency ablation for large hepatic hemangiomas: initial experience and lessons. *Surgery* 2013; 153: 78-85.
95. Ng KK, Lam CM, Poon RT, Shek TW, Ho DW, Fan ST. Safety limit of large-volume hepatic radiofrequency ablation in a rat model. *Arch Surg* 2006; 141: 252-8.
96. Homsy J, Daud AI. Spectrum of activity and mechanism of action of VEGF/PDGF inhibitors. *Cancer Control* 2007; 14: 285-94.
97. Folkman J, Merler E, Abernathy C, Williams G. Isolation of a tumor factor responsible for angiogenesis. *J Exp Med* 1971; 133: 275-88.
98. Berard M, Sordello S, Ortega N, Carrier JL, Peyri N, Wassef M, Bertrand N, et al. Vascular endothelial growth factor confers a growth advantage in vitro and in vivo to stromal cells cultured from neonatal hemangiomas. *Am J Pathol* 1997; 150: 1315-26.
99. Mitchell A, Adams LA, MacQuillan G, Tibballs J, vanden Driesen R, Delriviere L. Bevacizumab reverses need for liver transplantation in hereditary hemorrhagic telangiectasia. *Liver Transpl* 2008; 14: 210-3.
100. Biswal BM, Sandhu M, Lal P, Bal CS. Role of radiotherapy in cavernous hemangioma liver. *Indian J Gastroenterol* 1995; 14: 95-8.
101. Gaspar L, Mascarenhas F, da Costa MS, Dias JS, Afonso JG, Silvestre ME. Radiation therapy in the unresectable cavernous hemangioma of the liver. *Radiation Oncol* 1993; 29: 45-50.
102. Brady LW. Radiotherapy for non-malignant disorders. Contemporary concepts and clinical results. Berlin, Germany: Springer; 2008.
103. Ishak KG, Rabin L. Benign tumors of the liver. *Med Clin North Am* 1975; 59: 995-1013.
104. Cohen RC, Myers NA. Diagnosis and management of massive hepatic hemangiomas in childhood. *J Pediatr Surg* 1986; 21: 6-9.
105. Iyer CP, Stanley P, Mahour GH. Hepatic hemangiomas in infants and children: a review of 30 cases. *Am Surg* 1996; 62: 356-60.
106. Yamamoto T, Kawarada Y, Yano T, Noguchi T, Mizumoto R. Spontaneous rupture of hemangioma of the liver: treatment with transcatheter hepatic arterial embolization. *Am J Gastroenterol* 1991; 86: 1645-9.

107. Seo HI, Jo HJ, Sim MS, Kim S. Right trisegmentectomy with thoracoabdominal approach after transarterial embolization for giant hepatic hemangioma. *World J Gastroenterol* 2009; 15: 3437-9.
108. Giavroglou C, Economou H, Ioannidis I. Arterial embolization of giant hepatic hemangiomas. *Cardiovasc Intervent Radiol* 2003; 26: 92-6.
109. Srivastava DN, Gandhi D, Seith A, Pande GK, Sahni P. Transcatheter arterial embolization in the treatment of symptomatic cavernous hemangiomas of the liver: a prospective study. *Abdom Imaging* 2001; 26: 510-4.
110. Vassiou K, Rountas H, Liakou P, Arvanitis D, Fezoulidis I, Tepetes K. Embolization of a giant hepatic hemangioma prior to urgent liver resection. Case report and review of the literature. *Cardiovasc Intervent Radiol* 2007; 30: 800-2.
111. Zeng Q, Li Y, Chen Y, Ouyang Y, He X, Zhang H. Gigantic cavernous hemangioma of the liver treated by intra-arterial embolization with pingyangmycin-lipiodol emulsion: a multi-center study. *Cardiovasc Intervent Radiol* 2004; 27: 481-5.
112. Huang XQ, Huang ZQ, Duan WD, Zhou NX, Feng YQ. Severe biliary complications after hepatic artery embolization. *World J Gastroenterol* 2002; 8: 119-23.
113. Hashimoto M, Sugawara M, Ishiyama K, Sato T, Yamamoto Y, Nanjo H. Reduction in the size of a hepatic haemangioma after chemotherapy. *Liver Int* 2008; 28: 1043-4.
114. Okano H, Shiraki K, Inoue H, Ito T, Yamanaka T, Deguchi M, Sugimoto K, et al. Natural course of cavernous hepatic hemangioma. *Oncol Rep* 2001; 8: 411-4.
115. Longeville JH, de la Hall P, Dolan P, Holt AW, Lillie PE, Williams JA, Padbury RT. Treatment of a giant haemangioma of the liver with Kasabach-Merritt syndrome by orthotopic liver transplant a case report. *HPB Surg* 1997; 10: 159-62.
116. Unal E, Francis F, Aquino A, Xu R, Morgan G, Teperman L. Liver transplant for mixed capillary-cavernous hemangioma masquerading as hepatocellular carcinoma in a patient with hepatocellular carcinoma. *Exp Clin Transplant* 2011; 9: 344-8.