



Thoracoscopic resection of a lung keratocyst associated with pulmonary sequestration in a neonate



Prof. Mario Lima^a, Dr. Tommaso Gargano^{a,*}, Dr. Raed Al-Taher^a, Dr. Michela Maffi^a,
Dr. Nunzio Salfi^b

^a Department of Pediatric Surgery, S. Orsola Hospital, University of Bologna, Bologna, Italy

^b Department of Anatomy and Pathological Histology, S. Orsola Hospital, University of Bologna, Bologna, Italy

ARTICLE INFO

Article history:

Received 13 February 2017

Received in revised form

9 May 2017

Accepted 15 May 2017

Available online 16 May 2017

Keywords:

Lung keratocyst

Odontogenic keratocyst

Bronchogenic cyst

Pulmonary sequestration

Thoracoscopic resection

ABSTRACT

Introduction: Keratocyst is a cystic lesion that has distinctive histological features composed of an epithelial lining with surface keratinization that is corrugated and predominantly parakeratotic. Although rare, they present in human pathology mainly as odontogenic keratocysts and have occasional association with Gorlin-Goltz syndrome.

Case report: We present a neonate with an antenatal suspicion of congenital pulmonary airway malformation. While stable and imaging confirmation studies were accomplished, thoracoscopic exploration and excision of a bronchogenic-like lung cyst attached to an extralobar pulmonary sequestration superomedial to the left upper lobe were performed successfully at day 44 of life. Histopathological study has revealed keratocystic features of the lung cyst attached to a pulmonary sequestration tissue.

Discussion and conclusion: Up to our knowledge and review of the literature, this is the first case and occurrence of Lung Keratocyst. Long-term follow up is needed to assess the outcome of the thoracoscopic management of such lesion, and its hypothetic relation with odontogenic keratocysts or any associated syndromes.

© 2017 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The term “Lung Keratocyst” is not existing in literature. Keratocysts in literature are almost always of odontogenic origin and called “Keratocystic Odontogenic Tumors” that affect the mandible in most cases. We report an exceptional case of a lung cyst of histopathological keratocystic features associated with an extralobar pulmonary sequestration that underwent a thoracoscopic resection.

2. Case report

A 37 days old male baby, who was referred to our care from a

peripheral hospital with an antenatal diagnosis of congenital pulmonary airway malformation and a postnatal suspicion of a bronchogenic cyst according to postnatal chest radiographs and computed tomography images. The child was term and had a birth weight of 3700 g with stable hemodynamic and cardiorespiratory status after birth. He had good general health conditions with no any respiratory symptom. Chest radiograph has shown a well demarcated round radio-opaque structure occupying the left upper hemithorax with slight shifting of mediastinum toward the right side. Upper gastrointestinal contrast study has ruled out any communication with the gastrointestinal tract (Fig. 1). Computed tomography scan with contrast has shown a non-enhanced round cystic lesion of fluid density occupying the left paramediastinal region in corresponding to the apico-dorsal segment of the left upper lobe, 23 × 24 × 24 mm in size, in close contact with the descending thoracic aorta and pushing a what suspected to be the apical segment arterial branch anteriorly (Fig. 2). Our primary diagnosis was a bronchogenic cyst and therefore thoracoscopic resection was planned.

At day 44 of life, thoracoscopy was performed and we found a

* Corresponding author. Department of Pediatric Surgery, S. Orsola Hospital, University of Bologna, Via Giuseppe Massarenti, 11, Bologna (BO) 40138, Italy.

E-mail addresses: mario.lima@unibo.it (M. Lima), tommaso.gargano2@unibo.it (T. Gargano), raed_nt@hotmail.com (R. Al-Taher), michela.maffi@libero.it (M. Maffi), nunziocm.salfi@aosp.bo.it (N. Salfi).




Fig. 1. A) Chest radiograph showing the round radio-opaque lesion that occupies the left upper hemithorax with slight shifting of mediastinum toward the contralateral side. B) An upper GI contrast study showing normal esophageal contour with no evidence of communication with the cystic lesion.

bronchogenic-like cyst that was firmly attached to an extralobar pulmonary sequestration and to the adjacent normal lung tissue as well. It was difficult to dissect and isolate the cyst from the surrounding structures, that is why it was evacuated and opened where whitish creamy fluid came out of the cyst compatible with keratinaceous material. The part that was attached to the normal lung tissue was removed using Ligature™. The feeding vessel of the sequestration tissue was doubly ligated using Endoloops® and transected. The sequestration tissue with the attached remaining part of the lung cyst were removed and extracted out together (Fig. 3). Chest drain was left in place and operative time was 100 min. The patient had a smooth recovery postoperatively and he was discharged home at the tenth postoperative day. Follow up for 6 months does not show any complication.

Histological examination of the specimen has reported a pulmonary sequestration that is characterized by unaerated pulmonary parenchyma, serpiginous vessels of systemic type, and subpleural arterioles. The attaching cyst looks of bronchogenic origin due to the presence of sero-mucinous glands, cartilage and layers of smooth muscle cells, but surprisingly, it shows typical histologic characteristics of a keratocyst as well. These characteristics include the presence of few layers of squamous epithelium with incomplete



Fig. 2. Thoracic CT scan showing the cystic lesion at the left upper hemithorax (asterisk). The cyst is pushing a systemic vessel anteriorly (arrows) which was found intraoperatively to be the feeding vessel of an extralobar pulmonary sequestration.

keratinization, the corrugated and irregular luminal epithelial layer, and the complete absence of mucous-secreting goblet cells all through the wall of the lung cyst (Figs. 4 and 5).

3. Discussion

We describe the first case of an unusual occurrence of a lung cyst which has histological features of both a bronchogenic cyst and a keratocyst, with an unusual association with upper extralobar pulmonary sequestration. The lung cyst and sequestration were resected by thoracoscopy.

When mentioning a keratocyst, it has no occurrence in literature rather being an odontogenic one. Odontogenic keratocysts (OKC) are cystic lesions of the bone that are lined with keratinized epithelium and thought to originate from the dental lamina [1]. It is known for its high recurrence rate, aggressive behavior, and its occasional association with the nevoid basal cell carcinoma syndrome (NBCCS) or the Gorlin-Goltz syndrome [2]. Three-fourths of OKCs present in the mandible [3]. The term OKC was first coined by Philipsen in 1956 [4], until in recent years when the WHO has recommended that the term “keratocystic odontogenic tumor” (KCOT) replace the term “odontogenic keratocyst”, as it better reflects the neoplastic nature of the lesion [5]. Diagnosis is usually made with histopathology studies, where the main histologic features are composed of a thin lining epithelium of a parakeratinized surface which is typically corrugated, rippled or wrinkled, a remarkable uniformity of thickness of the epithelium, usually ranging from 6 to 10 cells thick, and a prominent palisaded, polarized basal layer of cells often described as having a ‘picket fence’ or ‘tombstone’ appearance [6].

Gorlin-Goltz syndrome is a rare inherited multisystem disorder, where affected patients have multiple developmental anomalies and early development of multiple basal cell carcinomas, odontogenic keratocysts, especially in the mandible, palmar and plantar pitting, ectopic intracranial calcification, and family history of NBCCS [7,8]. It has never been reported an association between any of the OKC or the Gorlin-Goltz syndrome with a lung sequestration or any other bronchopulmonary malformation.

On the other hand, bronchogenic cysts (BC) are congenital anomalies arising as an abnormal ventral budding of the tracheo-bronchial tree and so have an epithelium which is similar to bronchi [9]. Histologically, they are lined by ciliated pseudostratified respiratory epithelium or metaplastic squamous cells and have focal areas of hyaline cartilage, smooth muscle, and bronchial glands within their walls [10]. Furthermore, by definition and the histological components, our cyst is neither an enteric duplication cyst, an epidermal cyst, nor a dermal cyst.

An association of pulmonary sequestration (PS) and BC is unusual. We have reviewed the MEDLINE® database searching for the reported cases of PS that are associated with other pathologies, and specifically with BCs (Table 1). There is a wide range of associated anomalies and disorders but with few reported cases of each. There are 12 reported cases that describe an association between PS and BC. They include 7 cases of extralobar PS type [11–17], and 5 cases of intralobar type [18–22]. In addition, there are five other reported cases of association between PS and enteric duplication cysts.

Leaning on the aforementioned results, we suppose our case report to be the first occurrence of a “Lung Keratocyst”, of a bronchogenic origin, in association with an unusual left upper lobe extralobar PS. Hereby, we assume to introduce the new term of “Bronchogenic Keratocyst”, where the tissue of origin is not the usual odontogenic one. Could there be an association between bronchogenic and odontogenic keratocysts? Would bronchogenic keratocysts present aggressiveness and infiltrative behavior




Fig. 3. Thoracoscopic resection of the lung keratocyst in concurrence with the extralobar pulmonary sequestration.




Fig. 4. General microscopic view of the whole specimen showing the pulmonary sequestration tissue (arrows) with the attached opened lung cyst (asterisk).

resembling odontogenic keratocysts? Would we face a high recurrence rate? Questions to be answered with further histopathological analysis, long-term follow up, and possible occurrence of other similar cases in future.

4. Conclusion

Up to our knowledge and review of the literature, this is the first case and occurrence of a Lung Keratocyst, that is of bronchogenic origin, and in association with the uncommon left upper lobe extralobar PS on top. Long-term follow up is needed to assess the outcome of the thoracoscopic resection of such lesion. Further histopathological analysis and studying of possible future cases are needed to outline the doubtful aggressiveness and infiltrative behavior of bronchogenic keratocysts, and their possible relation with odontogenic keratocysts or the associated syndromes.




Fig. 5. Higher magnification microscopic views of the lung cyst showing features of both a bronchogenic cyst and a keratocyst.

Table 1
Literature review of reported cases of PS that is associated with other pathologies.

Associated pathologies	No. of cases	Details
Cystic lesions	17	12 with bronchogenic cysts 5 with duplication cysts
Congenital diaphragmatic hernia (CDH)	14	7 alone 1 with BEF 2 with CPAM & LTE cleft 1 with dermoid cyst & intestinal duplication 1 with gut malrotation 1 with pulmonary hypoplasia 1 with intrathoracic kidney
Congenital pulmonary airway malformation (CPAM)	11	8 alone 2 with CDH and LTE Cleft 1 with Prune-Belly Syndrome
Broncho-Esophageal Fistula (BEF)	6	5 alone 1 with CDH
Pulmonary Hypoplasia	4	3 alone 1 with CDH
Bronchial Stenosis/Atresia	3	
Laryngo-tracheo-esophageal cleft (LTEC)	3	2 with CDH & CPAM 1 with Taussing-Bing Syndrome
Vascular Anomalies	6	
Tumors	7	Adenoma-like-hamartoma Hepatoplastoma (Aicardi syndrome) Carcinoid tumourlets Hemangioma Lung cancer Lung sarcoma Thymoma (Lupus Syndrome)
Infection	18	12 Aspergillosis 3 TB 2 non-TB Mycobacterium 1 mixed
Other miscellaneous	13	Torsion of PS, vertebral anomalies, funnel chest, hemothorax, bronchiectasis, lung herniation, azygos lobe, Scimitar Syndrome, Achalasia, accessory diaphragm

References

- [1] Katase Naoki, Nagatsuka Hitoshi, Tsujigiwa Hidetsugu, Gunduz Mehmet, Tamamura Ryo, Pwint Han Phuu, et al. Analysis of the neoplastic nature and

- biological potential of sporadic and nevoid basal cell carcinoma syndrome-associated keratocystic odontogenic tumor. *J Oral Pathol Med* 2007;36(9): 550–4. <http://dx.doi.org/10.1111/j.1600-0714.2007.00578.x>.
- [2] Eryilmaz Tolga, Ozmen Selahattin, Findikcioglu Kemal, Kandal Sebahattin, Aral Mubin. Odontogenic keratocyst. *Ann Plast Surg* 2009;62(2):210–2. <http://dx.doi.org/10.1097/sap.0b013e31817dad9c>.
- [3] Myoung Hoon, Hong Sam-Pyo, Hong Seong-Doo, Lee Jae-Il, Lim Chang-Yun, Chung Pill-Hoon, et al. Odontogenic keratocyst: review of 256 cases for recurrence and clinicopathologic parameters. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2001;91(3):328–33. <http://dx.doi.org/10.1067/moe.2001.113109>.
- [4] Philipsen HP. Om keratocysts (kolesteatomer) I kaerberne. *Tandlaegebladet* 1956;60:963–80.
- [5] Philipsen HP. Keratocystic odontogenic tumor. In: Barnes L, Eveson JW, Reichart PA, Sidransky D, editors. *World health organization classification of tumours: pathology and genetics head and neck tumours*. Lyon, France: IARC Press; 2005. p. 306–7.
- [6] Rajendran A, Sivapathasundharam B. *Shafer's textbook of oral pathology*. Elsevier Health Sciences APAC; 2014.
- [7] Evans DG, Ladusans EJ, Rimmer S, Burnell LD, Thakker N, Farndon PA. Complications of the naevoid basal cell carcinoma syndrome: results of a population based study. *J Med Genet* 1993;30(6):460–4.
- [8] Rahbari Homayoon, Mehregan Amir H. Basal cell epithelioma (carcinoma) in children and teenagers. *Cancer* 1982;49(2):350–3.
- [9] Correia-Pinto Jorge, Gonzaga Silvia, Huang Yadi, Rottier Robbert. Congenital lung lesions—underlying molecular mechanisms. *Semin Pediatr Surg* 2010;19(3):171–9. <http://dx.doi.org/10.1053/j.sempedsurg.2010.03.003>.
- [10] Ali SZ, Yang GCH. *Lung and mediastinum cytohistology*. Cambridge University Press; 2012.
- [11] Ropion-Michaux H, Noel JB, Besnard M, Prevot M, Sauvage PJ. Extra-lobar pulmonary sequestration associated with a bronchogenic cyst: a case report. *J Radiol* 2010;91(11 Pt 1):1164–7.
- [12] Kim Kyoung Won, Kim W-S, Cheon Jung-Eun, Lee HJ, Kim Chong Jai, Kim I-O, et al. Complex bronchopulmonary foregut malformation: extralobar pulmonary sequestration associated with a duplication cyst of mixed bronchogenic and oesophageal type. *Pediatr Radiol* 2001;31(4):265–8. <http://dx.doi.org/10.1007/s002470000410>.
- [13] MacKenzie Tippi C, Guttenberg Marta E, Nisenbaum Harvey L, Johnson Mark P, Adzick N Scott. A fetal lung lesion consisting of bronchogenic cyst, bronchopulmonary sequestration, and congenital cystic adenomatoid malformation: the missing link? *Fetal Diagn Ther* 2001;16(4):193–5. <http://dx.doi.org/10.1159/000053907>.
- [14] Ikeda Y, Matsushita Y, Chimoto M, Kobayashi S, Sugita K, Ishihama H, et al. A case report of extralobar pulmonary sequestration associated with mediastinal bronchogenic cyst. *Kyobu Geka* 1998;51(2):154–7.
- [15] Lee S, Lee J, Park JH, Kim YW, Yang MH, Shin DW. Extralobar pulmonary sequestration with an associated cyst of mixed bronchogenic and esophageal type—a case report. *J Korean Med Sci* 1997;12(6):567–9. <http://dx.doi.org/10.3346/jkms.1997.12.6.567>.
- [16] Williams S, Burton EM, Day S, Hanley M, Mercado-Deane MG, Rogers DA. Combined sequestration, bronchogenic cyst, and dysgenetic lung simulating congenital lobar emphysema. *South Med J* 1996;89(12):1220–2.
- [17] Black Timothy L, Fernandes Eduardo T, Wrenn Earle L, Lynn Magill H. Extralobar pulmonary sequestration and mediastinal bronchogenic cyst. *J Pediatr Surg* 1988;23(11):999–1001.

- [18] Onuki T, Nakaoka K, Yamana T, Nishiyama N, Saitou K, Inagaki M. Independent occurrence of a mediastinal bronchogenic cyst and an intralobar pulmonary sequestration. *Kyobu Geka* 2015;68(11):947–9.
- [19] Traibi Akram, Strauss Christiane, Validire Pierre, Gossot Dominique. Intralobar pulmonary sequestration associated with bronchogenic cyst in adult. *Asian Cardiovasc Thorac Ann* 2012;20(5):597–9. <http://dx.doi.org/10.1177/0218492312440702>.
- [20] Barbut J, Fernandez C, Blanc F, Dubus JC, Guys JM, de Lagausie P. Pulmonary sequestration of the left upper lobe associated with a bronchogenic cyst: case report of an exceptional association. *Pediatr Pulmonol* 2011;46(5):509–11. <http://dx.doi.org/10.1002/ppul.21403>.
- [21] Grewal RG, Yip CK. Intralobar pulmonary sequestration and mediastinal bronchogenic cyst. *Thorax* 1994;49(6):615–6.
- [22] Croyle P, Estrera AS. Bronchogenic cyst and intralobar sequestration mimicking thoracic aortic aneurysm. *South Med J* 1982;75(10):1267–8.