

Flavonoids profile and antioxidant activity in flowers and leaves of hawthorn species (*Crataegus* spp.) from different regions of Iran

Abolfazl Alirezalu, Peyman Salehi, Nima Ahmadi, Ali Sonboli, Serena Aceto, Hamid Hatami Maleki & Mahdi Ayyari

To cite this article: Abolfazl Alirezalu, Peyman Salehi, Nima Ahmadi, Ali Sonboli, Serena Aceto, Hamid Hatami Maleki & Mahdi Ayyari (2018): Flavonoids profile and antioxidant activity in flowers and leaves of hawthorn species (*Crataegus* spp.) from different regions of Iran, International Journal of Food Properties, DOI: [10.1080/10942912.2018.1446146](https://doi.org/10.1080/10942912.2018.1446146)

To link to this article: <https://doi.org/10.1080/10942912.2018.1446146>



© 2018 The Author(s). Published by Taylor & Francis.



Accepted author version posted online: 11 Apr 2018.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

Flavonoids Profile and Antioxidant Activity in Flowers and Leaves of Hawthorn Species (*Crataegus* spp.) from Different Regions of Iran

Abolfazl Alirezalu^{a,c}, Peyman Salehi^{b*}, Nima Ahmadi^{c*}, Ali Sonboli^b, Serena Aceto^d, Hamid Hatami Maleki^e, and Mahdi Ayyari^c

^aDepartment of Horticultural Sciences, Faculty of Agriculture, Urmia University, Urmia, Iran

^bMedicinal Plants and Drugs Research Institute, Shahid Beheshti University, Tehran, Iran

*p-salehi@sbu.ac.ir

^cDepartment of Horticultural Sciences, Faculty of Agriculture, Tarbiat Modares University, Tehran, Iran

*ahmadin@modares.ac.ir

^dDepartment of Biology, University of Naples Federico II, Napoli, Italy

^eDepartment of Agronomy and Plant Breeding, Faculty of Agriculture, University of Maragheh, Iran

ABSTRACT

This study was undertaken to determine the total quantity of phenolic and flavonoids, as well as to find out about HPLC quantification of some individual phenolic compounds (*i.e.* chlorogenic acid, vitexin 2"-*O*-rhamnoside, vitexin, rutin, hyperoside, quercetin, and isoquercetin) in flower and leaves of 56 samples of different hawthorn species (*Crataegus* spp.) collected from different geographical regions of Iran. The amount of total phenolics ranges from 7.21 to 87.73 mg GAE/g in dry weight of the plant and total amount of flavonoids varied amongst species and in different plant organs ranging from 2.27 to 17.40 mg/g dry weight. Chlorogenic acid, vitexin and vitexin 2"-*O*-rhamnoside were found to be the most abundant phenolic compounds in the extracts of hawthorn leaves. Meanwhile, chlorogenic acid, hyperoside and rutin were the most abundant

phenolic compounds in the extracts of hawthorn flowers in the most genotypes. The antioxidant activity was widely varied in species and in different organs of each individual plant, ranging from 0.9 to 4.65 mmol Fe⁺⁺/g DW plant, calculated through FRAP method. Thus, this could provide valuable data for developing breeding strategies and plans, by the way it can help us in selecting genotypes with high phenolic contents for producing natural antioxidants and other bioactive compounds beneficial for food or the pharmaceutical industries.

Keywords: Antioxidant, *Crataegus* spp, Flavonoids, HPLC, Phenolics

INTRODUCTION

Wild edible plants including hawthorn have been an indispensable part of human life for ages. Ever since ancient times, their fruits, seeds, leaves, flowers even roots and branches have been used to meet personal and social needs such as severing food, curing diseases and beautifying the planet [1-5]. *Crataegus* which commonly named as hawthorn or thorn-apple is a genus with over 1000 species, belonging to the subfamily of Maloideae in family Rosaceae that is mainly distributed in Asia, Europe and North America [6]. Various species of hawthorn are capable of free hybridization, because they have possessed the base haploid chromosome number of $x=17$. The genus *Crataegus* comprises of a complex group of deciduous shrubs and small trees, which are native to northern temperate regions [7], mostly between latitudes of 30° and 50° N [8]. Hawthorn species are shrubs or small trees, with the height of about 15-18 feet. Various parts of hawthorn including fruits, leaves, flowers, and flowering tops have medicinal properties, which are mostly used as antispasmodic, cardiogenic, diuretic, hypotensive and anti-atherosclerotic

agents [9]. Flavonoids, oligomeric procyanidins and some phenolic acids are considered as the main active constituents of *Crataegus* species [10] with positive effects on heart function and blood circulation [11].

Food antioxidants are useful compounds to neutralize the negative effects of free radicals in the human body, through which the risk of some chronic diseases related to the redox state of the human body reduces [12]. Furthermore, the food industry has widely used natural antioxidants to extend the shelf life of food products [13]. Due to the limited sources of natural antioxidants and their high prices, finding new sources of safe and inexpensive natural antioxidants as substitutes for synthetic antioxidants could definitely be a plausible strategy for the food and pharmaceutical industries, with the purpose of avoiding potential health risks and toxicity [14, 15].

Various organs of hawthorn such as leaves, flowers, and fruits could be excellent source of antioxidants, due to the highly rich phenolic compositions and some well-known antioxidant compounds namely, hyperoside, isoquercetin, epicatechin, chlorogenic acid, quercetin, rutin, and protocatechuic acids. These compounds potentially protect human LDL from Cu^{++} -mediated oxidation. They are also believed to prevent the peroxy free radical-induced oxidation of α -tocopherol in human LDL. Structures of the main phenolic compounds that have already been identified from hawthorn species are shown in Fig. 1. [16-18].

Preharvest environmental conditions, postharvest conditions and processing techniques are key factors which may impacts on antioxidant activity and the chemical compositions of phenolic compounds in leaves and flowers [19]. In addition, level of flavonoids and the quantity of phenolic compounds in plant organs are also affected by genetic variations among different

species, even within the same species and also by the maturity of plant organs at harvest time [20]. Several studies have reported various ranges of phenolic compounds and antioxidant activities based on *Crataegus* accessions and collection regions [21-25].

Apparently, there is a growing interest in utilization of natural antioxidants and their application for nutritional and medical treatments [26, 27]. Iran is known as one of the primary centers of genetic diversity of *Crataegus*; however, few studies have been carried out on phytochemicals of this genus in Iran. The present study was undertaken to determine the total phenolic and flavonoid contents, antioxidant activity, and HPLC quantification of some individual phenolic compounds in the flower and leaves of 56 samples (including 14 species) taken from different hawthorn species (*Crataegus* spp.) that have been collected from different regions of Iran.

MATERIALS AND METHODS

Plant samples

A total of 112 leaves and flowers specimens (including 14 species) were collected from wild growing *Crataegus* genotypes from 11 provinces of Iran (Table 1), in 2014. Individual trees were selected from some genotypes based on their several distinct characteristics. The flowers and leaves were dried at room temperature (20–25°C) after sampling, and then were stored in dry and cool conditions until analysis.

Chemical reagents

The following materials, 2,4,6-Tripyridyl-s-triazine (TPTZ), Folin-Ciocalteu's reagent, aluminum chloride, standard antioxidants, phenolic compound standards (chlorogenic acid, vitexin 2"-*O*-rhamnoside, vitexin, rutin, hyperoside, quercetin, and isoquercetin) and other chemicals used for extraction were obtained from Sigma Co. (USA).

Preparation of the plant extracts

Leaves and flowers of each genotype were dried at room temperature and were grounded to homogenize particle size before extraction. Powdered samples (1 g) were extracted by ultrasound (for 30 min at 25 °C) using methanol/water (80 %, v/v), filtered.

Total phenolic content

The total content of phenolic compounds was determined by the Folin–Ciocalteu method [28]. The extracted samples (0.5 ml of different dilutions) were mixed with Folin Ciocalteu reagent (5 ml, 1:10 diluted with distilled water) for 5 min and aqueous Na₂CO₃ (4 ml, 1 M) was then added. The mixture was allowed to stand for 15 min and the phenolics were determined by spectrophotometer at 765 nm (Bio-Rad's Model). The standard curve ($y=0.0003x-0.0264$; $R^2=0.995$) was prepared by 50, 100, 150, 200, and 250 mg ml⁻¹ solutions of gallic acid in methanol:water (50:50). Total phenolic values are expressed in terms of gallic acid equivalent (mg g⁻¹ DW), which is a common reference compound.

Total flavonoid content

The total flavonoid content of the leaves and flowers extracts was determined using the aluminum chloride colorimetric method with slight modification using quercetin as standard ($y=0.028x-0.0123$; $R^2=0.997$) and the results were expressed as mg of quercetin equivalents per g dry weight of the plant (mg g^{-1} DW). Briefly, the extract solution (0.5 ml) was mixed with 1.5 ml of 80% methanol, 0.1 ml of 10% aluminum chloride hexahydrate (AlCl_3), 0.1 ml of 1 M potassium acetate (CH_3COOK), and 2.8 ml of deionized water. After incubation at room temperature for 30 min, absorbance of the reaction mixture was measured at 415 nm against deionized water blank [29].

Ferric-reducing antioxidant power (FRAP)

Diluted extracts from different organs of hawthorn (100 μl) and 3.0 ml of freshly prepared FRAP-reagent (containing 25 ml of 300 mM acetate buffer, pH 3.6 plus 2.5 ml of 10 mM TPTZ solution in 40 mM HCl plus 2.5 mL of 20 mM $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$) were mixed. The absorbance was recorded at 593 nm against a blank, containing 100 μl of resembling solvent, after 30 min incubation at 37 °C. The FRAP-value was calculated from the calibration curve of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ standard solutions, covering the concentration ranging 100–1000 $\mu\text{mol/L}$ and expressed as mmol Fe^{++} /g dry weight plant [17].

HPLC analysis

The separation of phenolic compounds (chlorogenic acid, vitexin 2"-*O*-rhamnoside, vitexin, rutin, hyperoside, quercetin, and isoquercetin) was performed on a Knauer reversed-phase liquid chromatography apparatus consisting of a 1000 Smartline pump, a 5000 smartline manager solvent organizer and a 2800 Smartline photo- diode array detector. Injection was performed through a 3900 Smartline autosampler injector equipped with a 100 μ l loop. The temperature control of the column was made with a jet stream 2 plus oven (Knauer, advanced scientific instrument, Berlin, Germany). Separation was achieved on an Eclipse XDB-C18 (4.6 mm \times 250 mm, 5 μ m), Agilent (USA) column. Data acquisition and integration was performed with EZChrome Elite software. The flow rate of the mobile phase was kept at 1 mL/min. Solvent A was water containing 0.05% formic acid, and Solvent B was acetonitrile/methanol (80:20, v/v). The gradient conditions were as follows: 0-5 min, 10% B; 5-15 min, 10-18% B; 15-25 min, 18% B; 25-30 min, 18-25% B; 30-35 min, 25% B; 35-40 min, 25-35% B; 40-45 min, 35-60% B; 45-50 min 60-10% B and 50-55 min with 10% B. The temperature of the column was controlled at 25°C. The partial loop injection volume was 10 μ L. The detection wavelengths of DAD were set at three selected positions: 320, 335 and 360 nm.

Preparation of standard solutions

The standard of each phenolic compound was weighed accurately (1 mg) and dissolved in 1:1 MeOH/water in a 10 mL volumetric flask to prepare the stock solution. For calibration curves, the stock solution was diluted by adding MeOH/water (1:4) to obtain the concentration sequence.

10 μ L of each solution was injected into HPLC. The linear range and the equations of linear regression were obtained through a sequence of 1000, 500, 250, 100, 50, 20, 10, 5, 2, and 1 mg/L. Mean areas (n=3) generated from the standard solutions were plotted against concentration to establish calibration equations.

Statistical analysis

All of the analyses were done in triplicate with a factorial experiment based on completely randomized design. SAS 9.1.3 software package (SAS Institute) was used for statistical data analysis. The multivariate ANOVA test and Fisher's Least Significant Difference (LSD) post hoc test were used for means comparison and determination of statistical significance at the $P < 0.05$ probability level. Also, principal component analysis (PCA) and Pearson correlation coefficients were performed by using Minitab 16.2.4 software.

RESULTS AND DISCUSSION

Total phenolic content

The total phenolic content of leaves and flower organs of hawthorn is presented in Table 2. The amount of total phenolic was significantly variable both amongst species and in different plant organs ranging from 7.21 to 87.73 mg GAE/g dry weight plant. Total phenolic content was in its highest value (87.73 mg GAE/g DW) in the flowers of G7 (*C. pseudomelanocarpa*), whereas the lowest level (7.21 mg GAE/g DW) was found in the flowers of G4 (*C. monogyna*). Furthermore, phenolic content reached the highest value (82.74 mg GAE/g DW) in leaves of G1 (*C.*

pentagyna), whereas leaves of G50 (*C. atrosanguinea*) ranked the lowest position (19.98 mg GAE/g DW). Both leaves and flower organs of G7 species (*C. pseudomelanocarpa*) exhibited a high level of total phenolic content which is worthy of consideration.

Results clearly show that total phenolic content is significantly under the influence of both the species and also the type of organs. Accordingly, some studies suggest that the polyphenolic content of plant organs is influenced by species and habitat conditions [30] as well as altitude, light, temperature, and the nutritive available in the soil, which may influence the metabolism of phenylpropanoid [31]. The time of harvest (the stage of maturity) is also a very important factor. Variation in total phenolic of hawthorn due to genetic and climatic factors has been reported in several other studies [21, 22]. Similar results have also been obtained in term of the total phenolic content i.e. 12.8 mg GAE/g DW for *C. monogyna*, [32], 2.9 mg GAE/g DW for *C. pinnatifida*, [33] and 26.4 mg GAE/g DW for *C. monogyna* [34]. In another study, total content of polyphenols in fruits of *C. pinnatifida* was $96.9 \pm 4.3 \text{ mg g}^{-1}$ [35].

Total flavonoid content

Table 2 shows the total flavonoids content in different organs of hawthorn. The amount of total flavonoids was significantly variable both amongst species and in different plant organs ranging from 2.27 to 17.40 mg/g dry weight. Differences between the species and also the parts of plants were highly significant ($p \leq 0.01$). Total flavonoids content was in its highest amount in the flowers (17.40 mg/g DW) of G10 (*C. songarica*), whereas the lowest level was found in the flower of G35 (2.27 mg/g DW, *C. orientalis*). Furthermore, the highest total flavonoids content in the leaves (9.90 mg/g DW) was found in G5 (*C. monogyna*), while the lowest content (3.34

mg/g DW) was measured in G56 (*C. meyeri*). These results showed that in most hawthorn species, flower organs possessed higher total flavonoid content than the leaf organs. In terms of total flavonoid content, flower organs of *C. songarica* contained higher content than the other species.

The total content of flavonoids is influenced by the interaction between varieties and parts of plants. Also environmental factors have a significant contribution to the total flavonoid content in plants [21]. Total flavonoids content found in the present study was similar to those reported from other hawthorn species in previous works i.e. 9.13 mg/g DW for *C. aronia* var. *aronia* leaves [30], 5.3 mg/g DW for *C. atrosanguinea* flowers, 11.8 mg/g DW for *C. curvisepala* flowers, 12.3 mg/g DW for *C. curvisepala* leaves [36] and 1.10 mg/g DW for *C. azarolus* leaves [37].

Antioxidant activity of the hawthorn

The evaluation of antioxidant activity of *Crataegus* species exhibited that these species possess considerable antioxidant potential due to the presence of polyphenolic compounds. The antioxidant activity was widely varied in species and in different organs of the individual organs, ranging from 0.9 to 4.65 mmol Fe⁺⁺/g DW plant (Table 2). The highest antioxidant activity was observed in the leaves of G1 (*C. pentagyna*) as 4.65 mmol Fe⁺⁺/g DW, whereas the lowest activity (0.9 mmol Fe⁺⁺/g DW) was found in the leaves of G18 (*C. azarolus* var. *aronia*). Furthermore, the highest (2.84 mmol Fe²⁺/g DW) and the lowest (0.96 mmol Fe⁺⁺/g DW) antioxidant activity in the flowers were found in G4 (*C. monogyna*) and G6 (*C. meyeri*), respectively.

In this study, several indigenous species of *Crataegus* of from Iran were compared in terms of their antioxidant activity using FRAP method. Results showed that the antioxidant activity through 56 specimens was significantly varied in terms of both different plants organs and species (Table 2).

Chlorogenic acid, hyperoside, rutin, spiraeoside, quercetin 3-glucoside (isoquercetin), quercetin, (-)-epicatechin and procyanidin B2 were suggested to be the compounds with strong radical-scavenging activity in floral bud extracts of hawthorn [38]. The ethanol extract of *C. monogyna* fruits contained higher levels of phenolic compounds and showed greater radical scavenging activities than the aqueous extract of the fruits [34]. Most of the reports regarding antioxidant activity of *Crataegus* species were dealing with fruits, aerial parts or flowers of the plant [39]. Only a recent report of Ozyurek et al. [22], describing antioxidant activity determination of different *Crataegus* species from Turkey, revealed FRAP and total phenols data regarding leaves and flowers separately. In addition to polyphenolic compounds, genetic, climatic conditions, other secondary metabolites such as vitamin C levels and carotenoids, are also involved in antioxidant activity [40]. Furthermore, environmental stresses such as cold and drought increase phenolic compounds and antioxidant activity [41].

Phenolic compounds analyses

The amount of seven phenolic compounds including chlorogenic acid, vitexin 2"-*O*-rhamnoside, vitexin, rutin, hyperoside, quercetin, and isoquercetin were simultaneously analyzed by high performance liquid chromatography. Fig. 2 represents the chromatograms of the above mentioned standards. Tables 3 and 4 summarize the contents of phenolic compounds in all 56

samples analyzed in this study. The amounts of phenolic compounds were significantly variable both amongst species and different plant organs. Chlorogenic acid, vitexin and vitexin 2"-O-rhamnoside were found to be the most abundant phenolic compounds analyzed in the extracts of hawthorn leaves. Meanwhile, chlorogenic acid, hyperoside and rutin were found to be the most abundant phenolic compounds in the extracts of hawthorn flowers in the most of the species. Quercetin was not detected in some species and in other species quercetin was also found in very low quantities both in leaves and flowers.

The G5 species (*C. monogyna*) had the highest level (17.69 mg/g DW) of chlorogenic acid and G17 (*C. azarolus* var. *aronia*) had the lowest level (0.28 mg/g DW) among the leaves of the studied species. *C. monogyna* species had the highest content and *C. azarolus* the lowest content of chlorogenic acid among the species studied. Vitexin was in the highest value (5.51 mg/g DW) in G46 (*C. atrosanguinea*) while the lowest level (0.2 mg/g DW) was found in G19 (*C. curvisepala*) among the leaves of the species. Vitexin was not detected in the leaves of G13 (*C. pseudomelanocarpa*). The G30 species (*C. turkestanica*) had the highest level (4.25 mg/g DW) of vitexin 2"-O-rhamnoside and G19 (*C. curvisepala*) the lowest level (0.03 mg/g DW) among the leaves of the studied species.

In the flowers of the studied species, the G13 species (*C. pseudomelanocarpa*) had the highest level (12.67 mg/g DW) of chlorogenic acid and G36 (*C. curvisepala*) and G55 (*C. pseudoheterophylla*) had the lowest level (0.49 mg/g DW). *C. pseudomelanocarpa* species had the highest content of chlorogenic acid among the all species. The highest amount (8.50 mg/g DW) of hyperoside has been observed in G56 species (*C. meyeri*) and G6 (*C. meyeri*) had the

lowest level (0.09 mg/g DW) among the flowers of the studied species. Rutin was in its highest value (3.64 mg/g DW) in G2 (*C. pseudomelanocarpa*) whereas the lowest level (0.02 mg/g DW) was found in G4 (*C. monogyna*) among the flowers of the studied species. Rutin was not detected in the flowers of G42 (*C. azarolus* var. *aronia*) nor in G52 (*C. atosanguinea*).

The present study shows that the amount of phenolic compounds is significantly under the influence of both the species and also the type of organs [42]. 122 genotypes of *Crataegus* have been investigated in China and it was found that the vitexin 2"-*O*-rhamnoside and rutin were the main flavonoids in hawthorn leaves. Vitexin and quercetin were in the minimum amount and quercetin was not found in some species, which it is similar to our findings. The difference in the amount and type of phenolic compounds in different organs has been observed in other species of hawthorn [16, 43].

Several environmental factors affect the concentration of phenolic compounds in plants. It is reported that higher growing temperatures and the level of CO₂ increase flavonoid content and concentrations of the phenolic compounds [44]. Furthermore, soil conditions affect plant phenolic composition. Soil fertilization factors (such as high level of nitrogen) and deficiency in soil moisture lead to the lower synthesis of phenolics and can decrease the levels of some certain phenolics [45]. Moreover, also light is one of the most effective environmental factors in the phenolic metabolism. Light stimulates the synthesis of phenolic compounds such as flavonoids and flavones, anthocyanins and PAL (phenylalanine ammonia-lyase) enzyme [46].

In general, variability in the contents of phenolic compound and flavonoid concentrations within one species could be mainly associated with differences in growth conditions [31], genetic backgrounds [47], and methodological differences [48].

Principal component analysis (PCA)

PCA multivariate analysis was performed, in order to classify the species studied based on the 20 traits (LTPC, FTFC, LTFC, FTFC, LFRAP, FFRAP, LCHA, LVOR, LVIT, LRUT, LHYP, LISOQ, LQUE, FCHA, FVOR, FVIT, FRUT, FHYP, FISOQ and FQUE). In fact, PCA was applied, to reduce the multidimensional structure of the data and providing a two-dimensional map to explain the variance observed. The first two components of the PCA shows 37% of the total variance (19% for component 1 and 18% for component 2). The first component (PC1) is highly positively correlated with FTFC, FTFC, FCHA, FRUT and FISOQ. The second principal component (PC2) separates the samples according to LTPC, LTFC, LRUT, LISOQ, FVIT and FVOR traits. Generally six genotypes of G1 (*C. pentagyna*), G2, G7, G8, G9 (*C. pseudomelanocarpa*) and G27 (*C. pseudoheterophylla*) formed a single group characterized by higher quantities of phytochemical components which can be considered. Results of PCA showed that the *Crataegus* species collected from different areas of Iran were successfully classified by their, TPC, TFC, antioxidant activity and flavonoids profile (Fig. 3).

Correlations among phytochemical compounds

The analysis of Pearson correlation coefficients showed the highest correlation coefficients between FVOR and FVIT (1.00**) as well as between FTFC and FISOQ (0.68**) followed by

FTFC and FCHA (0.57^{**}) (Table 5). There was a positive and significant correlation between TPC and TFC in both flower and leaves organs. Correlation analysis for phytochemical components with antioxidant activity by FRAP assay of the *Crataegus* species, the ISOQ and CHA, RUT and ISOQ compounds had exhibited positive and significant association with antioxidant activity of leaves and flower, respectively (Table 5). Other correlations are presented in Table 5.

CONCLUSION

To the best of our knowledge, this is the first report regarding antioxidant activity and determination of phenolic compounds (chlorogenic acid, vitexin 2"-*O*-rhamnoside, vitexin, rutin, hyperoside, quercetin, and isoquercetin) in flowers and leaves in *Crataegus* species grown in Iran. Different organs and various species of the genus *Crataegus* specially G1 (*C. pentagyna*), G2, G7, G8, G9 (*C. pseudomelanocarpa*) and G27 (*C. pseudoheterophylla*) showed a high level of total phenolic content as well as antioxidant activity. As a conclusion, our results clearly demonstrate that there are considerable variation in the antioxidant activity and phenolic compounds of hawthorn genotypes. Thus, this could provide valuable data for developing breeding strategies, as well as for selecting genotypes with high phenolic contents when it comes to producing natural antioxidants and other bioactive compounds beneficial in food or pharmaceutical industries.

REFERENCES

1. Ercisli, S. Apricot culture in Turkey. *Scientific Research and Essays*. **2009**, *4*, 715-719.
2. Hricova, A.; Fejer, J.; Libiakova, G.; Szabova, M.; Gazo, J.; Gajdosova, A. Characterization of phenotypic and nutritional properties of valuable *Amaranthus cruentus* L. mutants. *Turkish Journal of Agriculture and Forestry*. **2016**, *40*, 761-771.
3. Ipek, A.; Turkmen, O.; Fidan, S.; Ipek, M.; Karci, H. Genetic variation within the purple carrot population grown in Ereğli District in Turkey. *Turkish Journal of Agriculture and Forestry*. **2016**, *40*, 570-576.
4. Rop, O.; Ercisli, S.; Mlcek, J.; Jurikova, T.; Hoza, I. Antioxidant and radical scavenging activities in fruits of 6 sea buckthorn (*Hippophae rhamnoides* L.) cultivars. *Turkish Journal of Agriculture and Forestry*. **2014**, *38*, 224-232.
5. Zorenc, Z.; Veberic, R.; Stampar, F.; Koron, D.; Mikulic-Petkovsek, M. Changes in berry quality of northern highbush blueberry (*Vaccinium corymbosum* L.) during the harvest season. *Turkish Journal of Agriculture and Forestry*. **2016**, *40*: 855-867.
6. Zhao, H.C.; Tian, B.F. *Hawthorn Flora*, Press: Zhongguo Lin Ye, Beijing, China, **1996**; 366 pp.
7. Mabberley, D.J. 1997. *The plant-book: a portable dictionary of the vascular plants*, Cambridge Univ. Press: Cambridge, UK, **1997**; 858 pp.
8. Phipps, J.B. Biogeographic, taxonomic, and cladistic relationships between east Asiatic and North American *Crataegus*. *Annals of the Missouri Botanical Garden*. **1983**, *70*, 667–700.

9. Nabavi, S.F.; Habtemariam, S.; Ahmed, T.; Sureda, A.; Daglia, M.; Sobarzo-Sánchez, E.; Nabavi, S.M. Polyphenolic Composition of *Crataegus monogyna* Jacq: From Chemistry to Medical Applications. *Nutrients*. **2015**, *11*, 7708–28.
10. Hellenbrand, N.; Sendker, J.; Lechtenberg, M.; Petereit, F.; Hensel, A. Isolation and quantification of oligomeric and polymeric procyanidins in leaves and flowers of Hawthorn (*Crataegus* spp.). *Fitoterapia*. **2015**, *104*, 14–22.
11. Bernatoniene, J.; Kucinskaite, A.; Masteiková, R.; Kalveniene, Z.; Kasparaviciene, G.; Savickas, A. The comparison of antioxidative kinetics in vitro of the fluid extract from maidenhair tree, motherwort and hawthorn. *Acta Poloniae Pharmaceutica*. **2009**, *66*, 415–421.
12. Valko, M.; Leibfritz, D.; Moncol, J.; Cronin, M.T.D.; Mazur, M.; Telser, J. Free radicals and antioxidants in normal physiological functions and human disease. *The International Journal of Biochemistry & Cell Biology*. **2007**, *39*, 44–84.
13. Tlili, N.; Elfalleh, W.; Hannachi, H.; Yahia, Y.; Khaldi, A.; Ferchichi, A.; Nasri, N. Screening of Natural Antioxidants from Selected Medicinal Plants. *International Journal of Food Properties*. **2013**, *16*, 1117–1126.
14. Ortega-Ramirez, L.A.; Rodriguez-Garcia, I.; Leyva, J.M.; Cruz-Valenzuela, M.R.; Silva-Espinoza, B.A.; Gonzalez-Aguilar, G.A.; Siddiqui, W.; Ayala-Zavala, J.F. Potential of medicinal plants as antimicrobial and antioxidant agents in food industry: a hypothesis. *Journal of Food Science*. **2014**, *79*, 129–37.
15. Ramos, A.B.P.; Santos, S.A.O.; Guerra, A.R.; Guerreiro, O.; Freire, C.S.R.; Silva, A.M.S.; Duarte, M.F.; Silvestre, A.J.D. Phenolic composition and antioxidant activity of different

morphological parts of *Cynara cardunculus* L. var. *altilis* (DC). *Industrial Crops and Products*. **2014**, *61*, 460–471.

16. Liu, P.Z.; Kallio, H.; Lü, D.G.; Zhou, C.S.; Yang, B.R. Quantitative analysis of phenolic compounds in Chinese hawthorn (*Crataegus* spp.) fruits by high performance liquid chromatography-electrospray ionization mass spectrometry. *Food Chemistry*. **2011**, *127*, 1370–1377.

17. Zugic, A.; Dordevic, S.; Arsic, I.; Markovic, G.; Zivkovic, J.; Jovanovic, S.; Tadic, V. Antioxidant activity and phenolic compounds in 10 selected herbs from Vrujci Spa, Serbia. *Industrial Crops and Products*. **2014**, *52*, 519–527.

18. Barros, L.; Carvalho, A.M.; Ferreira, I.C. Comparing the composition and bioactivity of *Crataegus monogyna* flowers and fruits used in folk medicine. *Phytochemical Analysis*. **2011**, *22*, 181–8.

19. Kalt, W. Effects of production and processing factors on major fruit and vegetable antioxidants. *Journal of Food Science*. **2005**, *70*, 11–19.

20. Wang, C.Y.; Chen C.T.; Wang, S.Y. Changes of flavonoid content and antioxidant capacity in blueberries after illumination with UV-C. *Food Chemistry*. **2009**, *117*, 426–431.

21. García-Mateos, R.; Ibarra-Estrada, E.; Nieto-Angel, R. Antioxidant compounds in hawthorn fruits (*Crataegus* spp.) of Mexico. *Revista Mexicana de Biodiversidad*. **2013**, *84*, 1298–1304.

22. Ozyurek, M.; Bener, M.; Guclu, K.; Donmez, A.A.; Suzgec-Selcuk, S.; Pirildar, S.; Mericli, A.H.; Apak, R. Evaluation of antioxidant activity of *Crataegus* species collected from different regions of Turkey. *Records of Natural Products*. **2012**, *6*, 263–277.


23. Caliskan, O.; Gündüz, K.; Serçe, S.; Toplu, C.; Kamiloglu, O.; Şengül, M.; Ercişli, S. Phytochemical characterization of several hawthorn (*Crataegus* spp.) species sampled from the Eastern Mediterranean region of Turkey. *Pharmacognosy Magazine*. **2012**, *8*, 16-21.
24. Chang, C.L.; Chen, H.S.; Shen, Y.C.; Lai, G.H.; Lin, P.K.; Wang, C.M. Phytochemical composition, antioxidant activity and neuroprotective effect of *Crataegus pinnatifida* fruit. *South African Journal of Botany*. **2013**, *88*, 432–437.
25. Lee, Y.C.; Chuah, A.M.; Yamaguchi, T.; Takamura, H.; Matoba, T. Antioxidant activity of traditional Chinese medicinal herbs. *Food Science and Technology Research*. **2008**, *14*, 205–210.
26. Kim, S.J.; Min, S.C.; Shin, H.J.; Lee, Y.J.; Cho, A.R.; Kim, S.Y.; Han, J. Evaluation of the antioxidant activities and nutritional properties of ten edible plant extracts and their application to fresh ground beef. *Meat Science*. **2013**, *93*, 715–22.
27. García-Mateos, R.; Aguilar-Santelises, L.; Soto-Hernández, M.; Nieto-Angel, R. Flavonoids and antioxidant activity of flowers of Mexican *Crataegus* spp. *Natural Product Research*. 2013, *27*, 834–836.
28. Ebrahimzadeh, M.A.; Hosseinimehr, S.J.; Hamidinia, A.; Jafari, M. Antioxidant and free radical scavenging activity of *Feijoa sellowiana* fruits peel and leaves. *Pharmacologyonline*. **2008**, *1*, 7–14.
29. Chang, Q.; Zuo, Z.; Harrison, F.; Chow, M. S.S. Hawthorn. *The Journal of Clinical Pharmacology*. **2002**, *42*, 605–612.

30. Orhan, I.; Ozcelik, B.; Kartal, M.; Ozdeveci, B.; Duman, H. HPLC quantification of vitexine-2"-*O*-rhamnoside and hyperoside in three *Crataegus* species and their antimicrobial and antiviral activities. *Chromatographia*. . **2007**, *66*, 153–157.
31. Dixon, R.A.; Paiva, N.L. Stress-induced phenylpropanoid metabolism. *Plant Cell*. **1995**, *7*, 1085–1097.
32. Froehlicher, T.; Hennebelle, T.; Martin-Nizard, F.; Cleenewerck, P.; Hilbert, J.; Trotin, F.; Grec, S. Phenolic profiles and antioxidative effects of hawthorn cell suspensions, fresh fruits, and medicinal dried parts. *Food Chemistry*. **2009**, *115*, 897–903.
33. Zhang, Z.; Chang, Q.; Zhu, M.; Huang, Y.; Ho, W.K.K.; Chen, Z.Y. Characterization of antioxidants present in hawthorn fruits. *The Journal of Nutritional Biochemistry*. **2001**, *12*, 144–152.
34. Bernatoniene, J.; Masteikova, R.; Majiene, D.; Savickas, A.; Kevelaitis, E.; Bernatoniene, R.; Dvorackova, K.; Civinskiene, G.; Lekas, R.; Vitkevicius, K.; Peciura, R. Free radical-scavenging activities of *Crataegus monogyna* extracts. *Medicina*. **2008**, *44*, 706–712.
35. Liu, T.; Cao, Y.; Zhao, M. Extraction optimization, purification and antioxidant activity of procyanidins from hawthorn (*C. pinnatifida* Bge. var. *major*) fruits. *Food Chemistry*. **2010**, *119*, 1656–1662.
36. Amanzadeh, Y.; Khanavi, M.; Khatamsaz, M.; Rajabi, A.; Ebrahimi, S.E.S. Highperformance thin-layer chromatographic fingerprints of flavonoids and phenol carboxylic acids for standardization of Iranian species of the Genus *Crataegus* L. *Iranian Journal of Pharmaceutical Sciences*. **2007**, *3*, 143–152.

37. Bignami, C.; Paolocci, M.; Scossa, A.; Bertazza, G. Preliminary evaluation of nutritional and medicinal components of *Crataegus azarolus* fruits. *Acta Horticulturae*. **2003**, *597*, 95100.
38. Bahri-Sahloul, R.; Ammar, S.; Grec, S.; Harzallah-Skhiri, F. Chemical characterisation of *Crataegus azarolus* L. fruit from 14 genotypes found in Tunisia. *The Journal of Horticultural Science and Biotechnology*. **2009**, *84*, 23–28.
39. Simirgiotis, M.J. Antioxidant capacity and HPLC-DADMS profiling of Chilean Peumo (*Cryptocarya alba*) fruits and comparison with German Peumo (*Crataegus monogyna*) from Southern Chile. *Molecules*. **2013**, *18*, 2061–2080.
40. Materska, M.; Perucka, I. Antioxidant activity of the main phenolic compounds isolated from hot pepper fruit (*Capsicum annuum* L.). *Journal of Agricultural and Food Chemistry*. **2005**, *53*, 1750–1756.
41. Kirakosyan, A.; Seymour, E.; Kaufman, P.B.; Warber, S.; Bolling, S.; Chang, S.C. Antioxidant capacity of polyphenolic extracts from leaves of *Crataegus laevigata* and *Crataegus monogyna* (hawthorn) subjected to drought and cold stress. *Journal of Agricultural and Food Chemistry*. **2003**, *51*, 3973–3976.
42. Zhao, Y.; Su, K.; Wang, G.; Liu, Z.; Dong, W.; Guo, Y. Genetic diversity of flavonoid content in leaf of hawthorn resources. *Pakistan Journal of Botany*. **2014**, *46*, 1543–1548.
43. Merieli, A.H.; Melikoglu, G. Investigation on Turkish *Crataegus* Species. *Acta Pharmaceutica Turcica*. **2003**, *44*, 169–173.

44. Wang, S.Y.; Bunce, J.A.; Maas, J.L. Elevated carbon dioxide increases contents of antioxidant compounds in field-grown strawberries. *Journal of Agricultural and Food Chemistry*. **2003**, *51*, 4315–4320.
45. Keinanen, M.; Julkunen-Tiitto, R.; Mutikainen, P.; Walls, M.; Ovaska, J.; Vapaavuori, E. Trade-offs in phenolic metabolism of silver birch: effects of fertilization, defoliation, and genotype. *Ecology*. **1999**, *80*, 1970–1986.
46. Falcone Ferreyra M.L.; Rius S.P.; Casati P. Flavonoids: Biosynthesis, biological functions, and biotechnological applications. *Frontiers in Plant Science*. **2012**, *3*:222.
47. Prior, R.L.; Cao, G.; Martin, A.; Sofic, E.; McEwen, J.; O'Brien, C.; Lischner, N.; Ehlenfeldt, M.; Kalt, W.; Krewer, G.; Mainland, C.M. Antioxidant capacity as influenced by total phenolic and anthocyanin content, maturity, and variety of *Vaccinium* species. *Journal of Agricultural and Food Chemistry*. **1998**, *46*, 2686–2693.
48. Heinonen, I.M.; Meyer, A.S.; Frankel, E.N. Antioxidant activity of berry phenolics on human low-density lipoprotein and liposome oxidation. *Journal of Agricultural and Food Chemistry*. **1998**, *46*, 4107–4112.

Fig. 1. Structures of the main phenolic compounds identified in hawthorn; vitexin (1), vitexin 2''-*O*-rhamnoside (2), rutin (3), hyperoside (4), isoquercetin (5), quercetin (6), chlorogenic acid (7).



Accepted Manuscript

Fig. 2. HPLC chromatograms of seven phenolic standards (1. chlorogenic acid, 2. vitexin 2"-*O*-rhamnoside, 3. vitexin, 4. rutin, 5. hyperoside, 6. isoquercetin, 7. quercetin).




Fig. 3. Principal component analysis (PCA) of hawthorn species based on the 20 traits. (FTPC; Flower total phenolic content, LTFC; Leaf total phenolic content, FTFC; Flower total flavonoid content, LTFC; Leaf total flavonoid content, FFRAP; Flower ferric-reducing antioxidant power, LFRAP; Flower ferric-reducing antioxidant power, FCHA; Flower chlorogenic acid, FVOR; Flower vitexin 2-O-rhamnoside, FVIT; Flower vitexin, FRUT; Flower rutin, FHYP; Flower hyperoside, FISOQ; Flower isoquercetin, FQUE; Leaf quercitrin, LCHA; Leaf chlorogenic acid, LVOR; Leaf vitexin 2-O-rhamnoside, LVIT; Leaf vitexin, LRUT; Leaf rutin, LHYP; Leaf hyperoside, LISOQ; Leaf isoquercetin and LQUE; Leaf quercitrin).

Fig. 3




Table 1. Sampling locations of the different *Crataegus* specimens studied

Code	Province	Species	Height	Latitude	Longitude	Code	Province	Species	Height	Latitude	Longitude
G1	Semnan	<i>C. pentagyna</i>	1540	36° 02'N	53° 28'E	G2	East Azerbaijan	<i>C. sakranensis</i>	1694	38° 14'N	45° 42'E
G2	Golestan	<i>C. pseudomelanocarpa</i>	409	36° 50'N	54° 47'E	G3	East Azerbaijan	<i>C. turkestanica</i>	1690	38° 14'N	45° 42'E
G3	Golestan	<i>C. pseudomelanocarpa</i>	413	36° 50'N	54° 47'E	G3	East Azerbaijan	<i>C. pseudoheterophylla</i>	1427	38° 10'N	45° 42'E
G4	Mazandaran	<i>C. monogyna</i>	1081	36° 25'N	51° 52'E	G3	East Azerbaijan	<i>C. szovitisii</i>	1426	38° 10'N	45° 42'E
G5	Mazandaran	<i>C. monogyna</i>	119	36°	51°	G3	East Azerbaijan	<i>C. meyeri</i>	126	38°	47°

	aran		2	26°N	51°E	3	ijan		5	49°N	03°E
G6	Mazand aran	<i>C. meyeri</i>	154 1	36° 26°N	51° 51°E	G3 4	East Azerba ijan		128 1	38° 49°N	47° 03°E
G7	Mazand aran	<i>C. pseudomelanocarpa</i>	981	36° 25°N	51° 28°E	G3 5	East Azerba ijan		127 7	38° 49°N	47° 03°E
G8	Mazand aran	<i>C. pseudomelanocarpa</i>	132 0	36° 24°N	51° 33°E	G3 6	East Azerba ijan	<i>C. curvisepala</i>	119 6	38° 50°N	47° 02°E
G9	Mazand aran	<i>C. pseudomelanocarpa</i>	137 1	36° 23°N	51° 32°E	G3 7	East Azerba ijan	<i>C. monogyna</i>	152 5	38° 23°N	47° 14°E
G10	Mazand aran	<i>C. songarica</i>	138 9	36° 23°N	51° 32°E	G3 8	East Azerba ijan	<i>C. atrosanguinea</i>	149 0	38° 23°N	47° 14°E

G1 1	Mazand aran	<i>C. monogyna</i>	138 8	36° 23'N	51° 32'E	G3 9	East Azerba ijan	<i>C. meyeri</i>	149 0	38° 23'N	47° 14'E
G1 2	Mazand aran	<i>C. monogyna</i>	138 9	36° 23'N	51° 32'E	G4 0	East Azerba ijan	<i>C. meyeri</i>	152 4	36° 50'N	54° 47'E
G1 3	Mazand aran	<i>C. pseudomelan ocarpa</i>	139 4	36° 23'N	51° 32'E	G4 1	Kordes tan	<i>C. szovitisii</i>	160 3	35° 23'N	46° 55'E
G1 4	Mazand aran	<i>C. pseudomelan ocarpa</i>	139 5	36° 23'N	51° 32'E	G4 2	Kordes tan	<i>C. azarolus var. aronia</i>	163 2	35° 23'N	46° 55'E
G1 5	Mazand aran	<i>C. songarica</i>	112 3	36° 25'N	51° 31'E	G4 3	Kordes tan	<i>C. szovitisii</i>	163 4	35° 23'N	46° 55'E
G1 6	Mazand aran	<i>C. monogyna</i>	137 1	36° 23'N	51° 31'E	G4 4	Kordes tan	<i>C. atrosanguine</i>	163 3	35° 23'N	46° 55'E

								<i>a</i>			
G1 7	Kogilou ye	<i>C. azarolus</i> <i>var. aronia</i>	160 7	31° 20'N	51° 13'E	G4 5	Kordes tan	<i>C. persica</i>	163 7	35° 23'N	46° 55'E
G1 8	Bakhtiy ari	<i>C. azarolus</i> <i>var. aronia</i>	191 3	31° 33'N	51° 12'E	G4 6	Kordes tan	<i>C.</i> <i>atrosanguine</i> <i>a</i>	164 4	35° 23'N	46° 55'E
G1 9	Bakhtiy ari	<i>C.</i> <i>curvisepala</i>	189 0	31° 26'N	50° 58'E	G4 7	Kordes tan	<i>C.</i> <i>pseudohetero</i> <i>phylla</i>	164 9	35° 24'N	46° 55'E
G2 0	Bakhtiy ari	<i>C. azarolus</i> <i>var. pontica</i>	193 5	31° 22'N	51° 13'E	G4 8	Kordes tan	<i>C. szovitisii</i>	150 6	36° 06'N	46° 20'E
G2 1	Bakhtiy ari	<i>C.</i> <i>curvisepala</i>	185 3	31° 20'N	51° 13'E	G4 9	Kordes tan	<i>C. szovitisii</i>	150 6	36° 06'N	46° 20'E
G2 2	Qazvin	<i>C.</i> <i>pseudohetero</i> <i>phylla</i>	133 0	36° 24'N	50° 33'E	G5 0	West Azerba ijan	<i>C.</i> <i>atrosanguine</i> <i>a</i>	172 8	36° 42'N	45° 56'E

G2 3	Alborz	<i>C. monogyna</i>	181 4	36° 10'N	50° 41'E	G5 1	West Azerbaijan	<i>C. pseudoheterophylla</i>	148 8	37° 27'N	44° 56'E
G2 4	Alborz	<i>C. meyeri</i>	185 0	36° 09'N	50° 42'E	G5 2	West Azerbaijan	<i>C. atrosanguinea</i>	148 8	37° 27'N	44° 56'E
G2 5	Alborz	<i>C. azarolus</i> <i>var. pontica</i>	184 6	36° 09'N	50° 42'E	G5 3	West Azerbaijan	<i>C. azarolus</i> <i>var. aronia</i>	143 2	37° 18'N	45° 07'E
G2 6	Alborz	<i>C. pseudoheterophylla</i>	196 4	36° 10'N	50° 47'E	G5 4	West Azerbaijan	<i>C. monogyna</i>	144 0	37° 29'N	44° 58'E
G2 7	Alborz	<i>C. pseudoheterophylla</i>	198 0	36° 11'N	50° 54'E	G5 5	Lorestan	<i>C. pseudoheterophylla</i>	164 0	33° 56'N	48° 40'E
G2	East Azerbaijan	<i>C. meyeri</i>	143	38°	45°	G5	Lorestan	<i>C. meyeri</i>	164	33°	48°

8	jan		9	10'N	42'E	6	an		3	55'N	41'E
---	-----	--	---	------	------	---	----	--	---	------	------

Accepted Manuscript

Table 2. Level of total phenolic content, total flavonoids and antioxidant activity in flowers and leaves of different hawthorn (*Crataegus* spp.) species

Code	Species	Organ					
		Flower			Leaves		
		Total phenolic content (mg GAE/g DW)	Total flavonoids (mg/g DW)	Antioxidant activity (mmol Fe ²⁺ /g DW)	Total phenolic content (mg GAE/g DW)	Total flavonoids (mg/g DW)	Antioxidant activity (mmol Fe ²⁺ /g DW)
G1	<i>C. pentagyna</i>	52.56±1.85	7.27±0.21	0.45±0.04	82.74±1.17	8.03±0.24	0.56±0.04
G2	<i>C. pseudomelanocarpa</i>	46.94±0.14	10.83±0.32	0.47±0.06	45.21±0.94	7.31±0.25	0.70±0.05
G3	<i>C. pseudomelan</i>	34.28±0.4	12.81±0.2	0.69±0.10	28.72±0.3	5.91±0.11	0.24±0.12

	<i>ocarpa</i>	7	4		2		
G4	<i>C. monogyna</i>	7.21±0.08	4.01±0.36	0.33±0.06	12.41±0.6 3	3.34±0.15	0.24±0.11
G5	<i>C. monogyna</i>	6.79±0.26	4.59±0.17	0.37±0.05	48.90±1.1 8	9.90±0.32	0.23±0.14
G6	<i>C. meyeri</i>	7.61±0.13	3.25±0.26	0.24±0.07	49.16±0.3 6	6.02±0.16	0.75±0.03
G7	<i>C. pseudomelan ocarpa</i>	87.73±1.8 9	9.05±0.27	0.47±0.09	64.63±0.3 4	7.62±0.11	0.99±0.04
G8	<i>C. pseudomelan ocarpa</i>	46.39±2.1 6	8.12±0.11	0.39±0.07	79.41±1.7 3	7.53±0.12	1.16±0.12
G9	<i>C. pseudomelan ocarpa</i>	78.39±0.3 4	13.16±0.3 5	0.57±0.06	57.32±0.2 5	6.24±0.18	0.57±0.07

G1 0	<i>C. songarica</i>	37.56±1.5	17.40±0.2	0.45±0.10	36.07±0.6	4.95±0.31	0.27±0.06
G1 1	<i>C. monogyna</i>	47.78±0.3	6.52±0.25	0.55±0.08	33.88±0.2	6.46±0.28	0.23±0.09
G1 2	<i>C. monogyna</i>	12.23±0.1	6.15±0.18	0.29±0.06	76.74±0.8	4.68±0.16	0.61±0.06
G1 3	<i>C. pseudomelanocarpa</i>	62.89±2.6	12.94±0.1	0.71±0.11	42.12±0.8	5.88±0.12	0.55±0.04
G1 4	<i>C. pseudomelanocarpa</i>	43.76±0.1	7.89±0.25	0.48±0.09	55.17±0.3	8.03±0.14	0.56±0.08
G1 5	<i>C. songarica</i>	25.86±0.2	15.26±0.1	0.44±0.07	36.81±0.6	3.61±0.32	0.48±0.05
G1	<i>C. monogyna</i>	19.63±0.1	7.36±0.19	0.28±0.14	50.90±0.1	7.67±0.18	0.58±0.13

6		4			9		
G1 7	<i>C. azarolus</i> <i>var. aronia</i>	18.88±0.2 4	4.68±0.27	0.43±0.06	37.76±0.9 4	5.17±0.23	0.25±0.14
G1 8	<i>C. azarolus</i> <i>var. aronia</i>	20.29±0.1 3	4.81±0.24	0.61±0.14	36.67±0.2 5	5.75±0.27	0.23±0.05
G1 9	<i>C. curvisepala</i>	38.78±4.5 8	8.31±0.21	0.59±0.08	32.14±1.9 8	6.81±0.22	0.39±0.07
G2 0	<i>C. azarolus</i> <i>var. pontica</i>	18.66±0.1 7	5.47±0.28	0.51±0.07	70.30±3.3 5	5.75±0.19	0.23±0.05
G2 1	<i>C. curvisepala</i>	31.55±0.3 1	7.60±0.32	0.55±0.06	42.59±0.2 1	3.34±0.25	0.68±0.04
G2 2	<i>C. pseudohetero</i> <i>phylla</i>	57.89±1.1 6	6.21±0.25	0.45±0.07	36.70±0.5 4	6.82±0.14	0.25±0.08
G2	<i>C. monogyna</i>	28.98±0.1	6.12±0.19	0.56±0.06	32.34±0.6	6.87±0.18	0.49±0.07

3		0			4		
G2 4	<i>C. meyeri</i>	24.36±0.1 1	10.44±0.1 7	0.43±0.04	26.90±1.8 4	7.26±0.16	0.64±0.11
G2 5	<i>C. azarolus</i> <i>var. pontica</i>	24.08±0.3 2	3.84±0.15	0.32±0.09	31.47±0.3 2	4.86±0.12	0.39±0.16
G2 6	<i>C. pseudohetero</i> <i>phylla</i>	28.93±0.1 8	14.28±0.1 5	0.61±0.11	49.50±0.2 7	7.69±0.19	0.43±0.09
G2 7	<i>C. pseudohetero</i> <i>phylla</i>	43.18±0.2 4	8.88±0.14	0.40±0.08	80.52±2.5 6	6.40±0.32	0.46±0.07
G2 8	<i>C. meyeri</i>	23.38±0.0 9	11.95±0.2 4	0.61±0.07	28.52±0.2 7	6.06±0.15	0.48±0.06
G2 9	<i>C. sakranensis</i>	18.30±0.4 2	7.97±0.23	0.45±0.07	49.41±0.2 8	5.08±0.17	0.24±0.07

G3 0	<i>C. turkestanica</i>	27.54±0.2 4	5.21±0.19	0.41±0.10	28.45±0.5 7	8.92±0.29	0.53±0.08
G3 1	<i>C. pseudohetero phylla</i>	16.20±0.1 1	15.26±0.2 0	0.54±0.04	39.32±0.1 3	4.12±0.22	0.61±0.05
G3 2	<i>C. szovitisii</i>	18.28±0.1 3	7.62±0.25	0.47±0.03	39.72±0.3 4	4.14±0.18	0.24±0.06
G3 3	<i>C. meyeri</i>	28.68±0.1 5	7.42±0.33	0.57±0.09	37.96±0.2 4	5.40±0.22	0.65±0.13
G3 4	<i>C. meyeri</i>	29.82±0.0 3	5.39±0.23	0.61±0.08	44.87±0.1 8	6.61±0.19	0.24±0.11
G3 5	<i>C. orientalis</i>	47.14±0.4 7	2.27±0.19	0.46±0.06	48.23±0.5 5	4.51±0.20	0.66±0.04
G3 6	<i>C. curvisepala</i>	8.27±0.22	3.25±0.14	0.37±0.05	31.96±0.7 3	6.02±0.11	0.25±0.06

G3 7	<i>C. monogyna</i>	28.91±0.1 8	4.68±0.36	0.33±0.04	59.30±0.3 1	3.96±0.15	0.21±0.07
G3 8	<i>C. atrosanguinea</i>	58.89±0.4 4	6.22±0.16	0.50±0.06	26.79±0.3 7	6.21±0.14	0.42±0.06
G3 9	<i>C. meyeri</i>	28.02±0.2 0	4.10±0.19	0.58±0.07	51.90±0.3 6	4.95±0.32	0.49±0.06
G4 0	<i>C. meyeri</i>	60.23±0.7 6	6.69±0.24	0.53±0.06	42.59±0.3 2	5.48±0.22	0.63±0.12
G4 1	<i>C. szovitisii</i>	24.75±0.1 0	8.97±0.29	0.46±0.04	39.99±0.3 2	5.35±0.27	0.43±0.16
G4 2	<i>C. azarolus</i> <i>var. aronia</i>	27.59±0.2 9	8.96±0.31	0.57±0.05	26.65±0.3 0	7.09±0.15	0.46±0.09
G4 3	<i>C. szovitisii</i>	29.64±0.1 6	2.63±0.44	0.55±0.06	32.08±0.3 7	6.82±0.17	0.62±0.07

G4 4	<i>C. atrosanguinea</i>	28.54±0.2 4	10.71±0.3 7	0.41±0.08	62.08±0.1 4	6.93±0.29	0.45±0.06
G4 5	<i>C. persica</i>	27.09±0.2 3	5.13±0.14	0.43±0.07	25.98±0.2 9	5.90±0.31	0.37±0.05
G4 6	<i>C. atrosanguinea</i>	39.63±0.4 4	8.65±0.32	0.48±0.12	25.32±0.4 5	6.30±0.19	0.26±0.04
G4 7	<i>C. pseudoheterophylla</i>	33.58±0.2 0	2.54±0.27	0.39±0.11	44.56±0.2 6	4.88±0.12	0.23±0.08
G4 8	<i>C. szovitisii</i>	23.60±0.1 8	6.60±0.26	0.55±0.12	21.79±0.2 1	6.02±0.15	0.47±0.12
G4 9	<i>C. szovitisii</i>	20.04±0.2 3	3.32±0.11	0.38±0.14	37.18±0.7 3	4.34±0.16	0.64±0.13

G5 0	<i>C. atrosanguine a</i>	29.18±0.3 5	8.74±0.18	0.46±0.08	19.98±0.3 4	4.65±0.32	0.34±0.12
G5 1	<i>C. pseudohetero phylla</i>	12.52±0.3 8	7.49±0.23	0.57±0.06	31.50±0.4 0	4.86±0.21	0.69±0.08
G5 2	<i>C. atrosanguine a</i>	36.33±0.5 0	5.13±0.31	0.34±0.05	46.65±0.2 3	4.97±0.29	0.53±0.04
G5 3	<i>C. azarolus var. aronia</i>	19.26±0.1 7	3.43±0.22	0.48±0.11	31.27±0.2 9	5.68±0.15	0.44±0.11
G5 4	<i>C. monogyna</i>	17.41±0.3 4	3.43±0.16	0.30±0.16	28.74±0.8 0	5.35±0.31	0.23±0.05
G5 5	<i>C. pseudohetero phylla</i>	46.74±0.2 8	3.43±0.13	0.57±0.09	37.65±0.1 9	6.22±0.22	0.70±0.06

G5	<i>C. meyeri</i>	65.73±2.5	12.54±0.1		34.96±0.7		
6		1	9	0.43±0.07	7	3.34±0.18	0.47±0.08
LSD _{5%}		9.16	0.46	0.12	2.42	0.33	0.17

Accepted Manuscript

Table 3. Content of phenolic compounds in leaves of different hawthorn (*Crataegus* spp.) species

Code	Species	Leaves Phenolic Compounds						
		Chlorogenic acid	Vitexin 2"- O- rhamnoside	Vitexin	Rutin	Hyperoside	Isoquercetin	Quercetin
G1	<i>C. pentagyna</i>	1.86±0.03	0.11±0.04	0.47±0.05	2.48±0.02	1.39±0.04	1.36±0.03	0.05±0.01
G2	<i>C. pseudomelanocarpa</i>	1.13±0.03	0.28±0.05	0.70±0.04	1.13±0.06	0.36±0.05	0.64±0.02	0.02±0.00
G3	<i>C. pseudomelanocarpa</i>	1.45±0.04	0.17±0.04	0.39±0.05	0.54±0.03	0.30±0.05	0.45±0.04	0.02±0.00
G4	<i>C. monogyna</i>	5.39±0.03	0.67±0.03	0.22±0.06	0.36±0.03	1.66±0.03	0.35±0.04	-

G5	<i>C. monogyna</i>	17.69±0.06	0.04±0.05	1.61±0.03	1.28±0.05	3.20±0.03	0.75±0.03	0.02±0.00
G6	<i>C. meyeri</i>	11.59±0.05	0.69±0.06	0.78±0.02	0.65±0.06	0.53±0.06	2.37±0.02	0.02±0.00
G7	<i>C. pseudomelanocarpa</i>	1.72±0.03	0.18±0.03	0.41±0.04	0.82±0.04	0.89±0.05	1.53±0.04	0.02±0.00
G8	<i>C. pseudomelanocarpa</i>	2.90±0.04	0.46±0.05	0.29±0.07	0.36±0.04	1.34±0.05	0.85±0.03	0.02±0.00
G9	<i>C. pseudomelanocarpa</i>	1.68±0.05	0.15±0.02	0.60±0.04	0.66±0.05	0.28±0.06	0.38±0.03	-
G10	<i>C. songarica</i>	2.34±0.02	0.54±0.07	0.36±0.05	0.38±0.06	0.75±0.03	0.27±0.04	0.02±0.00

G1 1	<i>C. monogyna</i>	8.61±0.0 6	0.63±0.06	2.98±0.03	0.30±0.05	0.43±0.02	1.42±0.02	-
G1 2	<i>C. monogyna</i>	6.74±0.0 5	1.53±0.03	1.51±0.03	0.49±0.06	0.37±0.05	0.62±0.01	0.02±0.00
G1 3	<i>C. pseudomelanocarpa</i>	3.11±0.0 3	0.04±0.02	-	0.66±0.05	0.45±0.04	0.32±0.03	-
G1 4	<i>C. pseudomelanocarpa</i>	6.21±0.0 4	0.60±0.03	0.68±0.04	0.72±0.03	2.19±0.05	0.90±0.04	-
G1 5	<i>C. songarica</i>	2.14±0.0 5	0.33±0.07	0.38±0.05	0.35±0.01	0.52±0.07	0.22±0.04	-
G1 6	<i>C. monogyna</i>	11.55±0.0 04	0.20±0.05	3.93±0.04	0.47±0.04	1.01±0.03	1.92±0.03	0.02±0.00
G1	<i>C. azarolus var.</i>	0.28±0.0	1.14±0.06	0.35±0.0	0.25±0.0	1.81±0.0	0.67±0.0	0.03±0.0

7	<i>aronia</i>	4		04	05	4	5	1
G1	<i>C. azarolus var.</i>	0.55±0.0		4.08±0.	0.13±0.	2.51±0.0	0.43±0.0	0.02±0.0
8	<i>aronia</i>	5	0.64±0.07	07	06	4	6	0
G1	<i>C. curvisepala</i>	1.91±0.0		0.20±0.	0.45±0.	4.99±0.0	2.46±0.0	
9		5	0.03±0.05	05	03	5	4	-
G2	<i>C. azarolus var.</i>	1.43±0.0		3.00±0.	0.20±0.	3.09±0.0	0.45±0.0	0.02±0.0
0	<i>pontica</i>	4	0.54±0.04	03	06	3	5	0
G2	<i>C. curvisepala</i>	1.06±0.0		4.05±0.	0.12±0.	0.30±0.0	0.16±0.0	
1		6	3.62±0.05	03	05	6	6	-
G2	<i>C.</i>							
2	<i>pseudoheterophylla</i>	0.91±0.0		0.47±0.	0.61±0.	0.33±0.0	0.18±0.0	0.02±0.0
		3	0.78±0.05	05	03	6	4	0
G2	<i>C. monogyna</i>	2.45±0.0		2.52±0.	0.14±0.	0.92±0.0	0.38±0.0	
3		4	3.44±0.03	06	04	5	3	-
G2	<i>C. meyeri</i>	5.26±0.0	3.01±0.06	2.02±0.	0.24±0.	1.07±0.0	0.49±0.0	0.02±0.0

4		7		07	03	3	2	0
G2 5	<i>C. azarolus</i> var. <i>pontica</i>	0.65±0.0 5		2.20±0. 05	0.26±0. 08	0.39±0.0 4	0.19±0.0 2	0.02±0.0 0
G2 6	<i>C.</i> <i>pseudoheterophyll</i> <i>a</i>	6.03±0.0 6		2.77±0. 04	0.23±0. 05	0.99±0.0 5	0.35±0.0 6	0.02±0.0 0
G2 7	<i>C.</i> <i>pseudoheterophyll</i> <i>a</i>	4.07±0.0 6		0.99±0. 05	1.30±0. 07	0.98±0.0 3	0.49±0.0 4	0.03±0.0 0
G2 8	<i>C. meyeri</i>	4.69±0.0 5		0.83±0. 05	0.94±0. 04	0.82±0.0 3	0.46±0.0 3	-
G2 9	<i>C. sakranensis</i>	3.65±0.0 4		1.75±0. 06	0.38±0. 03	0.43±0.0 2	0.23±0.0 2	-
G3 0	<i>C. turkestanica</i>	3.85±0.0 3		2.79±0. 07	0.18±0. 02	1.27±0.0 4	0.63±0.0 4	0.02±0.0 0

G3 1	<i>C. pseudoheterophylla</i>	2.08±0.0 6	0.92±0.06	1.24±0.0 04	0.33±0.0 05	0.51±0.0 5	0.30±0.0 5	-
G3 2	<i>C. szovitisii</i>	0.89±0.0 1	0.99±0.05	1.80±0.0 06	0.62±0.0 05	0.55±0.0 3	0.43±0.0 4	0.02±0.0 0
G3 3	<i>C. meyeri</i>	5.72±0.0 7	0.73±0.04	3.23±0.0 06	0.11±0.0 08	1.50±0.0 5	0.57±0.0 3	0.02±0.0 0
G3 4	<i>C. meyeri</i>	3.91±0.0 6	0.24±0.03	3.51±0.0 04	0.31±0.0 04	1.70±0.0 2	0.98±0.0 4	-
G3 5	<i>C. orientalis</i>	1.54±0.0 5	0.07±0.06	2.67±0.0 04	0.31±0.0 03	0.90±0.0 3	0.54±0.0 3	-
G3 6	<i>C. curvisepala</i>	2.40±0.0 5	4.26±0.03	0.30±0.0 06	0.05±0.0 06	0.40±0.0 4	0.21±0.0 4	0.02±0.0 0
G3 7	<i>C. monogyna</i>	2.19±0.0 7	0.41±0.06	1.61±0.0 05	0.13±0.0 05	1.05±0.0 3	0.61±0.0 5	0.02±0.0 0

G3 8	<i>C. atrosanguinea</i>	3.77±0.0 3	0.24±0.04	4.41±0.03	0.11±0.04	1.44±0.03	0.59±0.03	-
G3 9	<i>C. meyeri</i>	1.96±0.0 6	0.46±0.05	1.95±0.04	0.22±0.03	1.88±0.05	0.98±0.06	0.02±0.00
G4 0	<i>C. meyeri</i>	6.07±0.0 5	0.11±0.08	2.03±0.05	0.28±0.05	1.30±0.08	0.64±0.04	0.02±0.00
G4 1	<i>C. szovitisii</i>	1.30±0.0 3	1.11±0.06	2.93±0.04	0.17±0.06	0.68±0.06	0.27±0.05	0.02±0.00
G4 2	<i>C. azarolus var. aronia</i>	0.81±0.0 8	2.47±0.05	2.08±0.03	0.18±0.04	2.09±0.07	0.57±0.04	0.02±0.00
G4 3	<i>C. szovitisii</i>	1.56±0.0 7	1.19±0.04	2.65±0.02	0.27±0.03	1.38±0.05	0.52±0.03	0.02±0.00
G4 4	<i>C. atrosanguinea</i>	4.51±0.0 6	0.49±0.06	4.02±0.01	0.35±0.06	1.47±0.07	0.69±0.02	0.02±0.00

G4 5	<i>C. persica</i>	1.55±0.0 5	0.29±0.03	4.70±0.07	0.07±0.07	0.39±0.05	0.19±0.04	-
G4 6	<i>C. atrosanguinea</i>	1.96±0.0 2	0.49±0.04	5.51±0.04	0.08±0.04	0.25±0.05	0.13±0.03	0.02±0.00
G4 7	<i>C. pseudoheterophylla</i>	0.78±0.0 5	2.05±0.08	2.34±0.03	0.08±0.03	0.28±0.04	0.14±0.05	-
G4 8	<i>C. szovitisii</i>	1.88±0.0 4	0.40±0.06	1.99±0.04	0.33±0.05	0.76±0.05	0.53±0.05	0.02±0.01
G4 9	<i>C. szovitisii</i>	1.15±0.0 5	0.65±0.03	1.54±0.05	0.22±0.06	0.89±0.04	0.34±0.04	0.02±0.00
G5 0	<i>C. atrosanguinea</i>	2.19±0.0 5	0.16±0.06	3.13±0.06	0.10±0.05	0.71±0.03	0.31±0.03	-
G5 1	<i>C. pseudoheterophylla</i>	5.53±0.0 3	1.20±0.05	0.57±0.07	0.15±0.04	0.72±0.04	0.31±0.05	-

	<i>a</i>							
G5 2	<i>C. atrosanguinea</i>	2.88±0.0 5	0.90±0.06	04	0.05±0.03	1.42±0.05	0.48±0.03	-
G5 3	<i>C. azarolus var. aronia</i>	1.34±0.0 4	0.80±0.07	03	0.30±0.02	1.28±0.06	0.51±0.03	-
G5 4	<i>C. monogyna</i>	1.47±0.0 3	3.31±0.03	04	0.03±0.08	0.22±0.04	0.14±0.04	0.02±0.00
G5 5	<i>C. pseudoheterophylla</i>	5.25±0.0 7	2.89±0.04	04	0.21±0.06	1.04±0.05	0.46±0.05	-
G5 6	<i>C. meyeri</i>	3.45±0.0 4	0.80±0.03	06	0.27±0.03	1.13±0.03	0.38±0.03	-
LSD _{5%}		0.06	0.05	0.04	0.06	0.06	0.09	0.02

Table 4. Content of phenolic compounds in flowers of different hawthorn (*Crataegus* spp.) species

Code	Species	Flower Phenolic Compounds						
		Chlorogenic acid	Vitexin 2"- <i>O</i> -rhamnoside	Vitexin	Rutin	Hyperoside	Isoquercetin	Quercetin
G1	<i>C. pentagyna</i>	2.82±0.04	0.10±0.04	0.32±0.06	2.44±0.03	1.34±0.04	1.25±0.02	-
G2	<i>C. pseudomelanocarpa</i>	6.06±0.03	0.22±0.06	0.55±0.05	3.64±0.04	2.00±0.03	1.96±0.05	0.03±0.01
G3	<i>C. pseudomelanocarpa</i>	7.34±0.04	0.26±0.06	0.63±0.04	3.03±0.03	1.95±0.04	2.28±0.06	0.04±0.01
G4	<i>C. monogyna</i>	1.01±0.03	0.28±0.05	0.68±0.05	0.02±0.05	1.59±0.05	0.24±0.05	0.08±0.00

G5	<i>C. monogyna</i>	1.31±0.0 4	-	0.03±0. 04	0.90±0. 06	2.99±0. 04	0.26±0. 05	-
G6	<i>C. meyeri</i>	4.25±0.0 6	0.35±0. 03	0.81±0. 03	0.43±0. 05	0.09±0. 05	0.44±0. 04	0.02±0. 00
G7	<i>C. pseudomelanocarpa</i>	12.24±0. 08	0.96±0. 06	2.00±0. 03	1.71±0. 04	1.49±0. 03	1.10±0. 06	0.06±0. 00
G8	<i>C. pseudomelanocarpa</i>	5.13±0.0 7	0.05±0. 01	0.21±0. 07	0.18±0. 05	2.70±0. 08	0.89±0. 03	0.07±0. 02
G9	<i>C. pseudomelanocarpa</i>	11.37±0. 09	-	0.09±0. 04	2.09±0. 04	1.26±0. 06	0.94±0. 04	0.09±0. 03
G10	<i>C. songarica</i>	5.22±0.0 7	0.03±0. 02	0.17±0. 06	0.78±0. 05	7.71±0. 05	0.71±0. 06	0.03±0. 01

G1 1	<i>C. monogyna</i>	5.94±0.0 3	0.03±0. 03	0.18±0. 03	0.20±0. 04	2.56±0. 04	0.39±0. 05	0.04±0. 01
G1 2	<i>C. monogyna</i>	3.96±0.0 3	0.02±0. 01	0.15±0. 04	0.24±0. 03	2.14±0. 05	0.39±0. 06	0.03±0. 02
G1 3	<i>C. pseudomelanoca rpa</i>	12.67±0. 10	0.18±0. 02	0.47±0. 03	2.86±0. 06	3.65±0. 05	0.91±0. 03	0.03±0. 01
G1 4	<i>C. pseudomelanoca rpa</i>	3.84±0.0 7	0.01±0. 02	0.15±0. 06	0.44±0. 05	3.94±0. 03	0.55±0. 06	0.03±0. 00
G1 5	<i>C. songarica</i>	5.40±0.0 3	0.05±0. 04	0.23±0. 07	0.88±0. 04	8.01±0. 04	1.01±0. 07	0.02±0. 00
G1 6	<i>C. monogyna</i>	9.00±0.0 8	0.27±0. 03	0.65±0. 04	0.91±0. 06	3.38±0. 03	0.43±0. 04	0.03±0. 01
G1	<i>C. azarolus var.</i>	5.76±0.0	0.04±0.	0.20±0.	1.49±0.	1.65±0.	0.59±0.	0.04±0.

7	<i>aronia</i>	9	04	03	05	05	03	03
G1 8	<i>C. azarolus var.</i> <i>aronia</i>	7.78±0.0 8	0.08±0. 03	0.28±0. 04	1.63±0. 05	2.84±0. 04	0.46±0. 02	0.02±0. 01
G1 9	<i>C. curvisepala</i>	4.04±0.0 3	0.33±0. 06	0.77±0. 03	1.73±0. 06	3.63±0. 03	0.76±0. 05	0.03±0. 01
G2 0	<i>C. azarolus var.</i> <i>pontica</i>	7.77±0.0 6	0.08±0. 06	0.27±0. 05	1.18±0. 04	2.99±0. 04	0.48±0. 08	0.06±0. 00
G2 1	<i>C. curvisepala</i>	2.12±0.0 7	0.27±0. 05	0.64±0. 03	1.17±0. 05	4.33±0. 05	0.67±0. 03	0.06±0. 03
G2 2	<i>C. pseudoheterophylla</i>	1.54±0.0 8	-	0.11±0. 04	1.24±0. 04	3.56±0. 03	0.48±0. 05	0.03±0. 02
G2 3	<i>C. monogyna</i>	2.57±0.0 6	0.26±0. 06	0.64±0. 03	0.65±0. 03	3.35±0. 02	0.70±0. 05	0.03±0. 01
G2	<i>C. meyeri</i>	7.52±0.0	0.44±0.	0.98±0.	1.35±0.	6.67±0.	1.03±0.	0.05±0.

4		5	07	04	05	06	04	01
G2 5	<i>C. azarolus var. pontica</i>	2.63±0.0 7	0.88±0. 06	1.05±0. 03	0.81±0. 04	0.41±0. 05	0.21±0. 05	0.03±0. 01
G2 6	<i>C. pseudoheterophylla</i>	7.99±0.0 7	0.93±0. 05	1.74±0. 04	2.42±0. 03	7.51±0. 03	1.23±0. 03	0.04±0. 00
G2 7	<i>C. pseudoheterophylla</i>	4.43±0.0 6	0.11±0. 06	0.34±0. 05	3.42±0. 05	4.20±0. 04	0.54±0. 05	0.05±0. 00
G2 8	<i>C. meyeri</i>	6.90±0.0 3	0.17±0. 07	0.46±0. 03	1.63±0. 04	7.38±0. 06	1.30±0. 06	0.11±0. 03
G2 9	<i>C. sakranensis</i>	3.16±0.0 4	0.06±0. 05	0.24±0. 07	0.22±0. 05	6.33±0. 05	1.08±0. 03	-
G3 0	<i>C. turkestanica</i>	1.54±0.0 5	0.10±0. 03	0.32±0. 05	0.87±0. 06	3.02±0. 04	0.79±0. 07	0.02±0. 02

G3 1	<i>C. pseudoheterophylla</i>	3.57±0.0 4	0.29±0. 05	0.69±0. 06	2.00±0. 05	8.33±0. 06	1.68±0. 05	0.04±0. 01
G3 2	<i>C. szovitisii</i>	6.09±0.0 3	0.16±0. 04	0.43±0. 04	0.61±0. 06	2.98±0. 05	0.32±0. 04	0.02±0. 01
G3 3	<i>C. meyeri</i>	4.52±0.0 5	0.33±0. 05	0.66±0. 06	0.50±0. 08	5.06±0. 04	0.82±0. 06	0.03±0. 02
G3 4	<i>C. meyeri</i>	2.46±0.0 6	0.13±0. 07	0.38±0. 04	0.23±0. 06	3.15±0. 06	0.71±0. 05	-
G3 5	<i>C. orientalis</i>	1.41±0.0 5	0.13±0. 05	0.37±0. 06	0.39±0. 03	0.85±0. 02	0.42±0. 04	0.02±0. 00
G3 6	<i>C. curvisepala</i>	0.49±0.0 4	0.05±0. 05	0.22±0. 05	0.09±0. 05	2.54±0. 05	0.37±0. 03	0.04±0. 00
G3 7	<i>C. monogyna</i>	1.87±0.0 3	0.04±0. 03	0.21±0. 03	0.12±0. 04	3.64±0. 05	0.61±0. 05	-

G3 8	<i>C. atrosanguinea</i>	2.51±0.0 5	0.19±0. 06	0.29±0. 03	0.39±0. 06	4.78±0. 07	0.57±0. 02	-
G3 9	<i>C. meyeri</i>	2.73±0.0 4	0.01±0. 07	0.15±0. 04	0.76±0. 05	2.43±0. 02	0.58±0. 07	0.04±0. 00
G4 0	<i>C. meyeri</i>	4.69±0.0 6	0.17±0. 08	0.46±0. 05	1.24±0. 03	3.42±0. 04	0.49±0. 05	0.05±0. 01
G4 1	<i>C. szovitisii</i>	7.32±0.1 1	0.16±0. 05	0.43±0. 06	3.16±0. 06	3.91±0. 03	0.70±0. 06	0.02±0. 00
G4 2	<i>C. azarolus var. aronia</i>	4.66±0.0 6	0.16±0. 07	0.43±0. 05	-	6.09±0. 06	0.96±0. 04	0.04±0. 02
G4 3	<i>C. szovitisii</i>	2.63±0.0 5	0.08±0. 03	0.28±0. 06	0.34±0. 03	1.57±0. 05	0.32±0. 06	0.02±0. 01
G4 4	<i>C. atrosanguinea</i>	4.70±0.0 3	0.21±0. 04	0.54±0. 03	0.97±0. 05	7.41±0. 04	1.23±0. 04	0.02±0. 00

G4 5	<i>C. persica</i>	5.05±0.0 4	0.43±0. 06	0.96±0. 05	0.40±0. 05	2.58±0. 05	0.45±0. 03	0.03±0. 01
G4 6	<i>C. atrosanguinea</i>	6.13±0.0 5	0.47±0. 07	1.04±0. 04	0.68±0. 06	5.08±0. 03	0.79±0. 02	0.03±0. 01
G4 7	<i>C. pseudoheterophylla</i>	2.15±0.0 6	0.45±0. 03	1.01±0. 05	0.31±0. 03	0.51±0. 05	0.23±0. 03	-
G4 8	<i>C. szovitisii</i>	6.47±0.0 4	0.27±0. 05	0.65±0. 04	0.23±0. 06	2.03±0. 03	0.85±0. 05	0.05±0. 02
G4 9	<i>C. szovitisii</i>	2.43±0.0 5	-	0.12±0. 05	0.45±0. 07	2.49±0. 04	0.33±0. 03	-
G5 0	<i>C. atrosanguinea</i>	7.09±0.0 3	0.54±0. 06	1.17±0. 06	0.69±0. 05	6.65±0. 06	0.65±0. 06	0.06±0. 03
G5 1	<i>C. pseudoheterophylla</i>	7.19±0.0 6	0.48±0. 05	1.07±0. 08	0.46±0. 06	0.88±0. 04	0.39±0. 06	-

	<i>lla</i>							
G5 2	<i>C. atrosanguinea</i>	1.77±0.0 5	0.02±0. 05	0.17±0. 04	-	4.97±0. 06	0.65±0. 03	-
G5 3	<i>C. azarolus var. aronia</i>	4.88±0.0 7	0.14±0. 08	0.40±0. 05	1.09±0. 05	1.39±0. 05	0.30±0. 07	0.04±0. 01
G5 4	<i>C. monogyna</i>	3.57±0.0 8	0.13±0. 05	0.37±0. 06	0.14±0. 06	2.31±0. 04	0.50±0. 06	0.03±0. 00
G5 5	<i>C. pseudoheterophylla</i>	0.49±0.0 4	0.23±0. 04	0.58±0. 05	0.26±0. 05	1.11±0. 03	0.25±0. 04	0.03±0. 01
G5 6	<i>C. meyeri</i>	11.05±0. 05	0.65±0. 06	1.40±0. 03	1.66±0. 02	8.50±0. 05	1.07±0. 03	0.05±0. 02
LSD5%		0.1	0.04	0.05	0.05	0.07	0.06	0.04

Table 5. Correlation coefficients between total phenolic and flavonoid contents, antioxidant activity and phenolic compounds on studied hawthorn (*Crataegus* spp.) species

Trait	LTPC	FTPC	LTPC	FTPC	LTPC	FTPC	LTPC	FTPC	LTPC	FTPC	LTPC	FTPC	LTPC	FTPC	LTPC	FTPC	LTPC	FTPC	LTPC	FTPC
TPC	1																			
FTPC	0.245	1																		
LTPC	0.338*	0.190	1																	

C	*																		
F T F C	0. 04 3	0. 28 6*	0. 02 3	1															
L F R A P	0. 34 6*	0. 33 3*	0. 16 0	0. 11 3	1														
F F R A P	- 0. 20 7	0. 30 6*	0. 05 2	0. 33 9*	- 0. 00 8	1													
L C H	0. 13	- 0. 21	0. 40 4*	- 0. 04	0. 09 25	1													

A	9	6	*	1	0	5														
L V O R	- 0. 19 5	- 0. 16 0	0. 03 3	- 0. 07 0	- 0. 01 5	- 0. 09 8	- 0. 00 8	1												
L VI T	- 0. 18 6	- 0. 02 6	- 0. 14 4	- 0. 09 2	- 0. 13 3	- 0. 00 6	0. 0 01 4 9	1												
L R U T	0. 51 1* *	0. 25 0	0. 53 0*	0. 20 3	0. 15 3	- 0. 02 7	- 0. 22 7 8 3	0. 38 3*	1											
L H Y	0. 07 9	- 0. 08	0. 34 5*	- 0. 19	- 0. 07	0. 16 8	0. 17 9	- 0. 0 9	0. 06 6	1										

P		4	*	7	3			0	2										
LI S O Q	0. 32 8*	0. 08 1	0. 43 0*	- 0. 12 8	0. 29 9*	- 0. 15 0	0. 45 1*	- 0. 1 9	2 0. 16 5		0. 32 9*	40 9*	1						
L Q U E	0. 35 6*	- 0. 09 6	0. 42 5*	- 0. 04 8	0. 02 7	- 0. 23 5	0. 02 1	0. 0 3	- 0. 07 7	0. 39 9*	0. 06 4	0. 17 0	1						
F C H A	0. 07 3	0. 37 8*	0. 07 1	0. 57 9*	0. 17 3	0. 32 5*	0. 04 3	- 0. 1 8	- 0. 03 0	0. 08 6	0. 12 5	0. 07 7	0. 1 3	1					
F V O	- 0. 22	0. 10 17	- 0. 17	0. 12	0. 13	0. 02	- 0. 04	0. 1 0	0. 27	- 0. 15	- 0. 15	- 0. 03	- 0. 0	0. 31	1				

R	7	7	7	9	2	2	2	7	9*	5	4	1	6	2						
													8							
F VI T	- 0. 22 8	0. 10 8	- 0. 18 5	0. 12 9	0. 13 5	0. 02 4	- 0. 05 4	0. 1 1 0	0. 28 1*	- 0. 16 2	- 0. 15 8	- 0. 03 1	- 0. 6 9	0. 31 2*	1. 00 0*	1				
F R U T	0. 19 2	0. 31 0*	0. 24 7	0. 50 9*	0. 05 9	0. 33 5*	- 0. 14 7	- 0. 0 8	- 0. 23 1	0. 47 2*	- 0. 07 3	- 0. 03 7	0. 2 1 8	0. 46 2*	0. 07 4	0. 0 7 2	1			
F H Y P	- 0. 12 0	- 0. 03 3	- 0. 12 9	0. 54 1*	- 0. 11 6	0. 19 4	0. 03 2	0. 2 0 5	0. 38 1*	- 0. 19 0	0. 08 8	- 0. 17 1	- 0. 1 4 2	0. 12 9 5	0. 23 5 7	0. 2 3 7	0. 13 8	1		

FI	0.	0.	0.	0.	0.	0.	-	-	-	0.	-	-	0.	0.	0.	0.	0.	0.
S	05	25	13	68	14	37	0.	0.	0.	0.	0.	0.	0.	34	11	1	52	42
O	3	4	8	5*	8	5*	18	02	27	15	04	6	4*	3	1	2*	0*	1
Q				*		*	8	3	0*	0	0	6	*	5		*	*	
F	-	0.	-	0.	0.	0.	-	0.	-	-	-	-	0.	0.	0.	0.	0.	0.
Q	0.	18	0.	29	20	21	0.	0.	0.	0.	0.	0.	42	14	1	27	07	18
U	03	8	01	6*	7	3	08	1	03	04	03	0	2*	4	6*	1	0	1
E	9		4				9	5	6	6	9	9	*	6	7			