

SPECIAL ARTICLE

Lung-protective ventilation for the surgical patient: international expert panel-based consensus recommendations

Christopher C. Young^{1,2,*}, Erica M. Harris², Charles Vacchiano^{1,3}, Stephan Bodnar³, Brooks Bukowy³, R. Ryland D. Elliott², Jaclyn Migliarese³, Chad Ragains², Brittany Trethewey³, Amanda Woodward⁴, Marcelo Gama de Abreu⁵, Martin Girard⁶, Emmanuel Futier⁷, Jan P. Mulier⁸, Paolo Pelosi^{9,10} and Juraj Sprung¹¹

¹Department of Anesthesiology, Duke University School of Medicine, Durham, NC, USA, ²Duke University Medical Center, Durham, NC, USA, ³Duke University School of Nursing, Durham, NC, USA, ⁴Duke University Medical Center Library, Durham, NC, USA, ⁵Pulmonary Engineering Group Dresden, Department of Anesthesiology and Intensive Care Medicine, University Hospital Carl Gustav Carus, TU Dresden, Dresden, Germany, ⁶Department of Anesthesiology, Centre Hospitalier de l'Université de Montréal, Montreal, QC, Canada, ⁷Department of Perioperative Medicine, Anesthesiology, and Critical Care Medicine, University Hospital of Clermont-Ferrand, Clermont-Ferrand, France, ⁸Department of Anesthesiology, Intensive Care and Reanimation, AZ Sint Jan Brugge-Oostende, Bruges, Belgium, ⁹Department of Surgical Sciences and Integrated Diagnostics, University of Genoa, Genoa, Italy, ¹⁰Anaesthesia and Intensive Care, San Martino Policlinico Hospital, IRCCS for Oncology and Neurosciences, Genoa, Italy and ¹¹Department of Anesthesiology and Perioperative Medicine, Mayo Clinic, Rochester, MN, USA

*Corresponding author. E-mail: christopher.young@duke.edu

Summary

Postoperative pulmonary complications (PPCs) occur frequently and are associated with substantial morbidity and mortality. Evidence suggests that reduction of PPCs can be accomplished by using lung-protective ventilation strategies intraoperatively, but a consensus on perioperative management has not been established. We sought to determine recommendations for lung protection for the surgical patient at an international consensus development conference. Seven experts produced 24 questions concerning preoperative assessment and intraoperative mechanical ventilation for patients at risk of developing PPCs. Six researchers assessed the literature using questions as a framework for their review. The modified Delphi method was utilised by a team of experts to produce recommendations and statements from study questions. An expert consensus was reached for 22 recommendations and four statements. The following are the highlights: (i) a dedicated score should be used for preoperative pulmonary risk evaluation; and (ii) an individualised mechanical ventilation may improve the mechanics of breathing and respiratory function, and prevent PPCs. The ventilator should initially be set to a tidal volume of 6–8 ml kg⁻¹ predicted body weight and positive end-expiratory pressure (PEEP) 5 cm H₂O. PEEP should be individualised thereafter. When recruitment manoeuvres are performed, the lowest effective pressure and shortest effective time or fewest number of breaths should be used.

Keywords: anaesthesia; adverse effects; Delphi method; intraoperative care; lung injury; perioperative; positive end-expiratory pressure; positive-pressure respiration; postoperative pulmonary complications; tidal volume

Editorial decision: 04 August 2019; Accepted: 4 August 2019

© 2019 The Authors. Published by Elsevier Ltd on behalf of British Journal of Anaesthesia. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

For Permissions, please email: permissions@elsevier.com

Editor's key points

- Expert consensus-based recommendations were produced to reduce pulmonary complications after surgery.
- Low tidal ventilation (6–8 ml kg⁻¹) and PEEP (5 cm H₂O) should be used initially.
- Alveolar recruitment manoeuvres are beneficial in reopening collapsed alveoli and improving lung mechanics.

Postoperative pulmonary complications (PPCs) account for substantial morbidity and mortality. The incidence of PPCs varies according to definition and type of surgery, and has been reported to range from 5% to 33%.^{1,2} The 30 day mortality rate for patients who develop PPCs can be as high as 20%.¹ Recent reviews have highlighted the growing evidence that lung-protective ventilation, consisting of low tidal volumes (V_T), application of PEEP, and use of alveolar recruitment manoeuvres (ARMs), can reduce PPCs.^{3,4} More recently, high ventilator driving pressure (ΔP =plateau pressure [P_{plat}]-PEEP) has been recognised as a significant determinant of lung injury⁵ and is linked to PPCs.⁶ Despite evidence of harm, a large proportion of patients continue to receive high V_T mechanical ventilation with a wide range of PEEP and frequently elevated ΔP .^{7,8}

Many factors may play a role in lung-protective ventilation, yet a consensus in the literature concerning the key clinical question of how to best provide lung protection during mechanical ventilation in surgical patients is lacking. For this reason, a multidisciplinary panel with expertise in perioperative care of mechanically ventilated patients was convened with the aim of developing consensus-based recommendations. As the practice of intraoperative mechanical ventilation varies widely in the published literature and amongst practitioners, a consensus-building approach from experts representing six countries in both Europe and North America was thought to best identify areas of agreement. The panel sought to first produce questions regarding preoperative pulmonary risk assessment and characteristics of intraoperative lung-protective ventilation. The current literature was then reviewed to provide evidence-based guidance in response to the identified questions and, in the absence of sufficient clinical data, an expert opinion was solicited. Subsequently, the panel convened and established consensus-based recommendations using the modified Delphi method. The Delphi method is a consensus-building method that is based upon a structured, iterative communication amongst content experts. The modified method allows for an expert discussion during the final round. Their combined contributions can help resolve complex clinical issues. It was used as a decision tool to efficiently identify best practices in protective lung ventilation whilst allowing for the experts to contribute their distinct perspectives.

Methods**Research/expert teams and main topics**

The president of the coordinating team (CCY) discussed the development of lung-protective-ventilation practice recommendations with the meeting sponsor (GE Healthcare). The meeting sponsor agreed to assist with establishing a consensus conference. The president and sponsor identified

individuals who were subsequently invited to participate in the consensus meeting. The selection criteria for the experts included previous publications in the field of intraoperative ventilation, demonstrated knowledge and interest in lung-protective strategies, and ability to participate in all pre-meeting teleconferences and a 1 day face-to-face meeting.

Seven experts (MGA, EF, MG, EMH, JPM, PP, and JS) from six countries agreed to serve on the panel for this consensus meeting. It has been suggested that between 5 and 10 experts are required for content validation,⁹ and that a 'suitable minimum size' for an expert panel is seven.¹⁰

The coordinating team (CCY and CV) and experts generated, reviewed, and approved 24 questions on perioperative mechanical ventilation (Supplementary Table S1). A content-validity universal agreement was not directly measured. However, the use of participants who have knowledge and interest in the topic increases the content validity of the Delphi method, and the use of successive rounds in the development of the questionnaire likewise improves validity.¹¹

A team of six researchers (SB, BB, RRDE, JM, CR, and BT) evaluated the existing literature for each question. A literature search was conducted in order to identify any additional topics of interest.

Processing literature

Research questions were used as guidance for literature searches conducted by a research librarian (AW). The search strategy combined subject headings and keywords for anaesthesia, surgery or perioperative care, and lung-protective ventilation in adults. A systematic literature search on each subject was performed by searching PubMed, Embase, and the Cochrane Central Register of Controlled Trials from inception to July 18, 2018 (Supplementary Table S2). Observational and experimental studies, and also literature reviews, systematic reviews, and meta-analyses written in English were included. The authors chose to include a clinically important, late-breaking randomised trial in the discussion even though it was not published until June 2019.¹²

All articles were screened and reviewed by teams of two for eligibility based on title and abstract (Fig. 1). Rayyan software (<https://rayyan.qcri.org>) was used as a screening tool to facilitate blind screening within the teams.¹³ Every citation was reviewed by two members of the research team using the same inclusion criteria. Any conflicts in including or excluding articles were resolved through a discussion within the research teams.

Eligible full-text articles were obtained and categorised according to sub-questions developed for each topic. They were summarised and evaluated according to the Grading of Recommendations Assessment, Development and Evaluation (GRADE) system,¹⁴ which systematically evaluates the available literature and focuses on the level of evidence based upon the types of studies included.

Each research team worked with one of the experts to formulate recommendations for their sub-questions based on the available literature and the input of their assigned expert. The quality of the evidence was evaluated according to the GRADE system, and assigned as 'high' (⊠⊠⊠⊠), 'moderate' (⊠⊠⊠), 'low' (⊠⊠), or 'very low' (⊠). The strength of the recommendation was based on judgement of the level of evidence, and reported as weak or strong. Expert and researcher teams produced recommendations for presentation at the face-to-face meeting. When the literature was insufficient to




Fig 1. PRISMA flow diagram. Each search term underwent a systematic review using PubMed, Embase, and the Cochrane Central Register of Controlled Trials from inception to July 18, 2018. Then, 651 eligible full-text articles were obtained and categorised. After question development for each topic, the relevant full-text articles were summarised and evaluated according to the GRADE system; 221 articles were included in the final development of the lung-protective ventilation recommendations. PRISMA, preferred reporting items for systematic reviews and meta-analyses.


provide a recommendation, the expert was asked to provide an opinion (Fig. 2).

Throughout the article, recommendations or statements are referred to by their main topic and sub-question. For example, Topic 1 (pulmonary risk assessment) and Question 1 (factors that increase risk of PPCs) are denoted as (Q1.1). The results of every question are displayed in Tables 1–3.

Consensus meeting

The consensus meeting, held in Frankfurt, Germany on October 1, 2018, was organised according to a modified Delphi methodology referred to as the ‘Amsterdam Delphi method’.¹⁵ The key components of the Delphi method include iteration

(two rounds), controlled acquisition of feedback, and aggregation of responses. The modified Delphi method was chosen because it allowed for expert interaction in the final round. This allowed members of the panel to provide further clarification on some matters and present arguments in order to justify their viewpoints. Anonymity, which is a component of the original Delphi method, was not feasible in this setting, and hence the ‘modified’ method was implemented. After displaying the recommendations, the experts voted their agreement or disagreement. Refraining from voting was not allowed. No discussion was allowed between the experts at this point. If 100% consensus was reached during the first round of voting, the recommendation was accepted without further voting or discussion. When the experts were not in full



agreement, the research team was given 2 min to present the underlying considerations. After this, 5 min of discussion amongst the experts was allowed and the recommendation could be reformulated. A given question could result in a statement rather than a recommendation at the discretion of the expert panel. A final round of voting was conducted using the revised recommendation or statement (Fig. 3). The 'consensus' level during the second round of voting was set at

70% agreement amongst experts. This level of agreement was validated and accepted at previous guideline development conferences, including the 2015 European Association for Endoscopic Surgery consensus meeting on appendicitis¹⁶ and the 2016 American Society for Metabolic and Bariatric Surgery consensus meeting on perioperative management of obstructive sleep apnoea in bariatric surgery.¹⁵

Results

Preoperative risk assessment

Preoperative assessment should include a dedicated score for pulmonary risk evaluation in order to identify patients with greater risk for PPCs (Table 1; Q1.1). Many scoring systems exist to quantify PPC risk, but most are too complex to be clinically useful or lack external validation confirming the accuracy of the score. Although the definition of PPCs was revisited recently with the goal of standardising the criteria, the consensus achieved in that publication does not differ substantially from previous ones.¹⁷ Despite the lack of evidence for the use of a specific prediction score, the patient factors and perioperative characteristics associated with increased PPC risk are well established. The panel agreed that the intraoperative ventilation strategy should be guided by an awareness of the factors that pose the greatest risk: age >50 yr, BMI >40 kg m⁻², ASA physical status >2, obstructive sleep apnoea, preoperative anaemia, preoperative hypoxaemia, emergency or urgent surgery, and ventilation duration >2 h (Table 1; Q1.1).

Intraoperative atelectasis, related changes in lung mechanics, and postoperative pulmonary complications

Atelectasis occurs in roughly 90% of all patients undergoing general anaesthesia and can persist for weeks after operation.^{18,19} Intraoperative atelectasis results in decreased functional residual capacity (FRC), increased heterogeneity of lung expansion, cyclic lung overstretch, and increased ΔP . ΔP is the pressure difference that generates V_T , and can be expressed as the ratio between V_T and respiratory system compliance (C_{RS}).²⁰ Lower intraoperative ΔP values have been associated with a reduction in PPCs,^{21,22} and high ΔP is considered a key mediator of lung injury during positive-pressure ventilation.²³ Therefore, intraoperative ventilation that avoids derecruitment without causing over-distension of alveoli may decrease postoperative pulmonary risk by improving perioperative oxygenation and respiratory mechanics,^{3,24,25} and reducing oxidative stress, inflammatory response, and lung injury.^{26,27}

Induction of anaesthesia

Patient positioning

Supine positioning during induction of anaesthesia causes cephalad displacement of abdominal contents, thereby forcing the diaphragm upwards and compressing dependent lung regions. These changes are attenuated by placing patients in a head-up or ramped position (Table 1; Q3.1). During induction of anaesthesia, particularly in obese individuals, the head-up method produces a longer non-hypoxic apnoea time compared with supine, allowing more time for laryngoscopy.^{28,29} The supine position should be avoided during

Table 1 Recommendations and statements concerning pulmonary risk assessment, case set-up, and ventilation management during anaesthesia induction. CPAP, continuous positive airway pressure; FIO₂, fraction of inspired oxygen; HOB, head of bed; I:E, inspiratory:expiratory; NIPPV, non-invasive positive-pressure ventilation; OSA, obstructive sleep apnoea; PBW, predicted body weight; PPC, postoperative pulmonary complication; P_{plat}, plateau pressure; SpO₂, peripheral oxygen saturation; V_T, tidal volume; ZEEP, zero end-expiratory pressure.

Question	Statement/recommendation	Consensus (%)	Quality of evidence	Strength of recommendation
1.1	A dedicated score should be used for risk evaluation.	100	☒☒☐☐	Strong
	The greatest risk factors for PPCs include age >50 yr, BMI >40 kg m ⁻² , ASA >2, OSA, preoperative anaemia, preoperative hypoxaemia, emergency or urgent surgery, ventilation duration >2 h, and intraoperative factors (such as haemodynamic impairment and low oxyhaemoglobin saturation).	100	☒☒☐☐	Statement
1.2	Use a low-tidal-volume protective-ventilation strategy (6–8 ml kg ⁻¹ PBW). ZEEP is not recommended. Appropriate PEEP and recruitment manoeuvres may improve intraoperative respiratory function and prevent PPCs.	86	☒☒☒☐	Strong
1.3	The formation of perioperative clinically significant atelectasis may be an important risk factor for the development of PPCs.	100	☒☒☒☒	Statement
2.1	Individualised mechanical ventilation should be used and may improve intraoperative respiratory function, but the beneficial effects are likely to disappear after extubation.	100	☒☒☐☐	Strong
2.2	The ventilator should initially be set to deliver V _T ≤6–8 ml kg ⁻¹ PBW and PEEP=5 cm H ₂ O. Evidence regarding I:E ratio settings is lacking.	86	☒☒☒☐	Strong
2.3	PEEP should be individualised to the patient in order to avoid increases in driving pressure (P _{plat} –PEEP) whilst maintaining a low V _T . To optimise intraoperative respiratory function in obese patients, during pneumoperitoneum insufflation, and during prone or Trendelenburg positioning, PEEP adjustment may be required.	100	☒☒☐☐	Strong
3.1	Before induction of anaesthesia, position the patient with the HOB elevated ≥ 30 deg (i.e. 'beach chair'); avoid flat supine position. If not contraindicated, before the loss of spontaneous ventilation, use NIPPV or CPAP to attenuate anaesthesia-induced respiratory changes.	100	☒☒☒☐	Strong
3.2	During induction, monitor for an obstructive breathing pattern and use a combination of appropriate techniques, including positioning, application of NIPPV or CPAP, or placement of a nasopharyngeal airway to avoid upper airway obstruction.	100	☒☐☐☐	Strong
3.3	After intubation, FIO ₂ should be set to ≤0.4. Thereafter, use the lowest possible FIO ₂ to achieve SpO ₂ ≥94%.	100	☒☐☐☐	Weak
3.4	No specific mode of controlled mechanical ventilation is recommended.	100	☒☐☐☐	Statement

anaesthesia induction, as 30 degree head-up and reverse Trendelenburg position is associated with less reduction of FRC.³⁰

Non-invasive ventilation during induction

Non-invasive positive-pressure ventilation (NIPPV) or continuous positive airway pressure (CPAP) should be considered as useful adjuncts during anaesthesia induction. Contraindications, such as altered mental status, certain procedures (face/nose/oesophageal resection), or emergency procedures, should be considered before applying NIPPV or CPAP (Table 1; Q3.1). Head-up positioning combined with NIPPV/CPAP³⁰ further attenuates FRC decrease with anaesthesia induction. Using NIPPV/CPAP during induction increases P_aO₂ and duration of non-hypoxic apnoea.^{29,31–33} Two meta-analyses of obese patients corroborated the finding that NIPPV/CPAP

during induction improved duration of non-hypoxic apnoea³⁴ and improved oxygenation.³⁵ A single study failed to demonstrate positive effects of NIPPV/CPAP on non-hypoxic apnoeic time.³⁶ NIPPV/CPAP was also noted to decrease venous admixture when compared with spontaneous breathing.³¹ Other methods, including monitoring of obstructive breathing, head positioning, and naso- or oropharyngeal airway insertion should be used to avoid upper airway obstruction during induction (Table 1; Q3.2).

Optimal intraoperative ventilator settings

Tidal volume

Low V_T ventilation, 6–8 ml kg⁻¹ predicted body weight (PBW), is a fundamental component of lung-protective ventilation (Table 1; Q1.2). Multiple studies have demonstrated a significant reduction in PPCs associated with low (<8 ml kg⁻¹) vs high

Table 2 Recommendations and statements concerning respiratory system monitoring and ventilation management during anaesthesia maintenance/surgery. ESA, European Society of Anaesthesiology; P_{plat} , plateau pressure; FIO_2 , fraction of inspired oxygen.

Question	Statement/recommendation	Consensus (%)	Quality of evidence	Strength of recommendation
4.1	In addition to standard monitoring (ASA/ESA), dynamic compliance, driving pressure ($P_{\text{plat}} - \text{PEEP}$), and P_{plat} should be monitored on all controlled mechanically ventilated patients.	100	☒☒☒☐	Strong
4.2	Decreasing compliance caused by surgical/ anaesthesia factors (i.e. pneumoperitoneum, positioning, and circuit disconnect) should be treated by appropriate interventions. Individualised PEEP can prevent progressive alveolar collapse. Recruitment manoeuvres can reverse alveolar collapse, but have limited benefit without sufficient PEEP. Statement: Increasing FIO_2 may be effective in increasing the oxygenation, but is not an effective intervention to improve dynamic compliance of the respiratory system.	86	☒☒☒☒	Strong
4.3	The effectiveness of interventions aimed at optimising respiratory system mechanics should be evaluated by measuring an improvement of the respiratory system compliance under a constant tidal volume.	100	☒☒☐☐	Strong

Table 3 Recommendations and statements concerning recruitment manoeuvres and ventilation management during anaesthesia emergence. ARM, alveolar recruitment manoeuvre; CPAP, continuous positive airway pressure; FIO_2 , fraction of inspired oxygen; HOB, head of bed; NIPPV, non-invasive positive-pressure ventilation; P_{plat} , plateau pressure; SpO_2 , peripheral oxygen saturation; ZEEP, zero end-expiratory pressure. *Consensus level <70%.

Question	Statement/recommendation	Consensus (%)	Quality of evidence	Strength of recommendation
5.1	High-quality supportive evidence is lacking to recommend a routine ARM for all patients after tracheal intubation. However, an ARM may be considered according to an individual risk-benefit assessment.	57*	☒☐☐☐	Weak
5.2	The bag-squeezing ARM should be avoided in favour of a ventilator-driven ARM.	100	☒☐☐☐	Weak
5.3	ARMs should be performed using the lowest effective P_{plat} (30–40 cm H_2O in non-obese; 40–50 cm H_2O in obese) and shortest effective time or fewest number of breaths.	100	☒☒☐☐	Weak
5.4	Continuous haemodynamic and oxygen saturation monitoring is recommended before and during an ARM. Ensure adequate haemodynamic stability before performing an ARM. Avoid ARMs when contraindicated.	100	☒☒☒☒	Strong
5.5	PEEP should be individualised after an ARM to avoid both alveolar overdistention and collapse.	71	☒☒☐☐	Weak
6.1	Optimise patient positioning and avoid ZEEP during emergence. Avoid tracheal tube suctioning immediately before tracheal extubation.	100	☒☐☐☐	Weak
6.2	Avoid apnoea with ZEEP before extubation.	100	☒☐☐☐	Weak
6.3	In the appropriate clinical scenario, the use of low FIO_2 (<0.4) during emergence from general anaesthesia can improve pulmonary function in the postoperative period.	71	☒☐☐☐	Weak
6.4	When high FIO_2 (>0.8) is used during emergence, the use of low FIO_2 (<0.3) CPAP immediately after tracheal extubation may reduce the risk of resorption atelectasis.	29*	☒☐☐☐	Weak
6.5	Administration of postoperative supplemental oxygen is recommended when room air SpO_2 decreases below 94%. Avoid routine application of supplemental oxygen without investigating and treating the underlying cause.	100	☒☐☐☐	Weak
6.6	Prophylactic NIPPV/CPAP should be considered after operation for patients with prior routine use of NIPPV/CPAP.	100	☒☒☐☐	Strong




Fig 3. Modified Delphi process flow chart. After the development of recommendations, the experts met in a face-to-face meeting to develop a consensus. All recommendations and statements underwent two rounds of voting as no recommendation achieved 100% consensus during the first round. The final round of voting was conducted using the revised recommendation or statement.

(>8 ml kg⁻¹) V_T ventilation.^{26,37,38} However, the use of a low V_T without adequate PEEP may increase the risk of atelectrauma as a result of cyclic lung de-recruitment.^{37,39}

End-expiratory pressure

A number of studies have suggested the negative effects associated with mechanical ventilation with zero end-expiratory pressure (ZEEP).^{4,40–43} These effects include a profound reduction in end-expiratory lung volume (EELV) after anaesthesia induction and an increased area of atelectasis. Loss of EELV and atelectasis contribute to decreased C_{RS} in de-recruited areas, and increase the propensity for overinflation of aerated lung tissue (volutrauma).^{40,41} Therefore, allowing airway/alveolar pressure to achieve ZEEP is not recommended (Table 1; Q1.2).

Individualised PEEP improves oxygenation, EELV, and respiratory system mechanics during ventilation; however, these improvements may disappear soon after extubation.^{44–51} Whilst the panel noted that many measurable effects of lung-protective ventilation may dissipate after extubation,

they agreed that mechanical ventilation should be targeted to optimise the respiratory function (Table 1; Q2.1), and that more studies are needed to quantify whether these positive intraoperative effects on ventilatory mechanics have a clinically meaningful impact on postoperative respiratory outcomes.

Although several studies of low V_T (6–8 ml kg⁻¹) have consistently shown improvement in pulmonary function and reduction of PPCs, the optimal level of PEEP remains a matter of debate.^{4,25,52,53} The panel agreed that lung-protective ventilation requires a combination of low V_T and some degree of PEEP (Table 1; Q2.2). Multiple studies demonstrate that the use of PEEP improves EELV; increases oxygenation; and improves dependent lung ventilation, C_{RS} , and postoperative pulmonary function when compared with ZEEP.^{39,54–58} Moreover, several large RCTs showed that intraoperative ventilation with reduced V_T (6–8 ml kg⁻¹) and increased levels of PEEP (6–10 cm H₂O) prevents PPCs^{38,59,60}; reduces atelectasis and recruitment/de-recruitment injury; and improves C_{RS} , EELV, PaO₂, and dependent lung ventilation with little-to-no overdistension.^{43,57} However, one large trial protective ventilation

during general anesthesia for open abdominal surgery: high versus low positive end-expiratory pressure (PROVHILO) showed no difference in the development of PPCs with low V_T and either high or low levels of PEEP (≤ 2 cm H₂O vs 12 cm H₂O).⁶¹ Whilst ZEEP is not recommended, the precise level of PEEP remains controversial.^{42,43,57,58,61–67}

Individualised PEEP has demonstrated many benefits to pulmonary function, and is especially important in obese patients, during abdominal insufflation, and during prone or Trendelenburg positioning (Table 1; Q2.2). One RCT of obese patients (BMI >35 kg m⁻²) undergoing laparoscopic surgery found that the average calculated individualised PEEP was 18.5 cm H₂O.⁴⁵ This trial also found that individualised PEEP decreased ΔP and increased PaO₂/FIO₂ ratios, EELV, C_{RS}, and ventilation to dependent lung regions. A recent large international trial, however, showed that, although higher PEEP with recruitment manoeuvres results in improved pulmonary function intraoperatively compared with a low PEEP without recruitment manoeuvres, it does not reduce the incidence of PPCs in obese surgical patients.¹²

The importance of individualised PEEP was further highlighted in a meta-analysis of individual patient data from RCTs comparing intraoperative protective ventilation with conventional ventilation, which found that the benefits of protective ventilation were related to reductions in ΔP rather than to changes in V_T or level of PEEP.⁶ The authors reported that only C_{RS} and ΔP were significantly associated with PPCs, and that their incidence was not affected by the level of PEEP unless it resulted in an increase in ΔP . Therefore, the panel recommends an initial PEEP setting of 5 cm H₂O and thereafter PEEP levels should be individualised (Table 1; Q2.2 and 2.3).

Inspiratory/expiratory ratio

Several studies have compared prolonged inspiratory-to-expiratory (I:E) ratios to the 1:2 ratio commonly used during mechanical ventilation. An I:E ratio of 1:1, which has been characterised as providing a 'balanced stress to time product',⁴ was associated with attenuation of lung damage. Prolonged I:E ratio increases mean airway pressure and concomitantly reduces peak airway pressure. Studies using prolonged inspiratory times have described beneficial effects, including increased C_{RS} and PaO₂, lower alveolar–arterial gradient, and reduced inflammatory markers.^{6,67–72} Given the lack of evidence for a clear benefit of a specific I:E ratio, no recommendation was offered by the panel (Table 1; Q2.2). However, the panel noted that optimisation of inspiratory time for individual patients can be achieved by monitoring parameters, such as oxygenation, C_{RS}, and ΔP .

Intraoperative FIO₂

Increased FIO₂ during mechanical ventilation is administered to prevent or correct hypoxaemia, but may result in hyperoxia.^{73,74} The negative effects of hyperoxia are not clear, but it has been suggested that it may increase oxidative stress, peripheral vascular and coronary artery vasoconstriction, decrease cardiac output, increase resorption atelectasis, and increase the rate of PPCs.^{75–81}

Recommendations for optimal use of oxygen and current evidence regarding the association between hyperoxaemia and clinically relevant outcomes during intraoperative mechanical ventilation are lacking. Few studies have revealed a protective effect of hyperoxaemia,⁸² some report an

association with mortality,⁸³ whilst others show no association with clinically relevant outcomes.⁸³ Therefore, in the absence of evidence, the most prudent course of action during mechanical ventilation is to maintain normoxaemia. SpO₂ monitoring can assist in the detection of hypoxaemia, but during oxygen therapy SpO₂ cannot detect hyperoxia.⁸⁴ Whilst SpO₂ monitoring reduces the incidence of hypoxaemia, it does not improve the overall patient outcomes and does not reduce morbidity and mortality.⁸⁵ Therefore, once the airway is secured, FIO₂ should be set to ≤ 0.4 with the goal of using the lowest possible FIO₂ to achieve normoxia (or SpO₂ $\geq 94\%$) (Table 1; Q3.3). Unnecessarily high FIO₂ should be avoided. Administering lower FIO₂ will not only decrease the risk of hyperoxia, but will also reduce the masking effect of oxygen therapy and allow for earlier diagnosis of gas-exchange impairment.⁸⁴

Modes of mechanical ventilation

A number of studies explored whether one mode of mechanical ventilation is better than others at reducing PPCs. When assessing pressure-controlled ventilation (PCV) vs volume-controlled ventilation (VCV), the results are mixed. VCV was associated with lower maximal plateau pressures, greater V_T , and less dead-space ventilation.⁸⁶ In an observational study, the risk of PPC was higher in patients who received PCV compared with VCV, particularly with PEEP <5 cm H₂O.⁸⁷ A meta-analysis regarding intraoperative ventilation mode in obese patients found VCV to be superior to PCV.⁸⁸

Pressure-controlled ventilation was superior to VCV on the basis of lower peak inspiratory pressure (PIP) or improved arterial blood gas (ABG) results in several studies. Four studies showed lower PIP with no change in arterial oxygenation.^{89–92} Another demonstrated improved ABG results in patients ventilated with PCV compared with VCV, with no change in airway pressures.⁹³ No significant differences between PCV and VCV were found in one randomised trial when assessing airway pressures, ABG results, or oxygenation.⁹⁴ VCV with an inspiratory pause does allow for measurement of P_{plat}, therefore allowing for a more accurate determination of ΔP . Given the heterogeneity of the published trials, no specific mode of controlled mechanical ventilation is recommended (Table 1; Q3.4).

Alveolar recruitment manoeuvres

General anaesthesia promotes the formation of atelectasis, which negatively impacts respiratory function and may be associated with subsequent PPCs.^{18,44} ARMs are beneficial in reopening collapsed alveoli and improving lung mechanics, suggesting that performing an ARM after intubation can combat anaesthesia-induced FRC changes.^{45,95–100} Even after an ARM, normal alveoli filled with 100% oxygen have a rapid tendency to collapse and form shunt.¹⁰¹ Therefore, resorption atelectasis can be attenuated with an ARM performed with FIO₂ <1.0 .¹⁸ After an ARM, C_{RS} and PaO₂ improved.^{24,60,102–104} ARMs are effective when applied after intubation and during any episodes of oxyhaemoglobin desaturations or release of positive pressure from the breathing circuit.

The period immediately after induction can often be a time of haemodynamic instability caused by medication and positive-pressure ventilation effects. Whilst ARMs are considered safe and effective,¹⁰⁵ some patients, such as those with hypovolaemia, severe emphysema, or chronic

obstructive lung disease, may be prone to hypotension during an ARM; therefore, the risk to benefit ratio of ARMs should be carefully considered. High-quality supportive evidence is lacking to recommend a routine ARM for all patients after tracheal intubation. However, an ARM may be considered according to an individual risk–benefit assessment (Table 3; Q5.1). Further research is needed to identify which patients would benefit from an ARM immediately after induction.

ARMs should be performed after a disconnection from the circuit and whenever the patient's SpO₂ is consistently $\leq 94\%$. The two primary methods are manual ARM and ventilator-driven ARM.

Manual alveolar recruitment manoeuvres

A manual ARM is performed by sustained lung inflation using the reservoir bag on the anaesthesia machine with the adjustable pressure-limiting valve set to the desired inflation pressure. The manual ARM can lead to brief loss of positive pressure when switching back to the ventilator circuit, which results in recollapse of alveoli. For this reason, the ventilator-driven ARM is favoured (Table 3; Q5.2).

Ventilator-driven alveolar recruitment manoeuvres

Ventilator-driven ARMs can be divided into three types: vital capacity, pressure-controlled, or volume-controlled cycling manoeuvres. The vital-capacity ARM resembles the manual ARM except that the V_T is delivered through the ventilator circuit. This requires a ventilator capable of providing CPAP or an inspiratory hold of 7–8 s.¹⁰⁶ The panel concurs that 7–8 s is an appropriate inspiratory time in patients with healthy lungs, but that individual patient characteristics (elevated BMI, Trendelenburg position, and abdominal insufflation) may require longer times and higher PIP. Studies that have evaluated intraoperative alveolar collapse have found that, in healthy patients with BMIs $< 35 \text{ kg m}^{-2}$, a PIP hold of 40 cm H₂O is required to improve PaO₂ and lung compliance.¹⁰⁷ For patients with BMIs $> 35 \text{ kg m}^{-2}$, pressures of up to 50 cm H₂O or multiple, successive ARMs have been recommended.^{51,62,90,102,108–113} The recently published effect of intraoperative high positive end-expiratory pressure (PEEP) with recruitment maneuvers vs low PEEP on postoperative pulmonary complications in obese patients (PROBESE) trial showed no reduction of PPCs when an ARM was performed after intubation and each hour afterwards as part of a non-individualised ventilator protocol in obese surgical patients.¹²

In pressure-controlled-mode ARM, recruitment airway pressure should be based upon patient BMI, as discussed previously, and this 'opening' pressure should be maintained for 10 breaths.^{21,45,114–118} The panel was unanimous in urging caution when using PIP $> 50 \text{ cm H}_2\text{O}$. When using volume-controlled mode for ARM,⁶⁰ one should start with a V_T of 6–8 ml kg⁻¹ PBW and I:E ratio of 1:1, and increase the V_T by 4 ml kg⁻¹ every three to six breaths until P_{plat} of 30–40 cm H₂O is reached. After an additional three to six breaths at this level, sufficient recruitment occurs and V_T settings can be reduced. PEEP adjustment after an ARM may be required to maintain alveolar recruitment. The panel further recommends that one should evaluate change in C_{RS} and ΔP after an ARM, and repeat the ARM with a longer inspiratory hold or higher pressure if recruitment is assessed as ineffective.

The panel recommends using the lowest FIO₂ during ARMs to aid in identifying the patient's opening and closing pressures, and sustain recruited alveoli by reducing the occurrence of resorption atelectasis.^{119,120} They also state that the method used to produce an ARM through the ventilator circuit is not as important as avoiding the use of manual ARMs. ARMs should be performed using the lowest effective PIP and shortest effective time or fewest number of breaths (Table 3; Q5.3). ARM effectiveness can be measured by improved oxygenation, C_{RS}, or ΔP . Further research is required, as there is currently little evidence linking ARMs to pulmonary outcomes.

Complications related to alveolar recruitment manoeuvres

Hypoxaemia and haemodynamic instability are reported complications of ARMs. No adverse effects of performing an ARM were found in 26 of 33 studies.^{24,51,57,59,60,62,65,95–100,102,104,108,111–114,116–118,121–123} Six studies identified transient haemodynamic instability requiring vasopressor treatment during ARMs.^{21,45,46,52,61,103} One study found more oxyhaemoglobin desaturation in the ARM group.¹²⁴ The panellists recommend continuous haemodynamic and SpO₂ monitoring before and during the ARM.¹²⁵ It is essential to ensure adequate haemodynamic stability before performing an ARM and avoid ARMs when contraindicated (Table 3; Q5.4).

Intraoperative monitoring of lung mechanics and oxygenation

Because the lung is a dynamic system, altered by both anaesthesia and surgery, the components of the mechanical breath should be continuously evaluated.²⁰ C_{RS}, ΔP , and P_{plat} should be monitored on all mechanically ventilated patients (Table 2; Q4.1), and interventions aimed at optimising respiratory system mechanics should be evaluated by measuring C_{RS} under constant V_T.⁶ (Table 2; Q4.3).

Current monitoring standards focus primarily on detecting hypoxaemia using SpO₂. Interventions tend to focus more on improving SpO₂, often by increasing FIO₂, rather than improving the underlying pulmonary system derangement. Whilst increasing FIO₂ may be effective in increasing oxygenation, it does not improve the underlying ventilation–perfusion mismatch (Table 2; Q4.2).

To minimise the risk associated with mechanical ventilation, the ventilator should be set to maintain the ΔP as low as possible.⁶ Appropriately set PEEP can maintain FRC without causing gross over-distension, and is evidenced by the lowest ΔP that achieves the desired V_T.¹²⁶ Surgical or anaesthesia factors that cause changes in C_{RS} or ΔP should be treated by interventions that restore physiological lung volume whilst avoiding both over- and under-distention (Table 2; Q4.2). During controlled mechanical ventilation, if the circuit is disconnected or switched from the ventilator to the manual mode, loss of lung volume will occur immediately, accompanied by a decrease in C_{RS} and an increase in ΔP .¹²⁷ In order to restore C_{RS} and prevent lung over-distension, FRC must be re-established by an increase in pressure sufficient to overcome the degree of lung collapse.^{50,127}

The FRC is maintained, not restored, by PEEP. Therefore, in order to prevent lung over-distension related to PEEP, FRC should be restored with an ARM before any increase in the

level of set PEEP.¹²⁷ Likewise, ARMs can reverse alveolar collapse, but the benefit will be of short duration without sufficient PEEP (Table 2; Q4.2). PEEP should be individualised after an ARM to avoid alveolar over-distension or collapse (Table 3; Q. 5.5).

Emergence from anaesthesia

Consideration should be given to avoiding conditions during emergence that negate the intraoperative efforts to recruit and maintain an open lung. Recommendations similar to those applied during induction include optimising patient positioning (head elevated ≥ 30 deg) and avoiding ZEEP (Table 3; Q6.1). Reduction of lung volume by routine suctioning of the tracheal tube just before extubation should be avoided. Other interventions likely beneficial include prevention of coughing and bucking on the tracheal tube, and avoiding upper airway obstruction after extubation. The common practice of turning off the ventilator allowing carbon dioxide to accumulate to stimulate spontaneous ventilation should also be avoided, as the period of apnoea is associated with ZEEP and collapse of alveoli (Table 3; Q6.2). Atelectasis that develops during general anaesthesia persists into the postoperative period. This finding argues for some methods of keeping recruited alveoli open, such as application of CPAP during the transition between mechanical ventilation and spontaneous breathing. However, applying an ARM followed by PEEP, and then maintaining positive airway pressure using CPAP from return of spontaneous breathing until extubation did not improve postoperative oxygenation.¹²²

FIO₂ during emergence

FIO₂ >0.8 during emergence significantly increases atelectasis formation.^{128–131} If clinically appropriate, FIO₂ ≤ 0.4 during emergence may be used to reduce atelectasis. Lower FIO₂ during emergence can improve postoperative pulmonary function¹³⁰ (Table 3; Q6.3). CPAP with low FIO₂ (<0.3) after extubation may decrease the area of atelectasis.^{31,123,130,132} However, current evidence regarding efficacy of this technique is lacking and cannot presently be universally recommended (Table 3; Q6.4). After extubation, supplemental oxygen should be administered for SpO₂ <94%; however, the underlying cause should be investigated and appropriate interventions should be used (Table 3; Q6.5).

Non-invasive ventilator support

A systematic review of CPAP administered after a major abdominal surgery found weak evidence that CPAP may reduce atelectasis, the rate of pneumonia, and the frequency of reintubation.¹³³ Prophylactic postoperative CPAP reduced the incidence of PPCs in patients undergoing abdominal surgery; however, the authors noted that the optimum CPAP in this setting is unknown and the administration of CPAP should be individualised.¹³⁴ Postoperative CPAP of 7.5 cm H₂O vs 6 L min⁻¹ flow of 50% oxygen by the Venturi mask may reduce reintubation rate, pneumonia, infection, and sepsis after a major abdominal surgery.¹³⁵ CPAP of 10 cm H₂O after thoracoabdominal surgery reduced PPCs and decreased the duration of ICU and hospital stay.¹³⁶

Administration of CPAP immediately post-extubation in the obese population has been shown to reduce atelectasis,

improve oxygenation and pulmonary function, and may minimise the risk of developing PPCs.^{66,137} The early postoperative use of NIPPV in obese patients promoted a more rapid recovery of lung function and improved oxygenation when compared with a 6 L min⁻¹ flow of 50% oxygen via Venturi mask.¹³⁸ In addition, the PaO₂ and PaO₂/FIO₂ ratio were significantly improved up to 24 h after operation when CPAP was applied immediately upon extubation in obese patients.¹³⁹ In obese patients undergoing laparoscopic surgery, NIPPV administration post-extubation improved pulmonary function and reduced the risk of respiratory complications; however, it did not reduce the risk of reintubation or unplanned ICU admission.³⁵

The postoperative prophylactic use of NIPPV or CPAP should be considered for patients who use these modalities to maintain adequate ventilation before operation (Table 3; Q6.6).

Discussion

A panel of experts produced consensus recommendations for intraoperative protective ventilation for the surgical patient. Those statements and recommendations that were of moderate to high quality and received strong support from the expert panel are presented in Table 4. We need to reiterate that two study questions did not achieve the consensus level of 70%. First, high-quality supportive evidence is lacking to recommend a routine ARM for all patients after tracheal intubation; however, 57% agreement was achieved that an ARM may be considered according to an individual risk-benefit assessment. Second, only 29% agreement was achieved that low FIO₂ (<0.3) with CPAP immediately after tracheal extubation may reduce the risk of resorption atelectasis. In both cases, published evidence was weak or non-existent, and the non-agreeing experts expressed concern about supporting potentially harmful interactions without more robust evidence.

Whilst these are the first published recommendations for the management of intraoperative mechanical ventilation, practice guidelines for mechanical ventilation in adult patients with acute respiratory distress syndrome (ARDS) strongly support the use of low V_T ventilation (4–8 ml kg⁻¹ PBW) and limiting P_{plat} to less than 30 cm H₂O.¹⁴⁰ The recommendations presented here are similar except for the use of ΔP instead of P_{plat}, as this appears better correlated with outcomes.^{5,6} In surgical patients, PEEP titration in conjunction with ARM is likely to be beneficial particularly during times when C_{RS} changes rapidly, such as during insufflation and steep Trendelenburg positioning. The use of higher levels of PEEP and ARM is only conditionally recommended in ARDS patients.¹⁴⁰ These differences likely reflect the different underlying pathophysiologies occurring in ARDS (inflammatory pulmonary oedema and cellular debris accumulation in alveoli) vs in the operating room (healthy lungs with a high degree of atelectasis). Whilst atelectatic alveoli during surgery can be reopened with ARM and incremental PEEP, the 'baby lung' of ARDS may not have a similar recruitable alveolar volume, and therefore, may not respond as favourably to ARM and PEEP.¹⁴¹

The modified Delphi method is recommended to determine a consensus for a defined clinical problem in the healthcare setting, and is an effective process for determining expert group consensus where there is little or no definitive evidence,

Table 4 Recommendations and statements with moderate- to high-quality and strong expert support. ARM, alveolar recruitment manoeuvre; CPAP, continuous positive airway pressure; ESA, European Society of Anaesthesiology; FIO₂, fraction of inspired oxygen; HOB, head of bed; I:E, inspiratory-to-expiratory ratio; NIPPV, non-invasive positive-pressure ventilation; PBW, predicted body weight; PPC, postoperative pulmonary complication; P_{plat}, plateau pressure; V_T, tidal volume; ZEEP, zero end-expiratory pressure.

Moderate- to high-quality recommendations with strong expert support:

- The ventilator should initially be set to deliver $V_T \leq 6-8 \text{ ml kg}^{-1}$ PBW and PEEP=5 cm H₂O. ZEEP is not recommended.
- Appropriate PEEP and recruitment manoeuvres may improve intraoperative respiratory function and prevent PPCs.
- Before the induction of anaesthesia, position the patient with the HOB elevated ≥ 30 deg (i.e. 'beach chair'); avoid flat supine position. If not contraindicated, before the loss of spontaneous ventilation, use NIPPV or CPAP to attenuate anaesthesia-induced respiratory changes.
- In addition to standard monitoring (ASA/ESA), dynamic compliance, driving pressure (P_{plat}-PEEP), and P_{plat} should be monitored on all controlled mechanically ventilated patients.
- Continuous haemodynamic and oxygen saturation monitoring is recommended before and during an ARM. Ensure adequate haemodynamic stability before performing an ARM. Avoid ARMs when contraindicated.

Moderate- to high-quality statements with strong expert support:

- The formation of perioperative clinically significant atelectasis may be an important risk factor for the development of PPCs.
- Decreasing compliance caused by surgical/anaesthesia factors (i.e. pneumoperitoneum, positioning, and circuit disconnect) should be treated by appropriate interventions.
- Individualised PEEP can prevent progressive alveolar collapse. Recruitment manoeuvres can reverse alveolar collapse, but have limited benefit without sufficient PEEP.
- Increasing FIO₂ may be effective in increasing the oxygenation, but is not an effective intervention to improve dynamic compliance of the respiratory system.

and where opinion is important.¹¹ The strengths of this method include the ability to bring a geographically dispersed and diverse group of expert panellists together, having an organised communication process in place, refining the content through repeated review, and the ability to condense expert opinion into clearly defined practice recommendations. Recognised limitations include the time required for expert participation and lack of anonymity during the face-to-face meeting. A limitation of our recommendations is that most of the literature focuses on surrogate endpoints, such as oxygenation or respiratory mechanics, and that relatively little published data support improvements in morbidity or mortality. By the same token, the recommendations are independent from the recently revised definition of PPCs.¹⁷ Interventions with associated costs or potential complications with no proven benefit in hard endpoints could not be recommended. Whilst the focus of this consensus conference was specifically to provide guidance for preoperative risk assessment and intraoperative mechanical ventilation for patients undergoing surgery, other factors not addressed in our review that may contribute to PPCs, such as incomplete reversal of neuromuscular block, postoperative opioid use, and surgical inflammation suppression, deserve further investigation. Future studies should continue to evaluate the roles of PEEP and ARM in the surgical patient. New imaging modalities, such as ultrasound and electrical impedance tomography, may help further elucidate their roles. Good-quality data on lung de-recruitment during emergence and possible mitigating methods are also needed. Finally, the role of FIO₂ in the development of PPCs requires further study.

Conclusion

In conclusion, this consensus meeting resulted in 26 recommendations and statements concerning the use of lung-protective ventilation in patients undergoing mechanical ventilation in the operating theatre. As the basic and clinical research focused on the application of mechanical ventilation in the surgical setting continues to emerge, it is likely that best practices to reduce or eliminate PPCs will likewise evolve. The

panel urges continued investigations and the adoption of proven interventions that will help optimise the perioperative care and safety of surgical patients. Further studies are needed to definitively confirm the beneficial effects of these interventions and manoeuvres on meaningful clinical outcomes.

Authors' contributions

Study design: CCY, EMH, CV

Literature search: AW

Literature review/compilation: SB, BB, RRDE, JM, CR, BT

Review of studies with research team: MGA, MG, EF, JPM, PP, JS, EMH

Writing of first draft: CCY, EMH, CV

Formatting of references: AW

Revising of final draft: all authors

CCY was the consensus conference president (coordinating team). CV was the consensus conference moderator (coordinating team). EMH, MGA, MG, EF, JPM, PP, and JS were the consensus conference experts. SB, BB, RRDE, JM, CR, and BT were the consensus conference participants.

Acknowledgements

GE Healthcare provided financial and logistical support for the development and implementation of the consensus panel meeting, including travel, lodging, and meals for the participants. GE Healthcare was not involved in the study data collection, analysis and interpretation of data, consensus panel deliberations, writing the report, or the decision to submit the report for publication. These were solely the decisions of the authors.

Declarations of interest

EF reports consulting fees from Dräger Medical, Edwards Lifesciences, GE Healthcare, and Orion Pharma, and lecture fees from Fresenius Kabi, Getinge, and Fisher & Paykel Healthcare. MGA has received financial support for research

and lecture fees from Dräger Medical AG, Ambu, GlaxoSmithKline, and GE Healthcare. MG and CCY are paid consultants for GE Healthcare. There are no other relationships or activities that could appear to have influenced the submitted work.

Funding

GE Healthcare (Anaesthesia and Respiratory Care)

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.bja.2019.08.017>.

References

- Canet J, Gallart L, Gomar C, et al. Prediction of postoperative pulmonary complications in a population-based surgical cohort. *Anesthesiology* 2010; **113**: 1338–50
- Fernandez-Bustamante A, Frenzl G, Sprung J, et al. Postoperative pulmonary complications, early mortality, and hospital stay following noncardiothoracic surgery: a multicenter study by the Perioperative Research Network Investigators. *JAMA Surg* 2017; **152**: 157–66
- Futier E, Constantin JM, Jaber S. Protective lung ventilation in operating room: a systematic review. *Minerva Anesthesiol* 2014; **80**: 726–35
- Guldner A, Kiss T, Serpa Neto A, et al. Intraoperative protective mechanical ventilation for prevention of postoperative pulmonary complications: a comprehensive review of the role of tidal volume, positive end-expiratory pressure, and lung recruitment maneuvers. *Anesthesiology* 2015; **123**: 692–713
- Amato MB, Meade MO, Slutsky AS, et al. Driving pressure and survival in the acute respiratory distress syndrome. *N Engl J Med* 2015; **372**: 747–55
- Neto AS, Hemmes SN, Barbas CS, et al. Association between driving pressure and development of postoperative pulmonary complications in patients undergoing mechanical ventilation for general anaesthesia: a meta-analysis of individual patient data. *Lancet Respir Med* 2016; **4**: 272–80
- Jaber S, Coisel Y, Chanques G, et al. A multicentre observational study of intra-operative ventilatory management during general anaesthesia: tidal volumes and relation to body weight. *Anaesthesia* 2012; **67**: 999–1008
- Hess DR, Kondili D, Burns E, Bittner EA, Schmidt UH. A 5-year observational study of lung-protective ventilation in the operating room: a single-center experience. *J Crit Care* 2013; **28**: 533.e9–e15
- Lynn MR. Determination and quantification of content validity. *Nurs Res* 1986; **35**: 382–5
- Linstone HA, Turoff M. *The Delphi method: techniques and applications*. Reading, MA: Addison-Wesley; 1975
- Goodman CM. The Delphi technique: a critique. *J Adv Nurs* 1987; **12**: 729–34
- Writing Committee for the PROBESE Collaborative Group of the PROtective VEntilation Network (PROVEnet) for the Clinical Trial Network of the European Society of Anaesthesiology, Bluth T, Serpa Neto A, Schultz MJ, Pelosi P, Gama de Abreu M. Effect of intraoperative high positive end-expiratory pressure (PEEP) with recruitment maneuvers vs low PEEP on postoperative pulmonary complications in obese patients: a randomized clinical trial. *JAMA* 2019; **321**: 2292–305
- Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan-a web and mobile app for systematic reviews. *Syst Rev* 2016; **5**: 210
- Atkins D, Eccles M, Flottorp S, et al. Systems for grading the quality of evidence and the strength of recommendations I: critical appraisal of existing approaches the GRADE Working Group. *BMC Health Serv Res* 2004; **4**: 38
- de Raaff CAL, de Vries N, van Wagenveld BA. Obstructive sleep apnea and bariatric surgical guidelines: summary and update. *Curr Opin Anaesthesiol* 2018; **31**: 104–9
- Gorter RR, Eker HH, Gorter-Stam MA, et al. Diagnosis and management of acute appendicitis. EAES consensus development conference 2015. *Surg Endosc* 2016; **30**: 4668–90
- Abbott TEF, Fowler AJ, Pelosi P, et al. A systematic review and consensus definitions for standardised end-points in perioperative medicine: pulmonary complications. *Br J Anaesth* 2018; **120**: 1066–79
- Hedenstierna G, Edmark L. Effects of anesthesia on the respiratory system. *Best Pract Res Clin Anaesthesiol* 2015; **29**: 273–84
- Gunnarsson L, Tokics L, Gustavsson H, Hedenstierna G. Influence of age on atelectasis formation and gas exchange impairment during general anaesthesia. *Br J Anaesth* 1991; **66**: 423–32
- Gattinoni L, Marini JJ, Collino F, et al. The future of mechanical ventilation: lessons from the present and the past. *Crit Care* 2017; **21**: 183
- Serpa Neto A, Juffermans NP, Hemmes SNT, et al. Interaction between peri-operative blood transfusion, tidal volume, airway pressure and postoperative ARDS: an individual patient data meta-analysis. *Ann Transl Med* 2018; **6**: 23
- Ladha K, Vidal Melo MF, McLean DJ, et al. Intraoperative protective mechanical ventilation and risk of postoperative respiratory complications: hospital based registry study. *BMJ* 2015; **351**: h3646
- Serpa Neto A, Amato MBP, Schultz MJ. Dissipated energy is a key mediator of VILI: rationale for using low driving pressures. In: Vincent J-L, editor. *Annual update in intensive care and emergency medicine* 2016. Cham: Springer; 2016. p. 311–21
- Weingarten TN, Whalen FX, Warner DO, et al. Comparison of two ventilatory strategies in elderly patients undergoing major abdominal surgery. *Br J Anaesth* 2010; **104**: 16–22
- Haliloglu M, Bilgili B, Ozdemir M, Umuroglu T, Bakan N. Low tidal volume positive end-expiratory pressure versus high tidal volume zero-positive end-expiratory pressure and postoperative pulmonary functions in robot-assisted laparoscopic radical prostatectomy. *Med Princ Pract* 2017; **26**: 573–8
- Wolthuis EK, Choi G, Densing MC, et al. Mechanical ventilation with lower tidal volumes and positive end-expiratory pressure prevents pulmonary inflammation in patients without preexisting lung injury. *Anesthesiology* 2008; **108**: 46–54
- Tang C, Li J, Lei S, et al. Lung-protective ventilation strategies for relief from ventilator-associated lung injury in patients undergoing craniotomy: a bicenter randomized, parallel, and controlled trial. *Oxid Med Cell Longev* 2017; **2017**: 6501248

28. Boyce JR, Ness T, Castroman P, Gleysteen JJ. A preliminary study of the optimal anesthesia positioning for the morbidly obese patient. *Obes Surg* 2003; **13**: 4–9
29. Dixon BJ, Dixon JB, Carden JR, et al. Preoxygenation is more effective in the 25 degrees head-up position than in the supine position in severely obese patients: a randomized controlled study. *Anesthesiology* 2005; **102**: 1110–5
30. Couture EJ, Provencher S, Somma J, Lellouche F, Marceau S, Bussieres JS. Effect of position and positive pressure ventilation on functional residual capacity in morbidly obese patients: a randomized trial. *Can J Anaesth* 2018; **65**: 522–8
31. Edmark L, Ostberg E, Scheer H, Wallquist W, Hedenstierna G, Zetterstrom H. Preserved oxygenation in obese patients receiving protective ventilation during laparoscopic surgery: a randomized controlled study. *Acta Anaesthesiol Scand* 2016; **60**: 26–35
32. Harbut P, Gozdzik W, Stjernfalt E, Marsk R, Hesselvik JF. Continuous positive airway pressure/pressure support pre-oxygenation of morbidly obese patients. *Acta Anaesthesiol Scand* 2014; **58**: 675–80
33. Rajan S, Joseph N, Tosh P, Paul J, Kumar L. Effects of preoxygenation with tidal volume breathing followed by apneic oxygenation with and without continuous positive airway pressure on duration of safe apnea time and arterial blood gases. *Anesth Essays Res* 2018; **12**: 229–33
34. Pang QY, Mo J, An R, Liu HL. Meta-analysis of the optimal ventilation strategies to improve perioperative oxygenation in obese patients. *Int J Clin Exp Med* 2017; **10**: 5883–91
35. Carron M, Zarantonello F, Tellaroli P, Ori C. Perioperative noninvasive ventilation in obese patients: a qualitative review and meta-analysis. *Surg Obes Relat Dis* 2016; **12**: 681–91
36. Cressey DM, Berthoud MC, Reilly CS. Effectiveness of continuous positive airway pressure to enhance preoxygenation in morbidly obese women. *Anaesthesia* 2001; **56**: 680–4
37. Yang D, Grant MC, Stone A, Wu CL, Wick EC. A meta-analysis of intraoperative ventilation strategies to prevent pulmonary complications: is low tidal volume alone sufficient to protect healthy lungs? *Ann Surg* 2016; **263**: 881–7
38. Serpa Neto A, Hemmes SN, Barbas CS, et al. Protective versus conventional ventilation for surgery: a systematic review and individual patient data meta-analysis. *Anesthesiology* 2015; **123**: 66–78
39. Cai H, Gong H, Zhang L, Wang Y, Tian Y. Effect of low tidal volume ventilation on atelectasis in patients during general anesthesia: a computed tomographic scan. *J Clin Anesth* 2007; **19**: 125–9
40. Futier E, Constantin JM, Petit A, et al. Positive end-expiratory pressure improves end-expiratory lung volume but not oxygenation after induction of anaesthesia. *Eur J Anaesthesiol* 2010; **27**: 508–13
41. Ostberg E, Thorisson A, Enlund M, Zetterstrom H, Hedenstierna G, Edmark L. Positive end-expiratory pressure alone minimizes atelectasis formation in non-abdominal surgery: a randomized controlled trial. *Anesthesiology* 2018; **128**: 1117–24
42. Wirth S, Baur M, Spaeth J, Guttmann J, Schumann S. Intraoperative positive end-expiratory pressure evaluation using the intratidal compliance-volume profile. *Br J Anaesth* 2015; **114**: 483–90
43. Wirth S, Kreysing M, Spaeth J, Schumann S. Intraoperative compliance profiles and regional lung ventilation improve with increasing positive end-expiratory pressure. *Acta Anaesthesiol Scand* 2016; **60**: 1241–50
44. Reis Miranda D, Gommers D, Struijs A, et al. Ventilation according to the open lung concept attenuates pulmonary inflammatory response in cardiac surgery. *Eur J Cardiothorac Surg* 2005; **28**: 889–95
45. Nestler C, Simon P, Petroff D, et al. Individualized positive end-expiratory pressure in obese patients during general anaesthesia: a randomized controlled clinical trial using electrical impedance tomography. *Br J Anaesth* 2017; **119**: 1194–205
46. Maisch S, Reissmann H, Fueellekrug B, et al. Compliance and dead space fraction indicate an optimal level of positive end-expiratory pressure after recruitment in anesthetized patients. *Anesth Analg* 2008; **106**: 175–81
47. Satoh D, Kurosawa S, Kirino W, et al. Impact of changes of positive end-expiratory pressure on functional residual capacity at low tidal volume ventilation during general anesthesia. *J Anesth* 2012; **26**: 664–9
48. Cinnella G, Grasso S, Spadaro S, et al. Effects of recruitment maneuver and positive end-expiratory pressure on respiratory mechanics and transpulmonary pressure during laparoscopic surgery. *Anesthesiology* 2013; **118**: 114–22
49. Karsten J, Heinze H, Meier T. Impact of PEEP during laparoscopic surgery on early postoperative ventilation distribution visualized by electrical impedance tomography. *Minerva Anestesiol* 2014; **80**: 158–66
50. Nieman GF, Satalin J, Andrews P, Aiash H, Habashi NM, Gatto LA. Personalizing mechanical ventilation according to physiologic parameters to stabilize alveoli and minimize ventilator induced lung injury (VILI). *Intensive Care Med Exp* 2017; **5**: 8
51. Ferrando C, Tusman G, Suarez-Sipmann F, et al. Individualized lung recruitment maneuver guided by pulse-oximetry in anesthetized patients undergoing laparoscopy: a feasibility study. *Acta Anaesthesiol Scand* 2018; **62**: 608–19
52. Asida SM, Badawy MS. Effect of low tidal volume during general anesthesia for urological procedures on lung functions. *Egypt J Anaesth* 2015; **31**: 127–34
53. Sato H, Nakamura K, Baba Y, Terada S, Goto T, Kurahashi K. Low tidal volume ventilation with low PEEP during surgery may induce lung inflammation. *BMC Anesthesiol* 2016; **16**: 47
54. Karsten J, Luepschen H, Grossherr M, et al. Effect of PEEP on regional ventilation during laparoscopic surgery monitored by electrical impedance tomography. *Acta Anaesthesiol Scand* 2011; **55**: 878–86
55. Ozkardesler Birlik S, Akan M, Atilla K, et al. The effects of the positive end expiratory pressure during laparoscopic cholecystectomy on postoperative respiratory function: a randomized controlled trial. *Gazz Med Ital* 2013; **172**: 11–9
56. Lee HJ, Kim KS, Jeong JS, Shim JC, Cho ES. Optimal positive end-expiratory pressure during robot-assisted laparoscopic radical prostatectomy. *Korean J Anesthesiol* 2013; **65**: 244–50
57. D'Antini D, Rauseo M, Grasso S, et al. Physiological effects of the open lung approach during laparoscopic

- cholecystectomy: focus on driving pressure. *Minerva Anesthesiol* 2018; **84**: 159–67
58. de Jong MAC, Ladha KS, Vidal Melo MF, et al. Differential effects of intraoperative positive end-expiratory pressure (PEEP) on respiratory outcome in major abdominal surgery versus craniotomy. *Ann Surg* 2016; **264**: 362–9
 59. Futier E, Constantin JM, Paugam-Burtz C, et al. A trial of intraoperative low-tidal-volume ventilation in abdominal surgery. *N Engl J Med* 2013; **369**: 428–37
 60. Severgnini P, Selmo G, Lanza C, et al. Protective mechanical ventilation during general anesthesia for open abdominal surgery improves postoperative pulmonary function. *Anesthesiology* 2013; **118**: 1307–21
 61. Hemmes SN, Gama de Abreu M, Pelosi P, Schultz MJ. High versus low positive end-expiratory pressure during general anaesthesia for open abdominal surgery (PROVHILO trial): a multicentre randomised controlled trial. *Lancet* 2014; **384**: 495–503
 62. D'Antini D, Huhle R, Herrmann J, et al. Respiratory system mechanics during low versus high positive end-expiratory pressure in open abdominal surgery: a sub-study of PROVHILO randomized controlled trial. *Anesth Analg* 2018; **126**: 143–9
 63. Treschan TA, Schaefer M, Kemper J, et al. Ventilation with high versus low PEEP levels during general anaesthesia for open abdominal surgery does not affect post-operative spirometry: a randomised clinical trial. *Eur J Anaesthesiol* 2017; **34**: 534–43
 64. Kim JY, Shin CS, Kim HS, Jung WS, Kwak HJ. Positive end-expiratory pressure in pressure-controlled ventilation improves ventilatory and oxygenation parameters during laparoscopic cholecystectomy. *Surg Endosc* 2010; **24**: 1099–103
 65. Spaeth J, Daume K, Goebel U, Wirth S, Schumann S. Increasing positive end-expiratory pressure (re-)improves intraoperative respiratory mechanics and lung ventilation after prone positioning. *Br J Anaesth* 2016; **116**: 838–46
 66. Neligan PJ, Malhotra G, Fraser M, et al. Continuous positive airway pressure via the Boussignac system immediately after extubation improves lung function in morbidly obese patients with obstructive sleep apnea undergoing laparoscopic bariatric surgery. *Anesthesiology* 2009; **110**: 878–84
 67. Jo YY, Kim JY, Park CK, Chang YJ, Kwak HJ. The effect of ventilation strategy on arterial and cerebral oxygenation during laparoscopic bariatric surgery. *Obes Surg* 2016; **26**: 339–44
 68. Kim WH, Hahm TS, Kim JA, et al. Prolonged inspiratory time produces better gas exchange in patients undergoing laparoscopic surgery: a randomised trial. *Acta Anaesthesiol Scand* 2013; **57**: 613–22
 69. Kim MS, Kim NY, Lee KY, Choi YD, Hong JH, Bai SJ. The impact of two different inspiratory to expiratory ratios (1:1 and 1:2) on respiratory mechanics and oxygenation during volume-controlled ventilation in robot-assisted laparoscopic radical prostatectomy: a randomized controlled trial. *Can J Anaesth* 2015; **62**: 979–87
 70. Mousa WF. Equal ratio ventilation (1:1) improves arterial oxygenation during laparoscopic bariatric surgery: a crossover study. *Saudi J Anaesth* 2013; **7**: 9–13
 71. Zhang WP, Zhu SM. The effects of inverse ratio ventilation on cardiopulmonary function and inflammatory cytokine of bronchoalveolar lavage in obese patients undergoing gynecological laparoscopy. *Acta Anaesthesiol Taiwan* 2016; **54**: 1–5
 72. Xu L, Shen J, Yan M. The effect of pressure-controlled inverse ratio ventilation on lung protection in obese patients undergoing gynecological laparoscopic surgery. *J Anesth* 2017; **31**: 651–6
 73. Suzuki S, Mihara Y, Hikasa Y, et al. Current ventilator and oxygen management during general anesthesia: a multicenter, cross-sectional observational study. *Anesthesiology* 2018; **129**: 67–76
 74. Stolmeijer R, Bouma HR, Zijlstra JG, Drost-de Klerck AM, Ter Maaten JC, JJM Ligtenberg. A systematic review of the effects of hyperoxia in acutely ill patients: should we aim for less? *Biomed Res Int* 2018; **2018**: 7841295
 75. Staehr AK, Meyhoff CS, Henneberg SW, Christensen PL, Rasmussen LS. Influence of perioperative oxygen fraction on pulmonary function after abdominal surgery: a randomized controlled trial. *BMC Res Notes* 2012; **5**: 383
 76. Haque WA, Boehmer J, Clemson BS, Leuenberger UA, Silber DH, Sinoway LI. Hemodynamic effects of supplemental oxygen administration in congestive heart failure. *J Am Coll Cardiol* 1996; **27**: 353–7
 77. Harten JM, Anderson KJ, Angerson WJ, Booth MG, Kinsella J. The effect of normobaric hyperoxia on cardiac index in healthy awake volunteers. *Anaesthesia* 2003; **58**: 885–8
 78. Austin MA, Wills KE, Blizzard L, Walters EH, Wood-Baker R. Effect of high flow oxygen on mortality in chronic obstructive pulmonary disease patients in pre-hospital setting: randomised controlled trial. *BMJ* 2010; **341**: c5462
 79. Stub D, Smith K, Bernard S, et al. Air versus oxygen in ST-segment-elevation myocardial infarction. *Circulation* 2015; **131**: 2143–50
 80. Kilgannon JH, Jones AE, Shapiro NI, et al. Association between arterial hyperoxia following resuscitation from cardiac arrest and in-hospital mortality. *JAMA* 2010; **303**: 2165–71
 81. Meyhoff CS, Jorgensen LN, Wetterslev J, Christensen KB, Rasmussen LS. Increased long-term mortality after a high perioperative inspiratory oxygen fraction during abdominal surgery: follow-up of a randomized clinical trial. *Anesth Analg* 2012; **115**: 849–54
 82. Qadan M, Akca O, Mahid SS, Hornung CA, Polk Jr HC. Perioperative supplemental oxygen therapy and surgical site infection: a meta-analysis of randomized controlled trials. *Arch Surg* 2009; **144**: 359–66
 83. Staehr-Rye AK, Meyhoff CS, Scheffenbichler FT, et al. High intraoperative inspiratory oxygen fraction and risk of major respiratory complications. *Br J Anaesth* 2017; **119**: 140–9
 84. Tusman G, Bohm SH, Suarez-Sipmann F. Advanced uses of pulse oximetry for monitoring mechanically ventilated patients. *Anesth Analg* 2017; **124**: 62–71
 85. Pedersen T, Moller AM, Pedersen BD. Pulse oximetry for perioperative monitoring: systematic review of randomized, controlled trials. *Anesth Analg* 2003; **96**: 426–31
 86. Aydin V, Kabukcu HK, Sahin N, et al. Comparison of pressure and volume-controlled ventilation in laparoscopic cholecystectomy operations. *Clin Respir J* 2016; **10**: 342–9
 87. Bagchi A, Rudolph MI, Ng PY, et al. The association of postoperative pulmonary complications in 109,360

- patients with pressure-controlled or volume-controlled ventilation. *Anaesthesia* 2017; **72**: 1334–43
88. Wang C, Zhao N, Wang W, et al. Intraoperative mechanical ventilation strategies for obese patients: a systematic review and network meta-analysis. *Obes Rev* 2015; **16**: 508–17
 89. Choi EM, Na S, Choi SH, An J, Rha KH, Oh YJ. Comparison of volume-controlled and pressure-controlled ventilation in steep Trendelenburg position for robot-assisted laparoscopic radical prostatectomy. *J Clin Anesth* 2011; **23**: 183–8
 90. Dion JM, McKee C, Tobias JD, et al. Ventilation during laparoscopic-assisted bariatric surgery: volume-controlled, pressure-controlled or volume-guaranteed pressure-regulated modes. *Int J Clin Exp Med* 2014; **7**: 2242–7
 91. Gupta SD, Kundu SB, Ghose T, et al. A comparison between volume-controlled ventilation and pressure-controlled ventilation in providing better oxygenation in obese patients undergoing laparoscopic cholecystectomy. *Indian J Anaesth* 2012; **56**: 276–82
 92. Tyagi A, Kumar R, Sethi AK, Mohta M. A comparison of pressure-controlled and volume-controlled ventilation for laparoscopic cholecystectomy. *Anaesthesia* 2011; **66**: 503–8
 93. Cadi P, Guenoun T, Journois D, Chevallier JM, Diehl JL, Safran D. Pressure-controlled ventilation improves oxygenation during laparoscopic obesity surgery compared with volume-controlled ventilation. *Br J Anaesth* 2008; **100**: 709–16
 94. De Baerdemaeker LE, Van der Hertten C, Gillardin JM, Pattyn P, Mortier EP, Szegedi LL. Comparison of volume-controlled and pressure-controlled ventilation during laparoscopic gastric banding in morbidly obese patients. *Obes Surg* 2008; **18**: 680–5
 95. Talab HF, Zabani IA, Abdelrahman HS, et al. Intraoperative ventilatory strategies for prevention of pulmonary atelectasis in obese patients undergoing laparoscopic bariatric surgery. *Anesth Analg* 2009; **109**: 1511–6
 96. El-Sayed KM, Tawfeek MM. Perioperative ventilatory strategies for improving arterial oxygenation and respiratory mechanics in morbidly obese patients undergoing laparoscopic bariatric surgery. *Egypt J Anaesth* 2012; **28**: 9–15
 97. He X, Jiang J, Liu Y, et al. Electrical impedance tomography-guided PEEP titration in patients undergoing laparoscopic abdominal surgery. *Medicine (Baltimore)* 2016; **95**, e3306
 98. Kostic P, LoMauro A, Larsson A, Hedenstierna G, Frykholm P, Aliverti A. Specific anesthesia-induced lung volume changes from induction to emergence: a pilot study. *Acta Anaesthesiol Scand* 2018; **62**: 282–92
 99. Soh S, Shim JK, Ha Y, Kim YS, Lee H, Kwak YL. Ventilation with high or low tidal volume with PEEP does not influence lung function after spinal surgery in prone position: a randomized controlled trial. *J Neurosurg Anesthesiol* 2018; **30**: 237–45
 100. Stankiewicz-Rudnicki M, Gaszynski W, Gaszynski T. Assessment of ventilation distribution during laparoscopic bariatric surgery: an electrical impedance tomography study. *Biomed Res Int* 2016; **2016**: 7423162
 101. Valenza F, Chevillard G, Fossali T, Salice V, Pizzocri M, Gattinoni L. Management of mechanical ventilation during laparoscopic surgery. *Best Pract Res Clin Anaesthesiol* 2010; **24**: 227–41
 102. Pang CK, Yap J, Chen PP. The effect of an alveolar recruitment strategy on oxygenation during laparoscopic cholecystectomy. *Anaesth Intensive Care* 2003; **31**: 176–80
 103. Whalen FX, Gajic O, Thompson GB, et al. The effects of the alveolar recruitment maneuver and positive end-expiratory pressure on arterial oxygenation during laparoscopic bariatric surgery. *Anesth Analg* 2006; **102**: 298–305
 104. Almarakbi WA, Fawzi HM, Alhashemi JA. Effects of four intraoperative ventilatory strategies on respiratory compliance and gas exchange during laparoscopic gastric banding in obese patients. *Br J Anaesth* 2009; **102**: 862–8
 105. Hartland BL, Newell TJ, Damico N. Alveolar recruitment maneuvers under general anesthesia: a systematic review of the literature. *Respir Care* 2015; **60**: 609–20
 106. Rothen HU, Neumann P, Berglund JE, Valtysson J, Magnusson A, Hedenstierna G. Dynamics of re-expansion of atelectasis during general anaesthesia. *Br J Anaesth* 1999; **82**: 551–6
 107. Rothen HU, Sporre B, Engberg G, Wegenius G, Hedenstierna G. Re-expansion of atelectasis during general anaesthesia: a computed tomography study. *Br J Anaesth* 1993; **71**: 788–95
 108. Ahn S, Byun SH, Chang H, Koo YB, Kim JC. Effect of recruitment maneuver on arterial oxygenation in patients undergoing robot-assisted laparoscopic prostatectomy with intraoperative 15 cmH₂O positive end expiratory pressure. *Korean J Anesthesiol* 2016; **69**: 592–8
 109. Cakmakkaya OS, Kaya G, Altintas F, Hayirlioglu M, Ekici B. Restoration of pulmonary compliance after laparoscopic surgery using a simple alveolar recruitment maneuver. *J Clin Anesth* 2009; **21**: 422–6
 110. Futier E, Constantin JM, Pelosi P, et al. Intraoperative recruitment maneuver reverses detrimental pneumoperitoneum-induced respiratory effects in healthy weight and obese patients undergoing laparoscopy. *Anesthesiology* 2010; **113**: 1310–9
 111. Park HP, Hwang JW, Kim YB, et al. Effect of pre-emptive alveolar recruitment strategy before pneumoperitoneum on arterial oxygenation during laparoscopic hysterectomy. *Anaesth Intensive Care* 2009; **37**: 593–7
 112. Reinius H, Jonsson L, Gustafsson S, et al. Prevention of atelectasis in morbidly obese patients during general anesthesia and paralysis: a computerized tomography study. *Anesthesiology* 2009; **111**: 979–87
 113. Shim JK, Chun DH, Choi YS, Lee JY, Hong SW, Kwak YL. Effects of early vital capacity maneuver on respiratory variables during multivessel off-pump coronary artery bypass graft surgery. *Crit Care Med* 2009; **37**: 539–44
 114. Aretha D, Fligou F, Kiekkas P, et al. Safety and effectiveness of alveolar recruitment maneuvers and positive end-expiratory pressure during general anesthesia for cesarean section: a prospective, randomized trial. *Int J Obstet Anesth* 2017; **30**: 30–8
 115. Choi ES, Oh AY, In CB, Ryu JH, Jeon YT, Kim HG. Effects of recruitment manoeuvre on perioperative pulmonary complications in patients undergoing robotic assisted radical prostatectomy: a randomised single-blinded trial. *PLoS One* 2017; **12**, e0183311
 116. Ferrando C, Suarez-Sipmann F, Tusman G, et al. Open lung approach versus standard protective strategies:

- effects on driving pressure and ventilatory efficiency during anesthesia—a pilot, randomized controlled trial. *PLoS One* 2017; **12**, e0177399
117. Tusman G, Bohm SH, Suarez-Sipmann F, Turchetto E. Alveolar recruitment improves ventilatory efficiency of the lungs during anesthesia. *Can J Anaesth* 2004; **51**: 723–7
 118. Tusman G, Bohm SH, Vazquez de Anda GF, do Campo JL, Lachmann B. 'Alveolar recruitment strategy' improves arterial oxygenation during general anaesthesia. *Br J Anaesth* 1999; **82**: 8–13
 119. Topuz U, Salihoglu Z, Gokay BV, Umutoglu T, Bakan M, Idin K. The effects of different oxygen concentrations on recruitment maneuver during general anesthesia for laparoscopic surgery. *Surg Laparosc Endosc Percutaneous Tech* 2014; **24**: 410–3
 120. Benoit Z, Wicky S, Fischer JF, et al. The effect of increased FIO₂ before tracheal extubation on postoperative atelectasis. *Anesth Analg* 2002; **95**: 1777–81
 121. Golparvar M, Mofrad SZ, Mahmoodieh M, Kalidarei B. Comparative evaluation of the effects of three different recruitment maneuvers during laparoscopic bariatric surgeries of morbid obese patients on cardiopulmonary indices. *Adv Biomed Res* 2018; **7**: 89
 122. Lumb AB, Greenhill SJ, Simpson MP, Stewart J. Lung recruitment and positive airway pressure before extubation does not improve oxygenation in the post-anaesthesia care unit: a randomized clinical trial. *Br J Anaesth* 2010; **104**: 643–7
 123. Neumann P, Rothen HU, Berglund JE, Valtysson J, Magnusson A, Hedenstierna G. Positive end-expiratory pressure prevents atelectasis during general anaesthesia even in the presence of a high inspired oxygen concentration. *Acta Anaesthesiol Scand* 1999; **43**: 295–301
 124. Park SJ, Kim BG, Oh AH, Han SH, Han HS, Ryu JH. Effects of intraoperative protective lung ventilation on postoperative pulmonary complications in patients with laparoscopic surgery: prospective, randomized and controlled trial. *Surg Endosc* 2016; **30**: 4598–606
 125. Tusman G, Groisman I, Fiolo FE, et al. Noninvasive monitoring of lung recruitment maneuvers in morbidly obese patients: the role of pulse oximetry and volumetric capnography. *Anesth Analg* 2014; **118**: 137–44
 126. Pereira SM, Tucci MR, Morais CCA, et al. Individual positive end-expiratory pressure settings optimize intraoperative mechanical ventilation and reduce postoperative atelectasis. *Anesthesiology* 2018; **129**: 1070–81
 127. Ball L, Costantino F, Orefice G, Chandrapatham K, Pelosi P. Intraoperative mechanical ventilation: state of the art. *Minerva Anesthesiol* 2017; **83**: 1075–88
 128. Hedenstierna G, Edmark L. Mechanisms of atelectasis in the perioperative period. *Best Pract Res Clin Anaesthesiol* 2010; **24**: 157–69
 129. Kleinsasser AT, Pircher I, Truebsbach S, Knotzer H, Loeckinger A, Trembl B. Pulmonary function after emergence on 100% oxygen in patients with chronic obstructive pulmonary disease: a randomized, controlled trial. *Anesthesiology* 2014; **120**: 1146–51
 130. Edmark L, Auner U, Lindback J, Enlund M, Hedenstierna G. Post-operative atelectasis—a randomised trial investigating a ventilatory strategy and low oxygen fraction during recovery. *Acta Anaesthesiol Scand* 2014; **58**: 681–8
 131. Edmark L, Auner U, Enlund M, Ostberg E, Hedenstierna G. Oxygen concentration and characteristics of progressive atelectasis formation during anaesthesia. *Acta Anaesthesiol Scand* 2011; **55**: 75–81
 132. Ostberg E, Auner U, Enlund M, Zetterstrom H, Edmark L. Minimizing atelectasis formation during general anaesthesia-oxygen washout is a non-essential supplement to PEEP. *Ups J Med Sci* 2017; **122**: 92–8
 133. Ireland CJ, Chapman TM, Mathew SF, Herbison GP, Zacharias M. Continuous positive airway pressure (CPAP) during the postoperative period for prevention of postoperative morbidity and mortality following major abdominal surgery. *Cochrane Database Syst Rev* 2014; **8**. Cd008930
 134. Singh PM, Borle A, Shah D, et al. Optimizing prophylactic CPAP in patients without obstructive sleep apnoea for high-risk abdominal surgeries: a meta-regression analysis. *Lung* 2016; **194**: 201–17
 135. Squadrone V, Cocha M, Cerutti E, et al. Continuous positive airway pressure for treatment of postoperative hypoxemia: a randomized controlled trial. *JAMA* 2005; **293**: 589–95
 136. Kindgen-Milles D, Muller E, Buhl R, et al. Nasal-continuous positive airway pressure reduces pulmonary morbidity and length of hospital stay following thoracoabdominal aortic surgery. *Chest* 2005; **128**: 821–8
 137. Hewidy AA, Suliman LA, El Hefnawy E, Hassan AA. Immediate continuous positive airway pressure (CPAP) therapy after sleeve gastrectomy. *Egypt J Chest Dis Tuberc* 2016; **65**: 701–6
 138. Zoremba M, Kalmus G, Begemann D, et al. Short term non-invasive ventilation post-surgery improves arterial blood-gases in obese subjects compared to supplemental oxygen delivery—a randomized controlled trial. *BMC Anesthesiol* 2011; **11**: 10
 139. Guimaraes J, Pinho D, Nunes CS, Cavaleiro CS, Machado HS. Effect of Boussignac continuous positive airway pressure ventilation on PaO₂ and PaO₂/FIO₂ ratio immediately after extubation in morbidly obese patients undergoing bariatric surgery: a randomized controlled trial. *J Clin Anesth* 2016; **34**: 562–70
 140. Fan E, Del Sorbo L, Goligher EC, et al. An Official American Thoracic Society/European Society of Intensive Care Medicine/Society of Critical Care Medicine clinical practice guideline: mechanical ventilation in adult patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2017; **195**: 1253–63
 141. Writing Group for the Alveolar Recruitment for Acute Respiratory Distress Syndrome Trial (ART) Investigators, Cavalcanti AB, Suzumura EA, et al. Effect of lung recruitment and titrated positive end-expiratory pressure (PEEP) vs low PEEP on mortality in patients with acute respiratory distress syndrome: a randomized clinical trial. *JAMA* 2017; **318**: 1335–45