



Nutrition behaviour and compliance with the Mediterranean diet pyramid recommendations: an Italian survey-based study

Renata Bracale¹ · Concetta M. Vaccaro² · Vittoria Coletta² · Claudio Cricelli³ · Francesco Carlo Gamaleri⁴ · Fabio Parazzini⁵ · Michele Carruba⁶

Received: 5 June 2019 / Accepted: 23 October 2019
© The Author(s) 2019, corrected publication 2019

Abstract

Purpose Adopting a Mediterranean-like dietary pattern may help in preventing several chronic diseases. We assessed the eating behaviour and compliance with the Mediterranean diet pyramid recommendations in Italy.

Methods This is a cross-sectional study conducted in subjects aged ≥ 20 years. A 14-question survey based on the updated Mediterranean diet pyramid was launched online from April 2015 to November 2016. At test completion, a personalized pyramid displaying the possible deficiencies and/or excesses was generated, that could be the basis to plan diet and lifestyle modifications.

Results Overall, 27,540 subjects completed the survey: the proportion of females (75.6%), younger subjects (20.7%) and people with a University degree (33.1%) resembled those of the Italian population of Internet users rather than of the general population. 37.8% of participants declared a sedentary lifestyle, including 29.6% of those aged 20–29 years. A lower-than-recommended intake of all food categories included in the Mediterranean diet pyramid, along with excess of sweets, red and processed meats, emerged, that may affect health in the long term. Low adherence to recommendations was observed especially among females and older people. Notably, a discrepancy surfaced between the responders' perceived and actual behaviour toward the regular consumption of fruits and vegetables (81.8% vs 22.7–32.8%, respectively).

Conclusions The nutritional habits and lifestyle of Italian participants are poorly adherent to the Mediterranean diet recommendations. The personalized pyramid tool may help in raising the awareness of individuals and their families on where to intervene, possibly with the support of healthcare professionals, to improve their behaviour.

Level of evidence Level V, cross-sectional descriptive study.

Keywords Compliance · Food pyramid · Italy · Lifestyle · Mediterranean diet · Nutrition

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s40519-019-00807-4>) contains supplementary material, which is available to authorized users.

✉ Michele Carruba
michele.carruba@unimi.it

¹ Department of Medicine and Sciences for Health, Molise University, Campobasso, Italy

² Fondazione Censis, Rome, Italy

³ Società Italiana di Medicina Generale e delle Cure Primarie Firenze, Florence, Italy

Introduction

Exhaustive evidence has established the central role of a healthy lifestyle, based on a balanced and diverse diet and on adequate physical activity, in preventing several chronic diseases, known to impose a substantial burden on health-care systems and communities.

⁴ Ordine dei Farmacisti delle Province di Milano, Lodi e Monza Brianza, Milan, Italy

⁵ Dipartimento di scienze Cliniche e di Comunità, Università di Milano, Milan, Italy

⁶ Department of Medical Biotechnology and Translational Medicine, Center for the Study and Research on Obesity, University of Milan, Milan, Italy

In particular, a number of studies have shown that adoption of a Mediterranean-like dietary pattern [1] improves the health status [2], decreases morbidity and mortality [3] and reduces the total lifetime costs [2]. Benefits have been demonstrated in terms of decreased risk of cardiovascular diseases (CVD) [4–6], obesity [7], metabolic syndrome [8, 9] and type-2 diabetes [10–13], as well as of certain types of neurodegenerative diseases and late-life cognitive disorders [14–17] and cancers [18–22]. Moreover, in the elderly, such dietary pattern has been reported to prolong survival [23] and reduce bone loss in subjects with osteoporosis supplemented with vitamin D [24, 25].

In the last decades, however, we have witnessed a change in dietary and lifestyle habits, that has led to the loss of the traditional Mediterranean diet pattern. Factors accounting for this shift include globalization, more poverty and sedentariness and increased intake of sugars and of energy-dense and processed foods. This is particularly relevant in Countries like Italy, where diabetes is rising (due to population aging, early diagnosis and prolongation of patient survival), and the rate of childhood obesity, although decreasing [26, 27], is one of the highest in Europe, with one child in five being obese [28]. As a consequence, the risk of adult obesity, diabetes and CVD later in life is expected to increase, rising a serious public health concern.

In this context, gaining insights into the actual nutrition habits of the Italians is the first step to plan simple and effective strategies aimed at improving diet and lifestyle or, where requested, to provide nutrient supplementation.

Here, we describe the nutrition behaviour of a large Italian population and their compliance with the Mediterranean diet pyramid recommendations [1].

Methods

Study design and participants

This population-based, cross-sectional study was undertaken to collect information about the nutrition behaviour of Italian subjects aged ≥ 20 years from across the Country. A 14-question survey was designed on the basis of the updated food pyramid of the Mediterranean diet, which establishes daily, weekly and occasional guidelines to follow a healthy and balanced diet [1]. Although not validated, the questionnaire employed was developed on the food frequency questionnaires used in published studies [1].

The survey was launched online on the Italian website www.curarelasalute.com, from April 2015 to November 2016. The venture was publicized mainly by pharmacists, informed through newsletters, activities of media relations on trade magazines and during events dedicated to pharmacists and nutritionists (e.g. Cosmofarma, Farmacista Più

and Pianeta Nutrizione). Moreover, pharmacies employed leaflets and informative materials distributed onsite, post on Facebook and activities of media relations on consumer-targeted magazines to publicize the website.

The survey collected demographics, general information on the lifestyle and nutrition habits, and information on the number of servings of each food category included in the Mediterranean pyramid (for the corresponding quantities in g/day refer to [29] and [30]). The complete questionnaire is provided as Supplementary Table 1. At test completion, a personalized pyramid was generated, displaying the possible food excesses and deficiencies. Moreover, a suggestion was made to discuss the resulting pyramid with the general practitioner (GP) or pharmacist to adopt measures to improve the individual's nutrition.

Participation was voluntary and anonymous and completing the survey was accepted as consent by the participants.

Statistical analysis

Qualitative variables were expressed as absolute and relative frequencies; comparisons between groups were made by the Chi-square test. A p value of < 0.05 was considered as statistically significant. All analyses were conducted using IBM SPSS Statistics v25.

Results

Participant characteristics

Overall, 27,540 subjects (females: 75.6%) completed the survey. The main demographic characteristics of the overall participant population are summarized in Table 1. Based on gender and age distribution, the sample population reflected the population of the Italian Internet users (i.e., 61.4% of people older than 20 years [31]) rather than fully represent the general Italian population, particularly with regard to the proportion of females, younger subjects and higher level of education (Table 1 and data not shown).

Lifestyle and nutrition habits

Overall, 62.1% of responders declared to have an active lifestyle, doing physical activity (including walking, biking and taking the stairs rather than using the lift) at least twice a week. As shown in Fig. 1, sedentariness was significantly more common in older people (from 29.6% of those aged 20–29 years to 40% of those aged ≥ 50 year, $p = 0.00$), in females (40% vs 35.3% in males, $p = 0.00$) and in responders from Central and Southern Italy compared to those from the Northwest and the Northeast (40.9% and 40.2% vs 36.4% and 34.4%, respectively, $p = 0.00$).

Table 1 Demographics of the responders and comparison with the population of Internet users in Italy

Characteristic	Responders <i>N</i> =27,540	Internet users ^a
Gender		
Females	75.6	47.6
Males	24.4	52.4
Age (years)		
20–29	20.7	18.6
30–39	19.2	22.9
40–49	22.9	25.8
50–59	24.1	19.6
≥60	13.1	13.1
Education		
Low	16.2	24.7
High school—Professional	47.1	50.2
University	33.1	22.2
Other	3.5	3.0
Geographical area		
Northwest	35.6	26.7
Northeast	20.0	19.1
Center	17.0	20.0
South and the islands	27.5	34.2

Data are expressed as frequencies (*N*, %)

^aData from the Italian Institute of Statistics

The regular consumption of fruits and vegetables was reported by 81.8% of participants and was significantly more common in females (85.0% vs 78.8% in males, $p=0.00$) and increased with age (from 72.1% of those aged 20–29 years to 92.1% of those older than 60 years, $p=0.00$) and education (from 77.3% of those with low education to 84.7% of those with a University degree, $p=0.00$) (Fig. 2a). Moreover,

75.0% of the responders declared to consume fruits and vegetables of different colors (data not shown) and 80.7% that they had products from the territory, with local (i.e., typical of the tradition of the territory) food intake increasing with age (from 72.6% of those aged 20–29 years to 90.0% of those older than 60 years, $p=0.00$) (Fig. 2b).

With regard to wine, 86.6% of the surveyed people declared to drink 0 or 1 glass of wine per day, while the remaining usually had more than one. The daily intake of wine was significantly lower among females (0–1 glass: 91.1% vs 82.6% of males; > 1 glass: 8.9% vs 17.4% in males $p=0.00$) and in those aged ≥ 60 years (78.7%, vs 92.2% among those of 20–29 years, $p=0.00$) (Fig. 2c).

Finally, 51.6% of the responders declared to drink at least 1.5 L of water daily; whereas, the remaining 48.4% had an insufficient intake. In particular, a level lower than recommended was reported significantly more frequently by females (56.7% vs 46.9% of males, $p=0.00$) and with increasing age (from 43.5% of those aged 20–29 years to 62.7% of those older than 60 years, $p=0.00$) (Fig. 2d).

Compliance to the Mediterranean food pyramid recommendations

Next, to assess the compliance to the Mediterranean diet recommendations, responders were asked to indicate how many servings of each food group included in the pyramid they consumed at every main meal, daily, weekly or occasionally (Table 2). Deficiencies and excesses are detailed in the pyramid depicted in Fig. 3. Overall, deficiencies were observed for all food categories, while excesses were observed solely in some cases.

At each main meal, only 18.9%, 26.8% and 34.8% of the responders adhered to the recommendations on the servings




Fig. 1 Sedentary lifestyle. Only the subgroups with statistically significant differences are displayed




Fig. 2 Nutrition habits in terms of regular intake of fruits and vegetables (a), territorial food (b), wine (c) and water (d). Only the subgroups with statistically significant differences are displayed

of fruits, pasta and cereals and vegetables, respectively (which constitute the pyramid base) (Table 2). A lower-than-recommended intake of these foods was reported significantly more frequently by age and gender (Fig. 4a), except for fruits, consumed at a similar extent by males and females. Notably, a discrepancy was observed between the % of participants who had declared a regular intake of fruits and vegetables (i.e. 81.8), and those stating to actually consume a recommended amount of these foods (22.7% for fruits and 32.8% for vegetables; $p = 0.00$).

A low level of compliance was reported also for the foods to be consumed on a daily basis, except for herbs and spices (normal consumption in 63.7% of cases, vs 36.3% of no or infrequent intake) (Table 2). The lowest rate of compliance was observed for milk and dairy (21.1%) and nuts (25.0%) (Table 2). Consumption was mostly deficient and varied by age (Figs. 3, 4b).

Most of the responders declared a normal weekly consumption of animal protein-enriched foods: 51.3% for red meats, 65.9% for white meats and 73.8% for eggs (Table 2). A scarce intake of red meats was reported

Table 2 Responder compliance to the Mediterranean food pyramid recommendations


	Compliant (%)	Non-compliant (%)
Every main meal		
Vegetables, ≥ 2 s	34.8	65.2
Fruits, ≥ 3 s	18.9	81.1
Bread, pasta etc., 4 s	26.8	73.2
Every day		
Milk and dairy, 2–3 s	21.1	78.9
Nuts, 1 s	25.0	75.0
Herbs and spices	63.7	36.3
Olive oil, 3 s	46.6	53.4
Hydration	48.4	51.6
Every week		
White meat, 1–2 s	65.9	34.1
Fish and seafood, ≥ 2 s	40.3	59.7
Legumes, ≥ 2 s	45.0	55.0
Eggs, 1–4	73.8	26.2
Red meat, 1 s	51.3	48.7
Processed meat, 1 s	44.0	56.0
Sweets, ≤ 2 s	59.1	40.9
Physical activity	62.1	37.9

Compliance rates of at least 50% are indicated in bold

Data are expressed as %

S serving

Fig. 3 Food pyramid built on the participants' responses to the questionnaire. The % of non-compliant responders in terms of excesses and deficiencies are displayed Modified from [1]



significantly more frequently among females (27.7% vs 18.4% of males), in Southern Italy (19.0% vs 25.4% in the Northeast) and with increasing age (from 19.6% in those aged 20–29 years to 29.1% in those older than 60 years), just like white meats (from 10.0% in those aged 20–29 years to 17.6% in those older than 60 years). In contrast, the weekly consumption of fish and seafood and of legumes was rather scarce (in 59.7% and 55.0% of responders, respectively; Table 2; Fig. 3), the latter being lower in Southern Italy (49.2% vs 59.2% in the Northeast) (Fig. 4c).

In the Mediterranean diet, sugar- and unhealthy fat-rich foods like sweets are allowed occasionally and in small amounts. In the present study, 59.1% of responders declared to have a normal consumption of sweets (52.9% of people between 20 and 29 years and 73.2% of those older than 60), whereas the remaining had excesses (Fig. 4d).

Discussion

This population-based study provides a snapshot of the nutritional habits and lifestyle of the Italian Internet users participating in the survey between April 2015 and November 2016. The tool employed is the food pyramid based on the updated food pyramid of the Mediterranean diet [1].

Overall, the population participating in the study was more representative of the Italian Internet users rather than

of the general population, as shown by the higher percentage of females, young subjects and those with a University degree. This result may be explained, at least partly, by the fact that the population of Internet users is younger than the general population, and, therefore, is more similar to the responders. Nevertheless, some interesting observations can be made. First, notwithstanding the clear interest of the responders in health, the younger age and the higher education level (adults with a high level of education are overrepresented in our sample population), nearly 40% of the participants declared to not conduct an active lifestyle. These data are in line with those collected by the Italian Institute of Statics, which, however, refer to a slightly different population (i.e., older than 6 years of age) [32]. It is not surprising that a significantly increased sedentariness was observed with increasing age, but the fact that 1 of 3

subjects between 20 and 29 years of age declared a sedentary behaviour is quite concerning. Indeed, sedentariness as well as suboptimal nutrition represents the modifiable risk factors for chronic diseases like obesity and type 2 diabetes [33], and adequate physical activity is recommended in the Mediterranean diet as a basic complement to nutrition for balancing the energy intake and maintaining a healthy body weight [1].

Although these data on the active lifestyle are in line with those observed in the general population (as reported by the Italian Institute of statistics [34]), the latter include also subjects between 3 and 20 years of age, who increase the rate of people with a non-sedentary behaviour and are instead excluded from the present study.

One of the main objectives of the study was to highlight possible food deficiencies and/or excesses. Accordingly, the




Fig. 4 Intake of foods to be consumed at each main meal (a), daily (b) weekly (c) and occasionally (d). Only the subgroups with statistically significant differences are displayed




Fig. 4 (continued)

overall picture emerged indicates a lower-than-recommended intake of all food categories included in the Mediterranean pyramid, together with some excesses that may impact on the subject's health status later in life. First, with regard to wine, the overall consumption was low to moderate: it is

worth noting that red wine contains flavonoids and resveratrol, known to play a role in the prevention of several diseases, including CDV and cancer [35–38]. Yet, it must be underlined that the International Agency for Research on Cancer has labeled alcohol as a Group-1 carcinogen, which

indicates that there is sufficient evidence to suggest that exposure in humans is carcinogenic [39].

More than half of the responders (especially females and those aged > 50 years) reported an insufficient intake of water per day. The risk of this behaviour is dehydration, which has been associated with increased morbidity, mortality and also with a high economic burden, particularly in the elderly, in whom the frequent coexistence of comorbidities requiring polypharmacy may exacerbate this status [40–43].

In addition to these findings, more than half of the responders declared a scarce consumption of all the basic elements of the pyramid (i.e., fruits, vegetables and cereals). An inadequate intake of these foods, which are central elements of the Mediterranean diet, exposes the whole population to specific micronutrient deficiencies that may compromise health [44]. Indeed, these foods are an important source of vitamins, minerals and, above all, antioxidants, which protect the body from oxidative damage and, thus, from cellular aging [44, 45].

On the other hand, the excessive intake of some of the foods at the top of the pyramid, that indeed should be consumed only occasionally, surfaced. A higher-than-recommended intake of sweets, red meats and processed meats increases the risk of CVD, type 2 diabetes, obesity and other metabolic diseases, due to the high content of simple sugars and saturated fats [11, 46]. In this regard, the consumption of fish and olive oil should be increased, as they are rich in mono and polyunsaturated fats that prevent the development of CVD [47]. Moreover, the intake of milk and dairy should be optimized, because an adequate consumption of these products, enriched in calcium and vitamin D, may improve bone health [48].

When the compliance with the recommendations of the Mediterranean diet pyramid was evaluated by subgroups (i.e., gender, age, geographical area and education), a rather low adherence emerged mainly among females and the elderly, followed by people from different Italian regions. In all, the differences observed by geographical area are likely indicative of the different regional culinary traditions. An important health concern regards the insufficient consumption of milk and dairy products in females and older age groups, who are at high risk for osteoporosis. Indeed, a Mediterranean-like dietary pattern has been reported to reduce bone loss in subjects with osteoporosis supplemented with vitamin D [24, 25]. Moreover, vitamin D insufficiency has been implicated in a wide spectrum of conditions, ranging from musculoskeletal diseases to cancer and CVD [49–56], and supplementation of calcium and vitamin D has shown to be beneficial in reducing the risk of fractures [57, 58]. Nevertheless, the effects and appropriate regimen of calcium and vitamin D supplementation remain highly debated [50, 56, 59, 60].

Another result that deserves attention is the fact that females and older people consume insufficient amounts of several key foods included in the pyramid: in fact, this

suggests a risk for suboptimal nutrition in children, because of the dependence from the parents' choices [61]. This is especially worrying considering that the rate of childhood obesity in Italy, although declining, is still one of the highest in Europe [28].

Finally, the present study allowed to highlight possible discrepancies between the responders' perceived and actual behaviour. Indeed, it is worth noting that 81.8% of participants had declared a regular consumption of fruits and vegetables, but only 22.7% and 32.8%, respectively, resulted compliant with the Mediterranean pyramid recommendations. This may be explained by the lack of information and awareness, by responders, on the correct intake of each food category.

The main limitation of the present study is the fact that responders do not fully represent the general Italian population. Moreover, the lack of data on BMI does not allow to correlate obesity with the compliance with the recommendations of the Mediterranean food pyramid. Finally, the food intake frequencies are self-reported, and this may lead to the actual misreporting of data. However, surveys similar to the one used here have been frequently employed.

On the other hand, the large sample size and the amount of information collected through the survey contribute to provide a reliable picture of the dietary pattern and lifestyle of participants.

From this population-based study, an overall low compliance of participants to the Mediterranean pyramid recommendations emerged, with several relevant food deficiencies and inconsistencies indicative of the low awareness of the subjects' own nutrition habits and lifestyle. Importantly, the personalized pyramid generated following survey completion should serve to the responders for a self-evaluation of their dietary pattern, to possibly raise their family's awareness on this matter, and to the healthcare professionals to plan simple intervention strategies to educate to a healthier behaviour and/or provide complete and balanced nutrient supplements. This is important in the long term, because of the positive impact that such measures may exert on public health in terms of chronic disease epidemiology and economic as well as social burden. In this context, GPs and pharmacists with their knowledge and expertise certainly play a key role and may support both health promotion and surveillance, which is required to monitor the situation and proactively evaluate the effectiveness of interventions.

Acknowledgements The authors would like to thank all participants who completed the online test. We also thank Scientific Committee members and contributors of Curare la Salute: Luigi Canciani, Alessandro Fornaro, Marcello Giovannini, Elisa Paganini, Annarosa Racca, Paolo Vintani. We thank Pfizer Consumer Healthcare, and especially Antonio Limitone, Ornella Parma and Domenico Giorgio Cassarà for supporting "Curare la salute". Editorial support was provided by Edra spa, and unconditionally funded by Pfizer Consumer Healthcare.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval All procedures performed in studies involving human participants were in accordance with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent Participation was voluntary and anonymous and completing the survey was accepted as consent by the participants.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Bach-Faig A, Berry EM, Lairon D et al (2011) Mediterranean diet pyramid today. Science and cultural updates. *Public Health Nutr* 14:2274–2284. <https://doi.org/10.1017/S1368980011002515>
- Martinez-Lacoba R, Pardo-Garcia I, Amo-Saus E, Escribano-Sotos F (2018) Mediterranean diet and health outcomes: a systematic meta-review. *Eur J Public Health* 28:955–961. <https://doi.org/10.1093/eurpub/cky113>
- Sofi F, Macchi C, Abbate R et al (2014) Mediterranean diet and health status: an updated meta-analysis and a proposal for a literature-based adherence score. *Public Health Nutr* 17:2769–2782. <https://doi.org/10.1017/S1368980013003169>
- Rees K, Hartley L, Flowers N et al (2013) “Mediterranean” dietary pattern for the primary prevention of cardiovascular disease. *Cochrane Database Syst Rev*. <https://doi.org/10.1002/14651858.CD009825.pub2>
- Panagiotakos DB, Georgousopoulou EN, Pitsavos C et al (2015) Exploring the path of Mediterranean diet on 10-year incidence of cardiovascular disease: the ATTICA study (2002–2012). *Nutr Metab Cardiovasc Dis NMCD* 25:327–335. <https://doi.org/10.1016/j.numecd.2014.09.006>
- Estruch R, Ros E, Salas-Salvadó J et al (2018) Primary prevention of cardiovascular disease with a mediterranean diet supplemented with extra-virgin olive oil or nuts. *N Engl J Med* 378:e34. <https://doi.org/10.1056/NEJMoa1800389>
- Zappalà G, Buscemi S, Mulè S et al (2018) High adherence to Mediterranean diet, but not individual foods or nutrients, is associated with lower likelihood of being obese in a Mediterranean cohort. *Eat Weight Disord EWD* 23:605–614. <https://doi.org/10.1007/s40519-017-0454-1>
- Kesse-Guyot E, Ahluwalia N, Lassale C et al (2013) Adherence to Mediterranean diet reduces the risk of metabolic syndrome: a 6-year prospective study. *Nutr Metab Cardiovasc Dis NMCD* 23:677–683. <https://doi.org/10.1016/j.numecd.2012.02.005>
- Babio N, Bulló M, Basora J et al (2009) Adherence to the Mediterranean diet and risk of metabolic syndrome and its components. *Nutr Metab Cardiovasc Dis NMCD* 19:563–570. <https://doi.org/10.1016/j.numecd.2008.10.007>
- Khalili-Moghadam S, Mirmiran P, Bahadoran Z, Azizi F (2018) The Mediterranean diet and risk of type 2 diabetes in Iranian population. *Eur J Clin Nutr*. <https://doi.org/10.1038/s41430-018-0336-2>
- Vitale M, Masulli M, Calabrese I et al (2018) Impact of a Mediterranean dietary pattern and its components on cardiovascular risk factors, glucose control, and body weight in people with type 2 diabetes: a real-life study. *Nutrients*. <https://doi.org/10.3390/nu10081067>
- Esposito K, Giugliano D (2014) Mediterranean diet and type 2 diabetes. *Diabetes Metab Res Rev* 30(Suppl 1):34–40. <https://doi.org/10.1002/dmrr.2516>
- Bonaccio M, Di Castelnuovo A, Costanzo S et al (2016) Adherence to the traditional Mediterranean diet and mortality in subjects with diabetes. Prospective results from the MOLI-SANI study. *Eur J Prev Cardiol* 23:400–407. <https://doi.org/10.1177/2047487315569409>
- Maraki MI, Yannakoulia M, Stamelou M et al (2018) Mediterranean diet adherence is related to reduced probability of prodromal Parkinson's disease. *Mov Disord*. <https://doi.org/10.1002/mds.27489>
- Solfrizzi V, Custodero C, Lozupone M et al (2017) Relationships of dietary patterns, foods, and micro- and macronutrients with Alzheimer's disease and late-life cognitive disorders: a systematic review. *J Alzheimers Dis JAD* 59:815–849. <https://doi.org/10.3233/JAD-170248>
- Solfrizzi V, Panza F (2014) Mediterranean diet and cognitive decline. A lesson from the whole-diet approach: what challenges lie ahead? *J Alzheimers Dis JAD* 39:283–286. <https://doi.org/10.3233/JAD-130831>
- Aridi YS, Walker JL, Wright ORL (2017) The association between the Mediterranean dietary pattern and cognitive health: a systematic review. *Nutrients*. <https://doi.org/10.3390/nu9070674>
- Bravi F, Spei M-E, Polesel J et al (2018) Mediterranean diet and bladder cancer risk in Italy. *Nutrients*. <https://doi.org/10.3390/nu10081061>
- Filomeno M, Bosetti C, Bidoli E et al (2015) Mediterranean diet and risk of endometrial cancer: a pooled analysis of three Italian case-control studies. *Br J Cancer* 112:1816–1821. <https://doi.org/10.1038/bjc.2015.153>
- Turati F, Bravi F, Polesel J et al (2017) Adherence to the Mediterranean diet and nasopharyngeal cancer risk in Italy. *Cancer Causes Control CCC* 28:89–95. <https://doi.org/10.1007/s10552-017-0850-x>
- Turati F, Carioli G, Bravi F et al (2018) Mediterranean diet and breast cancer risk. *Nutrients*. <https://doi.org/10.3390/nu10030326>
- Schwingshackl L, Schwedhelm C, Galbete C, Hoffmann G (2017) Adherence to Mediterranean diet and risk of cancer: an updated systematic review and meta-analysis. *Nutrients*. <https://doi.org/10.3390/nu9101063>
- Bonaccio M, Di Castelnuovo A, Costanzo S et al (2018) Mediterranean diet and mortality in the elderly: a prospective cohort study and a meta-analysis. *Br J Nutr* 120:841–854. <https://doi.org/10.1017/S0007114518002179>
- Jennings A, Cashman KD, Gillings R et al (2018) A Mediterranean-like dietary pattern with vitamin D3 (10 µg/day) supplements reduced the rate of bone loss in older Europeans with osteoporosis at baseline: results of a 1-y randomized controlled trial. *Am J Clin Nutr*. <https://doi.org/10.1093/ajcn/nqy122>
- Sahni S, Mangano KM, Kiel DP et al (2017) Dairy intake is protective against bone loss in older vitamin D supplement users: the framingham study. *J Nutr* 147:645–652. <https://doi.org/10.3945/jn.116.240390>

26. Bracale R, Milani L, Ferrara E et al (2013) Childhood obesity, overweight and underweight: a study in primary schools in Milan. *Eat Weight Disord EWD* 18:183–191. <https://doi.org/10.1007/s40519-013-0036-9>
27. Spinelli A, Nardone P, Buoncristiano M et al (2015) Italy 2014: childhood obesity is decreasing. *Epidemiol Prev* 39:63
28. Renzella J, Townsend N, Jewel J et al (2018) What national and subnational interventions and policies based on Mediterranean and Nordic diets are recommended or implemented in the WHO European Region, and is there evidence of effectiveness in reducing non-communicable diseases? http://www.euro.who.int/__data/assets/pdf_file/0011/365285/hen-58-eng.pdf?ua=1. Accessed 14 Oct 2019
29. Balzaretto CM, Ventura V, Ratti S et al (2018) Improving the overall sustainability of the school meal chain: the role of portion sizes. *Eat Weight Disord EWD*. <https://doi.org/10.1007/s40519-018-0524-z>
30. SINU—Società Italiana di Nutrizione Umana. <https://sinu.it/>. Accessed 14 Oct 2019
31. Aspetti della vita quotidiana (2016). <https://www.istat.it/it/archivio/186843>. Accessed 26 Sep 2019
32. Istat.it Multiscopo sulle famiglie (2018) Aspetti della vita quotidiana—parte generale. <https://www.istat.it/it/archivio/217037>. Accessed 14 Oct 2019
33. Mayer-Davis EJ, Costacou T (2001) Obesity and sedentary lifestyle: modifiable risk factors for prevention of type 2 diabetes. *Curr Diabetes Rep* 1:170–176
34. EUPATI (2017) Fattori di rischio per la salute: fumo, obesità, alcol e sedentarietà. <https://www.istat.it/it/archivio/202040>. Accessed 26 Sep 2019
35. Fernandes I, Pérez-Gregorio R, Soares S et al (2017) Wine flavonoids in health and disease prevention. *Mol Basel Switz*. <https://doi.org/10.3390/molecules22020292>
36. Haseeb S, Alexander B, Baranchuk A (2017) Wine and cardiovascular health: a comprehensive review. *Circulation* 136:1434–1448. <https://doi.org/10.1161/CIRCULATIONAHA.117.030387>
37. Bonnefont-Rousselot D (2016) Resveratrol and cardiovascular diseases. *Nutrients*. <https://doi.org/10.3390/nu8050250>
38. Farinetti A, Zurlo V, Manenti A et al (2017) Mediterranean diet and colorectal cancer: a systematic review. *Nutrition* (Burbank Los Angel City California) 43–44:83–88. <https://doi.org/10.1016/j.nut.2017.06.008>
39. International Agency for Research on Cancer, International Agency for Research on Cancer (1988) Alcohol drinking: views and experts opinions of an IARC Working Group on the Evaluation of Carcinogenic Risks to Humans, which met in Lyon 13–20 Oct. 1987. Lyon. <https://monographs.iarc.fr/wpcontent/uploads/2018/06/mono98.pdf>. Accessed 14 Oct 2019
40. Warren JL, Bacon WE, Harris T et al (1994) The burden and outcomes associated with dehydration among US elderly, 1991. *Am J Public Health* 84:1265–1269
41. Popkin BM, D’Anci KE, Rosenberg IH (2010) Water, hydration, and health. *Nutr Rev* 68:439–458. <https://doi.org/10.1111/j.1753-4887.2010.00304.x>
42. Frangeskou M, Lopez-Valcarcel B, Serra-Majem L (2015) Dehydration in the elderly: a review focused on economic burden. *J Nutr Health Aging* 19:619–627. <https://doi.org/10.1007/s12603-015-0491-2>
43. El-Sharkawy AM, Sahota O, Lobo DN (2015) Acute and chronic effects of hydration status on health. *Nutr Rev* 73(Suppl 2):97–109. <https://doi.org/10.1093/nutrit/nuv038>
44. Castro-Quezada I, Román-Viñas B, Serra-Majem L (2014) The Mediterranean diet and nutritional adequacy: a review. *Nutrients* 6:231–248. <https://doi.org/10.3390/nu6010231>
45. Manson JE, Bassuk SS (2018) Vitamin and mineral supplements: what clinicians need to know. *JAMA* 319:859–860. <https://doi.org/10.1001/jama.2017.21012>
46. DiNicolantonio JJ, Lucan SC, O’Keefe JH (2016) The evidence for saturated fat and for sugar related to coronary heart disease. *Prog Cardiovasc Dis* 58:464–472. <https://doi.org/10.1016/j.pcad.2015.11.006>
47. Widmer RJ, Flammer AJ, Lerman LO, Lerman A (2015) The Mediterranean diet, its components, and cardiovascular disease. *Am J Med* 128:229–238. <https://doi.org/10.1016/j.amjmed.2014.10.014>
48. van den Heuvel EGHM, Steijns JMJM (2018) Dairy products and bone health: how strong is the scientific evidence? *Nutr Res Rev* 31:164–178. <https://doi.org/10.1017/S095442241800001X>
49. Feldman D, Krishnan AV, Swami S et al (2014) The role of vitamin D in reducing cancer risk and progression. *Nat Rev Cancer* 14:342–357. <https://doi.org/10.1038/nrc3691>
50. Wintermeyer E, Ihle C, Ehnert S et al (2016) Crucial role of vitamin D in the musculoskeletal system. *Nutrients*. <https://doi.org/10.3390/nu8060319>
51. Pilz S, Trummer C, Pandis M et al (2018) Vitamin D: Current Guidelines and Future Outlook. *Anticancer Res* 38:1145–1151. <https://doi.org/10.21873/anticancer.12333>
52. Norman PE, Powell JT (2014) Vitamin D and cardiovascular disease. *Circ Res* 114:379–393. <https://doi.org/10.1161/CIRCRESAHA.113.301241>
53. Holick MF (2007) Vitamin D deficiency. *N Engl J Med* 357:266–281. <https://doi.org/10.1056/NEJMr070553>
54. Wacker M, Holick MF (2013) Vitamin D—effects on skeletal and extraskeletal health and the need for supplementation. *Nutrients* 5:111–148. <https://doi.org/10.3390/nu5010111>
55. Wimalawansa SJ (2018) Non-musculoskeletal benefits of vitamin D. *J Steroid Biochem Mol Biol* 175:60–81. <https://doi.org/10.1016/j.jsbmb.2016.09.016>
56. Chowdhury R, Kunutsor S, Vitezova A et al (2014) Vitamin D and risk of cause specific death: systematic review and meta-analysis of observational cohort and randomised intervention studies. *BMJ* 348:g1903
57. Weaver CM, Alexander DD, Boushey CJ et al (2016) Calcium plus vitamin D supplementation and risk of fractures: an updated meta-analysis from the National Osteoporosis Foundation. *Osteoporos Int J (Establ Result Coop Eur Found Osteoporos Natl Osteoporos Found USA)* 27:367–376. <https://doi.org/10.1007/s00198-015-3386-5>
58. Avenell A, Mak JCS, O’Connell D (2014) Vitamin D and vitamin D analogues for preventing fractures in post-menopausal women and older men. *Cochrane Database Syst Rev*. <https://doi.org/10.1002/14651858.CD000227.pub4>
59. Kahwati LC, Weber RP, Pan H et al (2018) Vitamin D, calcium, or combined supplementation for the primary prevention of fractures in community-dwelling adults: evidence report and systematic review for the us preventive services task force. *JAMA* 319:1600–1612. <https://doi.org/10.1001/jama.2017.21640>
60. Zhao J-G, Zeng X-T, Wang J, Liu L (2017) Association between calcium or vitamin d supplementation and fracture incidence in community-dwelling older adults: a systematic review and meta-analysis. *JAMA* 318:2466–2482. <https://doi.org/10.1001/jama.2017.19344>
61. Bracale R, Milani Marin LE, Russo V et al (2015) Family lifestyle and childhood obesity in an urban city of Northern Italy. *Eat Weight Disord EWD* 20:363–370. <https://doi.org/10.1007/s40519-015-0179-y>

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.