

Provided for non-commercial research and education use.  
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Contents lists available at SciVerse ScienceDirect

## Animal Reproduction Science

journal homepage: [www.elsevier.com/locate/anireprosci](http://www.elsevier.com/locate/anireprosci)

## Development of corpus luteum susceptibility to an analog of prostaglandin $F_{2\alpha}$ , throughout the luteal phase in llamas (*Lama glama*)

C.P. Bianchi\*, M.V. Cavilla, E. Jorgensen, M.A. Benavente, M.A. Aba

Área de Endocrinología, Facultad de Ciencias Veterinarias, U.N.C.P.B.A., Campus Universitario, Paraje Arroyo Seco, Tandil-7000-Buenos Aires, Argentina

## ARTICLE INFO

## Article history:

Received 15 July 2011

Received in revised form 2 March 2012

Accepted 8 March 2012

Available online 17 March 2012

## Keywords:

Corpus luteum

Luteolysis

Prostaglandin

Llamas

## ABSTRACT

The aim of the present study was to evaluate the susceptibility of the corpus luteum to d-cloprostenol (synthetic analog of  $PGF_{2\alpha}$ ) throughout the luteal phase in llamas. Female llamas ( $n=43$ ) were induced to ovulate by GnRH injection in the presence of an ovulatory follicle and randomly assigned into one of six groups: control and treated with an injection of d-cloprostenol on Day 3, 4, 5, 6 or 8 post GnRH. Blood samples were collected to determine plasma progesterone concentrations. There was no effect of treatment on animals injected on Day 3 or 4 post-GnRH. In animals treated on Day 5, different responses were observed. No effect of treatment was recorded in 27% of the animals whereas 55% of the llamas showed a transitory decrease followed by a recovery in plasma progesterone concentrations after d-cloprostenol injection, indicative of a resurgence of the corpus luteum, extending the luteal phase a day more than in control animals. In the remaining 18% of the animals injected on Day 5, (corresponding to those exhibiting the greatest plasma progesterone concentrations at the day of injection), complete luteolysis was observed. Plasma progesterone concentrations decreased to below  $1 \text{ ng ml}^{-1}$  24 h after d-cloprostenol in llamas injected on Day 6 or 8 post-GnRH. In conclusion, the corpus luteum of llamas is completely refractory to  $PGF_{2\alpha}$  until Day 4 after induction of ovulation, being partially sensitive by Day 5 and fully responsive to  $PGF_{2\alpha}$ , by Day 6 after induction of ovulation.

© 2012 Elsevier B.V. All rights reserved.

### 1. Introduction

Prostaglandin  $F_{2\alpha}$  ( $PGF_{2\alpha}$ ) has been recognized as the most important luteolysin in most domestic animals (McCracken et al., 1999). Luteolytic pulses of  $PGF_{2\alpha}$  are released from the endometrium at the end of the luteal phase to induce luteolysis. Nevertheless, the corpus luteum is sensitive to  $PGF_{2\alpha}$  earlier during the luteal phase although the time when a corpus luteum becomes susceptible to  $PGF_{2\alpha}$  varies between species (Cows: Louis et al., 1975; mares: Allen and Cooper, 1975; sheep: Rubianes et al., 2003; gilts: Estill et al., 1993).

In cows, the corpus luteum becomes sensitive to  $PGF_{2\alpha}$  on Day 4 following ovulation. Administration of  $PGF_{2\alpha}$

by either intrauterine or intramuscular route induces a decrease in the diameter of the corpus luteum and a decrease in plasma progesterone concentrations while the female returns to estrus 3–4 days later (Tervit et al., 1973; Louis et al., 1974). The corpus luteum of the ewe is susceptible to  $PGF_{2\alpha}$  at Day 3 post ovulation (Rubianes et al., 2003), which is earlier than had been previously known (Acritopoulou and Haresing, 1980). The corpus luteum of mares fully responds to  $PGF_{2\alpha}$  when it is administered by intrauterine, intramuscular or subcutaneous route after Day 4 post ovulation, inducing complete luteolysis (Allen and Cooper, 1975; Douglas and Ginther, 1975). In this species, when  $PGF_{2\alpha}$  is injected on Day 3 after ovulation an immediate functional and structural regression occurs followed by a later resurgence of the corpus luteum (Bergfelt et al., 2006). The period of luteal sensitivity seems to be shorter in pigs in comparison to other domestic species.  $PGF_{2\alpha}$  is effective to induce luteolysis around Day

\* Corresponding author. Tel.: +54 2293 439850; fax: +54 2293 439850.  
E-mail address: [cbianchi@vet.unicen.edu.ar](mailto:cbianchi@vet.unicen.edu.ar) (C.P. Bianchi).

12 after ovulation (Guthrie and Polge, 1976; Hallford et al., 1975). However, multiple injections between Day 5 and 10 are effective to cause premature luteolysis (Estill et al., 1993).

Llamas are induced ovulators, requiring copulation at the time when an ovulatory follicle is present to initiate the ovulatory process (San-Martín et al., 1968; Aba et al., 1995). The interval between mating and ovulation is approximately 30 h (range 24–48 h) (Adams et al., 1989; Vaughan and Tibary, 2006). Thereafter, plasma concentrations of progesterone increase on Day 4 after induction of ovulation, peak on Day 8 and, in non pregnant llamas, decrease by Days 10–11 in response to the release of PGF<sub>2α</sub> from the endometrium, attaining basal concentrations by Day 12 post mating (Aba et al., 1995). Small peaks of PGF<sub>2α</sub> are observed as early as Day 7 after mating, being the luteolytic pulses recorded between Days 8 and 13 when luteolysis occurs (Aba et al., 1995, 2000).

PGF<sub>2α</sub> and its synthetic analogs have been widely used in different species. PGF<sub>2α</sub> is considered the most potent luteolytic substance and it has been broadly used in different estrous synchronization protocols in cows (Hiers et al., 2003; Stevenson and Phatak, 2010), sheep (Menchaca et al., 2004) and mares (Loy et al., 1980) to ensure corpus luteum regression. Due to the capacity of PGF<sub>2α</sub> to induce myometrial contractions, it has also been used to treat uterine infections (Hirsbrunner et al., 2003; Kasimanickam et al., 2005) and to prevent post mating endometritis in mares (Combs et al., 1996). In llamas, cloprostenol, an analog of PGF<sub>2α</sub>, has been used as part of ovarian follicular superstimulation treatments (Bourke et al., 1995) and as abortifacient (Smith et al., 2000).

Information regarding the precise moment when the corpus luteum attains the capacity to respond to PGF<sub>2α</sub> in many domestic species has been known for many years, however, no reports are available regarding when the corpus luteum of llamas becomes susceptible to PGF<sub>2α</sub>. Thus, the aim of the present study was to investigate the susceptibility of the llama corpus luteum to an injection of d-cloprostenol (synthetic analog of PGF<sub>2α</sub>) on different days of the luteal phase.

## 2. Materials and methods

Field studies were performed in compliance with animal welfare regulations set by the Faculty of Veterinary Sciences, UNCPBA where activities were conducted. Facilities are located in Tandil, Argentina, at 37°S, 60°W. Animals were kept in pens isolated from males and fed pasture hay and water *ad libitum*. Female llamas ( $n=43$ ) were examined daily by transrectal ultrasonography to assess ovarian status (Mindray, DP 6600 Vet, with 5.0/7.5 variable traducer probe). When a follicle with a diameter  $\geq 8$  mm, considered ovulatory in this species (Bravo et al., 1991), was observed, ovulation was induced with a single injection of GnRH (16.8  $\mu\text{g}$  Receptal®, Intervet, Argentina). Occurrence of ovulation was assessed based on ultrasonographic visualization of the corpus luteum and further confirmed by the progesterone profiles. Thereafter, animals were randomly divided into six different groups: Control ( $n=10$ ; untreated) and G3 ( $n=4$ ); G4 ( $n=7$ ); G5 ( $n=11$ ); G6 ( $n=5$ )

and G8 ( $n=6$ ) which were treated with an injection of d-cloprostenol (112.5  $\mu\text{g}$ , Baker, Tecnofarm®, Argentina) into the hind leg on Days 3, 4, 5, 6 or 8 post-GnRH, respectively.


Blood samples for progesterone determinations were collected every second day until the day of d-cloprostenol injection and thereafter samples were obtained daily until the end of the study (Day 11 or 12 post-GnRH). Samples were centrifuged and plasma was stored at  $-20^\circ\text{C}$  until hormone assays were performed. Progesterone was measured using an RIA kit (COAT-A-COUNT®, Siemens Medical Solutions Diagnostics, Los Angeles, CA, USA) previously validated for use with llama plasma (Bianchi et al., 2007). The sensitivity of the assay was  $0.1 \text{ ng ml}^{-1}$  and the intra and inter-assay coefficient of variation were below 14% for concentrations between 0.1 and  $40 \text{ ng ml}^{-1}$ .

Data were analyzed using an ANOVA test to detect differences in the diameter of the largest follicle at the time of induction of ovulation between the different groups. Serial data were compared using the mixed model procedure of SAS (Statistical Analysis System Institute Inc., Cary, NC, USA) to determine main effect of treatment, days and their interactions. The compound symmetry model with a covariance structure was used. When the interactions were significant, the effect of days within treatment was compared by least squares means (adjustment Tukey–Kramer). Data are presented as mean  $\pm$  S.E.M., and differences were considered to be significant when  $P < 0.05$ .

## 3. Results

Mean diameters of the largest follicle at the time when GnRH was injected were similar in all groups ( $P=0.59$ ) and averaged  $9.6 \pm 0.42 \text{ mm}$ .

Plasma concentrations of progesterone were similar for llamas injected on Days 3 or 4 to controls and increased above  $1 \text{ ng ml}^{-1}$  on Day 5, had peaked on Day 8, started to decrease by Days 9 to 10 and had attained basal concentrations between Days 11 and 12 post induction of ovulation (Fig. 1A–C). In animals injected with d-cloprostenol on Day 5, diverse responses were observed. In 27% (3/11) animals, d-cloprostenol injection did not cause any effect, the pattern of progesterone concentrations were similar to that reported in the control group. A transient decrease (below  $1 \text{ ng ml}^{-1}$ ) in plasma progesterone concentrations was recorded 24 h after treatment in 55% (6/11) of the animals. One day later, circulating progesterone concentrations had increased and were maintained above  $1 \text{ ng ml}^{-1}$  until Day 11, one day longer than in animals from the control group. By Day 12, an important decrease was observed ( $0.17 \pm 0.07 \text{ ng ml}^{-1}$ ). Meanwhile, in 18% (2/11) of the animals in which plasma progesterone concentrations were above  $3 \text{ ng ml}^{-1}$  on Day 5, d-cloprostenol injection induced a complete luteolysis (Fig. 1D). In animals treated on Day 6 or 8 post GnRH, plasma progesterone concentrations had decreased to less than  $1 \text{ ng ml}^{-1}$  by 24 h after d-cloprostenol injection while in the control group plasma progesterone concentrations remained greater than  $2 \text{ ng ml}^{-1}$  on those days ( $P \leq 0.0001$ ; Fig. 1E and F).



**Fig. 1.** Mean plasma progesterone (P<sub>4</sub>) concentrations in control animals (A) and in animals injected with d-cloprostenol, synthetic analog of PGF<sub>2α</sub>, on Day 3 (B); Day 4 (C); Day 5 (D); Day 6 (E) or Day 8 (F) after induction of ovulation; (\*) indicates values statistically different between control and treated groups (P ≤ 0.0001).

#### 4. Discussion

Plasma concentrations of progesterone in the control group were similar to those reported in other studies. Progesterone reaches maximum concentrations at Day 8 and a decrease is observed between Days 10 and 12 after induction of ovulation that occurs subsequent to the luteolytic release of PGF<sub>2α</sub> (Aba et al., 1995).

The corpus luteum of llamas fully responds to d-cloprostenol, synthetic analog of PGF<sub>2α</sub>, by Day 6 after induction of ovulation. The observation that animals treated on Days 3 or 4 post induction of ovulation did not respond to d-cloprostenol injection suggests that the newly formed corpus luteum of llamas is refractory to PGF<sub>2α</sub> before Day 5. Many factors have been involved in the mechanism responsible for the refractoriness of the young corpus luteum to PGF<sub>2α</sub> in ruminants. However, clearly, it is not due to the lack of PGF<sub>2α</sub> receptors or receptor affinity (Cows: Tsai and Wiltbank, 1998; Sheep: Juengel et al., 1996). In cows, the existence of a positive feedback loop between PGF<sub>2α</sub> secretion and the expression of PG G/H synthase-2 (limiting enzyme in the synthesis of prostaglandin) in the mid-estrous cycle corpus luteum could be related to its responsiveness to PGF<sub>2α</sub> (Tsai and Wiltbank, 1998). In sheep, Silva et al. (2000) demonstrated that young and unresponsive corpora lutea have a greater capacity to catabolize PGF<sub>2α</sub> than more mature and sensitive corpora lutea. Several other factors, such as the development of endothelin-1 and its receptor (Levy et al., 2000); the differential expression of genes associated to PGF<sub>2α</sub> receptors (Goravanahally et al., 2009) and the regulation of many angiogenic factors (Shirasuna et al., 2010) has been involved in the achievement of responsiveness in other ruminant species. Nevertheless, the precise mechanisms responsible for the insensitivity of the newly developed corpus luteum to the luteolytic action of PGF<sub>2α</sub> are unknown.

The observation that animals within G5 showed different plasma progesterone concentrations at the time of d-cloprostenol injection could be in relation to a different time elapsed from induction of ovulation to ovulation within llamas (Adams et al., 1989). Thus, the corpus luteum could be some hours more mature in llamas with greater progesterone concentrations and fully responsive to PGF<sub>2α</sub> than in those with lesser concentrations at the time of PGF<sub>2α</sub> treatment. Similar results were recorded in mares where PGF<sub>2α</sub> treatment on Day 3 after ovulation provoked different responses within animals. An abrupt decrease in plasma progesterone concentrations was induced with a PGF<sub>2α</sub> injection in 25% of mares with greater plasma progesterone concentrations at the day of treatment while detectable corpus luteum resurgence was observed in those mares exhibiting lesser progesterone concentrations (75%). In this latter study, there is no information about plasma progesterone concentrations at the end of the luteal phase in mares with corpus luteum resurgences (Bergfelt et al., 2006). However, in the present study, 54% of llamas injected on Day 5 (*i.e.* around 3 days post ovulation) showed corpus luteum resurgence based on plasma progesterone concentrations and a day longer luteal phase than the control group.

Similar to the results observed in llamas injected on Day 5, a transient luteolytic effect was observed in ewes after the administration of a sub-luteolytic dose of PGF<sub>2α</sub> (Juengel et al., 2000). Nevertheless, the selected dose in the present study has demonstrated to be effective to induce a premature and complete luteolysis on Days 6 and 8 post induction of ovulation. In pigs, the administration of two doses of PGF<sub>2α</sub> per day was able to induce an earlier luteolysis (Estill et al., 1993). Thus, new studies will be necessary to test if a similar dose regimen on Day 5 could be able to induce a complete regression of the corpus luteum in llamas.

#### 5. Conclusion

In summary, the results of the present study indicate that the corpus luteum of llamas is refractory to PGF<sub>2α</sub> during the first 4 days after induction of ovulation, being partially sensitive by Day 5 depending on developmental status of the corpus luteum and becoming fully responsive by Day 6 post induction of ovulation.

#### Acknowledgments

This research was supported by grants from Agencia Nacional de Promoción Científica y Tecnológica (PICT N° 2074). We gratefully acknowledge Tecnofarm®, Argentina for providing d-cloprostenol. C.P. Bianchi and M.V. Cavilla are holders of fellowships from CONICET and M.A. Benavente from CICPBA.

#### References

- Aba, M.A., Forsberg, M., Kindahl, H., Sumar, J., Edqvist, L.-E., 1995. Endocrine changes after mating in pregnant and non-pregnant llamas and alpacas. *Acta Vet. Scand.* 36, 489–498.
- Aba, M.A., Kindahl, H., Forsberg, M., Quiroga, M., Auza, N., 2000. Levels of progesterone and changes in prostaglandin F<sub>2α</sub> release during luteolysis and early pregnancy in llamas and the effect of treatment with flunixin meglumine. *Anim. Reprod. Sci.* 59, 87–97.
- Acritopoulou, S., Haresing, N., 1980. Response of ewes to a single injection of an analogue of PGF – 2α given at different stages of the oestrous cycle. *J. Reprod. Fertil.* 58, 219–223.
- Adams, G.P., Griffin, P.G., Ginther, O.J., 1989. In situ morphologic dynamics of ovaries, uterus and cervix in llamas. *Biol. Reprod.* 41, 551–558.
- Allen, W.R., Cooper, J., 1975. The use of synthetic analogues of prostaglandins for inducing luteolysis in mares. *Ann. Biol. Anim. Biochim. Biophys.* 15, 461–469.
- Bergfelt, D.R., Pierson, R.A., Ginther, O.J., 2006. Regression and resurgence of the CL following PGF<sub>2α</sub> treatment 3 days after ovulation in mares. *Theriogenology* 65, 1605–1619.
- Bianchi, C.P., Meikle, A., Sartore, I., González, F., Aba, M.A., 2007. Uterine estrogen receptor alpha and progesterone receptor during the follicular and luteal phase in llamas (*Lama glama*). *Anim. Reprod. Sci.* 99, 117–126.
- Bourke, D.A., Kyle, C.E., McEvoy, T.G., Young, P., Adam, C.L., 1995. Superovulatory responses to eCG in llamas (*Lama glama*). *Theriogenology* 44, 255–268.
- Bravo, P.W., Stabenfeldt, G.H., Lasley, B.L., Fowler, M.E., 1991. The effect of ovarian follicle size on pituitary and ovarian responses to copulation in domesticated South American Camelids. *Biol. Reprod.* 45, 553–559.
- Combs, G.B., LeBlanc, M.M., Neuwirth, L., Tran, T.Q., 1996. Effects of prostaglandin F<sub>2α</sub>, Cloprostenol and Fenprostalene on uterine clearance of radiocolloid in the mare. *Theriogenology* 45, 1449–1455.
- Douglas, R.H., Ginther, O.J., 1975. Effects of prostaglandin F<sub>2α</sub> on estrous cycle or corpus luteum in mares and gilts. *J. Anim. Sci.* 40, 518–522.
- Estill, C.T., Britt, J.H., Gadsby, J.E., 1993. Repeated administration of prostaglandin F<sub>2α</sub> during the early luteal phase causes premature luteolysis in the pig. *Biol. Reprod.* 49, 181–185.

- Goravannahally, M.P., Salem, M., Yao, J., Inskip, E.K., Flores, J.A., 2009. Differential gene expression in the bovine corpus luteum during transition from early phase to midphase and its potential role in acquisition of luteolytic sensitive to prostaglandin F<sub>2</sub> alpha. *Biol. Reprod.* 80, 980–988.
- Guthrie, H.D., Polge, C., 1976. Luteal function and oestrus in gilts treated with a synthetic analogue of prostaglandin F<sub>2α</sub> (ICI 79,939) at various times during the oestrous cycle. *J. Reprod. Fertil.* 48, 423–425.
- Hallford, D.M., Wettemann, R.P., Turman, E.J., Omtvedt, I.T., 1975. Luteal function in gilts after prostaglandin F<sub>2α</sub>. *J. Anim. Sci.* 41, 1706–1710.
- Hiers, E.A., Barthle, C.R., Dahms, M.K.V., Portillo, G.E., Bridges, G.A., Rae, D.O., Thatcher, W.W., Yelich, J.V., 2003. Synchronization of *Bos indicus* × *Bos taurus* cows for timed artificial insemination using gonadotropin-releasing hormone plus prostaglandin F<sub>2α</sub> in combination with melengestrol acetate. *J. Anim. Sci.* 81, 830–835.
- Hirsbrunner, G., Knutti, B., K pfer, U., Burkhardt, H., Steiner, A., 2003. Effect of prostaglandin E<sub>2</sub>, DL – cloprostenol, and prostaglandin E<sub>2</sub> in combination with D – cloprostenol on uterine motility during diestrus in experimental cows. *Anim. Reprod. Sci.* 79, 17–32.
- Juengel, J.L., Wiltbank, M.C., Meberg, B.M., Niswender, G.D., 1996. Regulation of steady-state concentrations of messenger ribonucleic acid encoding prostaglandin F<sub>2α</sub> receptor in ovine corpus luteum. *Biol. Reprod.* 54, 1096–1102.
- Juengel, J.L., Haworth, J.D., Rollyson, M.K., Silva, P.J., Sawyer, H.R., Niswender, G.D., 2000. Effect of dose of prostaglandin F<sub>2α</sub> on steroidogenic components and oligonucleosomes in ovine luteal tissue. *Biol. Reprod.* 62, 1047–1051.
- Kasimanickam, R., Duffield, T.F., Foster, R.A., Gartley, C.J., Leslie, K.E., Walton, J.S., Johnson, W.H., 2005. The effect of a single administration of cephalirin or cloprostenol on the reproductive performance of dairy cows with subclinical endometritis. *Theriogenology* 63, 818–830.
- Levy, N., Kobayashi, S., Roth, Z., Wolfenson, D., Miyamoto, A., Meidan, R., 2000. Administration of prostaglandin F<sub>2α</sub> during early bovine luteal phase does not alter the expression of ET-1 and of its type A receptor: a possible cause for corpus luteum refractoriness. *Biol. Reprod.* 63, 377–382.
- Louis, T.M., Hafs, H.D., Morrow, D.A., 1974. Intrauterine administration of prostaglandin F<sub>2α</sub> in cows: progesterone, estrogen, LH, estrus and ovulation. *J. Anim. Sci.* 38, 347–353.
- Louis, T.M., Hafs, H.D., Stellflug, J.N., 1975. Control of ovulation, fertility and endocrine response after prostaglandin F<sub>2α</sub> in cattle. *Ann. Biol. Anim. Biochim. Biophys.* 15, 407–417.
- Loy, R.G., Pemstein, R., O’Canna, D., Douglas, R.H., 1980. Control of ovulation in cycling mares with ovarian steroids and prostaglandin. *Theriogenology* 15, 191–200.
- McCraken, J.A., Custer, E.E., Lamsa, J.A., 1999. Luteolysis: a neuroendocrine-mediated event. *Physiol. Rev.* 79, 263–324.
- Menchaca, A., Miller, V., Gil, J., Pinczak, A., Laca, M., Rubianes, E., 2004. Prostaglandin F<sub>2α</sub> treatment associated with timed artificial insemination in ewes. *Reprod. Domest. Anim.* 39, 352–355.
- Rubianes, E., Menchaca, A., Carbajal, B., 2003. Response of the 1–5 day-aged ovine corpus luteum to prostaglandin F<sub>2α</sub>. *Anim. Reprod. Sci.* 78, 47–55.
- San-Martin, M., Copaira, M., Z niga, R., Rodriguez, R., Bustinza, G., Acosta, L., 1968. Aspects in reproduction in alpacas. *J. Reprod. Fertil.* 16, 395–399.
- Shirasuna, K., Sasahara, K., Matsui, M., Shimizu, T., Miyamoto, A., 2010. Prostaglandin F<sub>2α</sub> differentially affects mRNA expression relating to angiogenesis, vasoactivation and prostaglandins in the early and mid corpus luteum in the cow. *J. Reprod. Dev.* 56, 428–436.
- Silva, P.J., Juengel, J.L., Rollyson, M.K., Niswender, G.D., 2000. Prostaglandin metabolism in the ovine corpus luteum: catabolism of prostaglandin F<sub>2α</sub> (PGF<sub>2α</sub>) coincides with resistance of the corpus luteum to PGF<sub>2α</sub>. *Biol. Reprod.* 63, 1229–1236.
- Smith, B.B., Timm, T.L., Reed, P.J., Christensen, M., 2000. Use of cloprostenol as an abortifacient in the llama (*Lama glama*). *Theriogenology* 54, 497–505.
- Stevenson, J.S., Phatak, A.P., 2010. Rates of luteolysis and pregnancy in dairy cows after treatment with cloprostenol or dinoprost. *Theriogenology* 73, 1127–1138.
- Tervit, H.R., Rowson, E.A., Brand, A., 1973. Synchronization of oestrus in cattle using a prostaglandin F<sub>2α</sub> analogue (ICI 79939). *J. Reprod. Fertil.* 34, 179–181.
- Tsai, S.-J., Wiltbank, M.C., 1998. Prostaglandin F<sub>2α</sub> regulates distinct physiological changes in early and mid-cycle bovine corpora lutea. *Biol. Reprod.* 58, 346–352.
- Vaughan, J.L., Tibary, A., 2006. Reproduction in female South American camelids: a review and clinical observations. *Small Rumin. Res.* 61, 259–281.