



## LJMU Research Online

**Franceschi, A, Robinson, M, Owens, D, Brownlee, T, Ferrari Bravo, D and Enright, K**

**Reliability and sensitivity to change of post-match physical performance measures in elite youth soccer players**

<http://researchonline.ljmu.ac.uk/id/eprint/20349/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Franceschi, A, Robinson, M, Owens, D, Brownlee, T, Ferrari Bravo, D and Enright, K (2023) Reliability and sensitivity to change of post-match physical performance measures in elite youth soccer players. *Frontiers in Sports and Active Living*. 5. ISSN 2624-9367**

LJMU has developed [LJMU Research Online](http://researchonline.ljmu.ac.uk/) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>



## OPEN ACCESS

## EDITED BY

James Malone,  
Liverpool Hope University, United Kingdom

## REVIEWED BY

Raphael Luiz Sakugawa,  
Federal University of Santa Catarina, Brazil  
Casey Watkins,  
Seattle University, United States

## \*CORRESPONDENCE

Thomas Brownlee  
✉ t.brownlee@bham.ac.uk

RECEIVED 24 February 2023

ACCEPTED 30 June 2023

PUBLISHED 13 July 2023

## CITATION

Franceschi A, Robinson MA, Owens D,  
Brownlee T, Ferrari Bravo D and Enright K  
(2023) Reliability and sensitivity to change of  
post-match physical performance measures in  
elite youth soccer players.  
Front. Sports Act. Living 5:1173621.  
doi: 10.3389/fspor.2023.1173621

## COPYRIGHT

© 2023 Franceschi, Robinson, Owens,  
Brownlee, Ferrari Bravo and Enright. This is an  
open-access article distributed under the terms  
of the [Creative Commons Attribution License  
\(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in  
other forums is permitted, provided the original  
author(s) and the copyright owner(s) are  
credited and that the original publication in this  
journal is cited, in accordance with accepted  
academic practice. No use, distribution or  
reproduction is permitted which does not  
comply with these terms.

# Reliability and sensitivity to change of post-match physical performance measures in elite youth soccer players

Alberto Franceschi<sup>1,2</sup> , Mark A. Robinson<sup>2</sup> , Daniel Owens<sup>2</sup> ,  
Thomas Brownlee<sup>3\*</sup> , Duccio Ferrari Bravo<sup>1</sup> and Kevin Enright<sup>2</sup>

<sup>1</sup>Sport Science and R&D Department, Juventus Football Club, Torino, Italy, <sup>2</sup>School of Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, United Kingdom, <sup>3</sup>School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham, Birmingham, United Kingdom

**Introduction:** To effectively monitor post-match changes in physical performance, valid, reliable and practical measures which are sensitive to change are required. This study aimed to quantify test-retest reliability and sensitivity to change of a range of physical performance measures recorded during an isometric posterior chain (IPC) lower-limb muscle test and a countermovement jump (CMJ) test.

**Methods:** Eighteen Italian Serie A academy soccer players performed three IPC repetitions per limb and five CMJ trials in 4 testing sessions. Test-retest reliability was evaluated between two testing sessions seven days apart using typical error of measurement, coefficient of variation and intraclass correlation coefficient. Sensitivity to change was assessed on two additional testing sessions performed before and immediately after a soccer match through Hedges' *g* effect size (*g*) and comparisons to typical error.

**Results:** Absolute reliability (coefficient of variations) ranged from 1.5 to 8.8%. IPC and CMJ measures demonstrated *moderate* to *excellent* relative reliability (intraclass correlation coefficients ranged from 0.70 to 0.98). A wide range of physical performance measures showed significant alterations post-match ( $p < 0.05$ ; *g*: *small* to *moderate*). IPC peak force and torque, CMJ reactive strength index modified, CMJ eccentric forces (mean breaking force, mean deceleration force, peak force, force at zero velocity) and CMJ mean power measures had post-match changes greater than their typical variation, demonstrating acceptable sensitivity in detecting performance changes at post-match.

**Discussion:** IPC peak force and torque, CMJ reactive strength index modified, CMJ eccentric phase forces and CMJ mean power were found to be both reliable and sensitive to change, and thus may be appropriate for monitoring post-match neuromuscular performance in youth soccer population.

## KEYWORDS

monitoring, neuromuscular fatigue, recovery, reliability, responsiveness, football

## 1. Introduction

Soccer match-play represents the most demanding stimulus of the competitive microcycle (1, 2) and has been shown to induce metabolic and mechanical fatigue acutely and in the days following the match in elite senior and youth soccer players (3–6). Nowadays, elite youth soccer players can be exposed to demanding periods of training and competition with limited between-match recovery time (7). The collective stresses

accumulated during the competitive season may lead players to a fatigued status, which may impact their preparedness for subsequent competition and increase the risk of non-functional overreaching, injury, and illness (8). This scenario has prompted a greater interest in monitoring players' post-match fatigue to inform training loads and recovery strategies within the weekly microcycle (9). Fatigue can be defined as a decline in an objective measure of performance over a discrete period (e.g., a decrease in force and power output), and as an increased perception of effort that regulates the integrity of an individual (10). A myriad of field-based physical performance tests is used to assess individual responses following training or competition (11), including isometric tests (e.g., squat, mid-thigh pull, posterior chain) and dynamic measures such as vertical jumps (12). Despite the widespread use of these player monitoring tools, evidence regarding the suitability of a comprehensive range of physical performance measures for monitoring post-match responses in a youth soccer population is limited.

A suitable measure used for effectively monitoring post-match response should possess high reproducibility between repeated tests performed under similar conditions (i.e., reliability) (13, 14) and should be capable of detecting changes induced by training or a match (i.e., sensitivity to change or responsiveness) (9). The reliability of a test refers to the degree of consistency between repeated tests within a practically relevant timeframe (13, 14), while sensitivity to change refers to the ability of a measure to change over a particular timeframe (15). Quantifying both test-retest reliability and sensitivity to change of post-match neuromuscular fatigue measures has been recommended to better interpret individual responses in high-performance sports (16, 17). Additionally, considering the challenges of team-sport settings, tests used for post-match response monitoring should be selected according to their relevance to performance and feasibility within the environment (18). Although numerous studies have reported the reliability of physical performance measures in youth soccer players including vertical jumps and isometric tests (19–21), less research has focused on examining their sensitivity to change following match-play (22).

The countermovement jump (CMJ) test is widely used in high-performance sports to indirectly monitor changes in neuromuscular fatigue and subsequent recovery in both team and individual sports (23, 24). Previous research in youth soccer has mainly investigated the reliability of outcome measures such as jump height (CVs: 4.3%–4.8%; ICCs: 0.83–0.88) (19, 20). Despite possessing acceptable levels of reliability, jump height has been demonstrated to have limited sensitivity to change in training loads in youth and senior soccer players (25, 26). Following a youth academy match play, decrements in CMJ height were observed immediately post and at +24 h, with a tendency towards recovery at +48 h post-match (6). Considering the metabolic, mechanical and neural factors associated with neuromuscular fatigue (10), measuring ground reaction forces during the CMJ may offer superior insights for indirectly detect neuromuscular fatigue (27). Previous research has examined the reliability and sensitivity to change of various CMJ measures in other sports, demonstrating their suitability for ongoing

monitoring (16, 22, 28, 29). In the context of youth soccer, additional research is required to identify reliable CMJ measures capable of detecting neuromuscular fatigue changes.

In team sports, physical performance monitoring also includes isometric tests, used in combination with vertical jumps (12) or as an alternative evaluation (30). Predominantly due to the elevated high-speed running demands (31) and the role of fatigue on hamstring strain injuries in soccer (32), hamstring assessment protocols were introduced as a tool to quantify muscle-specific neuromuscular fatigue during the competitive season. Soccer match results in substantial muscle function impairments until 48–72 h post-match with alterations in muscle contractile properties and central motor output (3). Larger and longer magnitudes of alterations have been observed for knee flexors compared to knee extensors, suggesting the importance of monitoring hamstring function at post-match (33). An isometric posterior chain (IPC) test has been proposed to measure the peak forces generated by the posterior chain musculature at 30° and 90° knee flexion in professional soccer players (34). IPC peak forces have been demonstrated to be reliable (CVs: 4%–6%; ICCs: 0.82–0.95) (34, 35) and sensitive to change following a competitive match-play (relative post-match changes from –20% to –10%) (34, 36). Despite its applicability, peak force does not account for individual differences in limb length and research examining the peak torque generated from the maximal voluntary contraction during the IPC is lacking. Therefore, it is important to demonstrate the reliability and sensitivity of IPC peak force and torque in the specific population to inform practitioners of the potential applications of this testing protocol.

This study extends previous research by quantifying the measurement characteristics of a range of physical performance measures used for monitoring post-match responses in elite youth soccer. Monitoring post-match neuromuscular fatigue using reliable and sensitive measures could potentially inform training prescription and recovery strategies at the individual level within the current or upcoming microcycle to ultimately optimize players' preparedness and development. Therefore, the aim of this study was to (1) quantify the reliability and (2) examine the sensitivity of a range of different physical performance measures from the IPC lower-limb muscle test and the CMJ test in a group of elite youth soccer players.

## 2. Materials and methods

### 2.1. Subjects

Eighteen elite youth soccer players (mean  $\pm$  SD, age: 16.7  $\pm$  0.3 years, height: 177.5 cm  $\pm$  5.9 cm, body mass: 70.4  $\pm$  4.4 kg, percentage adult height: 99.5  $\pm$  0.5%, maturity offset: 2.6  $\pm$  0.5 years) from the under-17 squad of a professional soccer academy in the Italian Serie A participated in this study. All players were free from injury and illness and completed on average 10 h of training (i.e., 5 training sessions) per week, plus a competitive match. Data collection was part of the club's monitoring system and written informed consent was provided by parents or legal

guardians. Ethics approval for the study was provided by the University Research Ethics Committee (22/SPS/006) and was conducted in accordance with the Declaration of Helsinki.

## 2.2. Design

The study was completed during the competitive season and consisted of a test-retest reliability and sensitivity to change assessment of physical performance measures commonly used for post-match monitoring (Figure 1). The IPC and the CMJ test were used for all test sessions. All players received a minimum of two familiarization sessions in the weeks before the experimental trials. Reliability assessment consisted of two testing sessions spaced 7 days apart using a test-retest design. Test-retest reliability sessions were conducted following a 72 hour rest period during which players refrained from participating in training and vigorous activity. Sensitivity to change was evaluated using an intra-club friendly match (2 × 40 min, 105 m × 68 m, artificial turf) as the intervention. Match demands were quantified using global navigation satellite systems (Apex Pro Series, 10 Hz GNSS, STATSports, Northern Ireland) and heart rate (HR) sensors (Polar H10, Polar, Finland). The average match demands were: total distance covered: 10,091 m ± 834 m; distance covered >20 km/h: 720 m ± 203 m; distance covered >25 km/h: 159 m ± 91 m; number of accelerations >3 m/s<sup>2</sup>: 60 ± 14; number of decelerations <-3 m/s<sup>2</sup>: 69 ± 15; time spent >85% HRmax: 29:48 ± 16:27 min; and time spent >90% HRmax: 9:13 ± 6:20 min. Each testing session was conducted at the same time of the day (i.e., early afternoon) to limit the influence of circadian variation and took place following a standardized 5-minute warm-up, except for the session performed immediately after the match. Testing sessions for sensitivity were performed pre- (i.e., 1 h and half before the commencement of the match) and immediately post-match (i.e., within 30 min of completing the match), including only the players who completed the full 80-minute duration. Players were instructed to maintain their normal dietary intake throughout the experimental period. Eighteen players completed the tests for reliability, while thirteen players completed the entire match and performed the tests for sensitivity.

## 2.3. Methodology


The IPC and the CMJ test were conducted using a force platform (ForceDecks Dual Force Plate System FD lite, VALD Performance, Australia). ForceDecks software was used to analyze and calculate the selected physical performance measures. The sampling rate of the force plate was 1,000 Hz.

### 2.3.1. Isometric posterior chain (IPC) lower-limb muscle test

Players lay in a supine position with their knee at 90° (IPC-90°) or 30° (IPC-30°) of flexion, with their calcaneus on the center of the force platform and the non-test leg extended alongside a box at an appropriate height for each participant (i.e., lower shank to be parallel to the floor) (32). Players performed a 3s maximal voluntary contraction by driving their heel down as hard as possible into the platform. The tester ensured a correct position of both legs and pressure was applied to the contralateral hip to control participant posture (i.e., keeping the buttocks, hips and head on the floor). Players were required to repeat trials if their hips raised off the ground. Three trials on each limb were executed with 30s rest between trials for both IPC-90° and IPC-30° lower-limb muscle tests. Peak force was quantified for each trial. Moment arm length was measured from the joint axis of rotation to the point of application of the force and peak torque was calculated by multiplying the peak force by the length of the moment arm. For both the dominant and non-dominant leg, the three-trial average of peak force and peak torque were used for the subsequent statistical analysis.

### 2.3.2. Countermovement jump (CMJ) test

Players performed five CMJ trials with 20s rest between trials. Before the commencement of each jump, players were advised to stand upright, with their arms akimbo and their feet hip-shoulder width apart. Once the starting position was adopted, players remained as still as possible for at least 3s before the start of the trial for the collection of the player's body weight. During the countermovement players were instructed to rapidly squat to their preferred depth and immediately jump as fast and as high as possible, with no knee or hip flexion during the flight phase, maintaining the



hands on the hips. Finally, players were encouraged to “absorb” the landing by flexing at the hips, knees, and ankles after impacting the force platform (37). The average of the three best trials was used for the subsequent statistical analyses (38) for the following CMJ measures: jump height (JH), contraction time (CT), reactive strength index modified (RSImod), concentric duration (ConcDur), concentric peak force (ConcPF), concentric peak velocity (ConcPV), concentric mean power (ConcMP), peak power (PP), eccentric duration (EccDur), eccentric braking phase duration (EccBrakPhDur), eccentric deceleration phase duration (EccDecPhDur), eccentric mean braking force (EccMBrakF), eccentric mean deceleration force (EccMDecF), eccentric peak force (EccPF), force at zero velocity (F@0 V) and eccentric mean power (EccMP) (Supplementary Material). These measures were selected to describe both outcome and movement strategy measures (28) and agree with previous works completed in other high-performance environments (16, 17).

### 2.4. Statistical analysis

Descriptive data are reported as mean ± standard deviation (SD). The assumption of normality was assessed using a Shapiro-Wilk test. Differences between the two reliability test sessions and between pre- and post-match for sensitivity were tested for each measure using a paired *t*-test with alpha ≤0.05. For the reliability analysis, the typical error (TE) of measurement and the coefficient of variation (CV)

were reported as measures of absolute reliability, while the intraclass correlation coefficient 3,1 (ICC; 2-way mixed-effects) with 95% confidence intervals was reported as a measure of relative reliability (13, 39). ICCs were interpreted in line with the lower 95% CI boundary based on previous recommendations: *excellent* (>0.90), *good* (0.75–0.90), *moderate* (0.50–0.75) and *poor* (<0.50) (40). For the sensitivity to change analysis, the magnitude of differences was assessed using Hedges’ *g* effect size (*g*) with 95% CI. Criteria for effect size statistics were interpreted as follows: *trivial* (<0.2), *small* (0.2–0.6), *moderate* (0.6–1.2), *large* (1.2–2.0) and *very large* (>2.0) (41). To determine the ability of each measure in detecting relevant post-match variations, changes were reported and assessed against TE values. If post-match changes were greater than 1.5 times the TE in the test, changes were deemed as sensitive to change (i.e., responsive) (13). The statistical analysis was performed using SPSS statistical software (IBM SPSS version 26, Chicago, IL, USA).

## 3. Results

### 3.1. Reliability of physical performance measures

Reliability statistics are shown in Table 1. Absolute reliability (CVs) ranged from 1.5 to 8.8%. Relative reliability (ICCs) ranged

TABLE 1 Reliability statistics of physical performance measures from the isometric posterior chain (IPC) and the countermovement jump (CMJ) test in elite youth soccer players.

	Test 1 Mean (SD)	Test 2 Mean (SD)	t test (p)	Change in mean (95% CI)	TE	CV	ICC (95% CI)
<b>Isometric posterior chain test</b>							
Peak Force (N)—Dominant leg at 90°	317.2 (54.5)	319.9 (57.1)	0.406	−2.7 (−9.6 to 4.1)	9.7	3.0	0.97 (0.92 to 0.99)
Peak Force (N)—Non-dominant leg at 90°	289.9 (49.2)	296.5 (47.2)	0.123	−6.6 (−15.3 to 2.0)	12.3	4.2	0.93 (0.84 to 0.97)
Peak Torque (Nm)—Dominant leg at 90°	128.3 (23.3)	129.6 (24.1)	0.307	−1.3 (−3.9 to 1.3)	3.7	2.9	0.98 (0.93 to 0.99)
Peak Torque (Nm)—Non-dominant leg at 90°	117.4 (21.7)	120.4 (20.6)	0.081	−3.0 (−6.3 to 0.4)	4.8	4.0	0.95 (0.89 to 0.98)
Peak Force (N)—Dominant leg at 30°	299.2 (49.1)	299.4 (45.8)	0.966	−0.2 (−9.8 to 9.4)	13.6	4.6	0.92 (0.79 to 0.97)
Peak Force (N)—Non-dominant leg at 30°	268.9 (47.2)	275.1 (45.6)	0.174	−6.2 (−15.3 to 3.0)	13.0	4.8	0.92 (0.80 to 0.97)
Peak Torque (Nm)—Dominant leg at 30°	121 (21.1)	121.1 (19.6)	0.965	−0.1 (−4.0 to 3.8)	5.6	4.6	0.93 (0.81 to 0.97)
Peak Torque (Nm)—Non-dominant leg at 30°	108.8 (19.7)	111.2 (20.1)	0.155	−2.7 (−6.4 to 1.1)	5.4	4.9	0.93 (0.82 to 0.97)
<b>Countermovement jump test</b>							
Jump Height (cm)	37.0 (3.9)	36.8 (3.7)	0.437	0.2 (−0.4 to 0.9)	1.0	2.6	0.93 (0.83 to 0.97)
Contraction Time (ms)	713.6 (73.1)	719.6 (84.6)	0.570	−6.0 (−27.7 to 15.8)	30.9	4.3	0.85 (0.64 to 0.94)
RSImodified (m/s)	0.53 (0.08)	0.52 (0.09)	0.296	0.01 (−0.01 to 0.03)	0.02	4.0	0.93 (0.83 to 0.97)
Concentric Duration (ms)	260.9 (28.7)	262.4 (34.7)	0.702	−1.5 (−9.9 to 6.8)	11.9	4.5	0.86 (0.66 to 0.95)
Concentric Peak Force (N)	1,757.4 (204.1)	1,749.7 (238.5)	0.693	7.7 (−32.8 to 48.2)	57.6	3.3	0.93 (0.83 to 0.97)
Concentric Peak Velocity (m/s)	2.77 (0.13)	2.77 (0.14)	0.578	0.01 (−0.02 to 0.04)	0.04	1.5	0.91 (0.77 to 0.96)
Concentric Mean Power (W)	2,087.5 (350.0)	2,083.6 (398.0)	0.847	3.8 (−37.4 to 45.1)	58.6	2.8	0.97 (0.94 to 0.99)
Peak Power (W)	3,702.6 (651.1)	3,723.0 (682.8)	0.524	−20.4 (−86.5 to 45.7)	94.0	2.5	0.98 (0.95 to 0.99)
Eccentric Duration (ms)	455.2 (48.6)	457.1 (54.8)	0.764	−2.0 (−15.6 to 11.6)	19.3	4.2	0.86 (0.67 to 0.95)
Eccentric Braking Phase Duration (s)	0.28 (0.44)	0.28 (0.43)	0.875	0.00 (−0.01 to 0.02)	0.02	7.6	0.77 (0.51 to 0.91)
Eccentric Deceleration Phase Duration (s)	0.16 (0.27)	0.16 (0.28)	0.584	0.00 (−0.01 to 0.01)	0.01	8.8	0.70 (0.37 to 0.88)
Eccentric Mean Braking Force (N)	896.8 (91.5)	900.5 (95.2)	0.614	−3.7 (−19.0 to 11.6)	21.8	2.4	0.95 (0.86 to 0.98)
Eccentric Mean Deceleration Force (N)	1,348.5 (142.8)	1,335.5 (167.1)	0.440	13.0 (−21.7 to 47.7)	49.4	3.7	0.90 (0.75 to 0.96)
Eccentric Peak Force (N)	1,739.9 (192.6)	1,737.2 (230.5)	0.926	−2.7 (−42.9 to 48.2)	64.8	3.7	0.91 (0.77 to 0.96)
Force at Zero Velocity (N)	1,729.5 (194.6)	1,725.5 (236.4)	0.903	4.0 (−42.6 to 50.6)	66.3	3.8	0.91 (0.77 to 0.97)
Eccentric Mean Power (W)	498.1 (76.6)	498.8 (61.6)	0.860	−0.7 (−16.8 to 15.4)	22.8	4.6	0.89 (0.73 to 0.96)

CI, confidence intervals; TE, typical error of measurement; CV, coefficient of variation; ICC, intraclass correlation coefficient, RSI, reactive strength index.

from *good* to *excellent* for IPC measures (ICC  $\geq 0.93$ ; 95% CI: 0.79 to 0.99), whilst it ranged from *moderate* to *excellent* for the CMJ measures (ICC  $\geq 0.85$ ; 95% CI: 0.61 to 0.99), except for Eccentric Braking Phase Duration (EccBrakPhDur) and Eccentric Deceleration Phase Duration (EccDecPhDur) (ICC: 0.70 and 0.77; 95% CI: 0.37 to 0.88, and 0.51 to 0.91, respectively).

### 3.2. Sensitivity of physical performance measures

A wide range of IPC and CMJ measures showed significant alterations at post-match ( $p < 0.05$ ) (Figures 2, 3, Table 2). The magnitude of change at post-match was *moderate* (from  $-10.0\%$  to  $-10.6\%$ ;  $g$ : from  $-0.58$  to  $-0.53$ ) for IPC measures, while it spanned from *trivial* to *moderate* (from  $-0.8$  to  $-10.7\%$ ;  $g$ : from  $-0.78$  to  $0.65$ ) for CMJ measures (Figures 2, 3, Table 2). Eleven measures (IPC-90° PF DL, IPC-90° PF NDL, IPC-90° PT DL, IPC-90° PT NDL, RSImod, EccMBF, EccMDecF, EccPF, F@0 V, EccMP, ConcMP) displayed mean changes greater than typical variation (i.e., 1.5 times the TE) (Figures 2, 3, Table 2). Conversely, the other nine measures (JH, CT, ConcPF, EccDur, EccBrakPhDur, EccDecPhDur, ConcDur, ConcPV, PP) displayed mean changes lower than their corresponding typical variation (Figures 2, 3, Table 2).

## 4. Discussion

The purpose of this study was to quantify the reliability and examine the sensitivity to change of a range of physical performance measures in a group of elite youth soccer players. The main findings from this study were: (1) IPC and CMJ neuromuscular measures demonstrated a high level of absolute reliability (CVs from 1.5 to 8.8%) and *moderate* to *excellent* level of relative reliability (ICCs from 0.70 to 0.98); (2) IPC peak force and torque, CMJ RSImod, CMJ eccentric forces (EccMBF, EccMDecF, EccPF, F@0 V) and CMJ mean power measures showed significant reductions of *small* to *moderate* effect and acceptable sensitivity by displaying post-match changes greater than their typical variation (Figures 2, 3); and (3) CMJ concentric phase measures and jump height showed *trivial* to *small* alterations by displaying post-match changes similar to their typical variation in elite youth soccer players (Figures 2, 3). These findings suggest that IPC peak force and torque, CMJ RSImod, CMJ eccentric phase forces and CMJ mean power may be considered more suitable measures for monitoring players' post-match neuromuscular performance in elite youth soccer players.

### 4.1. Reliability of physical performance measures


IPC lower-limb muscle peak force and torque were demonstrated to be reliable for both dominant and non-dominant leg at 90° and 30° knee flexion. There is no direct

comparative reliability data reporting peak torque recorded during the IPC test in youth soccer players. However, our findings are consistent with previous literature on professional soccer players which have reported better repeatability at 90° knee flexion compared to 30° knee flexion position (CV: 4.3% and 6.3%, respectively) (34). When comparing our reliability data with that of peak force from previous studies (34, 35), it appears that peak torque has greater absolute and relative reliability (CV  $< 5\%$ ; ICC  $> 0.90$ ) compared to peak force. One important consideration in reporting peak torque rather than peak force data is that it accounts for individual differences in limb length (42). Practitioners may therefore consider the IPC as a reliable neuromuscular performance test for monitoring post-match response in youth soccer players.

Most of the CMJ measures from the eccentric and concentric phases demonstrated a high level of absolute and relative reliability in our sample. Previous literature in youth soccer quantified the reliability of jump height only (19, 20). Our results showed greater absolute (CV: 2.6%) and relative (ICC: 0.93) reliability for jump height. These differences may be a result of the different technologies used between studies (i.e., force platform vs. contact mat vs. optical system). Additionally, we quantified the reliability of a wide range of eccentric and concentric phase measures which may provide deeper insights related to both jump outcome and movement strategy (28). Our CVs are lower than in previous literature that used a similar methodology (28, 29), which may be due to the differences in testing protocols, environment and population. However, CVs (1.7%–11.0%) and ICCs (0.70–0.99) were similar to that of a recent study conducted with professional rugby union players which quantified the reliability of 86 CMJ measures (16). Interestingly, CMJ concentric phase and jump outcome measures (e.g., jump height and peak power) displayed better reliability statistics when compared with time-based CMJ measures, suggesting greater variability of jump strategy measures. Despite this trend, several jump strategy measures (e.g., EccDur, RSImod, F@0 V) demonstrated an acceptable level of reliability and therefore may be valuable for monitoring post-match physical performance. In this light, this study uniquely provides reliability reference data of a comprehensive range of CMJ measures within an elite youth soccer academy environment.

### 4.2. Sensitivity of physical performance measures

Determining the inclusion or exclusion of neuromuscular performance measures based on reliability alone may lead to erroneous conclusions. A novel aspect of the present study was to establish the level of reliability in conjunction with the sensitivity to change for effectively monitoring post-match neuromuscular performance in youth soccer players. The significant reductions in IPC measures at 90° dominant and non-dominant leg ( $-10\%$  at post-match) suggest that soccer match play results in acute alterations of hamstring neuromuscular function in youth soccer players, in agreement with previous



**FIGURE 2** Post-match individual changes, mean difference (absolute and relative), *p*-value and Hedges' *g* effect size of physical performance measures in elite youth soccer players. (A) IPC-90° peak torque of dominant leg, (B) IPC-90° peak torque of non-dominant leg, (C) CMJ jump height, (D) CMJ peak power, (E) CMJ RSImodified, (F) CMJ contraction time. Light grey boxes represent the 1.5 times the TE of each measure to identify the players who exhibited changes greater than the typical variance in the IPC and CMJ test. \* indicates a significant mean difference at post-match ( $p \leq 0.05$ ), \*\* indicates a significant mean difference at post-match ( $p \leq 0.01$ ). \*\* indicates a significant mean difference at post-match ( $p \leq 0.001$ ). IPC-90° DL: isometric posterior chain 90° dominant leg; IPC-90° NDL: isometric posterior chain 90° non-dominant leg; JH: jump height; PP: peak power; RSImod: reactive strength index modified; CT: contraction time.

literature (36). However, the magnitude of post-match decline recorded in our research is smaller compared to previous studies (-15% at post-match) that assess hamstring neuromuscular

function following a 90 minute competitive match (34, 43). Despite the difference in the magnitude of change, IPC peak force and torque were demonstrated to be sensitive to change by




FIGURE 3

Post-match individual changes, mean difference (absolute and relative), *p*-value and Hedges' *g* effect size of physical performance measures in elite youth soccer players. (A) CMJ concentric duration, (B) CMJ eccentric duration, (C) concentric peak force, (D) eccentric peak force, (E) force at zero velocity, (F) eccentric mean deceleration force. Light grey boxes represent the 1.5 times the TE of each measure to identify the players who exhibited changes greater than the typical variance in the IPC and CMJ test. \* indicates a significant mean difference at post-match ( $p \leq 0.05$ ), \*\* indicates a significant mean difference at post-match ( $p \leq 0.01$ ), \*\*\* indicates a significant mean difference at post-match ( $p \leq 0.001$ ). ConcDur: concentric duration; EccDur: eccentric duration; ConcPF: concentric peak force; EccPF: eccentric peak force; F@0V: force at zero velocity; EccMDecF: eccentric mean deceleration force.

detecting post-match changes greater than the typical variation for both dominant and non-dominant legs. These moderate reductions in force and torque (−10% at post-match) likely reflect the high

eccentric involvement of the hamstring muscles during repeated sprinting, which could alter muscle function and lead to exercise-induced muscle damage in the days following the competition



TABLE 2 Sensitivity to change of physical performance measures from the isometric posterior chain (IPC) and the countermovement jump (CMJ) test in elite youth soccer players.

	Pre-match Mean (SD)	Post-match Mean (SD)	t test (p)	Change in mean (95% CI)	ES Hedges's g (95% CI)	TE × 1.5	Sensitivity to change	
<b>Isometric posterior chain test</b>								
Peak Force (N)—Dominant leg at 90°	302.2 (46.9)	272.6 (52.0)	0.002**	−29.6 (−46.2 to −12.9)	−0.58 (−1.44 to 0.29)	M	14.5	Yes
Peak Force (N)—Non-dominant leg at 90°	289.1 (42.6)	259.5 (54.0)	0.005**	−29.6 (−47.9 to −11.2)	−0.59 (−1.45 to 0.28)	M	18.4	Yes
Peak Torque (Nm)—Dominant leg at 90°	120.7 (19.9)	109.1 (22.6)	0.003**	−11.6 (−5.0 to −18.2)	−0.53 (−1.39 to −0.34)	M	5.5	Yes
Peak Torque (Nm)—Non-dominant leg at 90°	115.5 (18.8)	103.8 (23.5)	0.005**	−11.7 (−4.3 to −19.1)	−0.53 (−1.39 to −0.33)	M	7.2	Yes
Peak Force (N)—Dominant leg at 30°	–	–	–	–	–	–	–	–
Peak Force (N)—Non-dominant leg at 30°	–	–	–	–	–	–	–	–
Peak Torque (Nm)—Dominant leg at 30°	–	–	–	–	–	–	–	–
Peak Torque (Nm)—Non-dominant leg at 30°	–	–	–	–	–	–	–	–
<b>Countermovement jump test</b>								
Jump Height (cm)	39.5 (4.6)	38.4 (5.0)	0.005**	−1.1 (−1.9 to −0.4)	−0.22 (−1.03 to 0.59)	S	1.4	No
Contraction Time (ms)	692.1 (76.7)	720.9 (89.1)	0.236	28.8 (−21.5 to 79.2)	0.34 (−0.48 to 1.15)	S	46.4	No
RSImodified (m/s)	0.58 (0.11)	0.55 (0.11)	0.013*	−0.03 (−0.06 to 0.01)	−0.26 (−1.08 to 0.55)	S	0.03	Yes
Concentric Duration (ms)	244.2 (31.6)	245.9 (30.6)	0.756	1.7 (−9.8 to 13.1)	0.05 (−0.76 to 0.86)	T	17.8	No
Concentric Peak Force (N)	1,843.2 (185.1)	1,765.7 (181.5)	0.008**	−77.5 (−130.5 to −24.5)	−0.41 (−1.23 to 0.41)	S	86.3	No
Concentric Peak Velocity (m/s)	2.86 (0.15)	2.82 (0.17)	0.051	−0.04 (−0.07 to 0.00)	−0.24 (−1.05 to 0.57)	S	0.06	No
Concentric Mean Power (W)	2,239.3 (315.4)	2,141.2 (290.1)	0.000***	−98.1 (−142.2 to −54.0)	−0.31 (−1.13 to 0.50)	S	87.9	Yes
Peak Power (W)	3,981.9 (523.7)	3,943.9 (483)	0.282	−38.0 (−111.6 to 35.6)	−0.07 (−0.88 to 0.74)	T	141.0	No
Eccentric Duration (ms)	447.8 (56)	475.1 (66.2)	0.173	27.2 (−13.7 to 68.2)	0.43 (−0.39 to 1.25)	S	29.0	No
Eccentric Braking Phase Duration (s)	0.27 (0.05)	0.29 (0.05)	0.212	0.02 (−0.01 to 0.05)	0.35 (−0.47 to 1.16)	S	0.03	No
Eccentric Deceleration Phase Duration (s)	0.15 (0.03)	0.17 (0.03)	0.089	0.02 (−0.00 to 0.04)	0.65 (−0.19 to 1.48)	M	0.02	No
Eccentric Mean Braking Force (N)	888.6 (77.5)	850.3 (75.1)	0.025*	−38.3 (71.0 to −5.6)	−0.49 (−1.31 to 0.34)	S	32.7	Yes
Eccentric Mean Deceleration Force (N)	1,344.7 (151.1)	1,229.9 (133.2)	0.002**	−114.8 (−178.7 to −50.9)	−0.78 (−1.62 to 0.06)	M	74.0	Yes
Eccentric Peak Force (N)	1,769.8 (213.1)	1,621.2 (194.7)	0.000***	−148.6 (−205.4 to −91.8)	−0.71 (−1.54 to 0.13)	M	97.2	Yes
Force at Zero Velocity (N)	1,761 (215.2)	1,613.6 (194.4)	0.000***	−147.4 (−204.9 to −89.8)	−0.70 (−1.53 to 0.14)	M	99.4	Yes
Eccentric Mean Power (W)	487.5 (76.7)	442.1 (65.9)	0.005**	−45.5 (−74.0 to −16.9)	−0.61 (−1.44 to 0.22)	M	34.3	Yes

CI, confidence intervals; TE, typical error of measurement; ES, effect size; T, trivial; S, small; M, moderate, RSI, reactive strength index.

Sensitivity to change: absolute change at post-match >TE × 1.5 (Hopkins, 2000).

\*p ≤ 0.05.

\*\*p ≤ 0.01.

\*\*\*p ≤ 0.001.

(44). Considering that neuromuscular fatigue is one potential hamstring strain injury risk factor, our findings support the use of IPC test to indirectly monitor post-match neuromuscular function in youth soccer population.

A key finding of the present study was that CMJ eccentric phase measures had greater post-match changes compared with CMJ jump height and concentric phase measures. In line with previous studies on elite youth soccer players (6, 45), jump height showed a significant reduction but demonstrated limited sensitivity by displaying post-match changes similar to the typical variation. It has been reported that jump height may not be sensitive to training and match loads in elite soccer players (26, 46). Therefore, CMJ time-based measures may hold superior sensitivity to change following official matches and fatiguing intermittent protocol (22, 28). Our analysis of CMJ measures revealed that RSI<sub>mod</sub>, eccentric forces (EccMBF, EccMDecF, EccPF, F@0 V) and mean power (ConcMP, EccMP) demonstrated an acceptable level of sensitivity displaying changes greater than the typical variation (i.e., 1.5 times the TE). Gathercole and colleagues demonstrated a similar level of sensitivity to alternative CMJ measures in a group of collegiate team sports athletes showing acute changes greater than their typical variation and confidence intervals (28). The observed responses suggest that elite youth soccer players may experience

acute reductions in neuromuscular function which have been attributed to a reduced central drive and metabolic disturbances that impair excitation-contraction coupling and reduce stretch-reflex sensitivity, muscle stiffness, and force production (47). The greater magnitude of change of CMJ eccentric forces may be associated with the frequent intense acceleration and deceleration actions performed during the match (48, 49). Overall, the acute changes in time-based measures may reflect the altered stretch-shortening cycle (SSC) function and movement strategy employed over the downward phase of the CMJ (47). The current findings demonstrate CMJ eccentric force measures may reveal alterations in jump strategy which underline the presence of altered SSC and neuromuscular fatigue following match-play. As time, force and velocity used to perform soccer-specific actions are considered fundamental for success in elite soccer, these alterations in CMJ eccentric force measures may have relevant implications in monitoring players' physical performance.

This study provides practitioners with a range of physical performance measures suitable for indirectly monitoring players' post-match neuromuscular status in elite youth soccer players. It is important to highlight the main limitations inherent to this work. Despite the ecological validity of the study, sensitivity to change was evaluated using a single match of a shorter duration compared to the official match (i.e., 80 vs. 90 min), with match

demands inherently variable between players. Thus, match demands might have influenced the magnitude of alterations post-match. Another limitation of our study is the lack of investigation into the association between changes in physical performance and match demands. Examining this association may provide valuable insights for predicting post-match acute and residual fatigue (49). Additionally, it is important to emphasize that our study was conducted with players from one club only, limiting the generalizability of our results to other youth soccer academy settings. Future research should investigate the time course of recovery of these neuromuscular measures in the days following the match and examine the effects of different recovery interventions on these measures in the subsequent days following match play (i.e., match-day + 1 and match-day + 2) as little research exists in elite youth soccer (50).

## 5. Practical applications

The present study demonstrated that IPC and CMJ tests are reliable and sensitive tools for monitoring post-match changes in physical performance of elite youth soccer players. Our findings indicated that IPC peak force and torque, CMJ RSImod, CMJ eccentric measures (EccMBF, EccMDecF, EccPF, F@0 V) and CMJ mean power may be more appropriate to detect physical performance changes following match-play. Conversely, typically reported CMJ measures such as jump height and peak power showed limited sensitivity in detecting post-match changes. Performance practitioners are therefore encouraged to include such measures for monitoring neuromuscular performance following the match. This might allow coaches and scientists to optimize the scheduling of training across the microcycle, facilitating recovery whilst also optimizing the physical conditioning of players.

## Data availability statement

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

## Ethics statement

The studies involving human participants were reviewed and approved by University Research Ethics Committee, Liverpool John Moores University. Written informed consent to participate in this study was provided by the participants' legal guardian/

next of kin. Written informed consent was obtained from the minor(s)' legal guardian/next of kin for the publication of any potentially identifiable images or data included in this article.

## Author contributions

Authors contributed to this manuscript as follows: study conception and design (AF, MAR, KE); data collection (AF, DFB); data curation (AF, MAR); data analysis (AF, MAR, KE); supervision (DFB, KE); writing—original draft (AF); writing—review, editing and approval of final draft (AF, MAR, DO, TB, DFB, KE). All authors contributed to the article and approved the submitted version.

## Acknowledgments

We would like to thank the players and staff at Juventus Football Club for participating and cooperating in this research. Special thanks are acknowledged to Roberto Sassi, Darragh Connolly, Antonio Gualtieri and Francesco Lucia for their assistance with this research.

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fspor.2023.1173621/full#supplementary-material>

## References

- Kelly DM, Strudwick AJ, Atkinson G, Drust B, Gregson W. Quantification of training and match-load distribution across a season in elite English premier league soccer players. *Sci and Med in Football*. (2020) 4(1):59–67. doi: 10.1080/24733938.2019.1651934
- Hannon MP, Coleman NM, Parker LJF, McKeown J, Unnithan VB, Close GL, et al. Seasonal training and match load and micro-cycle periodization in male premier league academy soccer players. *J Sports Sci*. (2021) 00(00):1–12. doi: 10.1080/02640414.2021.1899610
- Silva JR, Rumpf MC, Hertzog M, Castagna C, Farooq A, Girard O, et al. Acute and residual soccer match-related fatigue: a systematic review and meta-analysis. *Sports Med*. (2018) 48(3):539–83. doi: 10.1007/s40279-017-0798-8
- Rampinini E, Bosio A, Ferraresi I, Petruolo A, Morelli A, Sassi A. Match-related fatigue in soccer players. *Med Sci Sports Exerc*. (2011) 43(11):2161–70. doi: 10.1249/MSS.0b013e31821e9c5c

5. Deely C, Tallent J, Bennett R, Woodhead A, Goodall S, Thomas K, et al. Etiology and recovery of neuromuscular function following academy soccer training. *Front Physiol.* (2022) 13:911009. doi: 10.3389/fphys.2022.911009
6. de Hoyo M, Cohen DD, Sañudo B, Carrasco L, Álvarez-Mesa A, del Ojo JJ, et al. Influence of football match time–motion parameters on recovery time course of muscle damage and jump ability. *J Sports Sci.* (2016) 34(14):1363–70. doi: 10.1080/02640414.2016.1150603
7. Castellano J, Martín-García A, Casamichana D. Most running demand passages of match play in youth soccer congestion period. *Biol Sport.* (2020) 37(4):367–73. doi: 10.5114/biolSport.2020.96853
8. Dupont G, Nedelec M, McCall A, McCormack D, Berthoin S, Wisløff U. Effect of 2 soccer matches in a week on physical performance and injury rate. *Am J Sports Med.* (2010) 38(9):1752–8. doi: 10.1177/0363546510361236
9. Thorpe RT, Atkinson G, Drust B, Gregson W. Monitoring fatigue status in elite team sports: implications for practice. *Int J Sports Physiol Perform.* (2017) 12: S227–34. doi: 10.1123/ijssp.2016-0434
10. Enoka RM, Duchateau J. Translating fatigue to human performance. *Med Sci Sports Exerc.* (2016) 48(11):2228–38. doi: 10.1249/MSS.0000000000000929
11. Alba-Jiménez C, Moreno-Doutres D, Peña J. Trends assessing neuromuscular fatigue in team sports: a narrative review. *Sports.* (2022) 10(3):33. Available at: <https://www.mdpi.com/2075-4663/10/3/33> doi: 10.3390/sports10030033
12. Norris D, Joyce D, Siegler J, Cohen D, Lovell R. Considerations in interpreting neuromuscular state in elite level Australian rules football players. *J Sci Med Sport.* (2021) 24(7):702–8. doi: 10.1016/j.jsams.2021.02.007
13. Hopkins WG. Measures of reliability in sports medicine and science. *Sports Med.* (2000) 30(1):1–15. doi: 10.2165/00007256-200030010-00001
14. Atkinson G, Nevill AM. Statistical methods for assessing measurement error (reliability) in variables relevant to sports medicine. *Sports Medicine.* (1998) 26(4):217–38. doi: 10.2165/00007256-199826040-00002
15. Impellizzeri FM, Marcora SM. Test validation in sport physiology: lessons learned from clinimetrics. Vol. 4, *Int J Sports Physiol Perform.* Human Kinetics Publishers Inc.; (2009). p. 269–77. doi: 10.1123/ijssp.4.2.269
16. Howarth DJ, Cohen DD, McLean BD, Coutts AJ. Establishing the noise: interday ecological reliability of countermovement jump variables in professional rugby union players. *J Strength Cond Res.* (2022) 36(11):3159–66. doi: 10.1519/JSC.0000000000004037
17. Mercer RAJ, Russell JL, Mcguigan LC, Coutts AJ, Strack DS, Mclean BD. Finding the signal in the noise: interday reliability and seasonal sensitivity of 84 countermovement jump variables in professional basketball players. *J Strength Cond Res.* (2023) 37(2):394–402. doi: 10.1519/JSC.0000000000004182
18. Bishop C, Turner AN, Jordan M, Harry J, Loturco I, Lake J, et al. A framework to guide practitioners when selecting metrics during the countermovement and drop jump tests. *Strength Cond J.* (2022) 44(4):95–103. doi: 10.1519/SSC.0000000000000677
19. Enright K, Morton J, Iga J, Lethian D, Roberts S, Drust B. Reliability of “in-season” fitness assessments in youth elite soccer players: a working model for practitioners and coaches. *Sci and Med in Football.* (2018) 2(3):177–83. doi: 10.1080/24733938.2017.1411603
20. Fitzpatrick JF, Hicks KM, Russell M, Hayes PR. The reliability of potential fatigue-monitoring measures in elite youth soccer players. *J Strength Cond Res.* (2021) 35(12):3448–52. doi: 10.1519/JSC.0000000000003317
21. Brown AM, Hader K, Guilhem G, Simpson BM, Buchheit M, Lacombe M. The ballistic hip thrust test: a potential tool to monitor neuromuscular performance. *Biol Sport.* (2022) 39(1):73–7. doi: 10.5114/biolSport.2021.102920
22. Rowell AE, Aughey RJ, Hopkins WG, Stewart AM, Cormack SJ. Identification of sensitive measures of recovery after external load from football match play. *Int J Sports Physiol Perform.* (2017) 12(7):969–76. doi: 10.1123/ijssp.2016-0522
23. Beattie CE, Fahey JT, Pullinger SA, Edwards BJ, Robertson CM. The sensitivity of countermovement jump, creatine kinase and urine osmolality to 90-min of competitive match-play in elite English championship football players 48-h post-match. *Sci Med Footb.* (2021) 5(2):165–73. doi: 10.1080/24733938.2020.1828614
24. Franceschi A, Conte D, Airale M, Sampaio J. Training load, neuromuscular readiness, and perceptual fatigue profile in youth elite long-jump athletes. *Int J Sports Physiol Perform.* (2020) 15(7):1034–8. doi: 10.1123/ijssp.2019-0596
25. Thorpe RT, Strudwick AJ, Buchheit M, Atkinson G, Drust B, Gregson W. Monitoring fatigue during the in-season competitive phase in elite soccer players. *Int J Sports Physiol Perform.* (2015) 10(8):958–64. doi: 10.1123/ijssp.2015-0004
26. Malone JJ, Murtagh CF, Morgans R, Burgess DJ, Morton JP, Drust B. Countermovement jump performance is not affected during an in-season training microcycle in elite youth soccer players. *J Strength Cond Res.* (2015) 29(3):752–7. doi: 10.1519/JSC.0000000000000701
27. Bishop C, Jordan M, Torres-Ronda L, Loturco I, Harry J, Virgile A, et al. Selecting metrics that matter: comparing the use of the countermovement jump for performance profiling, neuromuscular fatigue monitoring, and injury rehabilitation testing. *Strength Cond J.* (2023). doi: 10.1519/SSC.0000000000000772. [Epub ahead of Print].
28. Gathercole R, Sporer B, Stellingwerff T, Sleivert G. Alternative countermovement-jump analysis to quantify acute neuromuscular fatigue. *Int J Sports Physiol Perform.* (2015) 10(1):84–92. doi: 10.1123/ijssp.2013-0413
29. Heishman AD, Daub BD, Miller RM, Freitas EDS, Frantz BA, Bembem MG. Countermovement jump reliability performed with and without an arm swing in NCAA division I intercollegiate basketball players. *J Strength Cond Res.* (2020) 34(2):546–58. doi: 10.1519/JSC.0000000000002812
30. Brownlee TE, Murtagh CF, Naughton RJ, Whitworth-Turner CM, O’Boyle A, Morgans R, et al. Isometric maximal voluntary force evaluated using an isometric mid-thigh pull differentiates English premier league youth soccer players from a maturity-matched control group. *Sci and Med in Football.* (2018) 2(3):209–15. doi: 10.1080/24733938.2018.1432886
31. Barnes C, Archer DT, Hogg B, Bush M, Bradley PS. The evolution of physical and technical performance parameters in the English premier league. *Int J Sports Med.* (2014) 35(13):1095–100. doi: 10.1055/s-0034-1375695
32. Huygaerts S, Cos F, Cohen DD, Calleja-González J, Guitart M, Blazevich AJ, et al. Mechanisms of hamstring strain injury: interactions between fatigue, muscle activation and function. *Sports.* (2020) 8(5):65. doi: 10.3390/sports8050065
33. Trecroci A, Porcelli S, Perri E, Pedrali M, Rasica L, Alberti G, et al. Effects of different training interventions on the recovery of physical and neuromuscular performance after a soccer match. *J Strength Cond Res.* (2020) 34(8):2189–92. doi: 10.1519/JSC.0000000000003269
34. McCall A, Nedelec M, Carling C, le Gall F, Berthoin S, Dupont G. Reliability and sensitivity of a simple isometric posterior lower limb muscle test in professional football players. *J Sports Sci.* (2015) 33(12):1298–304. doi: 10.1080/02640414.2015.1022579
35. Cuthbert M, Comfort P, Ripley N, McMahon JJ, Evans M. Unilateral vs. Bilateral hamstring strength assessments: comparing reliability and inter-limb asymmetries in female soccer players. *J Sports Sci.* (2021) 39(13):1481–8. doi: 10.1080/02640414.2021.1880180
36. Constantine E, Taberner M, Richter C, Willett M, Cohen DD. Isometric posterior chain peak force recovery response following match-play in elite youth soccer players: associations with relative posterior chain strength. *Sports (Basel).* (2019) 7(10):218. doi: 10.3390/sports7100218
37. McMahon JJ, Suchomel TJ, Lake JP, Comfort P. Understanding the key phases of the countermovement jump force-time curve. *Strength Cond J.* (2018) 40(4):96–106. doi: 10.1519/SSC.0000000000000375
38. Claudino JG, Cronin J, Mezêncio B, McMaster DT, Mcguigan MR, Tricoli V, et al. The countermovement jump to monitor neuromuscular status: a meta-analysis. *J Sci Med Sport.* (2017) 20(4):397–402. doi: 10.1016/j.jsams.2016.08.011
39. Weir J. Quantifying test-retest reliability using the intraclass correlation coefficient and the SEM. *J Strength Cond Res.* (2005) 19(1):231–40. doi: 10.1519/15184.1
40. Koo TK, Li MY. A guideline of selecting and reporting intraclass correlation coefficients for reliability research. *J Chiropr Med.* (2016) 15(2):155–63. doi: 10.1016/j.jcm.2016.02.012
41. Hopkins WG, Marshall SW, Batterham AM, Hanin J. Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exerc.* (2009) 41(1):3–12. doi: 10.1249/MSS.0b013e31818cb278
42. Baltzopoulos B, King M, Gleeson N, de Ste Croix M. The BASES expert statement on measurement of muscle strength with isokinetic dynamometry. *The Sport and Exercise Scientist.* (2012) 31:12–3.
43. Wollin M, Thorborg K, Pizzari T. The acute effect of match play on hamstring strength and lower limb flexibility in elite youth football players. *Scand J Med Sci Sports.* (2017) 27(3):282–8. doi: 10.1111/sms.12655
44. Owens DJ, Twist C, Cogley JN, Howatson G, Close GL. Exercise-induced muscle damage: what is it, what causes it and what are the nutritional solutions?. *Eur J Sport Sci.* (2019) 19(1):71–85. doi: 10.1080/17461391.2018.1505957
45. Romagnoli M, Sanchis-Gomar F, Alis R, Risso-Ballester J, Bosio A, Graziani RL, et al. Changes in muscle damage, inflammation, and fatigue-related parameters in young elite soccer players after a match. *J Sports Med Phys Fitness.* (2016) 56(10):1198–205.
46. Thorpe RT, Strudwick AJ, Buchheit M, Atkinson G, Drust B, Gregson W. The influence of changes in acute training load on daily sensitivity of morning-measured fatigue variables in elite soccer players. *Int J Sports Physiol Perform.* (2017) 12(Suppl 2):S2107–13. doi: 10.1123/ijssp.2016-0433
47. Komi PV. Stretch-shortening cycle: a powerful model to study normal and fatigued muscle. *J Biomech.* (2000) 33(10):1197–206. doi: 10.1016/S0021-9290(00)00064-6
48. Harper DJ, Carling C, Kiely J. High-Intensity acceleration and deceleration demands in elite team sports competitive match play: a systematic review and meta-analysis of observational studies. *Sports Med.* (2019) 49(12):1923–47. doi: 10.1007/s40279-019-01170-1
49. Hader K, Rumpf MC, Hertzog M, Kilduff LP, Girard O, Silva JR. Monitoring the athlete match response: can external load variables predict post-match acute and residual fatigue in soccer? A systematic review with meta-analysis. *Sports Med Open.* (2019) 5(1):48. doi: 10.1186/s40798-019-0219-7
50. Calleja-González J, Mielgo-Ayuso J, Miguel-Ortega Á, Marqués-Jiménez D, del Valle M, Ostojic SM, et al. Post-exercise recovery methods focus on young soccer players: a systematic review. *Front Physiol.* (2021) 12:505149. doi: 10.3389/fphys.2021.505149