

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/161008/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Smith, Holly-Anne B., Mullins, Alex J. , Webster, Gordon , Sapsford, Devin , Gaskin, Paul, Monis, Paul T., Hoefel, Daniel, Saint, Christopher P., Weightman, Andrew J. and Maresca, Julia A. 2023. *Sphingopyxis* Species Isolated from Sand Filter Biofilm at an Australian Drinking Water Treatment Works. *Microbiology Resource Announcements* 10.1128/mra.00249-23 file

Publishers page: <http://dx.doi.org/10.1128/mra.00249-23>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.





# *Sphingopyxis* Species Isolated from Sand Filter Biofilm at an Australian Drinking Water Treatment Works

 Holly-Anne B. Smith,<sup>a</sup>  Alex J. Mullins,<sup>a</sup>  Gordon Webster,<sup>a</sup> Devin Sapsford,<sup>b</sup> Paul Gaskin,<sup>c</sup> Paul T. Monis,<sup>d</sup> Daniel Hoefel,<sup>d</sup> Christopher P. Saint,<sup>e,f</sup>  Andrew J. Weightman<sup>a</sup>

<sup>a</sup>Microbiomes, Microbes and Informatics Group, Organisms and Environment Division, School of Biosciences, Cardiff University, Cardiff, Wales, United Kingdom

<sup>b</sup>School of Engineering, Queen's Buildings, Cardiff University, Cardiff, Wales, United Kingdom

<sup>c</sup>Dwr Cymru Welsh Water, Coed Kernew, Newport, Wales, United Kingdom

<sup>d</sup>Australian Water Quality Centre, SA Water, Adelaide, Australia

<sup>e</sup>Future Industries Institute, University of South Australia, Adelaide, Australia

<sup>f</sup>UniSA STEM, University of South Australia, Adelaide, Australia

**ABSTRACT** Three strains isolated by geosmin enrichment from a sand filter in an Australian drinking water treatment works were genome sequenced to identify their taxonomic placement, and a bench-scale batch experiment confirmed their geosmin-degrading capability. Using the average nucleotide identity based on the MUMmer algorithm (ANIm), pairwise digital DNA-DNA hybridization (dDDH), and phylogenomic analyses, the strains were identified as *Sphingopyxis* species.

Proposed in 2001 (1), the genus *Sphingopyxis* currently comprises 21 validly published species (2), belonging to the family *Sphingomonadaceae* of the class *Alphaproteobacteria*. *Sphingopyxis* strains have been isolated from diverse natural environments, including volcanic rock (3), freshwater (4), soils (5), seawater (6), and some contaminated sites (7–9). *Sphingopyxis* species can utilize a broad range of carbon sources (1), including aromatic compounds such as tetralin (10) and microcystins (11). *Sphingopyxis* species have also been found to biotransform heavy metals and biodegrade polyethers, antibacterials, and geosmin (1, 12).

*Sphingopyxis* strains Geo24, Geo25, and Geo48 were originally isolated in the lab by geosmin enrichment from a sand filter from an Australian drinking water treatment works (13, 14). Geo24 and Geo25 were identified as part of a bacterial consortium able to biodegrade geosmin as the sole carbon source (13), and Geo48 was later identified as an isolate capable of geosmin biodegradation (14). The isolates were stored long term at  $-80^{\circ}\text{C}$ , shipped on charcoal transport swabs, streaked onto tryptic soy agar (TSA), and grown for 48 h at  $30^{\circ}\text{C}$ . All strains were streaked three times to ensure the purity of individual colonies.

For genome sequencing, the strains were grown in 5 mL tryptic soy broth (TSB) at  $30^{\circ}\text{C}$  for 48 h, and total DNA was extracted using the FastDNA spin kit for soil (MP Biomedicals). Sequencing was performed on a NovaSeq 6000 SP instrument using a NEBNext Ultra II DNA library prep kit. Between 7 and 9 million (150-bp) read pairs were generated for each genome. The read quality was checked using FastQC v0.11.9, trimming and adapter removal was performed using Fastp v0.20.1, and the genomes were assembled using Unicycler v0.4.7. The genome sizes and other metrics are as follows: Geo24 has a size of 3.86 Mbp, 23 contigs, an  $N_{50}$  value of 665,169 bp, and a GC content of 65.3%; Geo25 comprises 3.87 Mbp, 23 contigs, an  $N_{50}$  value of 672,543 bp, and a GC content of 65.3%; and Geo48 has a size of 3.96 Mbp, 25 contigs, an  $N_{50}$  value of 978,883 bp, and a GC content of 65.2%.

Phylogenetic analysis of *Sphingopyxis* strains Geo24, Geo25, and Geo48 was performed against all available *Sphingopyxis* type strain genomes, downloaded using the NCBI genome download tool v0.2.10 and annotated using Prokka v1.14.6. The average nucleotide identity

**Editor** Julia A. Maresca, University of Delaware College of Engineering


**Copyright** © 2023 Smith et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Holly-Anne B. Smith, smithhb@cardiff.ac.uk, or Andrew J. Weightman, weightman@cardiff.ac.uk.

The authors declare no conflict of interest.

**Received** 27 March 2023

**Accepted** 1 June 2023



**FIG 1** ANIm, phylogeny, and geosmin biodegradation batch experiment results for novel *Shingopyxis* species Geo24, Geo25, and Geo48, isolated from an Australian drinking water treatment works sand filter. (A) ANIm heatmap produced using PyANI v0.2.12, depicting the average (Continued on next page)

**TABLE 1** Basic metrics and pairwise dDDH values for novel *Sphingopyxis* species and the three most closely related type strains<sup>a</sup>

Strain	Genome size (Mbp)	GC content (%)	Pairwise dDDH (%) with strain:					
			Geo24	Geo25	Geo48	<i>Sphingopyxis soli</i> BL03	<i>Sphingopyxis lindanitolerans</i> W55A3p	<i>Sphingopyxis macrogoltabida</i> 203
<i>Sphingopyxis</i> sp. Geo24	3.86	65.3		100.0	76.4	50.0	27.5	25.8
<i>Sphingopyxis</i> sp. Geo25	3.87	65.3	100.0		76.4	50.0	27.5	25.8
<i>Sphingopyxis</i> sp. Geo48	3.96	65.2	76.4	76.4		51.2	27.1	26.2
<i>Sphingopyxis soli</i> BL03	3.63	65.8	50.0	50.0	51.2		27.5	25.9
<i>Sphingopyxis lindanitolerans</i> W55A3p	4.15	65.3	27.5	27.5	27.1	27.5		25.9
<i>Sphingopyxis macrogoltabida</i> 203	5.75	64.9	25.8	25.8	26.2	25.9	25.9	

<sup>a</sup> dDDH values were calculated using TYGS and according to the *Sphingopyxis* phylogenetic tree (Fig. 1A). Pairwise dDDH values of <70% indicate different species.

(ANI) was calculated using PyANI v0.2.12 (Fig. 1A), and a phylogenomic tree was constructed using OrthoFinder v2.5.4 (15, 16) and RAXML-NG v1.1 (17) (Fig. 1B). All three strains showed an ANI of <94% to the most closely related type strains and displayed phylogenomic distinction, indicating them as potentially novel species (Fig. 1B). Pairwise digital DNA-DNA hybridization (dDDH) values were determined using the Type Strain Genome Server (TYGS) with the most closely related type strains (18); the results indicated that all three *Sphingopyxis* strains represent different species (dDDH, <70%) than those previously described (Table 1).

Geosmin biodegradation capacity was confirmed in a microcosm batch experiment by analyzing the geosmin concentration over 7 days. Microcosms comprised of 10 mL bacteria, diluted to 0.1 optical density at 600 nm ( $OD_{600}$ ), in basal salts medium (BSM) (19) in vials with 20 mL headspace. Geosmin losses by volatilization were controlled with microcosms of 10 mL BSM. Geosmin was added at 100 ng/L to each microcosm and measured at 0, 4, and 7 days in triplicate using solid-phase microextraction and gas chromatography mass spectrometry (GCMS) analysis. Significant removal of geosmin was observed for all inoculated microcosms compared to the control (Fig. 1C), demonstrating that all three *Sphingopyxis* strains can degrade geosmin.

**Data availability.** The genome sequences and raw reads have been deposited in the European Nucleotide Archive (ENA) under the project/study number [PRJEB60073](https://www.ebi.ac.uk/ena/record/PRJEB60073). The accession numbers for the genome sequences are [ERS14837053](https://www.ebi.ac.uk/ena/record/ERS14837053), [ERS14837054](https://www.ebi.ac.uk/ena/record/ERS14837054), and [ERS14837055](https://www.ebi.ac.uk/ena/record/ERS14837055) for Geo24, Geo25, and Geo48, respectively.

## ACKNOWLEDGMENTS

This study was supported by funding from KESS2 East and Dŵr Cymru Welsh Water. All analyses were conducted using Cloud Infrastructure for Big Data Microbial Bioinformatics (CLIMB), funded by the Medical Research Council (MR/L015080/1). We thank South Australian Water (SA Water) for providing the bacterial strains investigated in this paper, and we acknowledge L. Ho, G. Newcombe, A. Keegan, and W. Aunkofer for their contributions to the initial isolation of the strains.

Conceptualization, H.A.B.S., G.W., and A.J.W.; Data curation, H.A.B.S. and A.J.M.; Formal analysis, H.A.B.S., G.W., and A.J.M.; Funding acquisition, A.J.W., P.G., and D.S.; Investigation, H.A.B.S.; Methodology, H.A.B.S., G.W., A.J.M., and A.J.W.; Project administration, H.A.B.S., A.J.W., and G.W.; Resources, P.T.M., D.H., C.P.S., and A.J.W.;

## FIG 1 Legend (Continued)

nucleotide identity, as specified by the color bar key. Novel species are shown in bold against *Sphingopyxis* type strains. (B) Phylogenetic tree created using OrthoFinder v2.5.4 and RAXML-NG v1.1 to show the relationship between novel *Sphingopyxis* species and available type strain species, rooted with the type strain from the sister genus *Novosphingopyxis*. The maximum likelihood method was used with the LG model and G4 distribution, with bootstrap support (100 replicates) shown next to each node. Novel *Sphingopyxis* species are shown in bold. (C) Geosmin concentrations for microcosms in batch experiment comparing geosmin removal of novel *Sphingopyxis* species over 7 days. Bacteria (1 mL) grown in TSB, washed, and controlled to an  $OD_{600}$  of 1 were added to 9 mL BSM and spiked with 100 ngL<sup>-1</sup> geosmin. Blank control microcosms with 10 mL BSM, spiked with 100 ngL<sup>-1</sup> geosmin and with no inoculum, were run simultaneously. Geosmin concentration analysis was performed using SPME with GCMS analysis. Asterisks indicate statistical significance from the blank control at each time point, determined using the Mann-Whitney test.

Software, H.A.B.S. and A.J.M.; Supervision, P.G., D.S., and A.J.W.; Validation, H.A.B.S. and G.W.; Visualization, H.A.B.S., A.J.M., and G.W.; Writing – original draft, H.A.B.S.; Writing – review & editing, G.W., A.J.M., A.J.W., C.P.S., and P.T.M.

## REFERENCES

1. Takeuchi M, Hamana K, Hiraishi A. 2001. Proposal of the genus *Sphingomonas* sensu stricto, and three new genera, *Sphingobium*, *Novosphingobium* and *Sphingopyxis*, on the basis of phylogenetic and chemotaxonomic analyses. *Int J Syst Evol Microbiol* 51:1405–1417. <https://doi.org/10.1099/00207713-51-4-1405>.
2. Sharma M, Khurana H, Singh DN, Negi RK. 2021. The genus *Sphingopyxis*: systematics, ecology, and bioremediation potential—a review. *J Environ Manage* 280:111744. <https://doi.org/10.1016/j.jenvman.2020.111744>.
3. Alias-Villegas C, Jurado V, Laiz L, Saiz-Jimenez C. 2013. *Sphingopyxis italica* sp. nov., isolated from Roman catacombs. *Int J Syst Evol Microbiol* 63:2565–2569. <https://doi.org/10.1099/ijs.0.046573-0>.
4. Baik KS, Choe HN, Park SC, Hwang YM, Kim EM, Park C, Seong CN. 2013. *Sphingopyxis rigui* sp. nov. and *Sphingopyxis wooonensis* sp. nov., isolated from wetland freshwater, and emended description of the genus *Sphingopyxis*. *Int J Syst Evol Microbiol* 63:1297–1303. <https://doi.org/10.1099/ijs.0.044057-0>.
5. Chaudhary DK, Dahal RH, Kim J. 2017. *Sphingopyxis solisilvae* sp. nov., isolated from forest soil. *Int J Syst Evol Microbiol* 67:1820–1826. <https://doi.org/10.1099/ijsem.0.001869>.
6. Kim B-S, Lim YW, Chun J. 2008. *Sphingopyxis marina* sp. nov. and *Sphingopyxis litoris* sp. nov., isolated from seawater. *Int J Syst Evol Microbiol* 58:2415–2419. <https://doi.org/10.1099/ijs.0.65614-0>.
7. Chaudhary DK, Kim J. 2018. *Sphingopyxis nepalensis* sp. nov., isolated from oil-contaminated soil. *Int J Syst Evol Microbiol* 68:364–370. <https://doi.org/10.1099/ijsem.0.002514>.
8. Kaminski MA, Sobczak A, Spolnik G, Danikiewicz W, Dziembowski A, Lipinski L. 2018. *Sphingopyxis lindanitolerans* sp. nov. strain W55A3pT enriched from a pesticide disposal site. *Int J Syst Evol Microbiol* 68:3935–3941. <https://doi.org/10.1099/ijsem.0.003094>.
9. Sharma P, Verma M, Bala K, Nigam A, Lal R. 2010. *Sphingopyxis ummariensis* sp. nov., isolated from a hexachlorocyclohexane dump site. *Int J Syst Evol Microbiol* 60:780–784. <https://doi.org/10.1099/ijs.0.008805-0>.
10. Ledesma-García L, Sánchez-Azqueta A, Medina M, Reyes-Ramírez F, Santero E. 2016. Redox proteins of hydroxylating bacterial dioxygenases establish a regulatory cascade that prevents gratuitous induction of tetralin biodegradation genes. *Sci Rep* 6:23848. <https://doi.org/10.1038/srep23848>.
11. Massey IY, Zhang X, Yang F. 2018. Importance of bacterial biodegradation and detoxification processes of microcystins for environmental health. *J Toxicol Environ Health B Crit Rev* 21:357–369. <https://doi.org/10.1080/10937404.2018.1532701>.
12. Yang F, Feng H, Massey IY, Huang F, Guo J, Zhang X. 2020. Genome-wide analysis reveals genetic potential for aromatic compounds biodegradation of *Sphingopyxis*. *Biomed Res Int* 2020:5849123. <https://doi.org/10.1155/2020/5849123>.
13. Hoefel D, Ho W, Aunkofer W, Monis PT, Keegan A, Newcombe G, Saint CP. 2006. Cooperative biodegradation of geosmin by a consortium comprising three Gram-negative bacteria isolated from the biofilm of a sand filter column. *Lett Appl Microbiol* 43:417–423. <https://doi.org/10.1111/j.1472-765X.2006.01974.x>.
14. Hoefel D, Ho L, Monis PT, Newcombe G, Saint CP. 2009. Biodegradation of geosmin by a novel Gram-negative bacterium; isolation, phylogenetic characterisation and degradation rate determination. *Water Res* 43:2927–2935. <https://doi.org/10.1016/j.watres.2009.04.005>.
15. Emms DM, Kelly S. 2017. STRIDE: species tree root inference from gene duplication events. *Mol Biol Evol* 34:3267–3278. <https://doi.org/10.1093/molbev/msx259>.
16. Emms DM, Kelly S. 2018. STAG: species tree inference from all genes. *bioRxiv*. <https://doi.org/10.1101/267914>.
17. Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. 2019. RAXML-NG: a fast, scalable, and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* 35:4453–4455. <https://doi.org/10.1093/bioinformatics/btz305>.
18. Meier-Kolthoff JP, Göker M. 2019. TYGS is an automated high-throughput platform for state-of-the-art genome-based taxonomy. *Nat Commun* 10:2182. <https://doi.org/10.1038/s41467-019-10210-3>.
19. Webster G, Jones C, Mullins AJ, Mahenthiralingam E. 2020. A rapid screening method for the detection of specialised metabolites from bacteria: induction and suppression of metabolites from *Burkholderia* species. *J Microbiol Methods* 178:106057. <https://doi.org/10.1016/j.mimet.2020.106057>.