

This is the final peer-reviewed accepted manuscript of:

Broeckel, I., Cicoli, M., Maharana, A., Singh, K., & Sinha, K. (2022). On the search for low W0. *Fortschritte Der Physik*, 70(6)

The final published version is available online at <https://dx.doi.org/10.1002/prop.202200002>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.



Fortschritte der Physik -
Progress of Physics

On the Search for Low W_0

Journal:	<i>Fortschritte der Physik - Progress of Physics</i>
Manuscript ID	prop.202200002.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Singh, Kajal; Harish-Chandra Research Institute, A CI of Homi Bhabha National Institute Broeckel, Igor; University of Bologna, Physics and Astronomy Cicoli, Michele; University of Bologna, Physics and Astronomy Maharana, Anshuman; Harish-Chandra Research Institute, A CI of Homi Bhabha National Institute Sinha, Kuver; University of Oklahoma
Keywords:	Flux Compactification, String Phenomenology, String Compactification, High Energy Physics, Superstring Vacua
Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.	
WOPlainReply.toc WOPlainReply.flx WOPlainReply.aux WOPlainReply.fdb_latexmk	

SCHOLARONE™
Manuscripts

On the Search for Low W_0

Igor Broeckel ^{a,b}, Michele Cicoli ^{a,b}, Anshuman Maharana ^c,
Kajal Singh ^c Kuver Sinha ^d

^a Dipartimento di Fisica e Astronomia, Università di Bologna, via Irnerio 46, 40126 Bologna, Italy

^b INFN, Sezione di Bologna, viale Berti Pichat 6/2, 40127 Bologna, Italy

^c Harish-Chandra Research Institute, A CI of Homi Bhabha National Institute, Allahabad 211019, India

^d Department of Physics and Astronomy, University of Oklahoma, Norman, OK 73019, USA

igor.broeckel@bo.infn.it, michele.cicoli@unibo.it, anshumanmaharana@hri.res.in,
kajalsingh@hri.res.in, kuver.sinha@ou.edu

Abstract

The magnitude of the vacuum expectation value of the Gukov-Vafa-Witten superpotential $|W_0|$ plays a central role in the phenomenology of type IIB flux compactifications. Recent analytical constructions have shown that perturbatively flat vacua can be used to obtain very low values of $|W_0|$. We present systematic algorithms to carry out exhaustive numerical searches for such vacua. We also analyse them in the statistical context, as part of the entire ensemble of type IIB flux vacua at low $|W_0|$. Our preliminary analysis indicates that these perturbatively flat vacua are statistically sparse in the whole set of vacua at low $|W_0|$ as calculated by Denef and Douglas. Two-moduli examples are used to illustrate these more general findings in specific settings. We find that these simple (**two moduli**) cases are good examples for existence proofs but they do not feature a large statistical tuning freedom for phenomenological applications.

Contents

1	Introduction	1
2	A brief review of perturbatively flat vacua	3
3	Expectations from statistics	5
4	Exhaustive search with two moduli	6
4.1	The $\mathbb{C}\mathbb{P}_{[1,1,1,6,9]}$ example	6
4.2	The algorithm	7
4.3	General treatment of 2-moduli case	10
4.4	Comparison with statistics	11
5	A more general search algorithm	11
6	Conclusion and discussion	14
A	2-moduli examples data	17

1 Introduction

Central to the venture of string phenomenology is to carry out explicit constructions of reliable vacua which are phenomenologically viable. In this light, particularly attractive are type IIB flux compactifications [1]. Here, the complex structure moduli and axio-dilaton can be stabilised by turning on background 3-form fluxes. There are instead various scenarios for stabilising the Kähler moduli (see for example [2–11]). The Standard Model can be realised on intersecting D-branes, branes at singularities or their F-theory generalisations [12]. Cosmic inflation can also be driven by either closed or open string moduli [13, 14]. Furthermore, the large number of possibilities for choosing flux quanta leads to a multitude of solutions which can provide a way to tune the parameters of the associated 4-dimensional effective theory [15, 16].

Our understanding of the physics of type IIB flux compactifications has been growing steadily. This often involves the discovery of a novel class of solutions which have some desirable property needed for the construction of string vacua. One such very interesting class has been discovered recently [17]. These vacua are in the large complex structure limit of the underlying Calabi-Yau of the compactification. They correspond to choices of flux quanta that yield a Gukov-Vafa-Witten superpotential [18] which, when computed using the perturbative part of the prepotential, is a degree-2 homogeneous polynomial in the complex structure moduli and the axio-dilaton. As a result, at this level, these vacua have a flat direction and the expectation value of the Gukov-Vafa-Witten superpotential vanishes along the flat direction. Therefore, these vacua have been dubbed ‘perturbatively flat’. The flat direction is lifted when non-perturbative corrections to the prepotential are incorporated. With this, the Gukov-Vafa-Witten superpotential acquires a value which is exponentially small (at weak string coupling).

This discovery is particularly interesting in the context of KKLT models [2]. Defining as in [17] the vacuum expectation value of the Gukov-Vafa-Witten superpotential as¹

$$W_0 \equiv \sqrt{\frac{2}{\pi}} \langle e^{\mathcal{K}} \int_X G_3 \wedge \Omega \rangle, \quad (1.1)$$

¹Unless otherwise stated, in this article we will follow all the conventions of [17].

where \mathcal{K} is the Kähler potential for the complex structure moduli and axio-dilaton, G_3 is the complexified 3-form flux, and Ω the holomorphic 3-form of the underlying (orientifolded) Calabi-Yau X , controlled KKLT vacua require exponentially small values of $|W_0|$.² This is effectively realised in perturbatively flat vacua which feature $|W_0| \sim e^{-2\pi/(cg_s)} \ll 1$ at small string coupling $g_s \ll 1$ (with $c \in \mathbb{Q}^+$). The paper [17] presented an explicit choice of flux quanta in an orientifold of the Calabi-Yau obtained by considering a degree-18 hypersurface in $\mathbb{C}\mathbb{P}_{[1,1,1,6,9]}$, which yielded $|W_0| \sim 10^{-8}$ (for earlier work on obtaining low values of $|W_0|$ see for example [20,21]). Not stopping at that, an explicit example with $|W_0|$ as low as 10^{-95} was presented in [22, 23]. Here, important advances were made in developing Kähler moduli stabilisation in this context. Furthermore, [24, 25] extended the method to settings with a shrinking conifold modulus, an essential ingredient of the KKLT construction for anti-brane uplifting. The generalisation to F-theory has been considered in [26].

Perturbatively flat vacua are important also for LVS recent explicit realisations of the Standard Model with D3-branes at an orientifolded dP_5 singularity [27]. In these constructions the cancellation of all D7-charges and Freed-Witten anomalies forces the presence of a hidden D7-sector with non-zero gauge fluxes which induce a T-brane background suitable for de Sitter uplifting [28]. As can be seen from equations (5.41) and (5.46) of [27] the T-brane contribution can give a leading order Minkowski vacuum if $|W_0|$ takes a form similar to the one typical of perturbative flat vacua since $|W_0| \sim \lambda_1 e^{-2\pi\lambda_2/g_s}$ where λ_1 and λ_2 are $\mathcal{O}(1)$ model-dependent coefficients which depend on microscopic quantities like the Calabi-Yau Euler characteristic and intersection numbers, the number of blow-up modes, gauge flux quanta and the rank of condensing gauge groups. Phenomenologically viable vacua with a de Sitter minimum and soft terms above the TeV scale can require³ $|W_0|$ as small as $|W_0| \sim 10^{-13}$. All of these are significant developments in the direction of explicit constructions of fully reliable de Sitter vacua in string theory.

Returning to a broader discussion, phenomenological requirements will invariably lead us to specific subclasses of flux vacua (such as perturbatively flat vacua for low $|W_0|$). As we continue to examine flux vacua in detail, we will certainly discover many other interesting subclasses. Given a subclass of flux vacua, there are two important questions that are natural to ask:

- How does the subclass fit within the larger ensemble of the full set of vacua? More specifically, what can we say about the set from the point of view of the statistical approach to string phenomenology [30–34] (see [35–49] for studies in various settings in this context)?
- How can one carry out exhaustive searches which will allow us to have a complete understanding of the vacua in this set (and their physics)?

The goal of this paper is to take the first steps and develop methods necessary to answer the above questions in the context of perturbatively flat vacua. In the process, we hope to learn some lessons which should be applicable to the study of any subclass. Apart from the general motivation, there are interesting reasons to address both questions in the context of perturbatively flat vacua.

Usually, finding flux vacua requires solving a coupled set of equations involving the flux quanta and the complex structure moduli. In the case of perturbatively flat vacua, there is considerable simplification. As we will review below, to find vacua one just needs to solve a set of diophantine equations involving the flux quanta (once solutions to this set are found, the vacuum expectation values of the complex structure moduli

²For a general discussion on the magnitude W_0 in the context of moduli stabilisation and phenomenological implications, see [19] and references therein.

³Notice that very small values of $|W_0|$ are not a necessary condition for T-brane uplifting since this depends crucially on the model-dependent values of λ_1 and λ_2 . In fact, [29] found explicit LVS de Sitter models with $|W_0| \sim \mathcal{O}(1 - 10)$.

are automatically determined by a simple analytic formula). Given this simplification, perturbatively flat vacua are the ideal set to look at to develop methods for exhaustive searches for flux vacua.

As already mentioned, perturbatively flat vacua provide a natural way to construct KKLT models and can be useful for effective T-brane uplifting in some LVS models. Thus, developing an understanding of how they fit into the full set of flux vacua (in the statistical context) is important for obtaining the statistical predictions for observables in these models. For instance, the analysis of [50, 51] implies that if $|W_0|$ is exponentially small in the dilaton in most vacua in a set, then the scale of supersymmetry breaking has a logarithmic distribution. The above property is true for all perturbatively flat vacua. Therefore, gaining an understanding of what fraction of the vacua at low $|W_0|$ are perturbatively flat is central to determining the distribution of the scale of supersymmetry breaking in KKLT models. The distribution of the scale of supersymmetry breaking in the landscape is of course of much interest [52–57].

Before closing the introduction, we would like to make some comments regarding the approach that this article takes. Work on the search for flux vacua and their properties is a two step process - development of methods and then extensive numerical scan through models. The focus of the present paper is on the former. While we will make use of specific models to illustrate the methods,⁴ we will not be carrying out any extensive numerical scans through models. In fact, we will often stop midway with the analysis of particular models when the necessary point regarding the methods is made. We leave detailed numerical scans of models for future work [58].

This paper is organised as follows. In Sec. 2 we review the main ingredients of perturbatively flat vacua, while in Sec. 3 we discuss their statistical significance. Sec. 4 provides all the details of an algorithm to perform exhaustive searches for perturbatively flat vacua for the case with 2 complex structure moduli. In Sec. 5 we outline instead a more general search algorithm which is valid in principle to obtain perturbatively flat vacua for examples with any number of complex structure moduli (although the requirement of computational resources grows with the number of complex structure moduli, this growth has the number of complex structure moduli in an exponent). We present our conclusions and discuss our results in Sec. 6. Some technical details regarding our numerical search for cases with 2 complex structure moduli are summarised in App. A.

2 A brief review of perturbatively flat vacua

In this section we first recapitulate some basic material on type IIB flux compactifications and then go on to review [17]. Our discussion in the first part shall be primarily to set notation and will be quite brief. We refer the reader to [1, 59–62] for further details.

Type IIB flux compactifications have an internal manifold that is conformally an orientifolded Calabi-Yau X . To describe these in the language of special geometry, one works with a symplectic basis for $H_3(X, \mathbb{Z})$, $\{A_a, B^a\}$ for $a = 1, \dots, h_-^{1,2}(X)$ with $A_a \cap A_b = 0$, $A_a \cap B^b = \delta_a^b$, and $B^a \cap B^b = 0$, and projective coordinates on the complex structure moduli, U^a (in what follows, we will take $U^0 = 1$). The central object is the prepotential \mathcal{F} , which is degree-2 and homogeneous in the projective coordinates. The period vector is given by

$$\Pi = \begin{pmatrix} \int_{B^a} \Omega \\ \int_{A_a} \Omega \end{pmatrix} = \begin{pmatrix} \mathcal{F}_a \\ U^a \end{pmatrix}. \quad (2.1)$$

The flux vectors F and H are obtained by integrating the 3-form field strengths of the type IIB theory over

⁴For this we will work with models with 2 complex structure moduli, keeping the numerics light.

the A_a and B_a cycles

$$F = \begin{pmatrix} \int_{B^a} F_3 \\ \int_{A_a} F_3 \end{pmatrix}, \quad H = \begin{pmatrix} \int_{B^a} H_3 \\ \int_{A_a} H_3 \end{pmatrix}. \quad (2.2)$$

Dirac quantisation conditions require that these are integer valued. The flux superpotential, which is perturbatively exact, is given by

$$W = \sqrt{\frac{2}{\pi}} (F - \tau H)^T \cdot \Sigma \cdot \Pi, \quad (2.3)$$

where

$$\Sigma = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad (2.4)$$

is the symplectic matrix. The tree-level Kähler potential (for the complex structure moduli and the axio-dilaton) is

$$\mathcal{K} = -\ln(-i\Pi^\dagger \cdot \Sigma \cdot \Pi) - \ln(-i(\tau - \bar{\tau})). \quad (2.5)$$

In the large complex structure limit,⁵ the prepotential is a sum of perturbative terms which are at most degree-3 and instanton corrections, i.e $\mathcal{F}(U) = \mathcal{F}_{\text{pert}}(U) + \mathcal{F}_{\text{inst}}(U)$ with

$$\mathcal{F}_{\text{pert}}(U) = -\frac{1}{3!} \mathcal{K}_{abc} U^a U^b U^c + \frac{1}{2} \mathbf{a}_{ab} U^a U^b + b_a U^a + \xi, \quad (2.6)$$

where \mathcal{K}_{abc} are the triple intersection numbers of the mirror Calabi-Yau, \mathbf{a}_{ab} and b_a are rational, and $\xi = -\frac{\zeta(3)\chi}{2(2\pi i)^3}$, with χ the Euler number of the Calabi-Yau. The instanton corrections are

$$\mathcal{F}_{\text{inst}}(U) = \frac{1}{(2\pi i)^3} \sum_{\vec{q}} A_{\vec{q}} e^{2\pi i \vec{q} \cdot \vec{U}}, \quad (2.7)$$

where the sum runs over effective curves in the mirror Calabi-Yau.

Supersymmetric vacua which have $W = 0$ at the perturbative level of the prepotential and also have a flat direction were termed as perturbatively flat in [17]. The basic idea of [17] is that, when the instanton corrections are incorporated, the flat direction is lifted and W acquires an exponentially small vacuum expectation value. Furthermore, the paper provides an explicit algorithm to obtain perturbatively flat vacua, which was stated in the form of a Lemma.

The statement of the Lemma is: if there is a pair $(\vec{M}, \vec{K}) \in \mathbb{Z}^n \times \mathbb{Z}^n$ satisfying $N_{\text{flux}} \equiv -\frac{1}{2} \vec{M} \cdot \vec{K} \leq Q_{\text{D3}}$ (Q_{D3} being the D3-charge tadpole bound), such that $N_{ab} \equiv \mathcal{K}_{abc} M^c$ is invertible, and $\vec{K}^T N^{-1} \vec{K} = 0$, and $\vec{p} \equiv N^{-1} \vec{K}$ lies in the Kähler cone of the mirror Calabi-Yau, and such that $\mathbf{a}_{ab} M^b$ and $b_a M^a$ take on values in integers; then there exists a choice of fluxes for which a perturbatively flat vacuum exists. The perturbative F-flatness conditions are satisfied along the 1-dimensional subspace $\vec{U} = \tau \vec{p}$, on which W_{pert} vanishes. The Lemma is easily verified by taking the flux vectors to be

$$F = (\vec{M} \cdot \vec{b}, \vec{M}^T \cdot \mathbf{a}, 0, \vec{M}^T) \quad \text{and} \quad H = (0, \vec{K}^T, 0, 0). \quad (2.8)$$

The above choice of the flux vectors is also the most general that leads to a superpotential that is a degree-2 homogenous polynomial in the $(h_{-}^{1,2} + 1)$ moduli.⁶ Note that this guarantees that the F-flatness conditions imply $W_{\text{pert}} = 0$, and also the existence of the flat direction.

⁵For detailed studies of flux vacua in the large complex structure limit see e.g. [63–69].

⁶Degree-2 homogeneous flux superpotentials and associated flat directions in toroidal compactifications were discussed in [70].

As mentioned earlier, the flat direction is lifted by the non-perturbative terms in \mathcal{F} . Choosing the axio-dilaton to be the coordinate along the flat direction, the superpotential which is effectively generated looks like

$$\frac{W_{\text{eff}}(\tau)}{\sqrt{2/\pi}} = M^a \partial_a \mathcal{F}_{\text{inst}} = \sum_{\vec{q}} \frac{A_{\vec{q}} \vec{M} \cdot \vec{q}}{(2\pi i)^2} e^{2\pi i \tau \vec{p} \cdot \vec{q}}. \quad (2.9)$$

The above superpotential can lead to a controlled racetrack stabilisation if the two dominant instantons (which we will call \vec{q}_1 and \vec{q}_2) satisfy $\vec{p} \cdot \vec{q}_1 \approx \vec{p} \cdot \vec{q}_2$. Furthermore, stabilisation at weak string coupling requires that there is a hierarchy between the prefactors of the instantons. This amounts to a hierarchy in the associated Gopakumar-Vafa invariants [71, 72].

3 Expectations from statistics

As discussed in the introduction, it is of much interest to develop an understanding of how perturbatively flat vacua fit in the larger ensemble of type IIB flux vacua in the statistical sense. The question is central to understanding the distribution of the scale of supersymmetry breaking for KKLТ vacua [50]. Perturbatively flat vacua are supersymmetric (even after the incorporation of instanton effects in the prepotential) and have low values of $|W_0|$. The statistical properties of such vacua were derived in [32]. The number of such vacua \mathcal{N} with the value of $|W_0|^2$ below λ_* is given by an integral of a density over the moduli space⁷

$$\mathcal{N}(N_{\text{flux}} \leq Q_{\text{D3}}, |W_0|^2 \leq \lambda_*) = \frac{(2\pi Q_{\text{D3}})^{2m} \pi \lambda_*}{2(2m)!} \int_{\mathcal{M}} d^{2m}z \sqrt{g} \rho(z), \quad (3.1)$$

where the density function is given by

$$\rho(z) = \frac{2\pi m}{\pi^{2m} Q_{\text{D3}}} I(\mathcal{F}) \quad \text{for} \quad I(\mathcal{F}) = \int d^{2h_-^{1,2}} Z e^{-|Z|^2} \left| \det \begin{pmatrix} 0 & Z_J \\ Z_I & e^{\mathcal{K}} \mathcal{F}_{IJK} \bar{Z}^K \end{pmatrix} \right|^2, \quad (3.2)$$

with $m = h_-^{1,2} + 1$ ($h_-^{1,2}$ being the number of complex structure moduli). The $d^{2m}z$ integration runs over the $2m$ -dimensional space of the complex structure moduli and the axio-dilaton and it involves its metric. \mathcal{F}_{IJK} are components of triple derivatives of the prepotential expressed in a local frame. The integration variables Z_I are related to derivatives of the flux superpotential, but can be thought of as dummy integration variables for the purposes of computation of $I(\mathcal{F})$.

Now, let us turn to examining perturbatively flat vacua in this context. For this, we will exploit universal properties of these vacua. A striking property of these vacua is that for all of them, at their minima

$$\vec{U} = \tau \vec{p}, \quad (3.3)$$

where the vector \vec{p} is real and has all positive entries. The real parts of \vec{U} are axionic. Thus, after the axions are brought to their fundamental domain, the relations in (3.3) will continue to hold modulo factors of integers. Therefore, the solutions under consideration are contained in a subspace of the moduli space which is isomorphic to $\mathcal{M}_\tau \times (\mathbb{R}^+)^{h_-^{1,2}}$. Given that perturbatively flat vacua are a subset of the set of all solutions satisfying (3.3), a necessary criterion for them to have statistical significance is that the set of all solutions on the subspace (3.3) have statistical significance. This question can be examined from the point of view of the densities of [32]. **Here, the continuous flux approximation was taken, correspondingly for the discussion at hand, one can imagine that the vector \vec{p} can be dialed continuously, sweeping out essentially the entirety**

⁷In the discussion below, we translate the results of [32] and report them in the conventions of [17].

of the Kähler cone of the mirror-CY. Notice that the entire moduli space is $2(h_-^{1,2} + 1)$ -dimensional, while the subspace described by (3.3) is only $(h_-^{1,2} + 2)$ -dimensional. Since the subspace (3.3) is of lower dimensionality than the entire moduli space, and the densities are smooth function on the entire moduli space, even the set of all vacua on this subspace are not expected to be of statistical significance, implying the same for perturbatively flat vacua. Hence perturbatively flat vacua are expected to be statistically sparse in the set of flux vacua with low $|W_0|$ as given by the distribution from [32] which already gives a much smaller number of vacua with respect to cases with $|W_0| \sim \mathcal{O}(1 - 10)$ since (3.1) is linear in λ_* , implying (at fixed Q_{D3}) $\mathcal{N}(|W_0|^2 \leq \lambda_*) / \mathcal{N}(|W_0|^2 \leq 1) \sim \lambda_*$.

Notice that comparisons in [32] of results of explicit searches to the predictions making use of densities did show local fluctuations (such as overdensities and voids) but these were local effects not having any effect on the overall statistical predictions. It is however important to check if the distributions of [32] are peaked along the space described by (3.3). Consider the subspace in which the axio-dilaton is purely imaginary. Being on (3.3), then implies that U^a are also purely imaginary. The densities of [32] can be expressed in terms of the Kähler potential, the metric on the moduli space and triple derivatives of the prepotential \mathcal{F}_{IJK} . For the perturbative part of the prepotential (which dominates in the large complex structure limit), all the above quantities are independent of the real part of U^a . Thus the densities are also independent of the real part of U^a . Therefore, at fixed purely imaginary axio-dilaton, moving away from the locus (3.3) by switching on a non-zero real part of U^a does not lead to a fall in the value of the density. Similar considerations also apply when the axio-dilaton is not purely imaginary.

We would like to close this section with a cautionary remark. **The diagnostics presented here relies on the fact that the basic reasoning of [32] is valid, i.e. that the space of flux vacua of a given compactification can be described by smooth density functions obtained by replacing sums over flux quanta by integrals. The reasoning of [32] is based on the general theorems in [88]. If for some reason this fails in some cases at hand, the diagnostics would be irrelevant and there could be a high density of vacua localised in the $(h_-^{1,2} + 2)$ -dimensional subspace of the moduli space for $|W_0|$ below a certain value (in contradiction with the general reasoning of [32]).** Next, we turn our discussion of setting up exhaustive searches for perturbatively flat vacua, which is crucial for developing a full understanding of their explicit properties.

4 Exhaustive search with two moduli

4.1 The $\mathbb{C}\mathbb{P}_{[1,1,1,6,9]}$ example

In this section we describe algorithms for carrying out exhaustive searches for perturbatively flat vacua in Calabi-Yau threefolds with 2 complex structure moduli. As mentioned in the introduction, even if this paper intends mainly to focus on methods for searches of flux vacua, for completeness we will present an explicit example in full detail. We do so by looking at the degree-18 hypersurface in $\mathbb{C}\mathbb{P}_{[1,1,1,6,9]}$ used in [17] (studied in the context of mirror symmetry in [73]).

We begin by recalling some basic facts about the Calabi-Yau and some details of the analysis of [17]. The Calabi-Yau has 272 complex structure moduli but has a $\mathcal{G} = \mathbb{Z}_6 \times \mathbb{Z}_{18}$ symmetry. By considering \mathcal{G} -invariant fluxes, a solution found in truncation to G-singlets is guaranteed to lift to a solution of the full theory (see [20]). Thus the stabilisation problem is effectively reduced to a 2-moduli one. The relevant geometric data are

$$\mathcal{K}_{111} = 9, \quad \mathcal{K}_{112} = 3, \quad \mathcal{K}_{122} = 1, \quad \mathbf{a} = \frac{1}{2} \begin{pmatrix} 9 & 3 \\ 3 & 0 \end{pmatrix}, \quad \vec{b} = \frac{1}{4} \begin{pmatrix} 17 \\ 6 \end{pmatrix}, \quad (4.1)$$

and the instanton corrections are

$$(2\pi i)^3 \mathcal{F}_{\text{inst}} = \mathcal{F}_1 + \mathcal{F}_2 + \dots, \quad (4.2)$$

$$\mathcal{F}_1 = -540 q_1 - 3 q_2, \quad (4.3)$$

$$\mathcal{F}_2 = -\frac{1215}{2} q_1^2 + 1080 q_1 q_2 + \frac{45}{8} q_2^2, \quad (4.4)$$

where $q_a = \exp(2\pi i U^a)$ with $a \in \{1, 2\}$. We consider the orientifold involution described in [74] which yields a D3-charge $Q_{\text{D3}} = 138$.

Making use of (4.1), the condition $\vec{K}^T N^{-1} \vec{K} = 0$ gives

$$M^1 = \frac{M^2 K^2 (2K^1 - 3K^2)}{(K^1 - 3K^2)^2}, \quad (4.5)$$

and the flat direction is

$$\vec{U} = \tau \begin{pmatrix} p^1 \\ p^2 \end{pmatrix} = \frac{\tau (K^1 - 3K^2)}{M^2} \begin{pmatrix} -K^2/K^1 \\ 1 \end{pmatrix}. \quad (4.6)$$

The following choice of the vectors (\vec{M}, \vec{K})

$$\vec{M} = \begin{pmatrix} -16 \\ 50 \end{pmatrix}, \quad \vec{K} = \begin{pmatrix} 3 \\ -4 \end{pmatrix}, \quad (4.7)$$

meets all the conditions of the Lemma and the flat direction can be lifted by the inclusion of non-perturbative terms.

In the large complex structure limit, the Kähler potential (for the complex structure moduli and axio-dilaton) is given by

$$K = -\ln \left(i \frac{1}{6} \mathcal{K}_{abc} (U^a - \bar{U}^a)(U^b - \bar{U}^b)(U^c - \bar{U}^c) + 4i\xi \right) - \ln(-i(\tau - \bar{\tau})). \quad (4.8)$$

We are interested in the locus $U^a = p^a \tau$. Furthermore, since in this limit $\text{Im}(U^a) > 1$, the term involving ξ is subdominant. Thus, along this locus one has

$$\begin{aligned} K &= -\ln \left(\frac{1}{6} \mathcal{K}_{abc} p^a p^b p^c (-i(\tau - \bar{\tau}))^3 + 4i\xi \right) - \ln(-i(\tau - \bar{\tau})) \\ &\sim -\ln \left(\frac{1}{6} \mathcal{K}_{abc} p^a p^b p^c \right) - 4 \ln(-i(\tau - \bar{\tau})). \end{aligned} \quad (4.9)$$

The effective superpotential for stabilising the perturbatively flat direction takes the form

$$W_{\text{eff}}(\tau) = c \left(e^{2\pi i p^1 \tau} + A e^{2\pi i p^2 \tau} \right) + \dots, \quad (4.10)$$

where $c = \sqrt{\frac{2}{\pi}} \frac{8640}{(2\pi i)^2}$ and $A = -\frac{5}{288}$. Making use of the fluxes in (4.7), it can be easily found that $|W_0| \simeq 2 \times 10^{-8}$.

4.2 The algorithm

Now, we describe an algorithm for finding all perturbatively flat vacua in the $\mathbb{CP}_{[1,1,1,6,9]}$ model, which can however be easily generalised to other 2-moduli examples. The F-flatness condition is $D_\tau W_{\text{eff}} = (\partial_\tau + \partial_\tau K) W_{\text{eff}} = 0$. Note that the form of the Kähler potential (4.9) implies that $\partial_\tau K \propto (\text{Im}\tau)^{-1} = g_s$.

Therefore, for consistent stabilisation at weak string coupling, the term involving $\partial_\tau K$ must be a small correction to the F-flatness condition. The F-flatness condition neglecting this term is

$$e^{2\pi i\tau(p^1-p^2)} = -\frac{Ap^2}{p^1}. \quad (4.11)$$

Let us start our search by considering the cases with $p^1 > p^2$. Now, by making use of (4.6) the condition $p^1 > p^2$ translates to

$$-\frac{K^2}{K^1} > 1. \quad (4.12)$$

Thus K^1 and K^2 have to be of opposite sign. Furthermore, the entire set of conditions in the Lemma have a symmetry:

$$\vec{K} \rightarrow -\vec{K} \quad \text{and} \quad \vec{M} \rightarrow -\vec{M}. \quad (4.13)$$

This in fact corresponds to an S-duality transformation with the centre of the group. Thus, without loss of generality, we will look at cases with $K^1 > 0$ and $K^2 < 0$. With this, the factor $(K^1 - 3K^2)$ in (4.6) is positive, implying that M^2 must be positive (so that p^2 is positive, as required by the Kähler cone condition in the Lemma). With these signs of K^1 , K^2 and M^2 , equation (4.5) gives the sign of M^1 to be negative. This implies that $A = M^2/(180M^1)$ has to be negative, which can be compatible with (4.11) if at the minimum $\text{Re}(\tau) = k/(p^1 - p^2) \pmod{\mathbb{Z}}$ with $k \in \mathbb{Z}$. The above suggests the following efficient algorithm to carry out an exhaustive search for vacua:

1. Consider a rational number x between 0 and 1, express this as $x = p/q$ such that p and q are positive and have no common factors. Define the vector

$$\vec{\tilde{K}} = \begin{pmatrix} \tilde{K}^1 \\ \tilde{K}^2 \end{pmatrix} = \begin{pmatrix} p \\ -q \end{pmatrix}. \quad (4.14)$$

The vector $\vec{\tilde{K}}$ will eventually be related to the vector \vec{K} being searched for.

2. Now, compute the ratio

$$y = \frac{\tilde{K}^2 (2\tilde{K}^1 - 3\tilde{K}^2)}{(\tilde{K}^1 - 3\tilde{K}^2)^2}. \quad (4.15)$$

Note that this is related to the ratio M^1/M^2 as given by (4.5). Express y as r/s , such that r and s have no common factors, and $s > 0$. Define

$$\vec{\tilde{M}} = \begin{pmatrix} r \\ s \end{pmatrix}. \quad (4.16)$$

The vector $\vec{\tilde{M}}$ will be eventually related to the vector \vec{M} being searched for.

3. Check if $K_{abc}\tilde{M}^c$ is invertible or not. If it is not invertible, discard x and start again with a new one. If it is invertible, then proceed further.

4. Compute the values

$$\alpha \mathbf{a} \cdot \vec{\tilde{M}} \quad \text{and} \quad \alpha \vec{b} \cdot \vec{\tilde{M}}, \quad (4.17)$$

for $\alpha = 1, 2, 4$. Determine the minimum value of α for which the above quantities are integer valued. Call this $\hat{\alpha}$. Note that they certainly must be integer valued for the case of $\alpha = 4$, given the form of \mathbf{a} and \vec{b} in equation (4.1).

\vec{M}^T	\vec{K}^T	$\vec{b} \cdot \vec{M}$	$(\mathbf{a} \cdot \vec{M})^T$	N_{flux}	τ	\vec{U}^T	$ W_0 $
(32, -98)	(-1, 2)	-11	(-3, 48)	114	10.255 i	(1.465, 0.7325) i	6.989×10^{-4}
(16, -50)	(-3, 4)	-7	(-3, 24)	124	6.855 i	(2.742, 2.0565) i	2.048×10^{-8}

 Table 1: All perturbatively flat vacua for the $\mathbb{C}\mathbb{P}_{[1,1,1,6,9]}$ example.

5. Consider the quantity

$$-\frac{1}{2} \hat{\alpha} \vec{M} \cdot \vec{K}. \quad (4.18)$$

If this does not satisfy the D3-tadpole bound, then discard x and move to another x . If it lies in the allowed range, we have a solution satisfying all conditions of the Lemma with

$$\vec{M} = \hat{\alpha} \vec{M} \quad \text{and} \quad \vec{K} = \vec{K}. \quad (4.19)$$

Also, for any positive integer β such that $-\frac{1}{2} \beta \hat{\alpha} \vec{M} \cdot \vec{K}$ satisfies the D3-tadpole bound, we have solutions

$$\vec{M} = \hat{\alpha} \beta_1 \vec{M} \quad \text{and} \quad \vec{K} = \beta_2 \vec{K}, \quad (4.20)$$

where β_1 and β_2 are positive and provide a factorisation of β .

6. To scan through all x , note that the signs of K^a and M^a (with our working assumption of $K^1 > 0$) are such that $M^1 K^1 < 0$ and $M^2 K^2 < 0$. Thus both terms contribute with a *positive* sign to the inner product $-\frac{1}{2} \vec{M} \cdot \vec{K}$. Therefore, *the maximum value* of $|K^2|$ necessary to carry out an exhaustive search is $2Q_{\text{D3}} = 2 \times 138$ (as higher values would violate the D3-tadpole condition). This bound on $|K^2|$ implies that we need to consider only those x for which $q \leq 2Q_{\text{D3}}$. Reduced rationals between 0 and 1 with a fixed upper bound on the denominator are given by the Farey sequence. Thus an exhaustive search is carried out by selecting x from the set $\text{Farey}_{2Q_{\text{D3}}}$.
7. Scan through the solutions obtained in this way, checking that non-perturbative effects lead to stabilisation at weak string coupling. Discard the ones that do not satisfy this condition.
8. Enlarge the solution list by considering the solutions obtained by the above process and then generating the solutions related to them by the S-duality symmetry

$$\vec{K} \rightarrow -\vec{K} \quad \text{and} \quad \vec{M} \rightarrow -\vec{M}. \quad (4.21)$$

9. Finally, run the same algorithm considering the possibility of $p^1 \leq p^2$.

Carrying out the search using the above algorithm, after S-duality identification, we find that there exist only 2 solutions which satisfy the conditions of the Lemma, although one of them is at the borderline for the validity of the large complex structure approximation. We report these in Tab. 1 along with the associated value of $|W_0|$, after stabilisation by non-perturbative effects. The second entry in the table is the solution reported in [17] with $|W_0| \sim 2 \times 10^{-8}$. **The first has a low value of W_0 , but the value of one of the complex structure moduli is slightly below one, in this case a detailed check validity of the large complex structure approximation can be carried out using the results of [73]. In any case, for our purposes (which is to gain an understanding of the statistics), we conclude that the $\mathbb{C}\mathbb{P}_{[1,1,1,6,9]}$ model essentially features only $\mathcal{O}(1)$ perturbatively flat solutions with very low $|W_0|$.**

Q_{D3}	Number of perturbatively flat vacua
50	37
100	128
250	531
500	1445

Table 2: Number of perturbatively flat vacua in the $\mathbb{CP}_{[1,1,2,2,2]}$ model taking Q_{D3} as a free parameter. The reported numbers are before imposing any of the following 3 requirements: stabilisation at weak string coupling, low $|W_0|$, S-duality identification.

4.3 General treatment of 2-moduli case

In this section we will present a general discussion of the cases with 2 complex structure moduli. A key-feature of the algorithm in Sec. 4.2 was the bound on the range of the elements of the vectors \vec{M} and \vec{K} . First we show that this follows from general considerations. The definition $\vec{p} = N^{-1}\vec{K}$, together with the equation $\vec{K}^T N^{-1}\vec{K} = 0$ implies

$$\vec{K}^T \vec{p} = 0. \quad (4.22)$$

The requirement that \vec{p} lies in the Kähler cone, then implies that K^1 and K^2 have opposite signs. By making use of the definition of \vec{p} again, the equation $\vec{K}^T N^{-1}\vec{K}$ can alternatively be written as

$$p^a p^b K_{abc} M^c = 0. \quad (4.23)$$

The requirement that \vec{p} lies in the Kähler cone implies also that $\tilde{p}_c \equiv p^a p^b K_{abc}$ has positive entries. Thus the vector \vec{M} satisfies an equation similar to \vec{K} , i.e.

$$\vec{M}^T \vec{p} = 0. \quad (4.24)$$

Therefore, M^1 and M^2 have to have different signs.

Now, if K^1 and M^1 have the same sign, then so would K^2 and M^2 . And this would imply a negative value for $N_{\text{flux}} = -\frac{1}{2}\vec{M}\cdot\vec{K}$, which is impossible for imaginary self dual fluxes.⁸ Thus viable solutions feature K^1 and M^1 of opposite sign⁹. This implies that both terms contributing to the N_{flux} inner product have to be positive. Thus, an exhaustive search can be carried out by considering the range

$$|M^a| \leq 2Q_{D3} \quad \text{and} \quad |K^a| \leq 2Q_{D3}, \quad (4.25)$$

which is the same as for the $\mathbb{CP}_{[1,1,1,6,9]}$ example, obtained by using slightly different considerations. As an example, we have carried out the analysis for the Calabi-Yau embedded in $\mathbb{CP}_{[1,1,2,2,2]}$ discussed in [75]. To gain a model-independent picture, we treat Q_{D3} as a free parameter. The results are summarised in Tab. 2.

All these solutions can potentially correspond to perturbatively flat vacua but these numbers would be reduced by the following 3 requirements which have still to be imposed: (i) dilaton stabilisation at weak string coupling by instanton effects; (ii) a value of $|W_0|$ which is indeed very small; (iii) possible equivalences between solutions via S-duality. Given that these numbers are still small to be attractive in the context of a landscape, we do not push the analysis further. This expectation has been confirmed by detailed scan of models in [87].

⁸ Any fluxes that solve the conditions being imposed are imaginary self dual from the 10-dimensional perspective (see e.g. [1]).

⁹ These sign correlations have been confirmed in the detailed numerical scans of [87]

It is important to note that the key-element in obtaining the bounds in (4.25) was the sign correlations between the elements in \vec{M} and \vec{K} . While the arguments in the first part of this section hold for any number of moduli, it is easy to see that the sign correlations need not hold when there are more than 2 moduli. To remedy this, we will discuss a more general method in Sec. 5.

4.4 Comparison with statistics

Let us compare our results with the statistical expectations of [32]. For $h_-^{1,2} = 2$, (3.1) yields

$$\mathcal{N}(N_{\text{flux}} \leq Q_{\text{D3}}, |W_0|^2 \leq \lambda_*) = \left(\frac{2^6 \pi^4}{5!} \right) Q_{\text{D3}}^5 \lambda_* \int_{\mathcal{M}} d^6 z \sqrt{g} e^{2\mathcal{K}} \mathcal{F}_{abc} \bar{\mathcal{F}}^{abc}, \quad (4.26)$$

where the indices of \mathcal{F} have been converted to tangent bundle ones. For the $\mathbb{C}\mathbb{P}_{[1,1,1,6,9]}$ example discussed in Sec. 4.1, carrying out the integration over the large complex structure patch one finds

$$\mathcal{N}(N_{\text{flux}} \leq Q_{\text{D3}} = 138, |W_0|^2 \leq \lambda_*) \simeq 3 \times 10^{12} \lambda_*. \quad (4.27)$$

As pointed out in [17], this predicts the lowest value of $|W_0|$ being of order 6×10^{-7} , close to what was found. On the other hand, the same formula predicts $\mathcal{O}(10^8)$ vacua for $|W_0| \lesssim 0.01$, even if our exhaustive search has shown that there are only $\mathcal{O}(1)$ vacua with such a feature, in agreement with the argument presented in Sec. 3.¹⁰ We therefore conclude that in the $\mathbb{C}\mathbb{P}_{[1,1,1,6,9]}$ model, perturbatively flat vacua are interesting examples to show explicitly the existence of vacua with very low $|W_0|$, but they do not possess any tuning freedom in the value of $|W_0|$. Given the argument presented in Sec. 4.3, we expect this conclusion to hold for all cases with 2 complex structure moduli. Notice, for example, that in the $\mathbb{C}\mathbb{P}_{[1,1,2,2,2]}$ model the number of perturbatively flat vacua summarised in Tab. 2 is also much less than as predicted by the Q_{D3}^5 scaling of (4.26). Models with more than 2 complex structure moduli require a refined analysis for exhaustive searches which we outline in the next section, although the analysis of Sec. 3 indicates that they should still be statistically sparse (of course, in this context the caveat discussed at the end of section 3 should be kept in mind).

Let us close this section by stressing that a key-assumption in the derivation of the results of [32], is a high density of flux vacua allowing for the sums over integer fluxes to be converted to integrals. Our results indicate that for the $\mathbb{C}\mathbb{P}_{[1,1,1,6,9]}$ model, under these circumstances, there are many more vacua at low $|W_0|$ that remain to be discovered.

5 A more general search algorithm

The key to carry out exhaustive searches is isolating the region in the flux vector space which contains all perturbatively flat vacua. Once such a region is obtained, one can carry out numerical searches in this region to obtain all solutions (if the region is not too large). In this section we present a general method to isolate such regions which is in principle valid for examples with an arbitrary large number of complex structure moduli. Here, we will discuss the method and leave its detailed numerical implementation for future work.¹¹

¹⁰In this context, we would like to mention that the values of $|W_0|$ obtained after stabilisation crucially depend on the hierarchy in the Gopakumar-Vafa invariants. However, the densities of [32] in the moduli space in the large complex structure limit have mild sensitivity to this. This is in keeping with the arguments of Sec. 3 which suggest that perturbatively flat vacua are a small fraction of the vacua at low $|W_0|$. Of course, the cautionary remark at the end of section 3 is a caveat that should be kept in mind.

¹¹Our preliminary analysis indicates that the numerics can be quite involved when one considers models with more than 2 moduli.

Central to our arguments will be certain properties of N_{flux} . Recall that the quantity $-\frac{1}{2}\vec{M}\cdot\vec{K}$ is equal to the contribution of the fluxes to the D3-charge

$$N_{\text{flux}} = -\frac{1}{2}\vec{M}\cdot\vec{K} = \frac{1}{(2\pi)^4\alpha'^2} \int_X H_3 \wedge F_3, \quad (5.1)$$

where the integration is over the Calabi-Yau X . The fluxes of interest to us correspond to an imaginary self dual G_3 , i.e.

$$*\frac{H_3}{g_s} = -(F_3 - C_0 H_3). \quad (5.2)$$

Thus (see e.g. [76])

$$\int_X H_3 \wedge F_3 = \frac{1}{3!g_s} \int_X d^6y \sqrt{g_6} H_3^2. \quad (5.3)$$

This is the usual argument given to show that N_{flux} is positive semi-definite. Here we list two consequences that are important for our arguments:

- (a) Equation (5.3) implies that the only way for N_{flux} to vanish is $H_3 = 0$. Equation (5.2) then implies that $F_3 = 0$. Translating this in terms of the vectors \vec{M} and \vec{K} , one learns that, for consistent solutions, $N_{\text{flux}} = 0$ only if $\vec{M} = \vec{K} = 0$.
- (b) The derivation of (5.3) does not make use of flux integrality. Thus, the conclusions of the above point remain valid even when one considers fluxes which do not obey the Dirac quantisation conditions (we will do so as an intermediate step in our analysis).

Now, returning to finding the solutions to the conditions of the Lemma, let us think of carrying out a search by scanning through the vectors \vec{M} and \vec{K} , by starting from the origin and progressively going through points with larger and larger $|\vec{M}|$ and $|\vec{K}|$. We would like to obtain upper bounds on the values of $|\vec{M}|$ and $|\vec{K}|$ which can possibly yield solutions to the conditions of the Lemma. For this, we write the D3-tadpole condition as

$$-\frac{1}{2}|\vec{M}||\vec{K}|\epsilon \leq Q_{\text{D3}}, \quad (5.4)$$

where ϵ is the cosine of the angle between the vectors \vec{M} and \vec{K} . Since both $|\vec{M}|$ and $|\vec{K}|$ are bounded from below, the only way $|\vec{M}|$ or $|\vec{K}|$ (or both) can be large is if $|\epsilon|$ is small. While in general the cosine of the angle between two vectors in \mathbb{Z}^n can be arbitrarily small, our interest is only in vectors that satisfy the conditions of the Lemma (i.e. provide consistent solutions to the type IIB equations of motion). We begin by defining

$$\hat{m} = \frac{\vec{M}}{|\vec{M}|}, \quad \hat{k} = \frac{\vec{K}}{|\vec{K}|} \quad \text{and} \quad \hat{\mathbf{n}}_{ab} = K_{abc} m^c. \quad (5.5)$$

The vectors \vec{m} and \vec{k} lie on the unit sphere and the integrality condition of the fluxes is now that the ratio of any two components of the vectors is rational. The equation constraining the vectors in the Lemma becomes

$$\hat{k}^T \hat{\mathbf{n}} \hat{k} = 0. \quad (5.6)$$

We will consider the equation (5.6) as an equation over *real* variables \vec{m} and \vec{k} (taking values on the unit sphere). Furthermore, we will demand that the vector $\hat{p} = \hat{\mathbf{n}}^{-1} \hat{k}$ lies in the Kähler cone of the mirror Calabi-Yau. With the variables taking on values over reals, the solution space can be studied using standard numerical methods. A lower bound on $|\epsilon|$ can be obtained by numerically searching for the minimum (or

infimum) of $|\hat{m} \cdot \hat{k}|$ in the solution space. Once such a bound is obtained, an exhaustive search can be carried out by scanning through

$$0 < |\vec{M}|, |\vec{K}| \leq \frac{2Q_{D3}}{|\epsilon|_{\text{inf}}} . \quad (5.7)$$

We note that a bound so obtained is conservative, due to the expansion of the domain of the variables to the reals.

Next, we would like to discuss some aspects of the minimisation problem at hand. As we have reviewed above, as long as one is in the physically allowed region of the moduli space, N_{flux} is always greater than zero, i.e. $|\epsilon| > 0$. Thus there are two possibilities for the infimum of $|\epsilon|$: either it is a positive number or it is equal to zero. In the former case, an exhaustive search can be carried out by considering $|\vec{M}|$ and $|\vec{K}|$ in the range (5.7).

The later case (in which $|\epsilon|$ takes on arbitrarily small values) is more subtle. In this case, there would be a point with $|\epsilon| = 0$ as a limit point of points in the solution space. Since all points in the physically allowed region must have $|\epsilon| > 0$, the limit point must lie in the boundary of the physical region. Typically, as one approaches the boundary, one loses control over the effective field theory (for example new light degree of freedom can arise) or encounters phenomenological challenges (as we will see in our discussion below). Taking this into consideration will lead to an effective $|\epsilon|_{\text{inf}}$ which can be used to determine a region to carry out exhaustive searches.¹² **A limitation of the algorithm is that for $|\epsilon|_{\text{inf}}$ of order one one would at large $h_-^{2,1}$ have to scan through at least of order $(Q_{D3})^{2h_-^{2,1}}$ candidate flux quanta, which is exponentially unfeasible at large $h_-^{2,1}$.**

To illustrate the method in a concrete setting, we consider the 39 Calabi-Yau threefolds with 2 Kähler moduli¹³ constructed by Kreuzer and Skarke in [77] and listed (along with the intersection numbers) in Table 11 of [11]. In all these cases we have followed the above described procedure to determine $|\epsilon|_{\text{inf}}$. For 22 of them, $|\epsilon|_{\text{inf}}$ does not take values close to zero, implying a strong bound on the region where all solutions are contained. We record the associated values of $|\epsilon|_{\text{inf}}$ for them in Tab. 3 in App. A.

On the other hand, in the remaining 17, the numerics yield very low values of $|\epsilon|_{\text{inf}}$. Thus these models might seem to be more promising to find a larger number of perturbatively flat vacua (from the perspective of the present algorithm). However, as we discuss below, most of the would-be solutions would not be ideal for phenomenological applications. In fact, in these cases we find a solution to the equations with $|\epsilon| = 0$ on the boundary of the Kähler cone, i.e. $\hat{p}^b = 0$ for some b . The definition of \hat{n} in (5.5) together with the definition of \hat{p} implies that $\vec{p} = \hat{p} \frac{|\vec{K}|}{|\vec{M}|}$. Thus we have the relation

$$\vec{U} = \hat{p} \frac{|\vec{K}|}{|\vec{M}|} \tau . \quad (5.8)$$

Being in the large complex structure limit requires $\text{Im}(U^c) > 1 \forall c$. **Considering the limit $\hat{p}^a \rightarrow 0$ for one of the $a \in 1, 2$, while maintaining $\text{Im}(U^a)$ in the large complex structure limit, implies that either $|\vec{K}|/|\vec{M}| \gg 1$ or $\text{Im}(\tau) \rightarrow \infty$.** However both cases are problematic for the following reasons. $|\vec{K}|/|\vec{M}| \gg 1$ will induce large hierarchies in the vector \vec{p} , making it unsuited for racetrack stabilisation at small string coupling¹⁴.

¹²In principle, there can be situations where there are no good reasons to exclude a region of arbitrary small ϵ . In such a case, one would need to carry out a more extensive search along vectors \vec{m} and \vec{k} in this region.

¹³The mirrors have 2 complex structure moduli. Of course, our analysis in Sec. 4 already provides regions for exhaustive searches for these. The goal here is to obtain the analogous regions from the algorithm presented in this section. We will proceed without worrying about issues that can arise from orientifolding.

¹⁴**We find by direction computation that in none of the 17 cases both \hat{p}^1 and \hat{p}^2 tend to zero simultaneously, thus the relation $\vec{p} = \hat{p} \frac{|\vec{K}|}{|\vec{M}|}$ leads to a large hierarchy in the vector \vec{p} if the value of $\frac{|\vec{K}|}{|\vec{M}|}$ is large. This of course is in the context of the 17 two moduli examples at hand.**

On the other hand, $\text{Im}(\tau) = g_s^{-1}$ cannot become too small without inducing phenomenological problems. In fact, in type IIB compactifications the Standard Model can either be realised on D3- or D7-branes. In the first case, the strength of the gauge couplings is set by g_s , and in the second by the Einstein frame volume of the 4-cycle wrapped by the D7-stack (which we denote as $\text{Re}(T_{\text{SM}})$). In the scenario at hand, however Kähler moduli stabilisation [23] gives

$$\frac{4\pi}{g_{\text{SM}}^2} = \text{Re}(T_{\text{SM}}) \simeq \frac{1}{2\pi} \ln |W_0|^{-1} \sim \frac{1}{g_s}. \quad (5.9)$$

Thus, irrespective of how the Standard Model is realised, in perturbatively flat vacua the strength of its gauge couplings is always determined by g_s .¹⁵ This effectively sets a lower bound on the range of interest for g_s (for instance one can demand $10^{-3} \lesssim g_s \lesssim 0.1$). Thus, the regions of small ϵ should be effectively avoided, implying that also the remaining 17 models are not expected to produce a large number of perturbatively flat vacua which are phenomenologically viable. **Here, we have sought KKLТ vacua with small g_s so as to accommodate the Standard Model sector. Let us mention another more general challenge in this context, the results of [79] imply in the case in which the Standard Model is realised on D7 branes, unless the complex phases of the non-perturbative corrections are tuned there is tension with having a small QCD θ angle. Combining both the conditions can lead to strong constraints.**

Before closing this section, we note that the key-aspect of the algorithm has been that, by determining the minimum value of the angle between the flux vectors, one can isolate a region by scanning through which exhaustive searches can be carried out. It will be interesting to see if the same considerations can be used in other settings.

6 Conclusion and discussion

In this article we have developed exhaustive search algorithms to find perturbatively flat vacua. The 2-moduli case has been discussed in detail and an algorithm applicable to any number of moduli has been presented. Detailed numerical scans going through specific models (including ones with higher number of complex structure moduli) will be presented elsewhere [58].

In Sec. 3 we have also examined perturbatively flat vacua as part of the entire ensemble of vacua at low $|W_0|$ from the point of view of a statistical approach. We found that they are statistically sparse when compared to the expectation from the distribution of low values of $|W_0|$ from [32]. This expectation has been confirmed in Sec. 4 by our numerical searches for cases with 2 complex structure moduli. In particular, for the $\mathbb{CP}_{[1,1,1,6,9]}$ model we found that there are only $\mathcal{O}(1)$ perturbatively flat solution with $|W_0| \lesssim 0.01$ (featuring $|W_0| \sim 10^{-8}$), while [32] would predict around $\mathcal{O}(10^8)$ flux vacua. We argued that similar considerations apply to all other 2-moduli cases. This has been confirmed in the detailed numerical scans of two moduli models [87]. We therefore conclude that this set by itself does not provide tuning freedom for phenomenological applications. Using the general algorithm outlined in Sec. 5, it would be interesting in the future to perform a detailed search for cases with more than 2 complex structure moduli [58], although one expects them to be statistically sparse from the analysis of Sec. 3. Let us just mention here that, as one goes to higher values of $h_-^{1,2}$, one can expect more solutions. However, the analysis of [33] implies that with higher values of $h_-^{1,2}$ the vacua in the large complex structure limit give a lower contribution

¹⁵Notice that in the string frame $\text{Re}(T_{\text{SM}})|_{\text{str}} = g_s \text{Re}(T_{\text{SM}}) \sim \mathcal{O}(1)$, implying that one should consider all perturbative and non-perturbative α' corrections at string tree-level, except for those which come from 10-dimensional terms proportional to G_3^{2n} with $n > 1$ when $|W_0| \ll 1$ [78]. However, as shown in [23], these α' effects should induce just a subdominant shift of the KKLТ minimum for $|W_0| \ll 1$.

to the statistics. This poses an interesting challenge for achieving statistical tuning in phenomenological applications. Furthermore, one can expect that the numerics required to obtain all vacua explicitly should become harder as one goes up in the number of complex structure moduli. **Of course, the cautionary remark at the end of section 3 is a caveat that should be kept in mind while making any conclusions regarding the full ensemble of vacua (as opposed to the ensemble captured by the analysis of [32].)**

Let us also stress that our analysis in Sec. 3 relied heavily on the specific form of the vacuum expectation values of the complex structure moduli (equation (3.3) which is specific to the vacua of [17]) but in principle there can be other families of vacua featuring $W = 0$ at perturbative level. An interesting question is to develop diagnostic methods to study the statistical significance of the vacua of [17] in general. Let us touch upon this briefly. For all such vacua, the instanton effects that give W_0 a non-zero value would also be responsible for giving the perturbatively flat direction a mass. Thus a universal property is a modulus (in the subspace spanned by the complex structure moduli and the axio-dilaton) with a low mass, more specifically a mass proportional to a positive power of $|W_0|$. Given this, one can ask whether there is a correlation between low $|W_0|$ and a modulus of low mass. This question can be addressed by examining the bosonic mass matrix of Sec. 3.2 of [32]. If one of the masses scales as $|W_0|^k$ (for some positive k), then the determinant of the mass matrix would scale as $|W_0|^{2k}$, i.e. it would vanish in the $W_0 \rightarrow 0$ limit. On the other hand, taking $W_0 \rightarrow 0$ (which is equal to X in the notation of [32]) is not a sufficient condition for the vanishing of the determinant. This indicates that the correlation is not universal, and so that there should exist another set of vacua with $W = 0$ at perturbative level but with no flat directions. This observation agrees with the analysis of [80] based on scale invariance of the 10-dimensional tree-level type IIB action. One family of the two original scaling symmetries is broken spontaneously by the vacuum expectation value of the dilaton, resulting in a massless Goldstone boson in 4 dimensions which can be identified with τ . Non-zero background fluxes can act as explicit symmetry breaking parameters (like non-zero quark masses in chiral perturbation theory), and can lift this flat direction. However, $W = 0$ is not enough to guarantee no explicit breaking, and so no flat direction, since also derivatives of W should vanish (see [81] for a study of flat directions in toroidal flux vacua in this context). Given a model, the statistical significance of any family of perturbatively flat vacua can be determined by the cut in the integration range of the flux variable Z^I (of [32]) put by the requirement of a low mass (at $|W_0|$ below a certain value). We hope to return to this question in the future.

Finally, let us summarise how our results can be crucial for addressing some important physics issues:

- We have provided explicit procedures to obtain all perturbatively flat vacua given an orientifolded Calabi Yau in IIB. As a proof of principle, all the vacua with $|W_0| < 10^{-2}$ have been enumerated for the $\mathbb{C}\mathbb{P}_{[1,1,1,6,9]}$ example. Complex structure moduli stabilisation is the first of many steps in the construction of de Sitter vacua. Thus detailed studies of understanding of complex structure moduli (along with the enumeration of all flux quanta that can lead to a low value of $|W_0|$) is a vital step in obtaining a rigorous understanding of the question of existence of de Sitter vacua in string theory. The methods developed in the present work are concrete advance in this direction.
- The question of statistical significance of perturbatively flat vacua has been examined and they have been found to be sparse in the ensemble of vacua at low $|W_0|$ as computed by Denef and Douglas. It has been found that perturbatively flat vacua can lead to very low values of the cosmological constant [23]. An anthropic solution to the cosmological constant problem via the landscape requires a large number of vacua with finely spaced values of the cosmological constant. In this light, understanding the statistics of perturbatively flat vacua is of utmost importance. Again, this paper takes concrete

1 steps in the direction.
2

3 Of course, addressing both questions will require the use of sophisticated numerical methods. For
4 explicit use of such methods in the context of string phenomenology see e.g [82–86]. They can be useful for
5 our future explorations.
6
7
8
9

10 **Acknowledgments**

11
12 We would like to thank Manki Kim, Sven Krippendorf, Liam McAllister, Ashoke Sen and Ravindranathan
13 Thangadurai for useful discussions. AM is supported in part by the SERB, DST, Government of India by
14 the grant MTR/2019/000267. The work of K. Sinha is supported in part by DOE Grant desc0009956.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Model	\vec{m}^T	\vec{k}^T	$ \epsilon _{\text{inf}}$
$M_{2,2}$	(-0.186, 0.982)	(-0.399, 0.917)	0.9751
$M_{2,3}$	(-0.966, 0.257)	(-0.746, 0.666)	0.8919
$M_{2,4}$	(-0.966, 0.257)	(-0.746, 0.666)	0.8919
$M_{2,5}$	(-0.966, 0.257)	(-0.746, 0.666)	0.8919
$M_{2,7}$	(-0.967, 0.253)	(-0.778, 0.628)	0.9114
$M_{2,13}$	(-0.143, 0.989)	(-0.297, 0.955)	0.9870
$M_{2,18}$	(-0.132, 0.991)	(-0.272, 0.962)	0.9897
$M_{2,19}$	(-0.186, 0.982)	(-0.399, 0.917)	0.9750
$M_{2,21}$	(0.966, -0.257)	(0.746, -0.666)	0.8919
$M_{2,22}$	(-0.896, 0.438)	(-0.695, 0.709)	0.9336
$M_{2,23}$	(-0.896, 0.438)	(-0.695, 0.709)	0.9336
$M_{2,24}$	(-0.896, 0.438)	(-0.695, 0.709)	0.9336
$M_{2,25}$	(-0.186, 0.982)	(-0.399, 0.917)	0.9752
$M_{2,26}$	(-0.969, 0.243)	(-0.816, 0.577)	0.9321
$M_{2,27}$	(-0.969, 0.247)	(-0.599, 0.801)	0.7784
$M_{2,28}$	(-0.969, 0.247)	(-0.599, 0.801)	0.7784
$M_{2,29}$	(-0.969, 0.247)	(-0.599, 0.801)	0.7784
$M_{2,35}$	(-0.993, 0.114)	(-0.972, 0.233)	0.9927
$M_{2,36}$	(-0.186, 0.982)	(-0.399, 0.917)	0.9752
$M_{2,37}$	(-0.186, 0.982)	(-0.399, 0.917)	0.9752
$M_{2,38}$	(-0.186, 0.982)	(-0.399, 0.917)	0.9752
$M_{2,39}$	(-0.970, 0.243)	(-0.577, 0.817)	0.7581

Table 3: 22 2-moduli examples from the Kreuzer-Skarke list in which $|\epsilon|_{\text{inf}}$ does not take values close to zero. For each model, we show the associated values of $|\epsilon|_{\text{inf}}$ and the values of \vec{m} and \vec{k} at which the infimum is attained.

A 2-moduli examples data

As discussed in Sec. 5, our numerical analysis has shown that in 22 of the 39 2-moduli examples in the Kreuzer-Skarke list, $|\epsilon|_{\text{inf}}$ does not take values close to zero. This by itself gives a strong bound on the region where all possible solutions to the Lemma are contained (without the need of imposing requirements such as validity of the effective field theory or phenomenological viability). We list these models in Tab. 3 together with the associated values of $|\epsilon|_{\text{inf}}$ and the values of \vec{m} and \vec{k} at which the infimum is attained.

References

- [1] S. B. Giddings, S. Kachru and J. Polchinski, “Hierarchies from fluxes in string compactifications,” Phys. Rev. D **66** (2002), 106006 doi:10.1103/PhysRevD.66.106006 [arXiv:hep-th/0105097 [hep-th]].
- [2] S. Kachru, R. Kallosh, A. D. Linde and S. P. Trivedi, “De Sitter vacua in string theory,” Phys. Rev. D **68** (2003), 046005 doi:10.1103/PhysRevD.68.046005 [arXiv:hep-th/0301240 [hep-th]].

- 1 [3] V. Balasubramanian, P. Berglund, J. P. Conlon and F. Quevedo, “Systematics of moduli stabilisation in Calabi-Yau flux compactifications,” *JHEP* **03** (2005), 007 doi:10.1088/1126-6708/2005/03/007 [arXiv:hep-th/0502058 [hep-th]].
- 2
3
4
5
6 [4] M. Berg, M. Haack and B. Kors, “On volume stabilization by quantum corrections,” *Phys. Rev. Lett.* **96** (2006), 021601 doi:10.1103/PhysRevLett.96.021601 [arXiv:hep-th/0508171 [hep-th]].
- 7
8
9 [5] A. Westphal, “de Sitter string vacua from Kahler uplifting,” *JHEP* **03** (2007), 102 doi:10.1088/1126-6708/2007/03/102 [arXiv:hep-th/0611332 [hep-th]].
- 10
11
12 [6] M. Cicoli, J. P. Conlon and F. Quevedo, “General Analysis of LARGE Volume Scenarios with String Loop Moduli Stabilisation,” *JHEP* **10** (2008), 105 doi:10.1088/1126-6708/2008/10/105 [arXiv:0805.1029 [hep-th]].
- 13
14
15
16 [7] M. Cicoli, A. Maharana, F. Quevedo and C. P. Burgess, “De Sitter String Vacua from Dilaton-dependent Non-perturbative Effects,” *JHEP* **06** (2012), 011 doi:10.1007/JHEP06(2012)011 [arXiv:1203.1750 [hep-th]].
- 17
18
19
20 [8] M. Cicoli, D. Ciupke, S. de Alwis and F. Muia, “ α' Inflation: moduli stabilisation and observable tensors from higher derivatives,” *JHEP* **09** (2016), 026 doi:10.1007/JHEP09(2016)026 [arXiv:1607.01395 [hep-th]].
- 21
22
23
24
25 [9] D. Gallego, M. C. D. Marsh, B. Vercnocke and T. Wrase, “A New Class of de Sitter Vacua in Type IIB Large Volume Compactifications,” *JHEP* **10** (2017), 193 doi:10.1007/JHEP10(2017)193 [arXiv:1707.01095 [hep-th]].
- 26
27
28
29
30 [10] I. Antoniadis, Y. Chen and G. K. Leontaris, “Perturbative moduli stabilisation in type IIB/F-theory framework,” *Eur. Phys. J. C* **78** (2018) no.9, 766 doi:10.1140/epjc/s10052-018-6248-4 [arXiv:1803.08941 [hep-th]].
- 31
32
33
34
35 [11] S. AbdusSalam, S. Abel, M. Cicoli, F. Quevedo and P. Shukla, “A systematic approach to Kähler moduli stabilisation,” *JHEP* **08** (2020) no.08, 047 doi:10.1007/JHEP08(2020)047 [arXiv:2005.11329 [hep-th]].
- 36
37
38
39
40 [12] A. Maharana and E. Palti, “Models of Particle Physics from Type IIB String Theory and F-theory: A Review,” *Int. J. Mod. Phys. A* **28** (2013), 1330005 doi:10.1142/S0217751X13300056 [arXiv:1212.0555 [hep-th]].
- 41
42
43
44
45 [13] M. Cicoli and F. Quevedo, “String moduli inflation: An overview,” *Class. Quant. Grav.* **28** (2011), 204001 doi:10.1088/0264-9381/28/20/204001 [arXiv:1108.2659 [hep-th]].
- 46
47
48
49 [14] D. Baumann and L. McAllister, “Inflation and String Theory,” doi:10.1017/CBO9781316105733 [arXiv:1404.2601 [hep-th]].
- 50
51
52 [15] R. Bousso and J. Polchinski, “Quantization of four form fluxes and dynamical neutralization of the cosmological constant,” *JHEP* **06** (2000), 006 doi:10.1088/1126-6708/2000/06/006 [arXiv:hep-th/0004134 [hep-th]].
- 53
54
55
56 [16] L. Susskind, “The Anthropic landscape of string theory,” [arXiv:hep-th/0302219 [hep-th]].
- 57
58
59
60

- [17] M. Demirtas, M. Kim, L. McAllister and J. Moritz, “Vacua with Small Flux Superpotential,” *Phys. Rev. Lett.* **124** (2020) no.21, 211603 doi:10.1103/PhysRevLett.124.211603 [arXiv:1912.10047 [hep-th]].
- [18] S. Gukov, C. Vafa and E. Witten, “CFT’s from Calabi-Yau four folds,” *Nucl. Phys. B* **584** (2000), 69-108 [erratum: *Nucl. Phys. B* **608** (2001), 477-478] doi:10.1016/S0550-3213(00)00373-4 [arXiv:hep-th/9906070 [hep-th]].
- [19] M. Cicoli, J. P. Conlon, A. Maharana and F. Quevedo, “A Note on the Magnitude of the Flux Superpotential,” *JHEP* **01** (2014), 027 doi:10.1007/JHEP01(2014)027 [arXiv:1310.6694 [hep-th]].
- [20] A. Giryavets, S. Kachru, P. K. Tripathy and S. P. Trivedi, “Flux compactifications on Calabi-Yau threefolds,” *JHEP* **04** (2004), 003 doi:10.1088/1126-6708/2004/04/003 [arXiv:hep-th/0312104 [hep-th]].
- [21] A. Cole, A. Schachner and G. Shiu, “Searching the Landscape of Flux Vacua with Genetic Algorithms,” *JHEP* **11** (2019), 045 doi:10.1007/JHEP11(2019)045 [arXiv:1907.10072 [hep-th]].
- [22] M. Demirtas, M. Kim, L. McAllister, J. Moritz and A. Rios-Tascon, “A Cosmological Constant That is Too Small,” [arXiv:2107.09065 [hep-th]].
- [23] M. Demirtas, M. Kim, L. McAllister, J. Moritz and A. Rios-Tascon, “Small Cosmological Constants in String Theory,” [arXiv:2107.09064 [hep-th]].
- [24] M. Demirtas, M. Kim, L. McAllister and J. Moritz, “Conifold Vacua with Small Flux Superpotential,” *Fortsch. Phys.* **68** (2020), 2000085 doi:10.1002/prop.202000085 [arXiv:2009.03312 [hep-th]].
- [25] R. Álvarez-García, R. Blumenhagen, M. Brinkmann and L. Schlechter, “Small Flux Superpotentials for Type IIB Flux Vacua Close to a Conifold,” doi:10.1002/prop.202000088 [arXiv:2009.03325 [hep-th]].
- [26] Y. Honma and H. Otsuka, “Small flux superpotential in F-theory compactifications,” *Phys. Rev. D* **103** (2021) no.12, 126022 doi:10.1103/PhysRevD.103.126022 [arXiv:2103.03003 [hep-th]].
- [27] M. Cicoli, I. G. Etxebarria, F. Quevedo, A. Schachner, P. Shukla and R. Valandro, “The Standard Model Quiver in de Sitter String Compactifications,” [arXiv:2106.11964 [hep-th]].
- [28] M. Cicoli, F. Quevedo and R. Valandro, “De Sitter from T-branes,” *JHEP* **03** (2016), 141 doi:10.1007/JHEP03(2016)141 [arXiv:1512.04558 [hep-th]].
- [29] M. Cicoli, I. García-Etxebarria, C. Mayrhofer, F. Quevedo, P. Shukla and R. Valandro, “Global Orientifolded Quivers with Inflation,” *JHEP* **11** (2017), 134 doi:10.1007/JHEP11(2017)134 [arXiv:1706.06128 [hep-th]].
- [30] M. R. Douglas, “The Statistics of string/M theory vacua,” *JHEP* **05** (2003), 046 doi:10.1088/1126-6708/2003/05/046 [arXiv:hep-th/0303194 [hep-th]].
- [31] S. Ashok and M. R. Douglas, “Counting flux vacua,” *JHEP* **01** (2004), 060 doi:10.1088/1126-6708/2004/01/060 [arXiv:hep-th/0307049 [hep-th]].
- [32] F. Denef and M. R. Douglas, “Distributions of flux vacua,” *JHEP* **05** (2004), 072 doi:10.1088/1126-6708/2004/05/072 [arXiv:hep-th/0404116 [hep-th]].

- 1 [33] F. Denef, M. R. Douglas and B. Florea, “Building a better racetrack,” JHEP **06** (2004), 034
2 doi:10.1088/1126-6708/2004/06/034 [arXiv:hep-th/0404257 [hep-th]].
3
4 [34] F. Denef and M. R. Douglas, “Distributions of nonsupersymmetric flux vacua,” JHEP **03** (2005), 061
5 doi:10.1088/1126-6708/2005/03/061 [arXiv:hep-th/0411183 [hep-th]].
6
7 [35] F. Marchesano, G. Shiu and L. T. Wang, “Model building and phenomenology of flux-
8 induced supersymmetry breaking on D3-branes,” Nucl. Phys. B **712** (2005), 20-58
9 doi:10.1016/j.nuclphysb.2005.01.046 [arXiv:hep-th/0411080 [hep-th]].
10
11 [36] K. R. Dienes, “Statistics on the heterotic landscape: Gauge groups and cosmological constants of four-
12 dimensional heterotic strings,” Phys. Rev. D **73** (2006), 106010 doi:10.1103/PhysRevD.73.106010
13 [arXiv:hep-th/0602286 [hep-th]].
14
15 [37] F. Gmeiner, R. Blumenhagen, G. Honecker, D. Lust and T. Weigand, “One in a billion: MSSM-like
16 D-brane statistics,” JHEP **01** (2006), 004 doi:10.1088/1126-6708/2006/01/004 [arXiv:hep-th/0510170
17 [hep-th]].
18
19 [38] M. R. Douglas and W. Taylor, “The Landscape of intersecting brane models,” JHEP **01** (2007), 031
20 doi:10.1088/1126-6708/2007/01/031 [arXiv:hep-th/0606109 [hep-th]].
21
22 [39] B. S. Acharya, F. Denef and R. Valandro, “Statistics of M theory vacua,” JHEP **06** (2005), 056
23 doi:10.1088/1126-6708/2005/06/056 [arXiv:hep-th/0502060 [hep-th]].
24
25 [40] A. Giryavets, S. Kachru and P. K. Tripathy, “On the taxonomy of flux vacua,” JHEP **08** (2004), 002
26 doi:10.1088/1126-6708/2004/08/002 [arXiv:hep-th/0404243 [hep-th]].
27
28 [41] A. Misra and A. Nanda, “Flux vacua statistics for two-parameter Calabi-Yau’s,” Fortsch. Phys. **53**
29 (2005), 246-259 doi:10.1002/prop.200410194 [arXiv:hep-th/0407252 [hep-th]].
30
31 [42] J. P. Conlon and F. Quevedo, “On the explicit construction and statistics of Calabi-Yau flux vacua,”
32 JHEP **10** (2004), 039 doi:10.1088/1126-6708/2004/10/039 [arXiv:hep-th/0409215 [hep-th]].
33
34 [43] O. DeWolfe, “Enhanced symmetries in multiparameter flux vacua,” JHEP **10** (2005), 066
35 doi:10.1088/1126-6708/2005/10/066 [arXiv:hep-th/0506245 [hep-th]].
36
37 [44] A. Hebecker and J. March-Russell, “The Ubiquitous throat,” Nucl. Phys. B **781** (2007), 99-111
38 doi:10.1016/j.nuclphysb.2007.05.003 [arXiv:hep-th/0607120 [hep-th]].
39
40 [45] D. Martinez-Pedrerera, D. Mehta, M. Rummel and A. Westphal, “Finding all flux vacua in an explicit
41 example,” JHEP **06** (2013), 110 doi:10.1007/JHEP06(2013)110 [arXiv:1212.4530 [hep-th]].
42
43 [46] J. Halverson and F. Ruehle, “Computational Complexity of Vacua and Near-Vacua in Field and String
44 Theory,” Phys. Rev. D **99** (2019) no.4, 046015 doi:10.1103/PhysRevD.99.046015 [arXiv:1809.08279
45 [hep-th]].
46
47 [47] P. Betzler and E. Plauschinn, “Type IIB flux vacua and tadpole cancellation,” Fortsch. Phys. **67** (2019)
48 no.11, 1900065 doi:10.1002/prop.201900065 [arXiv:1905.08823 [hep-th]].
49
50 [48] I. Bena, J. Blabäck, M. Graña and S. Lüst, “Algorithmically solving the Tadpole Problem,”
51 [arXiv:2103.03250 [hep-th]].
52
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- [49] S. Krippendorf, R. Kroepsch and M. Syvaeri, “Revealing systematics in phenomenologically viable flux vacua with reinforcement learning,” [arXiv:2107.04039 [hep-th]].
- [50] I. Broeckel, M. Cicoli, A. Maharana, K. Singh and K. Sinha, “Moduli Stabilisation and the Statistics of SUSY Breaking in the Landscape,” *JHEP* **10** (2020), 015 doi:10.1007/JHEP10(2020)015 [arXiv:2007.04327 [hep-th]].
- [51] I. Broeckel, M. Cicoli, A. Maharana, K. Singh and K. Sinha, “Moduli Stabilisation and the Statistics of Axion Physics in the Landscape,” [arXiv:2105.02889 [hep-th]].
- [52] L. Susskind, “Supersymmetry breaking in the anthropic landscape,” doi:10.1142/9789812775344_0040 [arXiv:hep-th/0405189 [hep-th]].
- [53] M. R. Douglas, “Statistical analysis of the supersymmetry breaking scale,” [arXiv:hep-th/0405279 [hep-th]].
- [54] M. Dine, E. Gorbatov and S. D. Thomas, “Low energy supersymmetry from the landscape,” *JHEP* **08** (2008), 098 doi:10.1088/1126-6708/2008/08/098 [arXiv:hep-th/0407043 [hep-th]].
- [55] N. Arkani-Hamed, S. Dimopoulos and S. Kachru, “Predictive landscapes and new physics at a TeV,” [arXiv:hep-th/0501082 [hep-th]].
- [56] R. Kallosh and A. D. Linde, “Landscape, the scale of SUSY breaking, and inflation,” *JHEP* **12** (2004), 004 doi:10.1088/1126-6708/2004/12/004 [arXiv:hep-th/0411011 [hep-th]].
- [57] M. Dine, D. O’Neil and Z. Sun, “Branches of the landscape,” *JHEP* **07** (2005), 014 doi:10.1088/1126-6708/2005/07/014 [arXiv:hep-th/0501214 [hep-th]].
- [58] I. Broeckel, M. Cicoli, A. Maharana, K. Singh and K. Sinha, “Work in progress”.
- [59] S. Hosono, A. Klemm and S. Theisen, “Lectures on mirror symmetry,” *Lect. Notes Phys.* **436** (1994), 235-280 doi:10.1007/3-540-58453-6_13 [arXiv:hep-th/9403096 [hep-th]].
- [60] A. Klemm, “Topological string theory on Calabi-Yau threefolds,” *PoS RTN2005* (2005), 002 doi:10.22323/1.019.0002
- [61] G. Curio, A. Klemm, D. Lust and S. Theisen, “On the vacuum structure of type II string compactifications on Calabi-Yau spaces with H fluxes,” *Nucl. Phys. B* **609** (2001), 3-45 doi:10.1016/S0550-3213(01)00285-1 [arXiv:hep-th/0012213 [hep-th]].
- [62] K. Dasgupta, G. Rajesh and S. Sethi, “M theory, orientifolds and G-flux,” *JHEP* **08** (1999), 023 doi:10.1088/1126-6708/1999/08/023 [arXiv:hep-th/9908088 [hep-th]].
- [63] T. D. Dimofte, “Type IIB Flux Vacua at Large Complex Structure,” *JHEP* **09** (2008), 064 doi:10.1088/1126-6708/2008/09/064 [arXiv:0806.0001 [hep-th]].
- [64] M. Cicoli, D. Klevers, S. Krippendorf, C. Mayrhofer, F. Quevedo and R. Valandro, “Explicit de Sitter Flux Vacua for Global String Models with Chiral Matter,” *JHEP* **05** (2014), 001 doi:10.1007/JHEP05(2014)001 [arXiv:1312.0014 [hep-th]].

- [65] M. C. D. Marsh and K. Sousa, “Universal Properties of Type IIB and F-theory Flux Compactifications at Large Complex Structure,” *JHEP* **03** (2016), 064 doi:10.1007/JHEP03(2016)064 [arXiv:1512.08549 [hep-th]].
- [66] Y. Honma and H. Otsuka, *Phys. Lett. B* **774** (2017), 225-228 doi:10.1016/j.physletb.2017.09.062 [arXiv:1706.09417 [hep-th]].
- [67] J. J. Blanco-Pillado, K. Sousa, M. A. Urkiola and J. M. Wachter, “Universal Class of Type-IIB Flux Vacua with Analytic Mass Spectrum,” *Phys. Rev. D* **103** (2021) no.10, 106006 doi:10.1103/PhysRevD.103.106006 [arXiv:2011.13953 [hep-th]].
- [68] F. Marchesano, D. Prieto and M. Wiesner, “F-theory flux vacua at large complex structure,” [arXiv:2105.09326 [hep-th]].
- [69] T. W. Grimm, C. Li and I. Valenzuela, *JHEP* **06** (2020), 009 [erratum: *JHEP* **01** (2021), 007] doi:10.1007/JHEP06(2020)009 [arXiv:1910.09549 [hep-th]].
- [70] A. Hebecker, P. Henkenjohann and L. T. Witkowski, “Flat Monodromies and a Moduli Space Size Conjecture,” *JHEP* **12** (2017), 033 doi:10.1007/JHEP12(2017)033 [arXiv:1708.06761 [hep-th]].
- [71] R. Gopakumar and C. Vafa, “M theory and topological strings. 1.,” [arXiv:hep-th/9809187 [hep-th]].
- [72] R. Gopakumar and C. Vafa, “M theory and topological strings. 2.,” [arXiv:hep-th/9812127 [hep-th]].
- [73] P. Candelas, A. Font, S. H. Katz and D. R. Morrison, “Mirror symmetry for two parameter models. 2.,” *Nucl. Phys. B* **429** (1994), 626-674 doi:10.1016/0550-3213(94)90155-4 [arXiv:hep-th/9403187 [hep-th]].
- [74] J. Louis, M. Rummel, R. Valandro and A. Westphal, “Building an explicit de Sitter,” *JHEP* **10** (2012), 163 doi:10.1007/JHEP10(2012)163 [arXiv:1208.3208 [hep-th]].
- [75] P. Candelas, X. De La Ossa, A. Font, S. H. Katz and D. R. Morrison, “Mirror symmetry for two parameter models. 1.,” *Nucl. Phys. B* **416** (1994), 481-538 doi:10.1016/0550-3213(94)90322-0 [arXiv:hep-th/9308083 [hep-th]].
- [76] S. Kachru, M. B. Schulz and S. Trivedi, “Moduli stabilization from fluxes in a simple IIB orientifold,” *JHEP* **10** (2003), 007 doi:10.1088/1126-6708/2003/10/007 [arXiv:hep-th/0201028 [hep-th]].
- [77] M. Kreuzer and H. Skarke, “Complete classification of reflexive polyhedra in four-dimensions,” *Adv. Theor. Math. Phys.* **4** (2002), 1209-1230 doi:10.4310/ATMP.2000.v4.n6.a2 [arXiv:hep-th/0002240 [hep-th]].
- [78] M. Cicoli, F. Quevedo, R. Savelli, A. Schachner and R. Valandro, “Systematics of the α' Expansion in F-Theory,” [arXiv:2106.04592 [hep-th]].
- [79] J. P. Conlon, *JHEP* **05** (2006), 078 doi:10.1088/1126-6708/2006/05/078 [arXiv:hep-th/0602233 [hep-th]].
- [80] C. P. Burgess, M. Cicoli, D. Ciupke, S. Krippendorff and F. Quevedo, “UV Shadows in EFTs: Accidental Symmetries, Robustness and No-Scale Supergravity,” *Fortsch. Phys.* **68** (2020) no.10, 2000076 doi:10.1002/prop.202000076 [arXiv:2006.06694 [hep-th]].

- 1 [81] M. Cicoli, M. Licheri, R. Mahanta, A. Maharana, “Work in progress”.
- 2
- 3 [82] L. B. Anderson, M. Gerdes, J. Gray, S. Krippendorf, N. Raghuram and F. Ruehle, “Moduli-
- 4 dependent Calabi-Yau and SU(3)-structure metrics from Machine Learning,” JHEP **05**, 013 (2021)
- 5 doi:10.1007/JHEP05(2021)013 [arXiv:2012.04656 [hep-th]].
- 6
- 7
- 8 [83] M. Bies, M. Cvetič, R. Donagi, L. Lin, M. Liu and F. Ruehle, “Machine Learning and Al-
- 9 gebraic Approaches towards Complete Matter Spectra in 4d F-theory,” JHEP **01**, 196 (2021)
- 10 doi:10.1007/JHEP01(2021)196 [arXiv:2007.00009 [hep-th]].
- 11
- 12
- 13 [84] W. Cui and J. Gray, “Numerical Metrics, Curvature Expansions and Calabi-Yau Manifolds,” JHEP **05**,
- 14 044 (2020) doi:10.1007/JHEP05(2020)044 [arXiv:1912.11068 [hep-th]].
- 15
- 16 [85] H. Otsuka and K. Takemoto, “Deep learning and k-means clustering in heterotic string vacua with line
- 17 bundles,” JHEP **05**, 047 (2020) doi:10.1007/JHEP05(2020)047 [arXiv:2003.11880 [hep-th]].
- 18
- 19 [86] F. Ruehle, Phys. Rept. **839** (2020), 1-117 doi:10.1016/j.physrep.2019.09.005
- 20
- 21 [87] F. Carta, A. Mininno and P. Shukla, [arXiv:2112.13863 [hep-th]].
- 22
- 23 [88] *Geometry of Numbers*, P. M. Gruber and C. G. Lekkerkerker, North-Holland, 1987.
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60