

University of Nebraska - Lincoln

## DigitalCommons@University of Nebraska - Lincoln

---

H. W. Manter Laboratory Library Materials

Parasitology, Harold W. Manter Laboratory of

---

2021

# Global Distribution, Host Range and Prevalence of Trypanosoma vivax: A Systematic Review and Meta-Analysis

Eyerusalem Fetene

*Addis Ababa University*

Samson Leta

*Addis Ababa University*

Fikru Regassa

*FDRE Ministry of Agriculture, Ethiopia*

Philippe Büscher

*Institute of Tropical Medicine, Belgium*

Follow this and additional works at: <https://digitalcommons.unl.edu/manterlibrary>



Part of the Parasitology Commons

---

Fetene, Eyerusalem; Leta, Samson; Regassa, Fikru; and Büscher, Philippe, "Global Distribution, Host Range and Prevalence of Trypanosoma vivax: A Systematic Review and Meta-Analysis" (2021). *H. W. Manter Laboratory Library Materials*. 102.

<https://digitalcommons.unl.edu/manterlibrary/102>

This Article is brought to you for free and open access by the Parasitology, Harold W. Manter Laboratory of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in H. W. Manter Laboratory Library Materials by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

RESEARCH

Open Access



# Global distribution, host range and prevalence of *Trypanosoma vivax*: a systematic review and meta-analysis

Eyerusalem Fetene<sup>1†</sup>, Samson Leta<sup>1\*†</sup>, Fikru Regassa<sup>1,2</sup> and Philippe Büscher<sup>3</sup>

## Abstract

**Background:** Trypanosomosis caused by *Trypanosoma vivax* is one of the diseases threatening the health and productivity of livestock in Africa and Latin America. *Trypanosoma vivax* is mainly transmitted by tsetse flies; however, the parasite has also acquired the ability to be transmitted mechanically by hematophagous dipterans. Understanding its distribution, host range and prevalence is a key step in local and global efforts to control the disease.

**Methods:** The study was conducted according to the methodological recommendations of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist. A systematic literature search was conducted on three search engines, namely PubMed, Scopus and CAB Direct, to identify all publications reporting natural infection of *T. vivax* across the world. All the three search engines were screened using the search term *Trypanosoma vivax* without time and language restrictions. Publications on *T. vivax* that met our inclusion criteria were considered for systematic review and meta-analysis.

**Result:** The study provides a global database of *T. vivax*, consisting of 899 records from 245 peer-reviewed articles in 41 countries. A total of 232,6277 tests were performed on 97 different mammalian hosts, including a wide range of wild animals. Natural infections of *T. vivax* were recorded in 39 different African and Latin American countries and 47 mammalian host species. All the 245 articles were included into the qualitative analysis, while information from 186 cross-sectional studies was used in the quantitative analysis mainly to estimate the pooled prevalence. Pooled prevalence estimates of *T. vivax* in domestic buffalo, cattle, dog, dromedary camel, equine, pig, small ruminant and wild animals were 30.6%, 6.4%, 2.6%, 8.4%, 3.7%, 5.5%, 3.8% and 12.9%, respectively. Stratified according to the diagnostic method, the highest pooled prevalences were found with serological techniques in domesticated buffalo (57.6%) followed by equine (50.0%) and wild animals (49.3%).

**Conclusion:** The study provides a comprehensive dataset on the geographical distribution and host range of *T. vivax* and demonstrates the potential of this parasite to invade other countries out of Africa and Latin America.

**Keywords:** *Trypanosoma vivax*, Global distribution, Host species, Meta-analysis, Pooled prevalence, Domestic animals, Wild fauna

\*Correspondence: samiwude@gmail.com

†Eyerusalem Fetene and Samson Leta equally contributed

<sup>1</sup>College of Veterinary Medicine and Agriculture, Addis Ababa University,

P.O. Box 34, Bishoftu, Ethiopia

Full list of author information is available at the end of the article



© The Author(s) 2021. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

## Introduction

Trypanosomes are protozoan parasites belonging to the family of *Trypanosomatidae* and the genus *Trypanosoma* (*T.*). The genus *Trypanosoma* comprises many species such as *T. brucei*, *T. congolense*, *T. equiperdum*, *T. evansi*, *T. simiae*, *T. suis* and *T. vivax*, which cause diseases called trypanosomoses in different mammalian hosts including humans [1]. Trypanosomoses are widely distributed in Africa, Latin America and Asia [2, 3].

*Trypanosoma vivax* is one of the most important *Trypanosoma* species known to infect both domestic and wild animals [4, 5]. *Trypanosoma vivax* is reported from cattle, dromedary camel, [6], goat, sheep, pig, dog [7], horse, donkey [8], both domesticated and wild buffalo, warthog, hippopotamus, reedbuck, waterbuck [9], antelope [10], giraffe [11], rhinoceros [12], rodents, pangolins, primates, reptiles and different wild ungulates and carnivores [13]. In Sub-Saharan Africa, *T. vivax* is mainly transmitted by tsetse flies (Diptera: *Glossinidae*) in which the parasite can multiply and remain infective throughout the insect's life [14]. The parasite has the ability to be transmitted mechanically by hematophagous flies such as *Tabanus* spp., *Stomoxys calcitrans* and *Haematobia irritans*, which are responsible for the spread of *T. vivax* in tsetse-free areas of Africa and in Latin America [4, 15–18].

*Trypanosoma vivax* infection can be suspected by clinical and/or serological evidence and can be confirmed by parasitological or molecular methods [19]. *Trypanosoma vivax* prevalence shows considerable variation with geography, abundance of tsetse or blood-sucking flies, and host species. In tsetse-infested areas of tropical Africa, the *T. vivax* prevalence is typically reported between 5–15% and often accounts for up to half of the total trypanosome prevalence. Outside of the tsetse belt, *T. vivax* prevalence is lower, between 2–10%, and it is related to local and seasonal variation in biting fly abundance [20].

Trypanosomosis caused by *T. vivax* is an important cause of economic losses related to morbidity, mortality, reproductive issues and decreased production [4]. For example, economic losses associated with bovine trypanosomosis have been estimated to be around US\$5 billion a year in Africa, and the continent spends at least \$30 million every year to control bovine trypanosomosis in terms of curative and prophylactic treatments [21]. Estimates outside Africa indicate that > 11 million head of cattle with a value of > US\$ 3 billion are at risk from *T. vivax* infection in the Brazilian Pantanal and Bolivian lowlands, with potential losses in excess of US\$ 160 million [16].

Many studies have been conducted on *T. vivax* over the past 100 years. Studies before the 1950s focused more on the morphology and taxonomy [22, 23], pathogenicity

[24] and treatment [25, 26]. However, since the 1950s, a considerable number of epidemiological studies have been conducted. Notwithstanding the excellent review on livestock trypanosomoses and their vectors in Latin America [18] and a recent general review on *T. vivax* [20], a systematic literature review on the global distribution, prevalence and host range of *T. vivax* is lacking. Moreover, no information on the global distribution of *T. vivax* is available at the World Animal Health Information System of the World Health Organization ([https://www.oie.int/wahis\\_2/public/wahid.php/Diseasedistributionmap](https://www.oie.int/wahis_2/public/wahid.php/Diseasedistributionmap)).

Thus, this study was conducted to provide the global distribution of *T. vivax* and to estimate the pooled prevalence of trypanosomosis caused by *T. vivax* in naturally infected domestic and wild animals.

## Methods

The systematic review and meta-analysis were conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist [27]. Screening and data extraction were performed by two authors (SL and EF) independently. All disagreements were discussed and resolved by consensus. A third author (PB) was also involved in the search for full-text papers to ensure that all relevant publications were included.

### Literature search

On 30 August 2019, a systematic literature search was conducted on three databases to identify all publications reporting natural infection of *T. vivax* across the world. PubMed, Scopus and CAB Direct were screened using the search term *Trypanosoma vivax* without time and language restrictions. All references found were imported into Mendeley Desktop reference manager software.

### Inclusion and exclusion criteria

To be considered, articles were required to meet the following inclusion criteria: (i) should be observational studies such as cross sectional, longitudinal, case report or outbreak investigation, published in indexed journals, reporting any natural infection of *T. vivax* using any diagnostic test or tests available; (ii) the study design, sample size, sample type, diagnostic methods and number of *T. vivax*-infected animals or prevalence, including 0%; (iii) species of animals with *T. vivax* infections must be provided. Experimental studies; publications which fail to describe diagnostic tools, study design and/or sample sources; and reports solely based on clinical signs were removed despite reporting the prevalence of the disease. In addition, studies reporting *T. vivax* from multiple species without stratifying the report at species level were removed.

## Data extraction

All relevant information such as author names, year of publication, study period, country, region, province, district, latitude, longitude (if provided or if they can be retrieved), host species, number of samples analyzed, type of samples collected, diagnostic method used, number of positives and prevalence or percentage were extracted to a pre-prepared Microsoft Excel spreadsheet (Microsoft Corp., Redmond, WA, USA). When publications only reported the number of animals tested and the prevalence, the numbers of positives were calculated. When publications only reported the number of animals tested and the number of positives, prevalence values were calculated. Publications in other languages than English were translated using Google Translate.

## Data analysis

Owing to heterogeneity within and between studies, random-effects meta-analysis was used to estimate the pooled prevalence and its 95% confidence interval (CI) in different hosts [28]. The estimation was carried out after categorization of the results according to the diagnostic tests used and the host species tested. Accordingly, diagnostic tests were categorized into three categories: (i) parasitological methods, including wet blood smear, stained blood smear and microhematocrit concentration; (ii) serological methods, including enzyme-linked immunosorbent assay (ELISA) both antigen and antibody based, indirect fluorescence antibody test (IFAT) and antigen detection LATEX agglutination; (iii) molecular methods, including reverse line blot hybridization assay, real-time and conventional polymerase chain reaction (PCR). Species-wise, sheep and goat were categorized into “small ruminants,” horse, donkey and mule into “equine” and all studied wild animals including Cape buffalo into “wild animals.” For cattle, domestic buffalo, dromedary camel, pig and dog, pooled prevalence was estimated without categorization.

Heterogeneity between studies was evaluated through the Cochran's Q test (reported as *p* value), and the inverse variance index ( $I^2$ ).  $I^2$  describes the percentage of observed total variation between studies due to heterogeneity rather than to random error (intra-study variation).  $I^2$  values < 25% correspond with low heterogeneity, up to 50% with moderate and up to 75% with high heterogeneity [29]. Sub-group analysis using the variable test method was performed to determine the potential sources of heterogeneity among studies. The across-study bias was examined by a funnel plot and Egger's regression asymmetry test. A funnel plot was used to visually examine the presence of publication bias, and Egger's regression asymmetry test was used to test whether the

bias was statistically significant or not [30]. The unbiased estimates were calculated using the Duval and Tweedie non-parametric ‘fill and trim’ linear random method [31].

The meta-analysis was done using ‘meta’ package of R statistical software version 3.6.2 (R Foundation for Statistical Computing). The map representing the global distribution of *T. vivax* was created, using Quantum GIS software version 3.4.5 (Open Source Geospatial Foundation, Boston, MA, USA).

## Results


### Literature search selection and data extraction

A total of 1691 publications were retrieved, 348 from PubMed, 1269 from Scopus and 74 from CAB Direct (Fig. 1). After removal of 390 duplicates, the remaining 1301 articles were screened based on their titles and abstracts. Reviews and articles reporting on laboratory and field experiments ( $n = 993$ ) were excluded of further analysis. Articles without an abstract or without sufficient information to make a decision were left for full text review. Of the remaining 308 articles, 11 of the full text files remained inaccessible [32–42]. Finally, one additional article, missed by the systematic literature search, was included manually. Full-text papers of 298 articles were retrieved online or via the library of the Institute of Tropical Medicine Antwerp and eligibility assessed according to the pre-established inclusion/exclusion criteria. Further 53 articles were excluded leaving 245 articles fulfilling all inclusion criteria for the qualitative analysis [4–10, 12–14, 43–277]. Among these 245 articles published between 1958 and 2019, 10 are case reports, 186 report on a cross-sectional study, 35 on a longitudinal study and 14 on an outbreak investigation. All relevant data from these articles were recorded, according to diagnostic method and host species, in a Microsoft Excel file, thus containing 899 records used in the meta-analysis (Additional file 1).

Of these 245 articles, 187 are conducted in 27 African countries, with Ethiopia taking the lead with 43 articles, followed by Nigeria with 29, Uganda with 21 and Kenya with 15 articles. In Latin America, 57 studies were conducted of which 32 were from Brazil, 9 from Venezuela and 6 from Colombia.

### Geographic distribution

All the studies conducted in the 27 African countries reported the presence of *T. vivax* in at least one host species; natural *T. vivax* infections were found in 12 of the 13 studied Latin American countries (Fig. 2 and Table 1). In Martinique, Alonso and co-workers did not find clinical or serological evidence of *T. vivax* in cattle on this island [50]. One article mentions a cross-sectional study on 300 equines in Pakistan, but all animals were negative in




molecular tests for *T. vivax* [231]. We could not find any other reports on the presence of *T. vivax* in Asia, Antarctica, Australia, Europe and North America.

#### Host range

A total of 232,627 tests were performed, and 24,420 of them were positive for natural infection of *T. vivax*. *Trypanosoma vivax* was reported from nine domestic animal species: cattle, domestic buffalo, dog, donkey, dromedary camel, goat, horse, pig and sheep. Among them, cattle were

the most studied species with 198,593 tests performed on cattle in 36 countries and two territories (192 publications) and 20,964 were positive for *T. vivax*. Next to cattle, goat, sheep, pig and donkey were the most frequently studied species. The protozoal parasite was also reported from wild animals including diverse species of antelopes, Cape buffalo, hippopotamus, black rhinoceros, pangolin and warthog. *Trypanosoma vivax* was reported from 39 wild fauna species, including many antelope species and Cape buffalo (Tables 2, 3).



**Fig. 2** The global distribution of *T. vivax* based on this systematic review including publications between 1958 and 2019

#### Pooled prevalence estimates according to host species and type of diagnostic test

Pooled prevalence estimates by test methods for different hosts are presented in Table 4, and forest plots of the meta-analysis and the subgroup analyses can be found in Additional files 2 and 3. Substantial heterogeneity was observed in the pooled estimate except for dog, which remained significant ( $P < 0.05$ ) even after sub-group analysis.

A total of 145 cross-sectional studies from 32 countries were included in estimation of natural infection of *T. vivax* in cattle. The random effect model indicates the pooled prevalence to be 6.4% (5.7–7.2, 95% CI). For small ruminants, pooled prevalence of *T. vivax* was estimated from 33 studies in 16 countries and found to be 3.8% (2.5–5.6, 95% CI). A total of 15 studies from 10 different countries were used to estimate the pooled prevalence of *T. vivax* in equines. The random effect model estimates the pooled prevalence to be 3.7% (2.0–6.8, 95% CI). Pooled prevalence of *T. vivax* in camels was estimated from four studies in three different countries. The model estimates a pooled prevalence of 8.4% (3.4–19.3, 95% CI). A total of 12 studies from 8 different countries were included in the estimation of pooled prevalence in pigs, which was found to be 5.5% (3.0–10.1, 95% CI). Five studies from five countries were used in the estimation pooled prevalence of *T. vivax* in dogs. The pooled prevalence was estimated

to be 2.6% (1.0–6.3% 95% CI). Three studies reported natural infection of *T. vivax* in domestic buffaloes from Venezuela, and the random effect model estimates a pooled prevalence of 30.6% (14.2–54.1, 95% CI). For wild animals, a pooled prevalence of 12.9% (9.9–16.6, 95% CI) was estimated from six studies in five countries. Subgroup pooled prevalences estimated according to the type of diagnostic test, as represented in Table 4, were lowest with parasitological techniques (from 1.1% in pigs to 13.2% in wild animals) and highest with serological techniques (from 13.8% in small ruminants to 57.6% in domestic buffalo).

#### Publication bias

The presence of publication bias was analyzed only in five species since there were not enough publications to discuss its possible influence in camel, domestic buffalo and dogs. Possible publication bias was demonstrated by visualization of asymmetry in funnel plots for cattle (Fig. 3a), small ruminants (Fig. 3b), equines (Fig. 3c), pigs (Fig. 3d) and wild animals (Fig. 3e). It was further confirmed by 'metabias' test (Egger's test) with  $p$ -value  $< 0.05$ . The 'trimfill' method imputed 170, 43, 30, 27 and 11 studies to obtain symmetry in funnel plots in cattle, wild animals, equines, small ruminants and pigs, respectively. The new estimated prevalence equals to 14.8% for cattle, 26.8% for wild animals, 21.6% for equines, 9.5% for small ruminants and 24.5% for pigs.

**Table 1** Countries with reported *T. vivax* infection in diverse host species tested with diverse methods

Country	Host species studied	Test method	Number of tests (total = 232,627)	Number of positives (total = 24,420)	References
Argentina	Cattle	Reverse line blot	186	16	[217]
Benin	Buffoon kob, cattle, hartebeest, roan antelope, warthog, waterbuck,	Thin and thick blood smears	312	205	[10, 99]
Bolivia	Cattle	Thin blood smear, Giemsa stained blood smear, PCR	1520	311	[123, 177, 243]
Botswana	Cape buffalo, cattle, donkey, goat, greater kudu, impala, lechwe, reedbuck, sable antelope, tsessebe	MH-CT/Woo test, Giemsa stained thick and thin blood smear, IFAT	3040	399	[100, 239]
Brazil	Cattle, donkey, goat, horse, pampas deer, sheep, domestic buffalo	MH-CT/Woo test, PCR, IFAT, thin and thick blood smear, buffy coat smear, Ab-ELISA, PCR	11468	4079	[4, 51, 58–62, 71, 72, 81, 83–85, 91, 94, 112, 116, 125–127, 154, 212, 219, 220, 226, 228, 242, 244, 250, 274, 277]
Burkina Faso	Cattle	Buffy coat smear, Ag-ELISA, Ab-ELISA, PCR	11095	1095	[65, 90, 151, 216, 241, 251–253]
Cameroon	African civet, black legged mongoose, black striped duiker, blue duiker, bosman potto, brush tailed porcupine, cattle, cloaked mangabey, crested mangabey, crocodile, dark mongoose, de Brazza's monkey, dog, dwarf guenon, giant forest squirrel, giant rat, goat, golden cat, golden potto, greater cane rat, greater white-nosed monkey, guereza white colobus, long-tailed pangolin, mandrill, mona monkey, monitor lizard, mustached monkey, ogilby's duiker, Peter's duiker, pig, red-legged sun squirrel, royal antelope, sheep, sitatunga, small-spotted genet, tree dassie, tree pangolin, two-spotted palm civet, water chevrotain, white-eye lid mangabey, yellow-backed duiker	Buffy coat smear, PCR	4176	406	[7, 13, 14, 170, 202, 245] [7, 13, 14, 170, 202, 245]
Chad	Cattle	Buffy coat smear, Ab-ELISA	1866	435	[93]
Colombia	Cattle, goat, sheep	Blood smear, PCR, IFAT	6712	1699	[135, 136, 215, 221, 232, 274]
Costa Rica	Cattle	Blood smear, IFAT	642	53	[210, 274]
Côte d'Ivoire	Cattle, goat, pig, sheep	MH-CT/Woo test, PCR	2185	195	[45, 148, 197]
Democratic Republic of the Congo	Cattle, dog, goat, pig, sheep	MH-CT/Woo test, ELISA	685	41	[167, 168]
Ecuador	Cattle	IFAT	310	70	[274]
El Salvador	Cattle	IFAT	100	15	[274]
Equatorial Guinea	Goat, sheep	PCR	559	10	[79]

**Table 1 (continued)**

Country	Host species studied	Test method	Number of tests (total = 232,627)	Number of positives (total = 24,420)	References
Ethiopia	Cattle, donkey, dromedary camel, goat, horse, mule, sheep	Giemsa stained blood smear, blood smear, buffy coat smear, MHCT/Woo test, thin and thick blood smear, Ab-ELISA, PCR Ag-ELISA	55196	2600	[6, 43, 44, 48, 64, 69, 70, 74–76, 86, 87, 92, 98, 101, 103, 107, 113–115, 122, 142, 144, 145, 152, 178–182, 184, 187, 191, 229, 230, 240, 248, 257, 258, 262, 264–266] [95]
French Guiana	Cattle	Buffy coat smear, Ag-ELISA, PCR	3000	870	[80, 157, 268]
Gabon	Cattle	Buffy coat smear, Giemsa stained blood smear, Ab-ELISA, PCR	442	26	[8, 96, 102, 173, 213, 218]
Gambia	Cattle, donkey, goat, horse, sheep	Buffy coat smear, PCR, Ag-latex agglutination test	5745	1329	[46, 117, 143, 198]
Ghana	Cattle, goat, pig, sheep	MHCT/Woo test, IFAT	1786	231	[55, 272]
Guyana	Goat, sheep	Thin and thick blood smear, Giemsa stained blood smear, buffy coat smear, MHCT/Woo test, Ag-ELISA, PCR	467	15	[66, 146, 172, 174, 183, 185, 196, 200, 201, 208, 209, 227, 261, 267, 275]
Kenya	Black rhinoceros, cattle, dromedary camel, goat, horse, pig, sheep	Thin and thick blood smear, Giemsa stained blood smear, buffy coat smear, MHCT/Woo test, Ag-ELISA, PCR	5156	845	[46, 117, 143, 198]
Liberia	Cattle	Giemsa stained blood smear, IFAT, Ab-ELISA	700	327	[155, 176]
Malawi	Cattle	Giemsa stained blood smear	9309	9	[271]
Mali	Cattle	Buffy coat smear	796	34	[192]
Martinique	Cattle	IFAT	227	0	[50]
Mozambique	Cattle	Blood smear	16895	1245	[254]
Namibia	Cattle	Giemsa stained thick and thin blood smear, MHCT	1481	15	[270]
Nigeria	Cattle, dog, goat, horse, sheep	Blood smear, Giemsa stained blood smear, MHCT/buffy coat smear, MHCT/Woo test, Ag-ELISA, PCR	20080	2926	[52, 53, 63, 88, 89, 104–106, 108–110, 134, 137–140, 153, 166, 206, 207, 211, 214, 234–237, 259, 269, 276]
Pakistan	Donkey, horse, mule	PCR	300	0	[231]
Paraguay	Cattle	IFAT	15	6	[274]
Peru	Cattle	Blood smear, MHCT/Woo test, Giemsa stained blood smear	985	119	[171, 177, 222, 274]
Rwanda	Cattle	Blood smear	3630	36	[12]
Senegal	Cattle, dog, donkey, goat, horse, sheep	Buffy coat smear, blood smears, Ab-ELISA, PCR	4890	365	[111, 128, 129, 225, 238]
South Africa	Cattle	PCR	143	30	[169]
Sudan	Cattle, donkey, dromedary camel, horse	Blood smear, Buffy coat smear, PCR	4426	366	[132, 186, 223, 233]

**Table 1 (continued)**

Country	Host species studied	Test method	Number of tests (total = 232,627)	Number of positives (total = 24,420)	References
Tanzania	African civet, bohor reedbuck, cattle, Coke's hartebeest, giraffe, Grant's gazelle, hunting dog, impala, Kirk's dikdik, klipspringer, Lichtenstein's hartebeest, oribi, oryx, ostrich, pig, roan antelope, southern reedbuck, steinbuck, Thomson's gazelle, tsessebe, warthog, wildebeest, zebra	Blood smear, thin and thick blood smear, Giemsa stained blood smear, buffy coat smear, PCR, PCR-LAMP	9974	431	[5, 56, 78, 130, 131, 133, 141, 147, 149, 188, 194, 203, 247, 256]
Togo	Cattle, dog, donkey, goat, pig, sheep	PCR-RFLP	354	27	[263]
Uganda	Cattle, horse, sheep, domestic buffalo	Giemsa stained blood smear, MHCT/Woo test, thick and thin blood smear, buffy coat smear, Ab-ELISA, PCR	28510	1932	[47, 49, 54, 57, 67, 68, 77, 82, 158–165, 189, 190, 195, 205, 273]
Venezuela	Cattle, horse, sheep, domestic buffalo	MHCT/Woo test, stained blood smear, IFAT, Ab-ELISA, PCR	6328	1373	[73, 118–121, 124, 224, 255, 260]
Zambia	African civet, baboon, bat, black rhinoceros, bushbuck, cane rat, Cape buffalo, cattle, crocodile, eland, elephant, genet, giraffe, goat, greater kudu, grey duiker, grysbok, hare, hartebeest, hippopotamus, hunting dog, hyena, impala, jackal, leopard, lion, mongoose, pig, porcupine, puku, reedbuck, roan antelope, serval, vervet monkey, warthog, waterbuck, wild cat, wildebeest, zebra	PCR, buffy coat smear	6936	234	[9, 97, 150, 175, 193, 199, 204, 246]

Ab-ELisa antibody enzyme-linked immunosorbent assay, Ag-ELisa antigen enzyme-linked immunosorbent assay, MHCT micro-hematocrit centrifugation technique, Ifat immuno-fluorescence antibody test, Pcr polymerase chain reaction, Pcr-Lamp polymerase chain reaction-loop mediated isothermal amplification, Pcr-Rflp polymerase chain reaction-restriction fragment length polymorphism

**Table 2** Domestic animal species tested for infection with *T. vivax*

Species	List of countries	Number of tests	Positive animals	References
Cattle	Argentina, Benin, Bolivia, Botswana, Brazil, Burkina Faso, Cameroon, Chad, Colombia, Costa Rica, Côte d'Ivoire, Democratic Republic of the Congo, Ecuador, El Salvador, Ethiopia, French Guiana, Gabon, Gambia, Ghana, Kenya, Liberia, Malawi, Mali, Martinique <sup>a</sup> , Mozambique, Namibia, Nigeria, Paraguay, Peru, Rwanda, Senegal, South Africa, Sudan, Tanzania, Togo, Uganda, Venezuela, Zambia	198593	20964	[4–6, 12, 14, 40, 44–54, 57–59, 61, 62, 64–66, 68–71, 73–78, 80–83, 85–88, 90–95, 99, 101–103, 105, 106, 108, 109, 111, 113, 114, 116, 118, 122, 123, 125–127, 129–132, 134–136, 138, 139, 141–143, 145, 147–153, 155–159, 161–167, 169–171, 173–177, 180, 182, 184, 185, 187–200, 202, 203, 205–212, 214–217, 219, 220, 222–227, 229, 230, 232, 235, 236, 238–243, 246–259, 261–271, 273, 274, 276–282]
Dromedary camel	Ethiopia, Kenya, Sudan	161	133	[6, 115, 172, 186]
Dog	Cameroon, Democratic Republic the Congo, Nigeria, Senegal, Uganda	574	1	[7, 137, 158, 168, 214, 225]
Donkey	Botswana, Brazil, Ethiopia, Pakistan <sup>a</sup> , Sudan, Uganda	2713	152	[6, 8, 43, 96, 107, 178, 189, 225, 228, 231, 233, 239, 248]
Goat	Botswana, Brazil, Cameroon, Colombia, Côte d'Ivoire, Democratic Republic of the Congo, Equatorial Guinea, Ethiopia, Gambia, Ghana, Guyana, Kenya, Nigeria, Senegal, Uganda, Zambia	9715	526	[6, 7, 55, 57, 60, 63, 68, 79, 98, 112, 128, 137, 140, 143, 144, 150, 158, 168, 179, 193, 197, 201, 204, 211, 213, 218, 221, 225, 234, 237, 239, 246, 248]
Horse	Brazil, Ethiopia, Gambia, Kenya, Nigeria, Pakistan <sup>a</sup> , Senegal, Sudan, Venezuela	3305	857	[8, 84, 96, 104, 118, 146, 181, 214, 225, 231, 233]
Mule	Ethiopia, Pakistan <sup>a</sup>	353	0	[43, 181, 231, 248]
Pig	Cameroon, Côte d'Ivoire, Democratic Republic the Congo, Ghana, Kenya, Tanzania, Uganda, Zambia	2650	233	[7, 57, 67, 68, 133, 158, 168, 197, 198, 201, 245, 246]
Sheep	Brazil, Cameroon, Colombia, Côte d'Ivoire, Democratic Republic of the Congo, Equatorial Guinea, Ethiopia, Gambia, Ghana, Guyana, Kenya, Nigeria, Senegal, Uganda, Venezuela	6447	455	[6, 7, 55, 57, 60, 79, 98, 116, 118, 120, 128, 134, 137, 140, 143, 144, 168, 197, 201, 213, 221, 225, 234, 248, 272]
Small ruminants	Kenya, Nigeria	988	69	[89, 110, 227]
Domestic buffalo	Brazil, Venezuela	2144	509	[116, 118, 119, 121, 260]

<sup>a</sup> *T. vivax* was not observed in Martinique and Pakistan

**Table 3** Wild animal species tested positive for *T. vivax* infection

Host species	Scientific name	Country	Number of tests	Positive tests	Positivity rate	References
Black rhinoceros	<i>Diceros bicornis</i>	Kenya	1	1	100	[283]
Black striped duiker	<i>Cephalophus dorsalis</i>	Cameroon	37	3	8.1	[13]
Blue duiker	<i>Cephalophus monticola</i>	Cameroon	290	24	8.3	[13]
Bosman potto	<i>Perodicticus potto</i>	Cameroon	8	3	37.5	[13]
Brush tailed porcupine	<i>Atherurus africanus</i>	Cameroon	106	7	6.6	[13]
Buffoon kob	<i>Kobus kob</i>	Benin	50	1	2	[10]
Bushbuck	<i>Tragelaphus scriptus</i>	Zambia	51	4	7.8	[97]
Cape buffalo	<i>Synacerus caffer</i>	Botswana, Zambia	1105	285	25.8	[9, 97, 100]
Cloaked mangabey	<i>Cercocebus albigena</i>	Cameroon	12	2	16.7	[13]
Crocodile	<i>Crocodylus niloticus</i>	Cameroon	3	1	33.3	[13]
De Brazza's Monkey	<i>Cercopithecus neglectus</i>	Cameroon	1	1	100	[13]
Dwarf guenon	<i>Miopithecus tlapoin</i>	Cameroon	55	5	9.1	[13]
Eland	<i>Taurotragus oryx</i>	Zambia	3	1	33.3	[97]
Giant rat	<i>Cricetomys gambianus</i>	Cameroon	135	4	2.9	[13]
Greater kudu	<i>Tragelaphus strepsiceros</i>	Botswana, Zambia	36	26	72.2	[97, 100]
Greater white-nosed monkey	<i>Cercopithecus nictitans</i>	Cameroon	155	22	14.2	[13]
Grey duiker	<i>Sylvicapra grimmia</i>	Zambia	7	1	14.3	[97]
Guereza white colobus	<i>Colobus guereza</i>	Cameroon	14	2	14.3	[13]
Hartebeest	<i>Alcelaphus buselaphus</i>	Benin	20	1	5	[10]
Hippopotamus	<i>Hippopotamus amphibius</i>	Zambia	29	1	3.4	[9]
Impala	<i>Aepyceros melampus</i>	Botswana	23	14	60.9	[100]
Lechwe	<i>Kobus leche</i>	Botswana	110	39	35.5	[100]
Long tailed pangolin	<i>Manis tetradactyla</i>	Cameroon	34	2	5.9	[13]
Mona monkey	<i>Cercopithecus mona</i>	Cameroon	46	8	17.4	[13]
Monitor lizard	<i>Varanus ornatus</i>	Cameroon	8	1	12.5	[13]
Moustached monkey	<i>Cercopithecus cephus</i>	Cameroon	101	11	10.9	[13]
Oryx	<i>Oryx beisa</i>	Tanzania	1	1	100	[56]
Puku	<i>Kobus vardonii</i>	Zambia	24	1	4.2	[97]
Reedbuck	<i>Redunca sp.</i>	Botswana, Zambia	3	3	100	[9, 100]
Sable antelope	<i>Hippotragus niger</i>	Botswana	22	7	31.8	[100]
Sitatunga	<i>Tragelaphus spekii</i>	Cameroon	5	1	20	[13]
Small-spotted genet	<i>Genetta servalina</i>	Cameroon	8	1	12.5	[13]
Southern reedbuck	<i>Redunca arundinum</i>	Tanzania	4	1	25	[56]
Tree pangolin	<i>Manis tricuspidis</i>	Cameroon	20	5	25	[13]
Tsessebe	<i>Damaliscus lunatus</i>	Botswana	15	6	40	[100]
Two-spotted palm civet	<i>Nandinia binotata</i>	Cameroon	32	3	9.4	[13]
Warthog	<i>Phacochoerus aethiopicus</i>	Zambia	56	1	1.8	[9]
Waterbuck	<i>Kobus ellipsiprymnus</i>	Zambia	30	19	63.3	[9, 97]
White-eyelid mangabey	<i>Cercocebus torquatus</i>	Cameroon	5	2	40	[13]

## Discussion

This study presents the first systematic review of published literature since the 1950s describing global distribution, host range and prevalence of trypanosomosis caused by *T. vivax*. Not surprisingly, most publications report on *T. vivax* infections in domestic mammalian species, in particular in cattle and small ruminants, while few publications describe natural infections in wildlife.

Looking at the *T. vivax* distribution map (Fig. 2), there is an evident data gap for some sub-Saharan African countries where tsetse flies are present and therefore *T. vivax* may be endemic. Although our formal search strategy could not retrieve any publication on these "missing" countries, conventional Google search confirms the presence of *T. vivax* in South Sudan and Zimbabwe [284, 285], and Genevieve et al. [286] reported on the presence

**Table 4** Sub-group meta-analysis for different species using different diagnostic methods


Host species	Diagnostic method	Number of publications	Number of tests	Number of positives	Pooled prevalence in %	95% CI
Domestic buffalo	Parasitological	1	316	36	11.4	8.3–15.4
	Molecular	2	609	127	20.9	17.4–25.0
	Serological	2	556	301	57.6	22.5–86.4
Camel	Molecular	4	1611	133	8.4	3.4–19.3
Cattle	Parasitological	92	102910	5414	4.6	4.0–5.3
	Molecular	51	31549	3140	7.4	6.2–8.7
	Serological	23	16469	4495	34.6	28.0–41.9
Dog	Parasitological	3	257	0	3.4	1.1–9.6
	Molecular	2	189	1	1.2	0.2–8.5
Domestic buffalo	Parasitological	1	316	36	11.4	8.3–15.4
	Molecular	2	609	127	20.9	17.4–25.0
	Serological	2	556	301	57.6	22.5–86.4
Equine	Parasitological	8	2471	20	1.5	0.9–2.6
	Molecular	7	1425	251	5.6	2.7–11.3
	Serological	1	6	3	50	16.8–83.2
Pig	Parasitological	4	799	3	1.1	0.4–2.9
	Molecular	10	1851	230	9	4.9–15.9
Small ruminant	Parasitological	19	8990	220	2.3	1.5–3.6
	Molecular	15	4045	327	5	2.7–9.3
	Serological	3	408	43	13.8	6.1–28.4
Wild animal	Parasitological	3	1093	75	11.8	7.1–16.9
	Molecular	3	1618	121	10.7	8.6–13.3
	Serological	1	748	318	49.3	37.5–61.2

of potential vectors in the Central African Republic. Since Angola, the Central African Republic and the Republic of Congo are endemic for human African trypanosomiasis, the presence of *T. vivax* in these countries is likely [287]. Due to its adaptation to mechanical transmission, *T. vivax* is also present outside the tsetse belt in Africa, e.g. in Ethiopia and Sudan [114, 288]. As a consequence, the trypanosomiasis control efforts with focus on tsetse eradication might have little effect on *T. vivax*. Also, economic impact assessments that are solely based on tsetse distribution alone could seriously underestimate the problem of trypanosomiasis because of *T. vivax*.

Out of Africa, *T. vivax* is present in Latin America but not in North America, Australia, Asia and the Pacific regions. *Trypanosoma vivax* is believed to be introduced into Latin America in cattle and horses imported from Africa, possibly in the sixteenth century, and spread to different Latin American countries including Brazil, Colombia, French Guiana, Guadeloupe, Guyana, Martinique, Panama, Suriname and Venezuela [18]. Stephen [289] reviewed the presence of the parasite in Costa Rica, Ecuador, El Salvador, Paraguay and Peru, and according to Gardiner et al. [15], *T. vivax* was present in the Caribbean thus posing a threat to the livestock industries.

From our literature search, we can only confirm *T. vivax* to be endemic in 12 Latin American countries of which 7 (Argentina, Bolivia, Brazil, Colombia, Guyana, Peru, Venezuela) are also endemic for *T. evansi* [290]. Although, our literature search provides information on the potential spread of *T. vivax* in Latin America, it is important to note that the distribution could be much wider, for example, *T. vivax* was only detected in Argentina in 2018; this is this due to the lack of previous studies. Apparently, *T. vivax* has never spread into Asia, unlike *T. evansi*, although similar to the latter; it can be mechanically transmitted by bloodsucking flies. Unless there is a particular biological or environmental factor preventing *T. vivax* from invading the Middle East and Asia, as well Northern Africa, North America and Europe, we must remain alert about the risk of importing *T. vivax* into non-endemic countries as happened in Latin America.

This review suggests that *T. vivax* has a very diverse host range, including 9 domestic mammals and almost 40 wild fauna species. Regarding the latter, however, data should be interpreted with caution. Diagnostic tests, whether parasitological, serological or even molecular, have their limitations. For examples, by sequencing of PCR amplicons, Auty and co-workers [11] clearly



**Fig. 3** Publication bias evidenced by funnel plots for cattle (a), small ruminants (b), equines (c), pigs (d) and wild animals (e)

demonstrate that wildlife may harbor a diversity of trypanosomes, including taxonomically undefined species. Therefore, it is likely that many reports on *T. vivax* infection in wildlife and tsetse in fact deal with other trypanosome species that are not necessarily pathogenic for domestic animals.

The pooled prevalence of trypanosomosis in different hosts varies significantly depending on the detection methods; significantly higher estimates were reported in publications using serological techniques. Higher estimates using a serological technique could be due to the persistence of the antibody over several months after curative treatment and the possibility of low undetectable

parasitemia in parasitological techniques [20, 93, 291, 292]. Moti et al. [187] compared the percentage positivity obtained with different diagnostic techniques and showed that relative to the microhematocrit centrifugation technique the percent positivity increased by 50 and 250% when using PCR-RFLP. Also, Garcia et al. [118] reported that for the detection of trypanosomes, PCR-based assays are twice as sensitive as parasitological techniques such as the microhaematocrit centrifugation.

The study has the following limitations. The literature search was almost exclusively based on electronic databases whereby some older literature must have been

missed. The data showed a large degree of heterogeneity among studies, which remain significant after subgroup analysis. There is a significant publication bias, which could be due to incomplete or inaccurate information provided in the publications. In addition, studies were conducted between 1956 and 2017, and the result may not accurately reflect the current epidemiological situation and therefore could limit interpretation of the result to some degree. Furthermore, we suspect numerous data gaps mainly because of two reasons. First, due to the lack of a country-level monitoring and reporting system for trypanosomosis, most of the data included in this analysis are from research activities. Second, trypanosomosis diagnosis in most endemic countries relied to a great extent on low-sensitivity parasitological methods, while more sensitive molecular tools are rarely used. In addition, the majority of studies analyzing trypanosome's presence in the field may not have a sampling strategy that allows a robust estimation of prevalence. This is for multiple and understandable reasons—samples can be difficult and expensive to collect, and many studies rely on purposive sampling, or sampling of, for example, animals presented to veterinary clinics. While these kinds of studies provide a rough idea of pathogen presence/absence, they may not provide an accurate estimate of prevalence. Thus, caution should be taken when interpreting the results presented here.

## Conclusion

With this study, we intended to provide comprehensive information on the geographical distribution, host range and prevalence of trypanosomosis caused by *T. vivax* worldwide. The results confirm the wide geographical distribution and a diverse host range of *T. vivax*. The parasite parasitizes almost all domestic mammals and many wild animal species, thus suggesting the potential to get established in other countries with favorable environmental conditions, e.g. in the Middle East, Asia and Australia. The meta-analysis showed a high degree of variability in estimated prevalence values. The variability can be attributed to diagnostic tests used and the species of the animal infected.

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13071-021-04584-x>.

**Additional file 1.** Global *Trypanosoma vivax* occurrence records.

**Additional file 2.** Forest plots showing an overview of studies reporting *Trypanosoma vivax* in different host species.

**Additional file 3.** Forest plots showing an overview of studies reporting *Trypanosoma vivax* grouped by test methods in different host species.

## Acknowledgements

We extend our gratitude to staff members of the library of the Institute of Tropical Medicine Antwerp for providing some articles.

## Authors' contributions

PB and FR conceived the research and SL designed the research; EF and SL drafted the manuscript. EF, SL and PB extracted and compiled the data. All authors read and approved the final manuscript.

## Funding

No funding was available for this study

## Availability of data and materials

All data analyzed in this paper are provided as supplementary file.

## Ethics approval and consent to participate

Not applicable.

## Consent for publication

Not applicable.

## Competing interests

The authors declare that they have no competing interests.

## Author details

<sup>1</sup> College of Veterinary Medicine and Agriculture, Addis Ababa University, P.O. Box 34, Bishoftu, Ethiopia. <sup>2</sup> FDRE Ministry of Agriculture, P.O.Box 62347/3735, Addis Ababa, Ethiopia. <sup>3</sup> Institute of Tropical Medicine, Department of Biomedical Sciences, Nationalestraat 155, 2000 Antwerp, Belgium.

Received: 14 October 2020 Accepted: 6 January 2021

Published online: 25 January 2021

## References

1. Stevens J, Brisse S. Systematics of trypanosomes of medical and veterinary importance. In: Maudlin I, Holmes P, Miles M., editors. Trypanos. CABI Publishing; 2004. p. 1–19.
2. Luckins AG, Dwingler RH. Non-tsetse-transmitted animal trypanosomiasis. In: Maudlin I, Holmes PH, Miles MA, editors. Trypanos. CABI Publishing; 2004. p. 269–79.
3. Cecchi G, Paone M, Feldmann U, Vreyen MJB, Diall O, Mattioli RC. Assembling a geospatial database of tsetse-transmitted animal trypanosomiasis for Africa. Parasit Vectors. 2014;7:1–10.
4. Vieira OLE, de Macedo LO, Santos MAB, Silva JABA, de Mendonca CL, da Faustino MAG, et al. Detection and molecular characterization of *Trypanosoma (Duttonella) vivax* in dairy cattle in the state of Sergipe, northeastern Brazil. Rev Bras Parasitol Vet Brazil. 2017;26:516–20.
5. Kassian EN, Simunza MC, Silayo RS, Moonga L, Ndebele J, Sugimoto C, et al. Prevalence and risk factors of bovine trypanosomosis in Kilwa district, Lindi region of southern Tanzania. Vet Parasitol Reg Stud Reports. 2017;9:1–5.
6. Birhanu H, Fikru R, Said M, Kidane W, Gebrehiwot T, Hagos A, et al. Epidemiology of *Trypanosoma evansi* and *Trypanosoma vivax* in domestic animals from selected districts of Tigray and Afar regions, Northern Ethiopia. Parasit Vectors. 2015;8:212.
7. Nimpaye H, Njokou F, Njine T, Njitchouang GR, Cuny G, Herder S, et al. *Trypanosoma vivax*, *T. congolense* "forest type" and *T. simiae*: prevalence in domestic animals of sleeping sickness foci of Cameroon. Parasite. 2011;18:171–9.
8. Pinchbeck GL, Morrison LJ, Tait A, Langford J, Meehan L, Jallow S, et al. Trypanosomosis in the gambia: prevalence in working horses and donkeys detected by whole genome amplification and PCR, and evidence for interactions between trypanosome species. BMC Vet Res. 2008;4:7.
9. Anderson NE, Mubanga J, Fevre EM, Picozzi K, Eisler MC, Thomas R, et al. Characterisation of the wildlife reservoir community for human and animal trypanosomiasis in the Luangwa Valley, Zambia. PLoS Negl Trop Dis. 2011;5:e1211.
10. Guedegbe B, Verhulst A, Van Meirvenne N, Pandey VS, Doko A. Serological evidence of the existence of a wild reservoir of *Trypanosoma brucei*

- gambiense* in the Pendjari biosphere reservation in the Republic of Benin. Ann Soc Belg Med Trop. 1992;72:113–20.
11. Auty H, Anderson NE, Picozzi K, Lembo T, Mubanga J, Hoare R, et al. Trypanosome diversity in wildlife species from the Serengeti and Luangwa Valley Ecosystems. PLoS Negl Trop Dis. 2012;6:e1828.
  12. Mihok S, Otieno LH, Tarimo CS. Trypanosome infection rates in tsetse flies (Diptera: Glossinidae) and cattle during tsetse control operations in the Kagera River region of Rwanda. Bull Entomol Res. 1992;82:361–7.
  13. Njiokou F, Simo G, Nkinin SW, Laveisseire C, Herder S. Infection rate of *Trypanosoma brucei* s.l., *T. vivax*, *T. congolense* "forest type", and *T. simiae* in small wild vertebrates in south Cameroon. Acta Trop. 2004;92:139–46.
  14. Suh PF, Njiokou F, Mamoudou A, Ahmadou TM, Mouhaman A, Garabed R. Bovine trypanosomiasis in tsetse-free pastoral zone of the far-North region, Cameroon. J Vector Borne Dis. 2017;54:263–9.
  15. Gardiner PR, Pearson TW, Clarke MW, Mutharia LM. Identification and isolation of a variant surface glycoprotein from *Trypanosoma vivax*. Science. 1987;235:774–7.
  16. Jones TW, Dávila AMR. *Trypanosoma vivax*-Out of Africa. Trends Parasitol. 2001;17:99–101.
  17. Davila AM, Silva RA. Animal trypanosomiasis in South America: Current status, partnership, and information technology. Ann NY Acad Sci. 2000;916:199–212.
  18. Desquesnes M. Livestock trypanosomoses and their vectors in Latin America CIRAD-EMVT publication. Paris: OIE; 2004.
  19. Osório ALAR, Madruga CR, Desquesnes M, Soares CO, Ribeiro LRR, Da Costa SCG, et al. *Trypanosoma (Duttonella) vivax*: Its biology, epidemiology, pathogenesis, and introduction in the New World—a review. Mem Inst Oswaldo Cruz. 2008;103:1–13.
  20. Dagnachew S, Bezie M. Review on *Trypanosoma vivax*. African J Basic Appl Sci. 2015;7:41–64.
  21. Angara TE, Ismail A, Ibrahim A. An overview on the economic impacts of animal trypanosomiasis. Glob J Res Anal. 2012;3:275–6.
  22. Hoare CA, Broom JC. Morphological and taxonomic studies on mammalian trypanosomes. Trans R Soc Trop Med Hyg. 1938;31:517–34.
  23. Hoare CA, Broom JC. Morphological and taxonomic studies on mammalian trypanosomes. vii.—differentiation of *Trypanosoma uniforme* and *T. vivax* in mixed infections. Trans R Soc Trop Med Hyg. 1939;32:629–32.
  24. Hornby HE. The pathogenicity to cattle of *Trypanosoma vivax*. Vet Rec. 1946;58:178.
  25. Schwetz J. Treatment of *Trypanosoma vivax* infection. Trans R Soc Trop Med Hyg. 1929;23:109.
  26. Fiennes RNTW. The treatment experiments with *Trypanosoma vivax* (Ziemann) disease of cattle. Vet Rec. 1948;60:302.
  27. Moher D, Liberati A, Tetzlaff J, Altman D, Group TP. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA Statement. PLoS Med. 2009;6:e1000097.
  28. Hedges L, Vevea J. Fixed- and random-effects models in meta-analysis. Psychol Methods. 1998;3:486–504.
  29. Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. Stat Med. 2002;21:1539–58.
  30. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple graphical test. BMJ. 1997;315:629–34.
  31. Duval S, Tweedie R. Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. Biometrics. 2000;56:455–63.
  32. Baptista LCF, de Fernandes ACC, da Silva TIB, de Souza ACM, Sandes HMM, Alves LC, et al. *Trypanosoma vivax* infection in dairy cattle raised in the State of Pernambuco: case report. Veterinária e Zootec. 2011;18:919–22.
  33. Bolívar AM, Rojas A, Rosales D, Torres Y, García-Lugo P. Detection of hemotropic agents in a livestock farm using PCR and DGGE. Rev Salud Anim. 2014;36:53–7.
  34. Carvalho AU, Abrão DC, Facury Filho EJ, Paes PRO, Ribeiro MFB. Occurrence of *Trypanosoma vivax* in Minas Gerais state, Brazil. Arq Bras Med Vet e Zootec. 2008;60:769–71.
  35. Fidelis Junio OL, Cadioli FA, de Barnabé PA, Machado RZ, Teixeira MMG, Marques LC. Trypanosomiasis in dairy cattle in São Paulo state. Veterinária e Zootec. 2011;18:879–82.
  36. Gonzales JL, Jones TW, Picozzi K, Cuellar HR. Evaluation of a polymerase chain reaction assay for the diagnosis of bovine trypanosomiasis and epidemiological surveillance in Bolivia. Kinetoplastid Biol Dis. 2003;2:1–4.
  37. Kalu AU, Uzoukwu M, Ikeme M. Prevalence of tsetse fly and ruminant trypanosomiasis in Katsina-Ala Local Government Area, Nigeria. Roum Arch Microbiol Immunol. 1996;55:341–52.
  38. Kamani J, Sannusi A, Egwu OK, Dogo GI, Tanko TJ, Kemza S, et al. Prevalence and significance of haemoparasitic infections of cattle in North-Central, Nigeria. Vet World. 2010;3:445–8.
  39. Katunguka-Rwakishaya E. The prevalence of trypanosomiasis in small ruminants and pigs in a sleeping sickness endemic area of Buikwe County, Mukono district, Uganda. Rev Elev Med Vet Pays Trop. 1996;49:56–8.
  40. Magona JW, Mayende JSP. Occurrence of concurrent trypanosomiasis, theileriosis, anaplasmosis and helminthosis in Friesian, Zebu and Sahi-wal cattle in Uganda. Onderstepoort J Vet Res. 2002;69:133–40.
  41. Plagemann O. Epidemiological survey of trypanosomiasis in cattle in Northern Uganda. Berl Munch Tierarztl Wochenschr. 1974;87:352–5.
  42. Suárez C, García F, Román D, Coronado A, Perrone T, Reyna A, et al. Factores de riesgo asociados a la tripanosomosis bovina en explotaciones ganaderas de Venezuela. Zootec Trop. 2010;27:363–72.
  43. Abebe R, Wolde A. A cross-sectional study of trypanosomiasis and its vectors in donkeys and mules in Northwest Ethiopia. Parasitol Res. 2010;106:911–6.
  44. Abebe R, Gute S, Simon I. Bovine trypanosomiasis and vector density in Omo-Ghibe tsetse belt, South Ethiopia. Acta Trop. 2017;167:79–85.
  45. Acapovi-Yao G, Cisse B, Koumba CRZ, Mavoungou JF. Trypanosome infections in cattle in farms of different departments in Côte d'Ivoire. Rev Med Vet. 2016;167:289–95.
  46. Adam Y, Marcotty T, Cecchi G, Mahama CI, Solano P, Bengaly Z, et al. Bovine trypanosomiasis in the Upper West Region of Ghana: Entomological, parasitological and serological cross-sectional surveys. Res Vet Sci. 2012;92:462–8.
  47. Ahmed HA, Picozzi K, Welburn SC, MacLeod ET. A comparative evaluation of PCR-based methods for species-specific determination of African animal trypanosomes in Ugandan cattle. Parasit Vectors. 2013;6:316.
  48. Ali D, Bitew M. Epidemiological study of bovine trypanosomiasis in Mao-Komo special district, Benishangul Gumuz regional state, Western Ethiopia. Glob Vet. 2011;6:402–8.
  49. Alingu RA, Muhanguzi D, MacLeod E, Waiswa C, Fyfe J. Bovine trypanosome species prevalence and farmers' trypanosomiasis control methods in south-western Uganda. J S Afr Vet Assoc. 2014;85:1–5.
  50. Alonso M, Camus E, Rodriguez Diego J, Bertaudiére L, Tatareau JC, Liabeuf JM. Current status of bovine haemoparasitic diseases in Martinique (French West Indies). Rev Elev Med Vet Pays Trop. 1992;45:9–14.
  51. Alves WP, Cuglovici DA, Furtado LFV, Da Silveira JAG, Facury-Filho EJ, Ribeiro MFB, et al. Comparison of three methods for diagnosis of *Trypanosoma (Duttonella) vivax* in cattle. Genet Mol Res. 2017;16:52.
  52. Anene BM, Chime AB, Jibike GI, Anika SM. Prevalence of trypanosomiasis in Zebu cattle at Obudu ranch—a tsetse-free zone in Nigeria. Prev Vet Med. 1991;10:257–60.
  53. Anene BM, Chime AB, Jibike GI, Anika SM. Comparative study of clinical signs, haematology and prevalence of trypanosomiasis in Holstein Friesian and White Fulani Zebu cattle exposed to natural infection in a rain forest zone of Nigeria. Angew Parasitol. 1991;32:99–104.
  54. Angwech H, Nyeko JHPP, Opiyo EA, Okello-Onen J, Opiro R, Echodu R, et al. Heterogeneity in the prevalence and intensity of bovine trypanosomiasis in the districts of Amuru and Nwoya, Northern Uganda. BMC Vet Res. 2015;11:1–8.
  55. Applewhite LM. Small ruminant trypanosomiasis in Guyana – a preliminary report. Br Vet J. 1990;146:93–4.
  56. Ashcroft MT. An attempt to isolate *Trypanosoma rhodesiense* from wild animals. Trans R Soc Trop Med Hyg. 1958;52:276–82.
  57. Balyeidhusa ASP, Kironde FAS, Enyaru JCK. Apparent lack of a domestic animal reservoir in Gambiense sleeping sickness in northwest Uganda. Vet Parasitol. 2012;187:157–67.
  58. Bastos TSA, Faria AM, de Madrid DMCM, de Bessa LC, Linhares GFC, Fidelis OLJ, et al. First outbreak and subsequent cases of *Trypanosoma vivax* in the state of Goiás, Brazil. Rev Bras Parasitol Vet. 2017;26:366–71.
  59. Batista JS, Riet-Correa F, Teixeira MMG, Madruga CR, Simões SDV, Maia TF. Trypanosomiasis by *Trypanosoma vivax* in cattle in the Brazilian

- semiarid: Description of an outbreak and lesions in the nervous system. *Vet Parasitol.* 2007;143:174–81.
60. Batista JS, Oliveira AF, Rodrigues CMF, Damasceno CAR, Oliveira IRS, Alves HM, et al. Infection by *Trypanosoma vivax* in goats and sheep in the Brazilian semiarid region: from acute disease outbreak to chronic cryptic infection. *Vet Parasitol.* 2009;165:131–5.
  61. Batista JS, Rodrigues CMF, Olinda RG, Silva TMF, Vale RG, Camara ACL, et al. Highly debilitating natural *Trypanosoma vivax* infections in Brazilian calves: epidemiology, pathology, and probable transplacental transmission. *Parasitol Res.* 2012;110:73–80.
  62. Batista JS, Freitas CIA, da Silva JB, Cavalcante TV, de Paiva KAR, Lopes FC, et al. Clinical evaluation and reproductive indices of dairy cows naturally infected with *Trypanosoma vivax*. *Semin Ciências Agrárias.* 2017;38:3031–8.
  63. Behnke JM, Chiejina SN, Musongong GA, Nnadi PA, Ngongeh LA, Abonyi FO, et al. Resistance and resilience of traditionally managed West African Dwarf goats from the savanna zone of northern Nigeria to naturally acquired trypanosome and gastrointestinal nematode infections. *J Helminthol.* 2011;85:80–91.
  64. Bejano S, Kifle T, Bireda W. Study on the prevalence of bovine trypanosomosis in Assosa district of the Benishangul Gumuz region, West Ethiopia. *Livest Res Rural Dev.* 2016; 28.
  65. Bengaly Z, Kanwe AB, Duvallet G. Evaluation of an antigen detection-ELISA test for the diagnosis of trypanosomiasis in naturally infected cattle. *Trop Med Parasitol.* 1995;46:284–6.
  66. Bett B, Orente C, Irungu P, Munga LK. Epidemiological factors that influence time-to-treatment of trypanosomosis in Orma Boran cattle raised at Galana Ranch, Kenya. *Vet Parasitol.* 2004;120:43–53.
  67. Biryomumaisho S, Melville SE, Atunguka-Rwakishaya E, Lubega GW. Detection of natural *Trypanosoma vivax* infections in pigs with microhaematocrit centrifugation and amplification of ITS1 rDNA. *Onderstepoort J Vet Res.* 2009;76:285–9.
  68. Biryomumaisho S, Rwakishaya E-K, Melville SE, Cailleau A, Lubega GW. Livestock trypanosomosis in Uganda: Parasite heterogeneity and anaemia status of naturally infected cattle, goats and pigs. *Parasitol Res.* 2013;112:1443–50.
  69. Bishaw Y, Temesgen W, Yideg N, Alemu S. Prevalence of bovine trypanosomosis in Wemberma district of West Gojam zone North West Ethiopia. *Ethiop Vet J.* 2012;16:41–8.
  70. Bitew M, Amedie Y, Abebe A, Tolosa T. Prevalence of bovine trypanosomosis in selected areas of Jabi Tehenan district, West Gojam of Amhara regional state, Northwestern Ethiopia. *African J Agric Res.* 2011;6:140–4.
  71. Bittar JFF, Bassi PB, Moura DM, Garcia GC, Martins-Filho OA, Vasconcelos AB, et al. Evaluation of parameters related to libido and semen quality in Zebu bulls naturally infected with *Trypanosoma vivax*. *BMC Vet Res.* 2015;11:261.
  72. Cadioli FA, de Barnabé PA, Machado RZ, Teixeira MCA, André MR, Sam-pão PH, et al. First report of *Trypanosoma vivax* outbreak in dairy cattle in São Paulo state, Brazil. *Rev Bras Parasitol Veterinária.* 2012;21:118–24.
  73. Camejo MI, Aso PM, Gonzatti MI, Pérez-Rojas Y. Relationship between asymptomatic infections with *Anaplasma marginale*, *Babesia* spp. and *Trypanosoma vivax* in bulls and testosterone levels. *Rev Cient la Fac Ciencias Vet la Univ del Zulia.* 2016;26:13–9.
  74. Chanie M, Arega C, Bogale B. Hematopathology and hematological parametric alterations in indigenous cattle due to trypanosomosis. *Glob Vet.* 2012;9:546–51.
  75. Cherenet T, Sani RA, Panandam JM, Nadzr S, Speybroeck N, van den Bossche P. Seasonal prevalence of bovine trypanosomosis in a tsetse-infested zone and a tsetse-free zone of the Amhara Region, north-west Ethiopia. *Onderstepoort J Vet Res.* 2004;71:307–12.
  76. Cherenet T, Sani RA, Speybroeck N, Panandam JM, Nadzr S, Van den Bossche P. A comparative longitudinal study of bovine trypanosomiasis in tsetse-free and tsetse-infested zones of the Amhara Region, north-west Ethiopia. *Vet Parasitol.* 2006;140:251–8.
  77. Clausens PH, Wiemann A, Patzelt R, Kakaire D, Poetzsch C, Peregrine A, et al. Use of a PCR assay for the specific and sensitive detection of *Trypanosoma* spp. in naturally infected dairy cattle in peri-urban Kampala, Uganda. *Ann NY Acad Sci.* 1998;849:21–31.
  78. Connor RJ, Halliwell RW. Bovine trypanosomiasis in Southern Tanzania: parasitological and serological survey of prevalence. *Trop Anim Health Prod.* 1987;19:165–72.
  79. Cordon-Obras C, Berzosa P, Ndong-Mabale N, Bobukasi L, Buatiche JN, Ndongo-Asumu P, et al. *Trypanosoma brucei gambiense* in domestic livestock of Kogo and Mbini foci (Equatorial Guinea). *Trop Med Int Heal.* 2009;14:535–41.
  80. Cossic BGA, Adjahoutonon B, Gloaguen P, Dibanganga GL, Maganga G, Leroy P, et al. Trypanosomiasis challenge estimation using the diminazene aceturate (Berenil) index in Zebu in Gabon. *Trop Anim Health Prod.* 2017;49:619–24.
  81. de Costa VMM, Ribeiro MFB, Duarte ALL, Mangueira JM, Pessoa AFA, Azevedo SS, et al. Seroprevalence and risk factors for cattle anaplasmosis, babesiosis, and trypanosomiasis in a Brazilian semiarid region. *Rev Bras Parasitol Vet.* 2013;22:207–13.
  82. Cox AP, Tosas O, Tilley A, Picozzi K, Coleman P, Hide G, et al. Constraints to estimating the prevalence of trypanosome infections in East African zebu cattle. *Parasites Vectors.* 2010;3:82.
  83. Cuglovici DA, Bartholomeu DC, Reis-Cunha JL, Carvalho AU, Ribeiro MFB. Epidemiologic aspects of an outbreak of *Trypanosoma vivax* in a dairy cattle herd in Minas Gerais state, Brazil. *Vet Parasitol.* 2010;169:320–6.
  84. Da SAS, Perez HAG, Costa MM, Da Silva AS, Garcia Perez HA, Costa MM, et al. Horses naturally infected by *Trypanosoma vivax* in southern Brazil. *Parasitol Res.* 2011;108:23–30.
  85. da Silva AS, Costa MM, Polenz MF, Polenz CH, Teixeira MMG, Lopes STA, et al. First report of *Trypanosoma vivax* in bovines in the State of Rio Grande do Sul, Brazil. *Cienc Rural.* 2009;39:2550–4.
  86. Dagnachew S, Girma H, Abebe G. A cross-sectional study on bovine trypanosomiasis in Jawi district of Amhara Region, Northwest Ethiopia. *Ethiop Vet J.* 2011;15:69–78.
  87. Dagnachew S, Tsegaye B, Awukew A, Tilahun M, Ashenafi H, Rowan T, et al. Prevalence of bovine trypanosomosis and assessment of trypanocidal drug resistance in tsetse infested and non-tsetse infested areas of Northwest Ethiopia. *Parasite Epidemiol Control.* 2017;2:40–9.
  88. Danbirni SO, Okaiyeto SO, Kudi AC, Pewan SB. Bovine trypanosomosis and tuberculosis in a nomadic herd in Sabon gari local government area of Kaduna state, Nigeria. *J Anim Vet Adv.* 2010;9:1285–8.
  89. Daniel AD, Joshua RA, Kalejaiye JO, Dada AJ. Prevalence of trypanosomiasis in sheep and goats in a region of northern Nigeria. *Rev Elev Med Vet Pays Trop.* 1994;47:295–7.
  90. Dayo G-KK, Bengaly Z, Messad S, Bucheton B, Sidibe I, Cene B, et al. Prevalence and incidence of bovine trypanosomosis in an agro-pastoral area of southwestern Burkina Faso. *Res Vet Sci.* 2010;88:470–7.
  91. De Araujo MS, Barros ACE, Costa FB, De Carvalho Neta AV, De Candanedo GRDMN, Abreu-Silva AL. Bovine trypanosomiasis an emerging disease in Maranhão State-Brazil. *Vector-Borne Zoonotic Dis.* 2011;11:853–6.
  92. Degneh E, Shibeshi W, Terefe G, Asres K, Ashenafi H. Bovine trypanosomosis: changes in parasitemia and packed cell volume in dry and wet seasons at Gidami district, Oromia Regional State, western Ethiopia. *Acta Vet Scand.* 2017;59:59.
  93. Delafosse A, Thebaud E, Desquesnes M, Michaux Y. Epidemiology of *Trypanosoma vivax* infection in cattle in the tse-tse free area of Lake Chad. *Prev Vet Med.* 2006;74:108–19.
  94. de Melo BJ, Blanco YAC, Bruhn FRP, Guimaraes AM. Seroprevalence of *Trypanosoma vivax*, anaplasma marginale, and babesia bovis in dairy cattle. *Cienc Anim Bras.* 2016;17:564–73.
  95. Desquesnes M, Gardiner PR. Epidemiology of bovine trypanosomiasis (*Trypanosoma vivax*) in French Guiana. *Rev Elev Med Vet Pays Trop.* 1993;46:463–70.
  96. Dhollander S, Jallow A, Mbodge K, Kora S, Sanneh M, Gaye M, et al. Equine trypanosomosis in the Central River Division of The Gambia: a study of veterinary gate-clinic consultation records. *Prev Vet Med.* 2006;75:152–62.
  97. Dillmann JSS, Townsend AJ. A trypanosomiasis survey of wild animals in the Luangwa Valley Zambia. *Acta Trop.* 1979;36:349.
  98. Dinka H, Abebe G. Small ruminants trypanosomosis in the southwest of Ethiopia. *Small Rumin Res.* 2005;57:239–43.
  99. Dokko A, Guedegbe B, Baelmans R, Demey F, N'Diaye A, Pandey VS, et al. Trypanosomiasis in different breeds of cattle from Benin. *Vet Parasitol.* 1991;40:1–7.
  100. Drager N, Mehlitz D. Investigations on the prevalence of trypanosome carriers and the antibody response in wildlife in Northern Botswana. *Tropenmed Parasitol.* 1978;29:223–33.

101. Duguma R, Tasew S, Olani A, Damena D, Alemu D, Mulatu T, et al. Spatial distribution of *Glossina* sp. and *Trypanosoma* sp. in south-western Ethiopia. *Parasites Vectors.* 2015;8:430.
102. Dwinger RH, Agyemang K, Kaufmann J, Grieve AS, Bah ML. Effects of trypanosome and helminth infections on health and production parameters of village N'Dama cattle in the Cambia. *Vet Parasitol.* 1994;54:353–65.
103. Efrem DB, Yacob HT, Hagos AT, Basu AK. Bovine trypanosomosis in Gimbi district of Western Oromia, Ethiopia. *Anim Biol.* 2010;60:123–31.
104. Ehizibolo DO, Kamani J, Ehizibolo PO, Egwu KO, Dogo GI, Salami-Shinaba JO. Prevalence and significance of parasites of horses in some states of northern Nigeria. *J Equine Sci.* 2012;23:1–4.
105. Enwezor FNC, Samdi SM, Ijabor O, Abenga JN. The prevalence of bovine trypanosomes in parts of Benue state, north-central Nigeria. *J Vector Borne Dis.* 2012;49:188–90.
106. Enwezor FNC, Bello B, Kalgo A, Zaria LT. Surveillance and management of trypanosomiasis in cattle herds in Kaura area, Kaduna State Nigeria. *Livest Rear Farm Pract Dis.* 2011;11:145–62.
107. Eyob A, Mekuria S, Regassa A, Abebe R. A cross-sectional study of equine trypanosomosis and its vectors in Wolayta zone, Southern Ethiopia. *J Vet Med Anim Heal.* 2011;3:21–6.
108. Ezeani MC, Okoro H, Anosa VO, Onyenekwe CC, Meludu SC, Dioka CE, et al. Immunodiagnosis of bovine trypanosomiasis in Anambra and Imo states, Nigeria, using enzyme-linked immunosorbent assay: Zoonotic implications to human health. *J Vector Borne Dis.* 2008;45:292–300.
109. Fajinmi AO, Faleke OO, Magaji AA, Daneji AI, Gweba M, Fajinmi AO, Faleke OO, Magaji AA, Daneji AI, Gweba M. Presence of *Trypanosome* species and determination of anaemia in trade cattle at Sokoto Abattoir, Nigeria. *Res J Parasitol.* 2011;6:31–42.
110. Fakae BB, Chiejinai SN. The prevalence of concurrent trypanosome and gastrointestinal nematode infections in West African Dwarf sheep and goats in Nsukka area of eastern Nigeria. *Vet Parasitol.* 1993;49:313–8.
111. Fall A, Diack A, Diaite A, Seye M, D'leteren GDM, Diaité A, et al. Tsetse challenge, trypanosome and helminth infection in relation to productivity of village Ndama cattle in Senegal. *Vet Parasitol.* 1999;81:235–47.
112. Fávero JF, Da Silva AS, Biazus AH, Volpato A. *Trypanosoma vivax* infection in goat in west of Santa Catarina state, Brazil. *Comp Clin Path.* 2016;25:497–9.
113. Fentahun T, Tekeba M, Mitiku T, Chanie M. Prevalence of bovine trypanosomosis and distribution of vectors in Hawa Gelan district, Oromia region, Ethiopia. *Glob Vet.* 2012;9:297–302.
114. Regassa F, Goddeeris BM, Delespauv X, Moti Y, Tadesse A, Bekana M, et al. Widespread occurrence of *Trypanosoma vivax* in bovines of tsetse-as well as non-tsetse-infested regions of Ethiopia: a reason for concern? *Vet Parasitol.* 2012;190:355–61.
115. Regassa F, Andualem Y, Getachew T, Menten J, Hasker E, Merga B, et al. Trypanosome infection in dromedary camels in eastern Ethiopia: Prevalence, relative performance of diagnostic tools and host related risk factors. *Vet Parasitol.* 2015;211:175–81.
116. Galiza GJN, Garcia HA, Assis ACO, Oliveira DM, Pimentel LA, Dantas AFM, et al. High mortality and lesions of the central nervous system in trypanosomosis by *Trypanosoma vivax* in Brazilian hair sheep. *Vet Parasitol.* 2011;182:359–63.
117. Ganyo EY, Boampong JN, Masiga DK, Villinger J, Turkson PK. Haematology of N'Dama and West African Short Horn cattle herds under natural *Trypanosoma vivax* challenge in Ghana. *F1000Research.* 2018;7:314.
118. Garcia H, Garcia M-E, Perez H, Mendoza-Leon A. The detection and PCR-based characterization of the parasites causing trypanosomiasis in water-buffalo herds in Venezuela. *Ann Trop Med Parasitol.* 2005;99:359–70.
119. García H, García M-EE, Pérez G, Bethencourt A, Zerpa É, Pérez H, et al. Trypanosomiasis in Venezuelan water buffaloes: Association of packed-cell volumes with seroprevalence and current trypanosome infection. *Ann Trop Med Parasitol.* 2006;100:297–305.
120. García H, Rangel-Rivas A, Contreras I, García M-E, García F, Perrone T. Molecular characterization of *Trypanosoma vivax* in naturally-infected sheep from two farms at San Fernando and Biruaca Counties, Apure State, Venezuela. *Rev Cient la Fac Ciencias Vet la Univ del Zulia.* 2009;19:230–7.
121. Garcia HA, Ramirez OJ, Rodrigues CMF, Sanchez RG, Bethencourt AM, Del M, Perez G, et al. *Trypanosoma vivax* in water buffalo of the Venezuelan Llanos: an unusual outbreak of wasting disease in an endemic area of typically asymptomatic infections. *Vet Parasitol.* 2016;230:49–55.
122. Girmay G, Arega B, Tesfaye D, Berkvens D, Muleta G, Asefa G. Community-based tsetse fly control significantly reduces fly density and trypanosomiasis prevalence in Metekel Zone, Northwest, Ethiopia. *Trop Anim Health Prod.* 2016;48:633–42.
123. Gonzales JL, Chacon E, Miranda M, Loza A, Siles LM. Bovine trypanosomosis in the Bolivian Pantanal. *Vet Parasitol.* 2007;146:9–16.
124. González JR, Meléndez RD. Seroprevalence of bovine trypanosomiasis and anaplasmosis by ELISA at Juan Jose Mora county, Carabobo State, Venezuela. *Rev Cient la Fac Ciencias Vet la Univ del Zulia.* 2007;17:449–55.
125. Guedes Junior DS, Araújo FR, Silva FJMM, Rangel CP, Barbosa Neto JD, Fonseca AH. Frequency of antibodies to Babesia bigemina, B bovis, Anaplasma marginale, *Trypanosoma vivax* and *Borrelia burgdorferi* in cattle from the Northeastern region of the State of Pará, Brazil. *Rev Bras Parasitol Vet.* 2008;17:105–9.
126. Guerra RDMSNDc, Feitosa AB, Santos HP, Abreu-Silva AL, Santos ACG, Dos Feitosa AB, et al. Biometry of *Trypanosoma vivax* found in a calf in the state of Maranhão Brazil. *Cienc Rural.* 2008;38:833–5.
127. Guerra NR, Monteiro MFM, Sandes HMM, Da Cruz NLN, Ramos CAN, De Assis Santana VL, et al. Detection of IgG antibodies against *Trypanosoma vivax* in cattle by indirect immunofluorescence test. *Pesqui Vet Bras.* 2013;33:1423–6.
128. Gueye A, Mbengue M, Diouf A. Ticks and hemoparasitoses of livestock in Senegal. III. The Northern Sudan area. *Rev Elev Med Vet Pays Trop.* 1989;42:411–20.
129. Gueye A, Mbengue M, Diouf A, Sonko ML. Ticks and hemoparasitoses in livestock in Senegal. V. The northern Guinea area. *Rev Elev Med Vet Pays Trop.* 1993;46:551–61.
130. Haji IJ, Malele I, Namangala B. Occurrence of haemoparasites in cattle in Monduli district, northern Tanzania. *Onderstepoort J Vet Res.* 2014;81:1–4.
131. Haji IJ, Sugimoto C, Kajino K, Malele I, Simukoko H, Chitambo H, et al. Determination of the prevalence of trypanosome species in cattle from Monduli district, northern Tanzania, by loop mediated isothermal amplification. *Trop Anim Health Prod.* 2015;47:1139–43.
132. Hall MJR, Kheir SM, Rahman AHA, Noga S. Tsetse and trypanosomiasis survey of southern darfur province, Sudan - I Bovine trypanosomiasis. *Trop Anim Health Prod.* 1983;15:191–206.
133. Hamill LC, Kaare MT, Welburn SC, Picozzi K. Domestic pigs as potential reservoirs of human and animal trypanosomiasis in Northern Tanzania. *Parasites Vectors.* 2013;6:1–7.
134. Idehen CO, Ishola OO, Adeyemi IG, Abongaby G, Olaleye OO, Aluma AL, et al. Prevalence of African trypanosomosis in cattle and sheep in Bassa local government area of Plateau State, Nigeria. *Sokoto J Vet Sci.* 2018;16:11–7.
135. Jaimes-Dueñez J, Triana-Chávez O, Mejía-Jaramillo AM. Spatial-temporal and phylogeographic characterization of *Trypanosoma* spp. in cattle (*Bos taurus*) and buffaloes (*Bubalus bubalis*) reveals transmission dynamics of these parasites in Colombia. *Vet Parasitol.* 2018;249:30–42.
136. Jaimes-Dueñez J, Zapata-Zapata C, Triana-Chávez O, Mejía-Jaramillo AM. Evaluation of an alternative indirect-ELISA test using in vitro-propagated *Trypanosoma brucei brucei* whole cell lysate as antigen for the detection of anti-*Trypanosoma evansi* IgG in Colombian livestock. *Prev Vet Med.* 2019;169:104712.
137. Joshua RA. Occurrence of human serum-resistant *Trypanosoma congoense* in goats and sheep in Nigeria. *Vet Parasitol.* 1989;31:107–13.
138. Kalu AU, Oboegbulem SI, Uzoukwu M. Trypanosomosis in small ruminants maintained by low riverine tsetse population in central Nigeria. *Small Rumin Res.* 2001;40:109–15.
139. Kalu AU. Prevalence of trypanosomiasis among Trypanotolerant cattle at the lower Benue River area of Nigeria. *Prev Vet Med.* 1995;24:97–103.
140. Kalu AU. Current status of tsetse fly and animal trypanosomiasis on the Jos Plateau, Nigeria. *Prev Vet Med.* 1996;27:107–13.
141. Karimuribo ED, Morrison LJ, Black A, Turner CMR, Kambarage DM, Ballingall KT. Analysis of host genetic factors influencing African trypanosome species infection in a cohort of Tanzanian *Bos indicus* cattle. *Vet Parasitol.* 2011;179:35–42.

142. Kassaye BK. Prevalence of bovine Trypanosomosis and apparent density of tsetse flies in Sayonole district Western Oromia, Ethiopia. *J Vet Sci Technol.* 2015;6:254.
143. Kayang BB, Bosompem KM, Assoku RKG, Awumbila B. Detection of *Trypanosoma brucei*, *T. congolense* and *T. vivax* infections in cattle, sheep and goats using latex agglutination. *Int J Parasitol.* 1997;27:83–7.
144. Kebede N, Fetene T, Animut A. Prevalence of Trypanosomosis of small ruminants in Guanqua district of Awil Zone, northwestern Ethiopia. *J Infect Dev Ctries.* 2009;3:245–6.
145. Kidanemariam A, Hadgu K, Sahle M. Parasitological prevalence of bovine trypanosomosis in Kindo Koisha district, Wolloa zone, south Ethiopia. *Onderstepoort J Vet Res.* 2002;69:107–13.
146. Kihurani DO, Nantulya VM, Mbuki SM, Mogoa E, Nguhiu-Mwangi J, Mbiti PM. *Trypanosoma brucei*, *T. congolense* and *T. vivax* infections in horses on a farm in Kenya. *Trop Anim Health Prod.* 1994;26:95–101.
147. Kimaro EG, Toribio J-AALML, Gwakisa P, Mor SM. Occurrence of trypanosome infections in cattle in relation to season, livestock movement and management practices of Maasai pastoralists in Northern Tanzania. *Vet Parasitol Reg Stud Reports.* 2018;12:91–8.
148. Kouadio IK, Sokouri D, Koffi M, Konaté I, Ahouty B, Koffi A, et al. Molecular characterization and prevalence of *Trypanosoma* species in cattle from a northern livestock area in Côte d'Ivoire. *Open J Vet Med.* 2014;4:314–21.
149. Laohasinnarong D, Thekiso OMM, Malele I, Namangala B, Ishii A, Goto Y, et al. Prevalence of *Trypanosoma sp.* in cattle from Tanzania estimated by conventional PCR and loop-mediated isothermal amplification (LAMP). *Parasitol Res.* 2011;109:1735–9.
150. Laohasinnarong D, Goto Y, Asada M, Nakao R, Hayashida K, Kajino K, et al. Studies of trypanosomiasis in the Luangwa valley, north-eastern Zambia. *Parasites Vectors.* 2015;8:1–8.
151. Lefrançois T, Solano P, De La Rocque S, Bengaly Z, Reifenberg JM, Kabore I, et al. New epidemiological features on animal trypanosomiasis by molecular analysis in the pastoral zone of Sideradougou, Burkina Faso. *Mol Ecol.* 1998;7:897–904.
152. Lelisa K, Shimeles S, Bekele J, Sheferaw D. Bovine trypanosomosis and its fly vectors in three selected settlement areas of Hawa-Gelan district, western Ethiopia. *Onderstepoort J Vet Res.* 2014;81:1–5.
153. Lema AA, Maigoro MA, Said M, Marwana AM, Nuraddeen W. Prevalence of bovine trypanosomosis in Katsina central abattoir, Katsina state. *Niger J Parasitol.* 2018;39:226–9.
154. Lopes STP, da Prado BS, Martins GHC, Beserra HEA, de Sousa Filho MAC, de Evangelista LSM, et al. *Trypanosoma vivax* in dairy cattle. *Acta Sci Vet.* 2018;46:287.
155. Luckins AG, Mehlitz D. Evaluation of an indirect fluorescent antibody test, enzyme-linked immunosorbent assay and quantification of immunoglobulins in the diagnosis of bovine trypanosomiasis. *Trop Anim Health Prod.* 1978;10:149–59.
156. Madruga CR, Araujo FR, Cavalcante-Goes G, Martins C, Pfeifer IB, Ribeiro LR, et al. The development of an enzyme-linked immunosorbent assay for *Trypanosoma vivax* antibodies and its use in epidemiological surveys. *Mem Inst Oswaldo Cruz.* 2006;101:801–7.
157. Maganga GD, Mavoungou JF, Ndilimabaka N, Kinga IC, Mve-Ondo B, Mombo IM, et al. Molecular identification of trypanosome species in trypanotolerant cattle from the south of Gabon. *Parasite.* 2017;24:4.
158. Magona JW, Kakaire DW, Mayende JSP. Prevalence and distribution of animal trypanosomosis on Buvuma Islands in Lake Victoria, Uganda. *Trop Anim Health Prod.* 1999;31:83–7.
159. Magona JW, Greiner M, Mehlitz D. Impact of tsetse control on the age-specific prevalence of trypanosomosis in village cattle in southeast Uganda. *Trop Anim Health Prod.* 2000;32:87–98.
160. Magona JW, Mayende JSP, Walubengo J. Comparative evaluation of the antibody-detection ELISA technique using microplates precoated with denatured crude antigens from *Trypanosoma congolense* or *Trypanosoma vivax*. *Trop Anim Health Prod.* 2002;34:295–308.
161. Magona JW, Mayende JSP, Olaho-Mukani W, Coleman PG, Jonsson NN, Welburn SC, et al. A comparative study on the clinical, parasitological and molecular diagnosis of bovine trypanosomosis in Uganda. *Onderstepoort J Vet Res.* 2003;70:213–8.
162. Magona JW, Walubengo J, Odimum JJ. Differences in susceptibility to trypanosome infection between Nkedi Zebu and Ankole cattle, under field conditions in Uganda. *Ann Trop Med Parasitol.* 2004;98:785–92.
163. Magona JW, Walubengo J, Odiit M, Okedi LA, Abila P, Katabazi BK, et al. Implications of the re-invasion of Southeast Uganda by *Glossina pallidipes* on the epidemiology of bovine trypanosomosis. *Vet Parasitol.* 2005;128:1–9.
164. Magona JW, Walubengo J, Odimum JT. Acute haemorrhagic syndrome of bovine trypanosomosis in Uganda. *Acta Trop.* 2008;107:186–91.
165. Magona JW, Walubengo J, Odimum JT. Differences in prevalence of trypanosomosis in Nkedi zebu, Ankole and crossbred cattle under tethering and open grazing management systems in Uganda. *Livest Res Rural Dev.* 2011;23:141.
166. Majekodunmi AO, Fajinmi A, Dongkum C, Picozzi K, Thrusfield MV, Welburn SC. A longitudinal survey of African animal trypanosomiasis in domestic cattle on the Jos Plateau, Nigeria: prevalence, distribution and risk factors. *Parasit Vectors.* 2013;6:239.
167. Makumyaviri AM, Ngarambe M. Parasitological and serological diagnosis of trypanosomiasis in cattle in the Northern-Kivu province, Congo. *Rev Med Vet.* 1997;148:809–12.
168. Makumyaviri A, Mehlitz D, Kageruka P, Kazyumba GL, Molisho D. Animal reservoir hosts of *Trypanosoma brucei gambiense* in Zaire: Trypanosome infections in two foci in Bas-Zaire. *Trop Med Parasitol.* 1989;40:258–62.
169. Mamabolo MV, Ntantiso L, Latif A, Majiwa PAO. Natural infection of cattle and tsetse flies in South Africa with two genotypic groups of *Trypanosoma congolense*. *Parasitology.* 2009;136:425–31.
170. Mamoudou A, Njanloga A, Hayatou A, Suh PF, Achukwi MD. Animal trypanosomosis in clinically healthy cattle of north Cameroon: epidemiological implications. *Parasites Vectors.* 2016;9:206.
171. Manuel Tafur T, Amanda Chávez V, Eva Casas A, Enrique SM. Prevalence of *Trypanosoma vivax* in cattle in high forest of the province of Chachapoyas, Amazonas. *Rev Investig Vet del Peru.* 2002;13:94–7.
172. Masiga RC, Nyang'ao JMN. Identification of trypanosome species from camel using polymerase chain reaction and procyclic transformation test. *J Camel Pract Res.* 2001;8:17–22.
173. Mattioli RC, Faye JA, Jaitner J. Estimation of trypanosomal status by the buffy coat technique and an antibody ELISA for assessment of the impact of trypanosomosis on health and productivity of N'Dama cattle in The Gambia. *Vet Parasitol.* 2001;95:25–35.
174. Mbahin N, Affognon H, Andoque J, Tiberius M, Mbuvi D, Otieno J, et al. Parasitological prevalence of bovine trypanosomosis in Kubo division of Kwale county of coastal: baseline survey. *Am J Anim Vet Sci.* 2013;8:28–36.
175. Mbewu NJ, Namangala B, Sitali L, Vorster I, Michelo C. Prevalence of pathogenic trypanosomes in anaemic cattle from trypanosomosis challenged areas of Itezhi-tezhi district in central Zambia. *Parasit Vectors.* 2015;8:638.
176. Mehlitz D. Trypanosome infections in domestic animals in Liberia. *Tropenmed Parasitol.* 1979;30:212–9.
177. Mekata H, Konnai S, Witola WH, Inoue N, Onuma M, Ohashi K. Molecular detection of trypanosomes in cattle in South America and genetic diversity of *Trypanosoma evansi* based on expression-site-associated gene 6. *Infect Genet Evol.* 2009;9:1301–5.
178. Mekibib B, Manegerew M, Tadesse A, Abuna F, Megersa B, Regassa A, et al. Prevalence of haemoparasites and associated risk factors in working donkeys in Adigudem and Kwiha districts of Tigray region, northern Ethiopia. *J Anim Vet Adv.* 2010;9:2249–55.
179. Mekonnen B, Regassa V, Kahsay AG. Epidemiology of trypanosomosis in goats in Abelti, Bede and Ghibe valley, south West Ethiopia. *Int J Trop Med.* 2014;9:10–4.
180. Mekuria S, Gadissa F. Survey on bovine trypanosomosis and its vector in Metekel and Awi zones of Northwest Ethiopia. *Acta Trop.* 2011;117:146–51.
181. Mekuria S, Eyob A, Regassa A, Tadesse A, Mekibib B, Abebe R. A cross-sectional study of equine trypanosomosis and its vectors in Wolayta zone, Southern Ethiopia. *J Anim Vet Adv.* 2010;9:2061–6.
182. Melese M, Alemu S, Kemal J, Muktar Y, Abraha A. Vector identification and bovine trypanosomosis in edja district. South Ethiopia *Livest Res Rural Dev.* 2017;29:5.
183. Mihok S, Olubayo RO, Moloo SK. Trypanosomiasis in the black rhinoceros (*Diceros bicornis Linnaeus, 1758*). *Rev Sci Tech.* 1992;11:1169–73.

184. Mihret A, Mamo G. Bovine trypanosomosis in three districts of East Gojam Zone bordering the Blue Nile River in Ethiopia. *J Infect Dev Ctries*. 2007;1:321–5.
185. Moll G, Lohding A, Young AS. Epidemiology of theilerioses in the Trans-Mara division, Kenya: Husbandry and disease background and preliminary investigations on theilerioses in calves. *Prev Vet Med*. 1984;2:801–31.
186. Mossaad E, Salim B, Saganuma K, Musinguzi P, Hassan MA, Elamin EA, et al. *Trypanosoma vivax* is the second leading cause of camel trypanosomosis in Sudan after *Trypanosoma evansi*. *Parasites and Vectors*. 2017;10:1.
187. Moti Y, Fikru R, Büscher P, Van Den Abbeele J, Duchateau L, Delespaux V. Detection of african animal trypanosomes: The haematocrit centrifugation technique compared to PCR with samples stored on filter paper or in DNA protecting buffer. *Vet Parasitol*. 2014;203:253–8.
188. Mugittu KN, Silayo RS, Majiwa PAO, Kimbita EK, Mutayoba BM, Maselle R. Application of PCR and DNA probes in the characterisation of trypanosomes in the blood of cattle in farms in Morogoro, Tanzania. *Vet Parasitol*. 2001;94:177–89.
189. Muhangazi D, Mugenyi A, Bigirwa G, Kamusime M, Kitibwa A, Akurut GG, et al. African animal trypanosomiasis as a constraint to livestock health and production in Karamoja region: a detailed qualitative and quantitative assessment. *BMC Vet Res*. 2017;13:1–3.
190. Muhangazi D, Picozzi K, Hattendorf J, Thrusfield M, Kabasa JD, Waiswa C, et al. The burden and spatial distribution of bovine African trypanosomes in small holder croplivestock production systems in Tororo district, south-eastern Uganda. *Parasites Vectors*. 2014;7:603.
191. Mulaw S, Addis M, Fromsa A. Study on the prevalence of major trypanosomes affecting bovine in tsetse infested Asosa district of Benishangul Gumuz Regional State, Western Ethiopia. *Glob Vet*. 2011;7:330–6.
192. Mungube EO, Vitouley HS, Allegye-Cudjoe E, Diall O, Boucoum Z, Diarra B, et al. Detection of multiple drug-resistant *Trypanosoma congolense* populations in village cattle of south-east Mali. *Parasites Vectors*. 2012;5:155.
193. Musinguzi SP, Saganuma K, Asada M, Laohassinrong D, Sivakumar T, Yokoyama N, et al. A PCR-based survey of animal African trypanosomosis and selected piroplasm parasites of cattle and goats in Zambia. *J Vet Med Sci*. 2016;78:1819–24.
194. Mwambu PM, Mayende JSP. Salivarian trypanosome infections in cattle in Ikoma, South Mara district, Tanzania. *Parasitology*. 1973;66:381–5.
195. Mwambu PM. Prevalence of *Trypanosoma vivax* infection in cattle in Teso district, Eastern Uganda. *Bull Epizoot Dis Afr*. 1969;17:395–402.
196. Mwangi EK, Stevenson P, Gettinby G, Reid SW, Murray M. Susceptibility to trypanosomosis of three *Bos indicus* cattle breeds in areas of differing tsetse fly challenge. *Vet Parasitol*. 1998;79:1–17.
197. N'Djetchi MK, Ilboudo H, Koffi M, Kaboré J, Kaboré JW, Kaba D, et al. The study of trypanosome species circulating in domestic animals in two human African trypanosomiasis foci of Côte d'Ivoire identifies pigs and cattle as potential reservoirs of *Trypanosoma brucei gambiense*. *PLoS Negl Trop Dis*. 2017;11:1–16.
198. Nakayima J, Nakao R, Alhassan A, Mahama C, Afakye K, Sugimoto C. Molecular epidemiological studies on animal trypanosomiasis in Ghana. *Parasites Vectors*. 2012;5:1–7.
199. Nakayima J, Nakao R, Alhassan A, Hayashida K, Namangala B, Mahama C, et al. Genetic diversity among *Trypanosoma (Duttonella) vivax* strains from Zambia and Ghana, based on cathepsin L-like gene. *Parasite*. 2013;20:24.
200. Nantulya VM, Lindqvist KJ, Stevenson P, Mwangi EK. Application of a monoclonal antibody-based antigen detection enzyme-linked immunosorbent assay (antigen ELISA) for field diagnosis of bovine trypanosomiasis at Nguruman, Kenya. *Ann Trop Med Parasitol*. 1992;86:225–30.
201. Ngayo MO, Njiru ZK, Kenya EU, Muluvi GM, Osir EO, Masiga DK. Detection of trypanosomes in small ruminants and pigs in western Kenya: Important reservoirs in the epidemiology of sleeping sickness? *Kinetoplastid Biol Dis*. 2005;4:1–7.
202. Ngomtcho SCH, Weber JS, Ngo Bum E, Gbem TT, Kelm S, Achukwi MD. Molecular screening of tsetse flies and cattle reveal different Trypanosoma species including *T. grayi* and *T. theileri* in northern Cameroon. *Parasites Vectors*. 2017;10:1–16.
203. Nonga HE, Kambarage DM. Prevalence of Bovine trypanosomosis in Morogoro, Tanzania. *Pakistan J Nutr*. 2009;8:208–13.
204. Nyimba PH, Komba EVG, Sugimoto C, Namangala B. Prevalence and species distribution of caprine trypanosomosis in Sinazongwe and Kalomo districts of Zambia. *Vet Parasitol*. 2015;210:125–30.
205. Ocaido M, Otieno CP, Okuna NM, Erume J, Ssekitto C, Wafula RZOO, et al. Socio-economic and livestock disease survey of agropastoral communities in Serere county, Soroti district, Uganda. *Livest Res Rural Dev*. 2005;17.
206. Ode S, Adamu M, Taioe M, Thekiso O, Adamu S, Saror DI. Molecular occurrence of trypanosomes, erythrocyte and serum sialic acid concentrations of Muturu and Bunaji cattle in Benue State, Nigeria. *Vet Parasitol*. 2017;242:10–3.
207. Odeniran PO, Macleod ET, Ademola IO, Welburn SC. Molecular identification of bovine trypanosomes in relation to cattle sources in southwest Nigeria. *Parasitol Int*. 2019;68:1–8.
208. Odongo S, Delespaux V, Ngoro M, Bekkele SM, Magez S. Comparative evaluation of the nested ITS PCR against the 18S PCR-RFLP in a survey of bovine trypanosomiasis in Kwale County, Kenya. *J Vet Diagn Invest*. 2016;28:589–94.
209. Okech G, Dolan RB, Stevenson P, Alushula H, Watson ED, Luckins AG, et al. The effect of trypanosomosis on pregnancy in trypanotolerant Orma Boran cattle. *Theriogenology*. 1996;46:441–7.
210. Oliveira JB, Hernández-Gamboa J, Jiménez-Alfaro C, Zeledón R, Blandón M, Urbina A, et al. First report of *Trypanosoma vivax* infection in dairy cattle from Costa Rica. *Vet Parasitol*. 2009;163:136–9.
211. Omoogun GA, Akinboade OA. Tsetse and bovine trypanosomosis incidence at Egbe in the derived savanna zone of Nigeria. *Insect Sci its Appl*. 2000;20:215–9.
212. Ono MSB, Souto PC, Cruz JA, Guerra NR, Guimarães JA, Dantas AC, et al. *Trypanosoma vivax* outbreak in cattle in the 'Zona da Mata' of the state of Pernambuco'. *Med Vet*. 2017;11:96–101.
213. Osaer S, Goossens B, Kora S, Gaye M, Darboe L. Health and productivity of traditionally managed Djallonke sheep and West African dwarf goats under high and moderate trypanosomosis risk. *Vet Parasitol*. 1999;82:101–19.
214. Osiyemi TIOO, Agbonlahor DEAA. Incidence of protozoan blood parasites in livestock in northern Nigeria. *Trop Anim Health Prod*. 1980;12:115.
215. Otte MJ, Abuabara JY, Nieto MI, Gutierrez JR. Incidence of *Trypanosoma vivax* infection on three cattle farms on the north coast of Colombia. *Acta Vet Scand Suppl*. 1988;84:104–6.
216. Pagabeleguem S, Sangaré M, Bengaly Z, Akoudjini M, Belem AMGG, Bouyer J. Climate, cattle rearing systems and African animal trypanosomosis risk in Burkina Faso. *PLoS ONE*. 2012;7:e49762.
217. Paolletta MS, López Arias L, de la Fournière S, Guillemi EC, Luciani C, Sarmiento NF, et al. Epidemiology of *Babesia*, *Anaplasma* and *Trypanosoma* species using a new expanded reverse line blot hybridization assay. *Ticks Tick Borne Dis*. 2018;9:155–63.
218. De Pereira Almeida PJLL, Nda M, Goossens B, Osaer S. PCR primer evaluation for the detection of trypanosome DNA in naturally infected goats. *Vet Parasitol*. 1998;80:111–6.
219. Pereira HD, Simões SVDD, Souza FAL, Silveira JAG, Ribeiro MFB, Cadioli FA, et al. Clinical and epidemiological aspects and diagnosis of *Trypanosoma vivax* infection in a cattle herd, state of Maranhão, Brazil. *Pesqui Vet Bras*. 2018;38:896–901.
220. de Souza Pimentel D, do Nascimento Ramos CA, Ramos RADNRA, de Araújo FR, Borba ML, da Gloria Faustino MA, et al. First report and molecular characterization of *Trypanosoma vivax* in cattle from state of Pernambuco, Brazil. *Vet Parasitol*. 2012;185:286–9.
221. Ávila Pulgarín LS, Acevedo Restrepo A, Jurado Guevara JA, Polanco Echverry D, Velásquez Vélez R, Zapata Salas R, et al. Hemoparasite infection in goats and sheep at five municipalities in north and northeastern Antioquia (Colombia). *Rev CES Med Vet y Zootec*. 2013;8:11–21.
222. Patricia Quispe A, Amanda Chávez V, Eva Casas A, Antonio Trigueros V, Francisco SA. Prevalence of *Trypanosoma vivax* in cattle from the Coronel Portillo province, Ucayali. *Rev Investig Vet del Peru*. 2003;14:161–5.
223. Rahman AHA. Observations on the trypanosomosis problem outside the tsetse belts of Sudan. *Rev Sci Tech*. 2005;24:965–72.
224. Ramírez-Iglesias JR, Elejalde MC, Reyna-Bello A, Mendoza M. Molecular diagnosis of cattle trypanosomes in Venezuela: evidences of *Trypanosoma evansi* and *Trypanosoma vivax* infections. *J Parasit Dis*. 2017;41:450–8.

225. Ravel S, Mediannikov O, Bossard G, Desquesnes M, Cuny G, Davoust B. A study on African animal trypanosomosis in four areas of Senegal. *Folia Parasitol.* 2015;62:2015.044.
226. de Reis MO, Souza FR, Albuquerque AS, Monteiro F, Oliveira LFDS, Raymundo DL, et al. Epizootic Infection by *Trypanosoma vivax* in cattle from the State of Minas Gerais, Brazil. *Korean J Parasitol.* 2019;57:191–5.
227. Robson J, Rickman LR. Results of a field trial for the improved detection of *Trypanosoma vivax* in domestic animals. *Bull Epizoot Dis Afr.* 1972;20:297–9.
228. Rodrigues CMF, Batista JS, Lima JM, Freitas FJC, Barros IO, Garcia HA, et al. Field and experimental symptomless infections support wandering donkeys as healthy carriers of *Trypanosoma vivax* in the Brazilian Semiarid, a region of outbreaks of high mortality in cattle and sheep. *Parasit Vectors.* 2015;8:564.
229. Roeder PL, Scott JM, Pogram RG, Prod H. Acute *Trypanosoma vivax* infection of Ethiopian cattle in the apparent absence of tsetse. *Trop Anim Health Prod.* 1984;16:141–7.
230. Rowlands GJ, Leak SG, Peregrine AS, Nagda SM, Mulatu W, D'leteren GD. The incidence of new and the prevalence and persistence of recurrent trypanosome infections in cattle in southwest Ethiopia exposed to a high challenge with drug-resistant parasites. *Acta Trop.* 2001;79:149–63.
231. Sabir N, Chaudhry ZI, Aslam A, Muhammad K, Shahid M, Hussain A, et al. A study on prevalence and molecular characterization of trypanosomal species infecting equines in Lahore region, Pakistan. *J Parasit Dis.* 2018;42:96–101.
232. Salas RZ, Zuluaga EAC, Vélez JR, Chávez OT, García VHP, Osorio LAR, et al. Bovine trypanosomiasis in dairy farming in the high tropics: first report of Haematobia irritans as the main vector for *T. vivax* and *T. evansi* in Colombia. *Rev Med Vet.* 2017;33:21–34.
233. Salim B, Bakheit MA, Sugimoto C. Molecular detection of equine trypanosomes in the Sudan. *Vet Parasitol.* 2014;200:246–50.
234. Samdi S, Abenga JN, Fajinmi A, Kalgo A, Idowu T, Lawani F. Seasonal variation in trypanosomosis rates in small ruminants at the Kaduna abattoir, Nigeria. *African J Biomed Res.* 2008;11:229–32.
235. Samdi SM, Abenga JN, Wayo B, Mshelia WP, Musa D, Haruna MK, et al. The complementary roles of biting flies and reservoirs of infection: In the resurgent of African animal trypanosomosis in Keffi local Government area of Nassarawa state, Nigeria. *Asian J Anim Vet Adv.* 2010;6:316–21.
236. Samdi SM, Fajinmi AO, Kalejaye JO, Wayo B, Haruna MK, Yarnap JE, et al. Prevalence of trypanosomosis in cattle at slaughter in Kaduna central Abattoir. *Asian J Anim Sci.* 2011;5:162–5.
237. Sanni TM, Onasanya GO, Adefenwa MA, Yakubu A, Ikeobi CON, Adelbambo OA, et al. Molecular diagnosis of subclinical African *Trypanosoma vivax* infection and association with physiological indices and serum metabolites in extensively managed goats in the tropics. *Open J Vet Med.* 2013;3:39–45.
238. Seck MT, Bouyer J, Sall B, Bengaly Z, Vreyen MJB. The prevalence of African animal trypanosomoses and tsetse presence in Western Senegal. *Parasite.* 2010;17:257–65.
239. Sharma SP, Losho TC, Malau M, Mangate KG, Linchwe KB, Amanfu W, et al. The resurgence of trypanosomosis in Botswana. *J S Afr Vet Assoc.* 2001;72:232–4.
240. Sheferaw D, Birhanu B, Asrade B, Abera M, Tusse T, Fikadu A, et al. Bovine trypanosomosis and Glossina distribution in selected areas of southern part of Rift Valley, Ethiopia. *Acta Trop.* 2016;154:145–8.
241. Silbermayr K, Li F, Soudré A, Müller S, Sölkner J. A Novel qPCR assay for the detection of African animal trypanosomosis in trypanotolerant and trypanosusceptible cattle breeds. *PLoS Negl Trop Dis.* 2013;7:e2345.
242. Silva RA, da Silva JA, Schneider RC, de Freitas J, Mesquita D, Mesquita T, et al. Outbreak of trypanosomiasis due to *Trypanosoma vivax* (Ziemann, 1905) in bovines of the Pantanal, Brazil. *Mem Inst Oswaldo Cruz.* 1996;91:561–2.
243. Silva RA, Morales G, Eulert E, Montenegro A, Ybanez R. Outbreaks of trypanosomosis due to *Trypanosoma vivax* in cattle in Bolivia. *Vet Parasitol.* 1998;76:153–7.
244. Silveira JAG, Rabelo TML, Lacerda ACR, Borges PAL, Tomás WM, Pellegrin AO, et al. Molecular detection and identification of hemoparasites in pampas deer (*Ozotoceros bezoarticus* Linnaeus, 1758) from the Pantanal Brazil. *Ticks Tick Borne Dis.* 2013;4:341–5.
245. Simo G, Asonganyi T, Nkinin SW, Njiokou F, Herder S. High prevalence of *Trypanosoma brucei gambiense group 1* in pigs from the Fontem sleeping sickness focus in Cameroon. *Vet Parasitol.* 2006;139:57–66.
246. Simukoko H, Marcotty T, Phiri I, Geysen D, Vercruyse J, Van den Bossche P. The comparative role of cattle, goats and pigs in the epidemiology of livestock trypanosomiasis on the plateau of eastern Zambia. *Vet Parasitol.* 2007;147:231–8.
247. Simwango M, Ngonyoka A, Nnko HJ, Salekwa LP, Ole-Neselle M, Kimera SI, et al. Molecular prevalence of trypanosome infections in cattle and tsetse flies in the Maasai Steppe, northern Tanzania. *Parasites Vectors.* 2017;10:1.
248. Sinshaw A, Abebe G, Desquesnes M, Yoni W. Biting flies and *Trypanosoma vivax* infection in three highland districts bordering lake Tana Ethiopia. *Vet Parasitol.* 2006;142:35–46.
249. Sinyangwe L, Delespaux V, Brandt J, Geerts S, Mubanga J, Machila N, et al. Trypanocidal drug resistance in eastern province of Zambia. *Vet Parasitol.* 2004;119:125–35.
250. Snak A, Lara AA, Garcia FG, Pieri EM, Da Silveira JAG, Osaki SC, et al. Prevalence study on *Trypanosoma vivax* in dairy cattle in the western region on the State of Paraná, Brazil. *Semin Ciências Agrárias.* 2018;39:425–30.
251. Solano P, Michel JF, Lefrancois T, de La Rocque S, Sidibe I, Zoungrana A, et al. Polymerase chain reaction as a diagnosis tool for detecting trypanosomes in naturally infected cattle in Burkina Faso. *Vet Parasitol.* 1999;86:95–103.
252. Sow A, Sidibe I, Bengaly Z, Marcotty T, Sere M, Diallo A, et al. Field detection of resistance to isometamidium chloride and diminazene aceturate in *Trypanosoma vivax* from the region of the Boucle du Mouhoun in Burkina Faso. *Vet Parasitol.* 2012;187:105–11.
253. Sow A, Ganaba R, Percoma L, Sidibé I, Bengaly Z, Adam Y, et al. Baseline survey of animal trypanosomosis in the region of the Boucle du Mouhoun, Burkina Faso. *Res Vet Sci.* 2013;94:573–8.
254. Specht EJK. Prevalence of bovine trypanosomosis in Central Mozambique from 2002 to 2005. *Onderstepoort J Vet Res.* 2008;75:73–81.
255. Suárez C, García F, Román D, Coronado A, Perrone T, Reyna A, et al. Risk factors associated with the bovine trypanosomosis of livestock farms in Venezuela. *Zootec Trop.* 2010;27:363–72.
256. Swai ES, Kaaya JE. A parasitological survey for bovine trypanosomosis in the livestock/wildlife ecozone of Northern Tanzania. *Vet World.* 2012;5:459–64.
257. Tadesse A, Tsegaye B. Bovine trypanosomosis and its vectors in two districts of Bench Maji zone, South Western Ethiopia. *Trop Anim Health Prod.* 2010;42:1757–62.
258. Tafese W, Melaku A, Fentahun T. Prevalence of bovine trypanosomosis and its vectors in two districts of East Wollega Zone Ethiopia. *Onderstepoort J Vet Res.* 2012;79:E1–4.
259. Takeuti MI, Fagbemi BO, De Donato M, Yakubu A, Rodulfo HE, Peters SO, et al. Molecular survey of pathogenic trypanosomes in naturally infected Nigerian cattle. *Res Vet Sci.* 2013;94:555–61.
260. Tamasaukas R, Roa N, Cobo M. Trypanosomosis due to *Trypanosoma vivax* in two buffalo (*Bubalus bubalis*) farms of Guárico state, Venezuela. *Rev Cient la Fac Ciencias Vet la Univ del Zulia.* 2006;16:575–8.
261. Tarimo-Nesbitt RA, Golder TK, Dransfield RD, Chaudhury MF, Brightwell R. Trypanosome infection rate in cattle at Nguruman, Kenya. *Vet Parasitol.* 1999;81:107–17.
262. Taseew S, Duguma R. Cattle anaemia and trypanosomiasis in western Oromia State, Ethiopia. *Rev Med Vet.* 2012;163:581–8.
263. Tchamdja E, Kulo AE, Vitouley HS, Batawui K, Bankolé AA, Adomefa K, et al. Cattle breeding, trypanosomosis prevalence and drug resistance in Northern Togo. *Vet Parasitol.* 2017;236:86–92.
264. Terefe E, Haile A, Mulatu W, Dessie T, Mwai O. Phenotypic characteristics and trypanosome prevalence of Mursi cattle breed in the Bodi and Mursi districts of South Omo Zone, southwest Ethiopia. *Trop Anim Health Prod.* 2015;47:485–93.
265. Tesfaheyewet Z, Getnet F, Ayana M. A cross-sectional study on the prevalence of bovine trypanosomosis in Amhara region, Northwest Ethiopia. *Livest Res Rural Dev.* 2012;24.
266. Tesfaye D, Ibrahim N. Prevalence of bovine trypanosomosis in Assosa district of Benishangul Gumuz. *Adv Biol Res.* 2017;11:13–7.
267. Thumbi SM, Jung'A JO, Mosi RO, McOdimba FA. Spatial distribution of African animal trypanosomiasis in suba and teso districts in Western Kenya. *BMC Res Notes.* 2010;3:6.

268. Trail JC, D'leteren GD, Viviani P, Yangari G, Nantulya VM. Relationships between trypanosome infection measured by antigen detection enzyme immunoassays, anaemia and growth in trypanotolerant N'Dama cattle. *Vet Parasitol.* 1992;42:213–23.
269. Ukpai OM, Obasi NI. Prevalence of trypanosomiasis in relation to some haematological parameters in cattle, Ohafia LGA, Abia State Nigeria. *Niger J Parasitol.* 2017;38:250–2.
270. Van Den Bossche P, Mudenge D, Mubanga J, Norval A. The parasitological and serological prevalence of tsetse-transmitted bovine trypanosomosis in the Eastern Caprivi (Caprivi District, Namibia). *Onderstepoort J Vet Res.* 1999;66:103–10.
271. Van den Bossche P, Shumba W, Makhambera P. The distribution and epidemiology of bovine trypanosomosis in Malawi. *Vet Parasitol.* 2000;88:163–76.
272. Vokaty S, McPherson VO, Camus E, Applewhaitte L. Ovine trypanosomosis: a seroepidemiological survey in coastal Guyana. *Rev Elev Med Vet Pays Trop.* 1993;46:57–9.
273. Waiswa C, Katunguka-Rwakishaya E. Bovine trypanosomiasis in southwestern Uganda: packed-cell volumes and prevalences of infection in the cattle. *Ann Trop Med Parasitol.* 2004;98:21–7.
274. Wells EA, Betamcourt A, Ramirez LE. Serological evidence for the geographical distribution of *Trypanosoma vivax* in the new world. *Trans R Soc Trop Med Hyg.* 1977;71:448–9.
275. Wijers DJB. The complex epidemiology of Rhodesian sleeping sickness in Kenya and Uganda. Part II: observations in Samia (Kenya). *Trop Geogr Med.* 1974;26:182–97.
276. Yesufu HM, Mshelbwala AS. Trypanosomiasis survey in cattle and tsetse flies along a trade cattle route in southwestern Nigeria. *Ann Trop Med Parasitol.* 1973;67:307–12.
277. Zanatto DCS, Gatto IRH, Labruna MB, Jusi MMG, Samara SI, Machado RZ, et al. *Coxiella burnetii* associated with BVDV (Bovine Viral Diarrhea Virus), BoHV (bovine herpesvirus), *Leptospira* spp., *Neospora caninum*, *Toxoplasma gondii* and *Trypanosoma vivax* in reproductive disorders in cattle. *Rev Bras Parasitol Vet.* 2019;28:245–57.
278. Cadioli FA, de Barnabe PA, Machado RZ, Teixeira MCA, Andre MR, Sampaio PH, et al. First report of *Trypanosoma vivax* outbreak in dairy cattle in Sao Paulo state Brazil. *Rev Bras Parasitol Vet.* 2012;21:118–24.
279. Ganyo EY, Boampong JN, Masiga DK, Villinger J, Turkson PK. Haematology of N'Dama and West African Shorthorn cattle herds under natural *Trypanosoma vivax* challenge in Ghana. *F1000Research.* 2018;7:314.
280. González JR, Meléndez RD. Seroprevalencia de la tripanosomosis y anaplasmosis bovina en el municipio Juan José Mora del estado Carabobo, Venezuela, Mediante la técnica de ELISA. *Rev Cient la Fac Ciencias Vet la Univ del Zulia.* 2007;17:449–55.
281. Lopes STP, da Prado BS, Martins GHC, Beserra HEA, de Sousa Filho MAC, de Evangelista LSM, et al. *Trypanosoma vivax* em bovino leiteiro. *Acta Sci Vet.* 2018;46:1–5.
282. Wijers DJB. The complex epidemiology of Rhodesian sleeping sickness in Kenya and Uganda. 1. The absence of the disease on Mfangano island (Kenya). *Trop Geogr Med.* 1974;26:58–64.
283. Mihok S, Mnyoki E, Brett RA, Jonyo JF, Röttcher D, Majiwa PAO, et al. Trypanosomiasis and the conservation of black rhinoceros (*Diceros bicornis*) at the Ngulia Rhino Sanctuary, Tsavo West National Park Kenya. *Afr J Ecol.* 1992;30:103–15.
284. Thon MGA. Prevalence of trypanosomosis in cattle in Juba Area, Central Equatoria State, Sudan. MTAH thesis, Department of Preventive Medicine Faculty of Veterinary Medicine University of Khartoum. Department of Preventive Medicine Faculty of Veterinary Medicine University of Khartoum; 2009.
285. Shereni W, Anderson NE, Nyakupinda L, Cecchi G. Spatial distribution and trypanosome infection of tsetse flies in the sleeping sickness focus of Zimbabwe in Hurungwe District. *Parasites Vectors.* 2016;9:605.
286. Genevieve A-Y, Bakary C, Mavoungou JF, Silas Lendzele S, Abdallah NE. Preliminary study on vectors of bovine trypanosomosis in the central African republic one decade after the socio-military crisis. *J Anim Plant Sci.* 2019;39:6487–94.
287. Franco JR, Cecchi G, Priotto G, Paone M, Diarra A, Grout L, et al. Monitoring the elimination of human African trypanosomiasis: update to 2014. *PLoS Negl Trop Dis.* 2017;11:1–26.
288. Ahmed SK, Rahman AH, Hassan MA, Salih SEM, Paone M, Cecchi G. An atlas of tsetse and bovine trypanosomosis in Sudan. *Parasit Vectors.* 2016;9:194.
289. Stephen LE. Trypanosomiasis A Veterinary Perspective. Oxford: Pergamon Press; 1986.
290. Arengwi WG, Agga GE, Abdi RD, Büscher P. Systematic review and meta-analysis on the global distribution, host range, and prevalence of *Trypanosoma evansi*. *Parasit Vectors.* 2019;12(67):1–25.
291. Gardiner PR. Recent studies of the biology of *Trypanosoma vivax*. *Adv Parasitol.* 1989;28:229–317.
292. Desquesnes M. Evaluation of a simple PCR technique for the diagnosis of *Trypanosoma vivax* infection in the serum of cattle in comparison to parasitological techniques and antigen-enzyme-linked immuno sorbent assay. *Acta Trop.* 1997;65:139–48.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](http://biomedcentral.com/submissions)

