

1 **Palynostratigraphic correlation of the Sardhai Formation (Permian) of Pakistan**

2 **Irfan U. Jan^{a&c,*}, Michael H. Stephenson, Fazli R. Khan^c**

3 **^aDepartment of Geology, University of Leicester, University Road Leicester, LE1**
4 **7RH, UK**

5 **E-mail: ij21@le.ac.uk**

6 **Fax: +44 (0)1162523918**

7 **^bBritish Geological Survey, Keyworth, Nottingham, NG12 5GG, UK**

8 **^cNational Centre of Excellence in Geology, University of Peshawar, Pakistan.**

9

10 **Abstract**

11 Palynological assemblages from the Sardhai Formation shale (Permian), lying between
12 the red-bed Warchha Formation and the Amb Formation limestones in the Salt and
13 Khisor ranges of Pakistan contain abundant bisaccate pollen grains and few spores. In
14 particular, well-preserved specimens of *Florinites ?balmei*, a bilaterally symmetrical
15 monosaccate pollen grain, are common. The presence of this pollen and the stratigraphic
16 context suggest that the Sardhai Formation correlates with the Khuff transition beds of
17 Oman and the basal Khuff clastics of central Saudi Arabia. *Florinites ?balmei* was first
18 described by Stephenson and Filatoff in 2000 from the basal Khuff clastics of Saudi
19 Arabia, and it has since been reported from Oman, Kuwait, southeastern Turkey, Iraq,
20 United Arab Emirates and Qatar. This suggests that the plant that produced *Florinites*
21 *?balmei* had a rather limited palaeogeographic distribution in the Mid-Permian which

22 may be useful in reconstructing the problematic tectonic and palaeogeographic history of
23 this complex region.

24

25 **Keywords:** Permian; Palynology; Sardhai Formation; Salt Range; Khisor Range;
26 Nilwahan Group; Zaluch Group.

27

28 **Introduction**

29 The Permian succession of Pakistan crops out in the Salt Range and Trans-Indus Khisor
30 and Marwat ranges and partly in the Surghar Range (Fig. 1), which represent the southern
31 side of a rift flank basin, along the northern Gondwanan coastal margin (Wardlaw and
32 Pogue, 1995). The Salt Range and the Trans-Indus ranges of Pakistan are regarded as
33 important reference areas for Permian strata; however despite considerable research in the
34 area, few studies have dealt with biostratigraphy (e.g. Pakistani-Japanese Research
35 Group, 1985; Wardlaw and Pogue, 1995; Mertmann, 1999). Most studies have
36 concentrated on taxonomy of various groups, and Permian-Triassic boundary problems
37 (e.g. see Waagen, 1882-1885; Noetling, 1901; Diener, 1912; Grabau, 1931; Balme,
38 1970; Kummel and Teichert, 1970; Rowell, 1970; Grant, 1970; Glenister and Furnish,
39 1970; Kummel, 1970; Sohn, 1970; Sweet, 1970; Sarjeant, 1970).

40 The Permian succession is divided into two groups (Fig. 2), representing two different
41 depositional settings: the largely terrestrial Gondwana succession, represented by the
42 Nilwahan Group, and the shallow marine Tethyan succession, represented by the
43 overlying Zaluch Group (Wardlaw and Pogue, 1995). The base of the Nilawahan Group
44 is characterized by the Tobra Formation, showing glacially-influenced sedimentation

45 (Ghauri et al., 1977). It is overlain by the Dandot Formation, in the Salt Range. But this
46 formation is absent in the Khisor Range and western Salt Range. The Dandot Formation
47 consists of pale grey to olive green sandstone having occasionally scattered pebbles of up
48 to 10 cm in diameter or pebbly beds with subordinate dark grey and greenish splintery
49 shales (Shah, 1977) and containing the bivalve *Eurydesma* and the conularid, *Conularia*
50 (Reed, 1936; Pascoe, 1959). Many species of Bryozoa and Ostracoda along with a few
51 brachiopod taxa have also been described from the formation..

52 Arid conditions are indicated by the succeeding Warchha Formation, which consists of
53 medium- to coarse-grained, purple, arkosic sandstone, conglomeratic in places with
54 interbeds of reddish shale. The conglomerate clasts are mostly granitic in nature, though
55 quartzitic clasts are also present.

56 More humid conditions are indicated by the overlying Sardhai Formation (Sultan, 2004).
57 The name Sardhai Formation, as approved by the Stratigraphic Committee of Pakistan,
58 comes from Gee (written comm. 1964). Prior to which, Gee (in Pascoe, 1959) called it
59 “Lavender clay stage”. Earlier, Wynne (1878) called it “Lavender clay” whereas Noetling
60 (1901) called the formation “upper part of Warchha Group” (Shah, 1977). The type
61 locality of the formation has been suggested by Gee, as the Sardhai Gorge in the eastern
62 Salt Range (Shah, 1977). The formation is composed of bluish to greenish-grey claystone
63 with subordinate sandstone and siltstone interbeds. It is also reported to contain minor
64 carbonaceous clays. These clays are lavender in colour and contain copper minerals, as
65 well as jarosite, chert and gypsum (Shah, 1977). Shah (1977) observed lateral facies
66 changes from the lavender- coloured clays in the Salt Range to black shale and brownish
67 argillaceous limestone in the Khisor Range. Generally the formation is reported to be

68 unfossiliferous with occasional plant remains in the exposures in the Salt Range (Shah,
69 1977); however well-developed limestone interbeds from the Khisor Range, have been
70 reported by Shah (1977) to have yielded determinable brachiopods and bryozoans.
71 Hussain (1967) reported the fossils, *Anastomopora* sp., *Fenestella* sp., *Athyris* sp.,
72 *Spirifer* sp. from the sandy limestone beds exposed at the Saiyiduwali in the Khisor
73 Range and assigned an Early Permian age to the formation (Alam, 2008). Moreover a
74 unit recognized as Sardhai from the Jang drill core (Fig. 1; Alam et al., 1987), following
75 Wardlaw and Pogue (1995) contained common fish debris. One of the samples also
76 yielded common paleoniscoid fish teeth and a single *Hindeodus* conodont, the presence
77 of which by Wardlaw and Pogue (1995) has been attributed to the deposition of the
78 formation in a very shallow marine settings. Variable thicknesses of the formation have
79 been observed in eastern and western Salt Range and Trans-Indus Khisor Range. At the
80 type locality i.e Sardhai Gorge, it is reported to be 42m thick, in the western Salt Range,
81 it is 65m thick, whereas in the Khisor Range, it makes up 50m (Shah, 1977). Its lower
82 contact with the Warchha Formation has been assigned as transitional, whereas the upper
83 contact, following Shah (1977) is conformable with the overlying Amb Formation.
84 Wardlaw and Pogue (1995) have mentioned the presence of the significant hiatus
85 between the Nilawahan and Zaluch Group. Based on the presence of the conodont in the
86 upper part of the Nilawhan Group (Sardhai Formation) Wardlaw and Pogue (1995)
87 suggested the climatic amelioration from the significantly cold, non- existent conodont
88 glacial waters to cool, conodont-bearing waters. The Nilwahan Group, indicates marginal
89 marine fluvial deposition, that was glacially controlled, especially in the Tobra and
90 Dandot formations, thus reflecting the combined effects of the overall Permian climatic

91 amelioration, i.e. the southward expansion of the warmer climatic zones and the
92 northward continental movement into the milder climatic zones (Wardlaw and Pogue,
93 1995).

94 The overlying Zaluch Group is a well-exposed succession of the shallow marine to inter-
95 tidal carbonate facies of the Amb, Wargal and Chhidru formations (Fig. 2), that were
96 deposited when a large carbonate platform developed on the Gondwana continental crust
97 close to the Indian Shield (Mertmann, 2003). The contrast between the largely cool
98 climate continental “Gondwanan” deposits of the Nilawahen Group and the richly
99 fossiliferous, marine “Tethyan” deposits of the Zaluch Group appears great, showing
100 gradual change from the marginally glacial sedimentation to marginally warm water
101 deposition (Wardlaw and Pogue, 1995).

102 Thus the Permian succession of Pakistan provides information on the changing
103 palaeoclimate and palaeogeography of the region, comprising a record of warming as the
104 Carboniferous-Permian glaciations waned and northern Gondwana drifted northwards
105 (Stephenson et al., 2007; 2008).

106 Biostratigraphical dating has suggested a range of ages for Salt Range units. The
107 brachiopods of the Amb Formation were considered by Waterhouse (1976; 1981) to be
108 Late Baigendzinian (upper Artinskian). The fusulinid fauna from the formation was also
109 assigned as Baigendzinian by the Pakistani-Japanese Research Group (1985). Pollen and
110 spores investigated by Balme (1970), suggested an Artinskian age. The most recent age
111 determination is that of Wardlaw and Pogue (1995), who used conodonts, which indicate
112 a Wordian (middle Guadalupian, Kazanian) age.

113 The Pakistani-Japanese Research Group (1985) reported a number of megafossils from
114 the Wargal Formation, including commonly occurring brachiopods. Gastropods and
115 corals have also been reported, along with common occurrences of small foraminifera
116 throughout the formation. The fusulinids and foraminifers indicate a late Murghabian,
117 Tethyan equivalent of Capitanian, through early Dzhulfian, Tethyan equivalent of
118 Wuchiapingian. The brachiopods indicate an early Dzhulfian (Wuchiapingian) age
119 (Pakistani-Japanese Research Group, 1985). The conodonts have been assigned to
120 Capitanian through Wuchiapingian age by Wardlaw and Pogue (1995). The overlaying
121 Chhidru Formation is reported to include small foraminifera, along with brachiopods,
122 molluscan fossils, Bryozoa and rare ammonoids (*Cycolobus*). The conodonts are
123 commonly dominated by the near-shore *Hindeodus* (Wardlaw and Pogue, 1995). The
124 foraminifers from the formation support correlation with the Wachiaping Formation of
125 South China and thus it has been assigned to the late Dzhulfian (Wuchiapingian) by the
126 Pakistani-Japanese Research Group (1985).

127 Palynological study of the Permian succession is confined to the Salt Range; there are no
128 reports of palynology from the Trans-Indus ranges. Virkki (1946) and Venkatachala and
129 Kar (1966, 1968) studied samples from a horizon 20-25 feet above the Tobra Formation
130 (see Balme, 1970). Balme (in Teichert, 1967) also described assemblages from the Tobra
131 Formation at Zaluch Nala, eastern Salt Range and assigned them to the Permian
132 (Teichert, 1967). Kemp (1975) examined two samples from the Tobra Formation at
133 Zaluch Nala and reported the presence of *Brevitriletes* sp. cf. *B. unicus*, *Lophotriletes* sp.
134 cf. *L. scotinus*, *Horriditriletes*- *Lophotriletes* sp. *Potonieisporites neglectus*,
135 *Dentatisporites* sp. along with acritarchs, referable to the genus *Cymatiosphaera*, while

136 Khan et al., (2001) reported Tobra Formation assemblages from Nilawahan Gorge,
137 central Salt Range.

138 There is no palynological work on the Sardhai formation, but Balme (1970) made a
139 detailed taxonomic survey of the carbonate dominated succession of the overlying Amb,
140 Wargal and Chhidru formations. He recovered pollen and spores from the plant-bearing
141 horizons of the Amb Formation at three localities in the Salt Range: Zaluch Nala, Dhodha
142 Wahan, and near Warchha Water Tank. Trilete spores represented were: *Acanthotriletes*
143 *tereteangulatus*, *Camptotriletes warchianus*, *Leiotriletes cf. adnatus*, *Lophotriletes*
144 *novicus* and *Verrucosisporites* sp. cf. *V. planiverrucatus*, while *Reticuloidosporites*
145 *warchianus* was the only monolete spore taxon. The monosaccate pollen included
146 *Plicatipollenites indicus* and *Potonieisporites novicus*. Balme (1970) reported a high
147 diversity of bisaccate pollen including the taeniate taxa *Corisaccites alutas*,
148 *Guttulapollenites hannonicus*, *Hamiapollenites insolitus*, *Lueckisporites singhii*,
149 *Protohaploxylinus limpidus*, *P. goraiensis*, *P. diagonalis*, *P. varius*, *Striatopodocarpites*
150 *cancellatus*, *S. rarus* and *S. pantii*. Non-taeniate bisaccate taxa included *Alisporites*
151 *tenuicarpus*, *Falcisporites nuthallensis*, *Pinuspollenites thoracatus*, *Sulcatisporites*
152 *ovatus*, *S. nilssoni* and *Vitreisporites pallidus*. Balme (1970) noticed similarities between
153 the palynological assemblages from the Zaluch Group of Pakistan with those of the
154 Madagascar, Australia and the then USSR.

155 For the present study, two samples (Figs. 3 and 4) were collected from a 22 meter-thick
156 exposure of the Sardhai Formation in the Khisor Range at N32° 11' 52.1" E 70° 59' 18.0".
157 One more sample, 25 meters above base of the Sardhai Formation was collected from an

158 approximately 30 meter-thick exposure of the same formation at Zaluch Nala, Salt Range
159 at N32° 46' 58.4" E 71° 38' 49.4".

160 Assuming that the Salt and Khisor ranges comprise a potential Permian reference section
161 for the South Tethys, because of their good outcrop exposures, palyniferous lithologies
162 and the presence of abundant marine fauna for age calibration, yet the aim of this paper is
163 to only document and describe the assemblages of the Sardhai Formation and to correlate
164 them with the other reported assemblages from the southern Tethyan region including the
165 standard palynostratigraphic scheme of Arabia. Our long term intention would be to
166 develop such palynological succession and studies on the Tobra, Dandot and Warchha
167 formations.

168

169 **Materials and methods**

170 The preparation of strew mounts for palynological analysis involved established
171 procedures of crushing followed by hydrofluoric and hydrochloric acid treatments (Wood
172 et al., 1996). Post-hydrofluoric acid organic residues were oxidized with Schulze's
173 Solution and dilute nitric acid. The photography was done with a DP11 Olympus digital
174 camera mounted on a Zeiss Universal microscope. The samples collected and
175 palynological slides prepared are housed at the laboratories of the British Geological
176 Survey, UK.

177

178 **Description of assemblages**

179 The yield of the samples was mainly poor, however it was possible in most cases to count
180 at least two hundred specimens per slide. Thirty five taxa were identified from these
181 samples, including the palynostratigraphically important *Camptotriletes warchianus*,
182 *Florinites ?balmei* and *Lueckisporites virkkiae*. A list of all palynomorph species with
183 author citation is given in Appendix 2 and selected taxa are displayed in [Plates I and II](#).

184 The quantitative character of assemblages from the Sardhai Formation at Zaluch Nala and
185 Khisor Range is shown in [Tables 1 and 2](#).

186 **Correlation with Arabia**

187 The most extensively studied Tethyan Permian sections are those of Oman and Saudi
188 Arabia (Stephenson and Filatoff, 2000a,b; Stephenson, 2008). Stephenson et al., (2003)
189 and Stephenson (2006) established eight palynological biozones (OSPZ 1 to OSPZ6); and
190 OSPZ5 and OSPZ6 are considered Mid- and Late Permian in age. OSPZ5, associated
191 with the lower to middle parts of the Upper Gharif member in Oman, is dominated by
192 distally-taeniate bisaccate pollen including *Distriatites insolitus* and *Hamiapollenites*
193 *dettmannae*, but also contains *Densiopollenites indicus*, *Platysaccus cf. queenslandi*,
194 *Playfordiaspora cancellosa* and *Thymospora opaqua*.

195 The base of the succeeding biozone, OSPZ6, represents a considerable palynological
196 change because a number of taxa appear for the first time at this level, and because
197 diversity increases. The base of OSPZ6 is defined by the first occurrence of *Florinites*
198 *?balmei* which is usually very common. Other taxa of OSPZ6 include *Camptotriletes*
199 *warchianus*, *Pyramidosporites cyathodes* and *Protohaploxylinus uttingii* though many
200 taxa such as *Alisporites nuthallensis*, *Laevigatosporites callosus*, *Lueckisporites virkkiae*,

201 *Thymospora opaqua* and *Reduviasporonites chalastus* persist from OSPZ5 (Stephenson
202 et al., 2003; Stephenson, 2006, 2008).

203 The base of OSPZ6 occurs in the highest parts of the Upper Gharif member in Oman a
204 few meters below the base of the succeeding carbonate Khuff Formation in beds
205 sometimes referred to as the Khuff Transition beds (see Stephenson, 2006, 2008) and the
206 biozone extends into the Khuff Formation. In central Saudi Arabia, assemblages assigned
207 to OSPZ6 (i.e. containing *Florinites ?balmei*) also occur in clastic sedimentary rocks
208 below the base of the Khuff Formation.

209 The lower age limit of OSPZ6 is difficult to constrain since no independent
210 palaeontological data are available from the clastic sedimentary rocks of the Upper
211 Gharif member. In Oman the base of OSPZ6 occurs consistently a few metres below the
212 base of the carbonate Khuff Formation (see Stephenson, 2006, 2008) and the lower beds
213 of the Khuff Formation are dated as early Wordian in age (Angiolini et al., 2003). Since
214 no significant hiatus is present between the Upper Gharif member and the lower Khuff
215 Formation the lower limit of the age of OSPZ6 is likely to be Wordian. The upper age
216 limit of OSPZ6 in Oman and Saudi Arabia is yet to be defined but the assemblages that
217 characterise it are not known to extend into the Triassic.

218 The presence in the Sardhai Formation of *Florinites ?balmei* in addition to *Alisporites*
219 *nuthallensis*, *Corisaccites alutas*, *Camptotriletes warchianus*, *Laevigatosporites callosus*
220 and *Thymospora opaqua* suggests a correlation with the OSPZ6 Biozone of Arabia. The
221 Amb Formation above the Sardhai Formation is similar palaeontologically and
222 lithologically to the Arabian Khuff Formation and is widely considered to be its temporal

223 and sedimentological equivalent (see Angiolini and Bucher, 1999) having been formed
224 by essentially the same marine transgression associated with neo-Tethyan sea floor
225 spreading (Angiolini et al., 2003; Mertmann, 2003). The evidence thus suggests that the
226 Sardhai Formation correlates with the immediate pre-carbonate clastic sedimentary rocks
227 of Arabia, including the basal Khuff clastics and the Khuff transition beds. In the light of
228 this correlation, a Wordian age is tentatively suggested for the Sardhai Formation.

229

230 **Tethyan correlation**

231 Recently Stolle (pers. comm) investigated the Permian Kas and Gomaniibrik formations
232 in southeast Turkey. The Kas Formation has been dated as Wordian by foraminifera
233 (Stolle, pers. comm.) The assemblages in the Kas Formation are dominated by spores,
234 particularly monolete taxa, including *Punctatisporites* spp., *Spinospores* sp., *Torispora*
235 spp., and *Thymospora opaqua*, but also contain common *Camptotriletes warchianus*,
236 *Distriatites insolitus*, *Florinites ?balmei* (up to 23% of assemblages) and *Hamiapollenites*
237 *dettmannae*. Stolle (pers. comm.) correlated the Kas Formation assemblages with OSPZ6,
238 and noted similarities with northern Iraqi subsurface assemblages described by Nader et
239 al., (1993) and Singh (1964). Stolle's figure 5 illustrates a correlation of OSPZ6
240 assemblages across the Tethyan region, including Turkey, Iraq, Saudi Arabia and Oman
241 and this chart is here modified to include the Salt and Khisor ranges Sardhai Formation
242 OSPZ6 assemblages (Fig. 5).

243

244 **Palaeogeographic distribution of *Florinites ?balmei***

245 This correlation shows that the distinctive pollen *Florinites ?balmei* is present in
246 approximately coeval rocks in an area of the southern Neotethys which is now
247 represented by southeast Turkey and northern Iraq. In addition *Florinites ?balmei* has
248 recently been described from the basal Khuff clastics in Kuwait (Tanoli et al., 2008) and
249 is known to occur in the same unit in the United Arab Emirates and Qatar (BGS
250 unpublished reports; Fig. 6). Its occurrence in the Salt and Khisor ranges and apparent
251 absence from Middle Permian rocks elsewhere in Gondwana, Euramerica and Cathaysia
252 suggests that the plant that produced *Florinites ?balmei* had a rather restricted
253 palaeogeographic distribution along the palaeotropical coast of the Tethys Ocean (Fig. 6).
254 It also tends to support the palaeogeographic reconstructions of Ricou in Dercourt et al.,
255 (1993), Ziegler et al., (1998) and Gaetani et al., (2000) showing the Salt Range area in
256 contiguity with the southern part of the Arabian Plate.

257 The complex palaeogeography and palaeotectonics of the Tethyan margin from the Early
258 to Mid Permian has been discussed by amongst others Sengör (1979), Ricou (in Dercourt
259 et al., 1993), Ziegler et al., (1998), Gaetani et al., (2000) and Angiolini (2001).

260 Angiolini's (2001) reconstruction of the Wordian shows the Mega Lhasa Block or
261 Cimmerian blocks (comprising Iran, Afghanistan, Karakorum and Sibumasu, Thailand)
262 in contiguity and in relative proximity to the Gondwanan margin, however the form and
263 position of the Mega Lhasa Block is generally considered uncertain

264 (Gaetani, 1997; Muttoni et al., personal communication), thus further work to establish
265 whether *Florinites ?balmei* occurs in the Mega Lhasa Block; especially Thailand might
266 shed more light on such reconstructions.

267 The well known Oman Gharif palaeoflora (e.g. Broutin et al., 1995; Berthelin et al.,
268 2003) was described from the uppermost Gharif Formation in the Huqf area in interior
269 Oman, and is believed to consist of a mixture of Gondwanan, Cathaysian and
270 Euramerican fossil plant taxa. Plant taxa of the Permian Cathaysian Paleokingdom
271 present in Oman were considered by Berthelin et al., (2003) to indicate a close
272 relationship between the Neotethys realm and south China. Further work should aim to
273 establish whether *Florinites ?balmei* occurs in the Cathaysian Paleokingdom.

274

275 **Conclusion**

276 The presence of *Florinites ?balmei* together with other stratigraphically important taxa,
277 e.g. *Camptotriletes warchianus*, suggest that the Sardhai Formation correlates with the
278 Khuff transition beds of Oman and the basal Khuff clastics of central Saudi Arabia and
279 can be likewise assigned to the Arabian OSPZ6 biozone, indicating a tentative Wordian
280 age. Overall the Salt Range Permian succession is also similar in lithological character to
281 that of the Arabian Peninsula: both have successions of glacial diamictites at the base,
282 overlain by the red beds, followed by distinctive dark shale-sandstone interbeds and
283 conclude with thick limestones.

284 This work has also shown that monosaccate pollen grain *Florinites ?balmei* had a limited
285 palaeogeographic distribution in the Mid-Permian across most of the southern Tethys and
286 Arabia, whereas it is apparently not reported elsewhere in Gondwana and Euramerica. If
287 this distribution can be more precisely delineated in regions in the wider Middle East and
288 parts of present day southeast Asia, it would help reconstruct this region's complex
289 palaeogeography and tectonics.

290

291

292

293 **Appendix 1**

294 **Systematic Palynology**

295 *Florinites* Schopf, Wilson and Bentall 1944

296 Type Species: *Florinites pellucidus* (Wilson and Coe) Wilson, 1958.

297 ***Florinites ?balmei* Stephenson and Filatoff 2000**

298 Plate I, figures 1-15.

299 *Florinites ?balmei* Stephenson and Filatoff 2000: plate 2, figs j-m, p. 208-212

300 *Florinites millotti* Butterworth and Williams 1954 – Nader et al., 1993: plate 13, figures

301 7-8 [no description].

302 *Florinites millotti* Butterworth and Williams 1954 - Akyol, 1975: plate 9, figures 12-16
303 [no description].

304 *Florinipollenites millotti* (Butterworth and Williams) Coquel 1966 - Agrali and Akyol,
305 1967: plate 8, figures 21-24 [*Florinipollenites* Laveine 1965 is an obligate junior
306 synonym of *Florinites* Schopf, Wilson and Bentall 1944].

307 *Description:* Pollen, monosaccate, bilaterally symmetrical; amb oval. Corpus almost
308 imperceptible, though its presence is suggested by a narrow, oval fold structure in the
309 saccus. Long axis of corpus parallel to the long axis of the grain; diameter of the corpus
310 approximately half that of the grain overall. The detachment of the saccus from the
311 corpus is imperceptible. Saccus coarsely infrareticulate; brochi 1-2 μ m in diameter, muri
312 width <1 μ m. equatorial margin of the saccus is slightly thickened in some specimens.

313

314 **Appendix 2. List of taxa recorded**

315 *Alisporites* sp.

316 *Alisporites indarraensis* Segroves, 1969

317 *Alisporites* cf. *nuthallensis* Clarke, 1965

318 *Barakarites rotatus* (Balme and Hennelly) Bharadwaj and Tiwari, 1964

319 *Brevitriletes parmatus* (Balme and Hennelly) Backhouse, 1991

320 *Brevitriletes* sp.

321 *Camptotriletes warchianus* Balme, 1970

- 322 *Corisaccites alutas* Venkatachala and Kar, 1966
- 323 *Cedripites* sp.
- 324 *Cannanoropollis janakii* Potonié and Sah, 1960
- 325 *Complexisporites polymorphus* Jizba, 1962
- 326 *Distriatites* sp.
- 327 *Florinites ?balmei* Stephenson & Filatoff, 2000
- 328 *Guttulapollenites hannonicus* Goubin, 1965
- 329 *Hamiapollenites* sp.
- 330 *Hamiapollenites dettmannae* Segroves, 1969
- 331 *Hamiapollenites karrooensis* (Hart 1963) Hart, 1964
- 332 *Horriditriletes tereteangulatus* (Balme and Hennelly) Backhouse, 1991
- 333 *Kingiacolpites subcircularis* Tiwari and Moiz, 1971
- 334 *Laevigatosporites callosus* Balme, 1970
- 335 *Lueckisporites virkkiae* Potonié and Klaus emended Clarke, 1965
- 336 *Lundbladispota* sp.
- 337 *Punctatisporites* spp.
- 338 *Protohaploxypinus uttingii* Stephenson and Filatoff, 2000

- 339 *Protohaploxypinus* sp.
- 340 *Plicatipollenites* sp.
- 341 *Potonieisporites* sp.
- 342 *Retusotriletes* sp.
- 343 *Striatopodocarpites cancellatus* (Balme and Hennelly) Bharadwaj, 1962
- 344 *Striatopodocarpites fusus* (Balme & Hennelly) Potonié, 1958
- 345 *Strotersporites indicus* Tiwari, 1965
- 346 *Spelaeotriletes* sp.
- 347 *Thymospora opaqua* Singh, 1964
- 348 *Taeniasporites* sp.
- 349 *Verrucosisporites andersonii* Backhouse, 1988

350

351 **References**

- 352 Agrali, B., Akyol, E., 1967. Étude palynologique de charbons de Hazro et considérations
353 sur L'âge des horizons lacustres de Permo-Carbonifère. Mineral Res. Expl. Inst. Turkey,
354 Bull. (Foreign Edition), 68, 1-26.

355 Akyol, E., 1975. Palynologie du Permien inférieur de Sariz (Kayseri) et de Pamucak
356 Yaylasi (Antalya-Turquie) et contamination jurassique observée, due aux ruisseaux
357 “Pamucak” et “Göynük”. Pollen et Spores, Muséum National d’Histoire Naturelle, Paris
358 17, 141-179.

359 Alam, I., 2008. Structural and Stratigraphic Framework of the Marwat-Khisor Ranges,
360 N.W.F.P., Pakistan. Unpublished Ph.D thesis. National Centre of Excellence in Geology,
361 University of Peshawar, Pakistan.

362 Alam, G.S., Mashhadi, S. T. A., Khan, M. A., 1987. Interim report on Permian coal
363 exploration in Punjab Plains (drilling at Ashaba, District Jhang). Geol. Surv. Pakistan inf.
364 Release 286, 30.

365 Angiolini, L., Bucher, H., 1999. Taxonomy and quantitative biochronology of
366 Guadalupian brachiopods from the Khuff Formation, southeastern Oman. Geobios 32,
367 665-699.

368 Angiolini, L., 2001. Lower and Middle Permian brachiopods from Oman and Peri-
369 Gondwanan palaeogeographical reconstructions. In: Brunton, C. H. C., Cocks, L. R. M.,
370 Long, S. L. (Eds.), Brachiopods past and present. Systematics Association, Special
371 Volume 63, pp. 352-362.

372 Angiolini, L., Balini, M., Garzanti, E., Nicora, A., Tintori, A., Crasquin, S., Muttoni, G.,
373 2003. Permian climatic and palaeogeographic changes in Northern Gondwana: the Khuff
374 Formation of Interior Oman. Palaeogeography, Palaeoclimatology, Palaeoecology 191,
375 269 -300.

376 Balme, B. E., 1970. Palynology of Permian and Triassic strata in the Salt Range and
377 Surghar Range, West Pakistan. In: Kummel, B., Teichert, C. (Eds.), Stratigraphic
378 Boundary Problems: Permian and Triassic of West Pakistan. University Press of Kansas,
379 Department of Geology Special Publication 4, pp. 305-453.

380 Berthelin, M., Broutin, J., Kerp, H., Crasquin-Soleau, S., Platel, J.-P., Roger, J., 2003.
381 The Oman Gharif mixed paleoflora: a useful tool for testing Permian Pangea
382 reconstructions. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 196, 85-98.

383 Broutin, J., Roger, J., Platel, J.-P., Angiolini, L., Baud, A., Bucher, H., Marcoux, J., Al
384 Hasmi, H., 1995. The Permian Pangea. Phylogeographic implications of new
385 paleontological discoveries in Oman (Arabian Peninsula). *C. R. Acad. Sci. Paris, Série*
386 *Ila* 321, 1069-1086.

387 Butterworth, M. A., Williams, R. W., 1954. Descriptions of nine species of small spores
388 from the British Coal Measures. *The Annals and Magazine of Natural History* 7, 753-
389 764.

390 Dercourt, J., Ricou, L. E., Vrielinck, B., 1993. Atlas Tethys Palaeoenvironmental Maps.
391 Explanatory Notes. (Paris Ed BEICIP), pp. 1-307.

392 Diener, C., 1912. The Trias of the Himalayas. *Geol. Survey India, Memoir* 36, pp. 202-
393 360.

394 Gaetani, M., 1997. The Karakoram Block in central Asia, from Ordovician to Cretaceous.
395 *Sedimentary Geology* 109, 339-359.

396 Gaetani, M., et al., 2000. Wordian. In : Dercourt, J., Gaetani, M., et al. (Eds.), Atlas Peri-
397 Tethys, Palaeogeographical Maps CCGM/CGMW, Paris, map 3.

398 Gee, E. R., 1980. Salt Range Maps. Geological Survey of Pakistan, Lahore.

399 Gee, E. R., 1989. Overview of the geology and structure of the Salt Range, with
400 observation on related areas of northern Pakistan. In: Malinconico, L. L., Lillie, R. J.
401 (Eds.), Tectonics of the Western Himalayas. Geological Society of America Special
402 paper 232, 95-112.

403 Ghauri, A. A. K., Anwar, M., Kamel, K. S., Issa, I. M., 1977. The study of nature and
404 origin of the Tobra Formation in the eastern part of the Punjab Salt Range. Geological
405 Bulletin University of Peshawar 9, 67-78.

406 Glenister, B. F., Furnish, W. M., 1970. Permian Ammonoid *Cyclolobus* from the Salt
407 Range, West Pakistan. In: Kummel, B., Teichert, C. (Eds.), Stratigraphic Boundary
408 Problems : Permian and Triassic of West Pakistan. University Press of Kansas,
409 Department of Geology Special Publication 4, pp. 153-175.

410 Grabau, A. W., 1931. The Permian of Mongolia. Amer. Museum Natural History, Natural
411 History of Central Asia 4, pp. 665.

412 Grant, R. E., 1970. Brachiopods from Permian -Triassic Boundary Beds and Age of
413 Chhidru Formation, West Pakistan. In: Kummel, B., Teichert, C. (Eds.), Stratigraphic
414 Boundary Problems: Permian and Triassic of West Pakistan. University Press of Kansas,
415 Department of Geology Special Publication 4, pp. 117-151.

416 Hussain, B. R., 1967. Saiyiduwali member, a new name for the lower part for the
417 Permian Amb Formation, West Pakistan: Pakistan Geol. Surv. Mem. Palaeont.
418 Pakistanica. Vol. 5, pp. 88-95.

419 Kemp, E. M., 1975. The palynology of Late Palaeozoic Glacial Deposits of
420 Gondwanaland. In: Campbell, K. S. W. (Eds.), Gondwana geology: Canberra. Australian
421 University Press. pp. 397-413.

422 Khan, Q, M., Sarfraz, A., Kamran, M., Fazli, R. K., 2001. Some monosaccate pollen from
423 the Tobra Formation of the Nilawahan Gorge, Central Salt Range, Pakistan. Geological
424 Bulletin Punjab University 36, 95-102.

425 Kummel, B., Teichert, C., 1970. Stratigraphy and Paleontology of the Permian-Triassic
426 Boundary Beds, Salt Range and Trans-Indus Ranges, West Pakistan. In: Kummel, B.,
427 Teichert, C. (Eds.), Stratigraphic Boundary Problems: Permian and Triassic of West
428 Pakistan. University Press of Kansas, Department of Geology Special Publication 4, pp.
429 2-110.

430 Kummel, B., 1970. Ammonoids from the Kathwai Member, Mianwali Formation, Salt
431 Range, West Pakistan. In: Kummel, B., Teichert, C. (Eds.), Stratigraphic Boundary
432 Problems: Permian and Triassic of West Pakistan. University Press of Kansas,
433 Department of Geology Special Publication 4, pp. 177-192.

434 Mertmann, D., 1999. Das marine Perm in der Salt Range und in den Trans Indus Ranges,
435 Pakistan. Berliner Geowissensch. Abh. A. 204, 1– 94.

436 Mertmann, D., 2003. Evolution of the marine Permian carbonate platform in the Salt
437 Range (Pakistan), *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 191, 373–384.

438 Nader, A. D., Khalaf, F.H., Hadid, A. A., 1993. Palynology of the Permo -Triassic
439 boundary in Borehole Mityaha-1, south west Mosul City – Iraq. *Mu'tah Journal for*
440 *Research and Studies, Mu'tah University, Jordan* 8, 223-280.

441 Noetling, F., 1901. Beiträge zur Geologie der Salt Range, insbesondere der permischen
442 und triassischen Ablagerungen: *Neues Jahrb. Mineralogie, Geologie, Palaeontologie* 14,
443 369-471.

444 Pakistani-Japanese Research Group. 1985. Permian and Triassic in the Salt Range and
445 Surghar Range, Pakistan. In: Nakazawa, K., Dickins, J. M. (Eds.), *The Tethys- Her*
446 *Paleogeography and Paleobiogeography from Paleozoic to Mesozoic.* Tokai University
447 Press, Tokyo. pp. 221-312.

448 Pascoe, E. H., 1959. A manual of the geology of India and Burma. India Govt. Press
449 Calcutta. II 484-1338.

450 Reed, F. R. C., 1936. Some fossils from the *Eurydesma* and *Conularia* beds (Punjabian)
451 of the Salt Range. *India Geological Survey Memoirs Palaeont. Indica New Series* 1, 1-
452 36.

453 Rowell, A.J., 1970. *Lingula* from the Basal Triassic, Kathwai Member, Mianwali
454 Formation, Salt Range and Surghar Range, West Pakistan. In: Kummel, B., Teichert, C.
455 (Eds.), *Stratigraphic Boundary Problems: Permian and Triassic of West Pakistan.*
456 University Press of Kansas, Department of Geology Special Publication 4, pp. 111-116.

457 Sarjeant, W. A. S., 1970. Acritarchs and Tasmanitids from the Chdidru Formation,
458 Uppermost Permian of West Pakistan. In: Kummel, B., Teichert, C. (Eds.), *Stratigraphic*

459 Boundary Problems: Permian and Triassic of West Pakistan. University Press of Kansas,
460 Department of Geology Special Publication 4, pp. 277-304.

461 Sengör, A. M. C., 1979. Mid-Mesozoic closure of Permo-Triassic Tethys and its
462 implications. *Nature* 279, 590-593.

463 Shah, S. M. I., 1977. Stratigraphy of Pakistan. Geological Survey of Pakistan Memoir 12.

464 Singh, H.P., 1964. A miospore assemblage from the Permian of Iraq. *Palaeontology*,
465 *Palaeontological Association, London* 7, 240-265.

466 Sohail, W., Shakeel U. R., Sadaf, B., 2004. Structural Analysis of Central Khisor Range,
467 North west of Dakki, D.I. Khan NWFP Pakistan. unpublished MSc. Thesis. Department
468 of Geology, University of Peshawar, Pakistan.

469 Sohn, I. G., 1970. Early Triassic Marin Ostracodes from the Salt Range and Surghar
470 Range, West Pakistan. In: Kummel, B., Teichert, C. (Eds.), *Stratigraphic Boundary*
471 *Problems: Permian and Triassic of West Pakistan*. University Press of Kansas,
472 Department of Geology Special Publication 4, pp. 193-206.

473 Stephenson, M. H., Filatoff, J., 2000a. Correlation of Carboniferous-Permian
474 Palynological Assemblages from Oman and Saudi Arabia. *GeoArabia Special*
475 *Publication- Stratigraphic Palynology of the Palaeozoic of Saudi Arabia*. Gulf Petrolink,
476 Bahrain 1, 168-191.

477 Stephenson, M. H., Filatoff, J., 2000b. Description and Correlation of Late Permian
478 Palynological Assemblages from the Khuff Formation, Saudi Arabia and Evidence for

479 the Duration of the pre-Khuff Hiatus. *GeoArabia Special Publication- Stratigraphic*
480 *Palynology of the Palaeozoic of Saudi Arabia*. Gulf PetroLink, Bahrain 1, 192-215.

481 Stephenson, M. H., Osterloff, P.L., Filatoff, J., 2003. Palynological biozonation of the
482 Permian of Oman and Saudi Arabia; progress and challenges. *GeoArabia* 8, 467-496.

483 Stephenson, M. H., 2006. Stratigraphic Note: Update of the standard Arabian Permian
484 palynological biozonation; definition and description of the OSPZ5 and 6. *GeoArabia* 11,
485 173-178.

486 Stephenson, M. H., Angiolini, L., Leng, M. J., 2007. The Early Permian fossil record of
487 Gondwana and its relationship to deglaciation: a review In: Williams, M., Heywood, A.
488 (Eds.), *Climate Change in Deep Time*. Special Publication of the Geological Society,
489 London. pp. 45-85.

490 Stephenson, M. H., Angiolini, L., Leng, M.J., Brewer, T.S., Berra, F., Jadoul, F.,
491 Gambacorta, G., Verna, V., Al Beloushi, B., 2008. Abrupt environmental and climatic
492 change during the deposition of the Early Permian Haushi limestone, Oman.
493 *Palaeogeography, Palaeoclimatology, Palaeoecology* 270, 1- 18.

494 Stephenson, M. H., 2008. Spores and Pollen from the Middle and Upper Gharif members
495 (Permian) of Oman. *Palynology* 32, 157-182.

496 Stolle, E., (pers. comm.). Regional Permian Palynological correlation: southeast Turkey –
497 northern Iraq.

498 Sultan, S. A., 2004. Mineralogical investigations of the phosphatic nodules from
499 Warchha sandstone, salt range with emphasis on replacement process. Geological
500 Bulletin University of Peshawar 37, 65-75.

501 Sweet, W. C., 1970. Uppermost Permian and Lower Triassic Conodonts of the Salt
502 Range and Trans-Indus Range, West Pakistan. In: Kummel, B., Teichert, C. (Eds.),
503 Stratigraphic Boundary Problem between Permian and Triassic of West Pakistan.
504 University Press of Kansas, Department of Geology Special Publication 4, pp. 207-275.

505 Tanoli, S. K., Husain, R., Sajer, A. A., 2008. Facies in the Unayzah Formation and the
506 Basal Khuff Clastics in subsurface, northern Kuwait. GeoArabia 13, 15-40.

507 Teichert, C., 1967. Nature of the Permian glacial record, Salt Range and Khisor Range,
508 West Pakistan. N. Jb. Geol. Paläont. Abh. 129, 167-184.

509 Venkatachala, B. S., Kar, R. K., 1966. *Corisaccites* gen. nov., a new saccate pollen genus
510 from the Permian of the Salt Range, West Pakistan. Palaeobotanist 15, 107-109.

511 Venkatachala, B. S., Kar, R. K., 1968. Palynology of the Kathwai Shales, Salt Range,
512 West Pakistan. 1. Shales 25ft above the Talchir Boulder Bed. Palaeobotanist 16, 156-166.

513 Virkki, C., 1946. Spores from the Lower Gondwanas of India and Australia. Proc. Indian
514 Acad. Sc. India 15, pp. 93-176.

515 Waagen, W, H., 1882-1885. Salt Range fossils. Part 4: Brachiopoda. Palaeonologia
516 Indica 13, pp. 329- 770.

517 Wardlaw, B. R., Pogue, K. R., 1995. The Permian of Pakistan. In: Scholle, P. A., Peryt,
518 T. M., Ulmer-Scholle, P. M. (Eds.), The Permian of Northern Pangea vol. 1
519 Palaeogeography, Palaeoclimates, Stratigraphy. Springer Verlag. pp. 215-224.

520 Waterhouse, J.B., 1976. World correlations for Permian Marine Faunas. Univ.
521 Queensland papers, Dept. Geol 7(2) 232p.

522 Waterhouse, J. B., 1981. Age of the Rat Buri Limestone of Southern Thailand. In:
523 Waterhouse , J. B., Pitakpaivan, K., Mantajit, N. (Eds.), The Permian stratigraphy and
524 palaeontology of the Southern Thailand. Dep. Minerals Res. Geol. Surv. Mem. 4, pp. 2-
525 42.

526 Wood, G. D., Gabriel, A. M., Lawson, J. C., 1996. Palynological techniques-processing
527 and microscopy. In: Jansonius, J., McGregor, D.C. (Eds.), Palynology: Principles and
528 Applications. AASP Foundation. vol. 1, pp. 29-50.

529 Wynne, A. B., 1878. On the geology of the Salt Range in the Punjab: Geol. Surv. India,
530 Mem. vol. 14, pp. 1-313.

531 Ziegler, A. M., Gibbs, M. T., Hulver, M. L., 1998. A mini-atlas of oceanic water masses
532 in the Permian period. Proceedings of the Royal Society of Victoria 110, pp. 323–343.

533

534

535 **Figure Captions**

536 Figure 1. Location map of the study area. Showing Salt and Khisor ranges of Pakistan
537 (Modified after Gee 1980, 1989).

538 Figure 2. Simplified Permian stratigraphy of the Salt Range and central part of Khisor
539 Range. (Modified after Gee, 1989; Sohail et al., 2004).

540 Figure 3. Vertical beds of the Sardhai Formation underlain by red beds of the Warchha
541 Formation (to the right in the photo). Samples studied were from levels 1.8m and 16.2m
542 above base of the Sardhai Formation. Rock outcrop to the left is 5m high.

543 Figure 4. Measured section of the Sardhai Formation in the Khisor Range Pakistan.

544 Plate I. The monosaccate pollen grain *Florinites ?balmei* identified from Sardhai
545 Formation Khisor Range, Pakistan. The slide number followed by the England finder
546 coordinates are given as follows,

547 (1). MPA-57533. W 12/3. (2). MPA-57533. Y11. (3). MPA-57533. W25. (4). MPA-
548 57533. S19. (5). MPA-57533. V8/1. (6). MPA-57533. V29/1. (7). MPA-57533. V33/2.
549 (8). MPA-57533. W 20/3. (9). MPA-57533. T10/3. (10). MPA-57533. S17/4. (11). MPA-
550 57533. U3/3. (12). MPA-57533. W27/4. (13). MPA-57533. L 29/3. (14). MPA-57533.
551 O12. (15). MPA-57533. P5.

552 Plate II. Palynomorphs from the Sardhai Formation, Pakistan. Each with slide number
553 followed by England finder coordinates is given as follows,

554 (1). *Lueckisporites virkkiae*, MPA-57528, T23/3 (proximal focus). (2). *Lueckisporites*
555 *virkkiae*, MPA-57528, T23/3 (distal focus). (3). *Corisaccites alutas*, MPA-57528, Q33/3.

556 (4). *Striatopodocarpites fusus*, MPA-57528, N24. (5). *Camptotriletes warchianus*, MPA-
557 57528, S21. (6). *Protohaploxypinus uttingii*, MPA-57528, S35. (7). *Alisporites*
558 *nuthallensis*, MPA-57528, W32/1. (8). *Guttulapollenites hannonicus*, MPA-57528,
559 X29/1. (9). *Taeniaesporites* sp, MPA-57528, D31/4. (10). *Thymospora opaqua*, MPA-
560 57528, S22. (11). *Laevigatosporites callosus*, MPA-57528, U4/1. (12). *Lundbladispora*
561 sp., MPA-57528, R4/1. (13). *Alisporites nuthallensis*, MPA-57528, Q19/2. (14).
562 *Protohaploxypinus* sp., MPA-57528, Q19/2. (15). *Guttulapollenites hannonicus*, MPA-
563 57528, E17.

564 Table 1. Percentage abundance of taxa, Zaluch Nala. Data from sample 25m (MPA-
565 57528) above base of the Sardhai Formation.

566 Table 2. Percentage abundance of taxa, Khisor Range. Data from sample 1.8m (MPA-
567 57533) and 16.2m (MPA-57532) above base of the Sardhai Formation.

568 Figure 5. Correlation of the OSPZ6 biozone between Southeast Turkey, northern Iraq,
569 central Saudi Arabia, Oman and Pakistan (Modified after Stolle, pers. comm.).

570 Figure 6. The mid-Permian (Roadian-Wordian) continental configuration. Solid circles
571 indicate the location of *Florinites ?balmei* across, 1- Oman, 2- Salt Range, 3- UAE, 4-
572 Kuwait, 5- Saudi Arab, 6- Qatar and 7- southeast Turkey (Modified after Angiolini et al.,
573 2001).

574


575

576

577

578

579 Figure1



593

594

595 Figure2

596

597

598

599

600

601


602

603

604

605

606



607

608

609

610

611

612 Figure3



619

620

621

622

623

624

625

626

627

628

629 Figure4

630

631

632

633

634


635

636

637

638

639



640

641


642













643

644

645

646 Plate I





648

649

650

651 Table 1

Taxon	% age Abundance (MPA 57528)
<i>Alisporites indarraensis</i>	15
<i>Alisporites nuthallensis</i>	1
<i>Brevitriletes parmatus</i>	2
<i>Barakarites rotatus</i>	2
<i>Cannanoropollis janakii</i>	1
<i>Corisaccites alutas</i>	14
<i>Cedripites</i> sp.	1.5
<i>Complexisporites polymorphus</i>	0.5
<i>Camptotriletes warchianus</i>	1
<i>Distriatites</i> sp.	3
<i>Florinites ?balmei</i>	0.5
<i>Guttulapollenites hannonicus</i>	2.5
<i>Hamiapollenites</i> sp.	0.5
<i>Horriditriletes tereteangulatus</i>	0.5
<i>Kingiacolpites subcircularis</i>	4.5
<i>Lundbladispota</i> sp.	1
<i>Laevigatosporites callosus</i>	3
<i>Lueckisporites virkkiae</i>	1
<i>Plicatipollenites</i> sp.	0.5
<i>Protohaploxylinus</i> sp.	1
<i>Protohaploxylinus uttingii</i>	0.5
<i>Punctatisporites</i> spp.	3
<i>Potonieisporites</i> sp.	2
<i>Retusotriletes</i> sp.	0.5
<i>Striatopodocarpites cancellatus</i>	0.5
<i>Striatopodocarpites fusus</i>	1
<i>Strotersporites indicus</i>	1
<i>Spelaeotriletes</i> sp.	0.5
<i>Thymospora opaqua</i>	2
<i>Taeniaesporites</i> sp.	1
<i>Verrucosisporites andersonii</i>	3
Indeterminate monosaccate pollen	2
Indeterminate bisaccate pollens	27

664

665

666

667 Table2

Taxon	% age Abundance (MPA-57533)	% age Abundance (MPA-57532)
<i>Alisporites indarraensis</i>	3.5	22.5
<i>Alisporites nuthallensis</i>	1	
<i>Alisporites</i> sp.	3	
<i>Brevitriletes</i> sp.		0.5
<i>Corisaccites alutas</i>	1	1
<i>Distriatites</i> sp.		0.5
<i>Florinites ?balmei</i>	60	1.5
<i>Hamiapollenites dettmannae</i>	0.5	
<i>Hamiapollenites karrooensis</i>		0.5
<i>Kingiacolpites subcircularis</i>		0.5
<i>Laevigatosporites callosus</i>	22	1
<i>Punctatisporites</i> spp.	3.5	
<i>Protohaploxylinus uttingii</i>		55.5
<i>Strotersporites indicus</i>		0.5
<i>Thymospora opaqua</i>	2	2
<i>Verrucosisporites andersonii</i>	1	
Indeterminate monosaccate pollen	1.5	
Indeterminate bisaccate pollens	1	14

675

676

677

678


679

680

681

682

683 Figure5

Chronostratigraphy		Palynological Biozonation	Lithostratigraphy				
			Southeast Turkey	North Iraq	Central Saudi Arabia	Oman	Pakistan
Middle Permian Guadalupian	Capitanian	OSPZ6	Gomaniibrik Formation (part) B	Chia Zairi Formation (part) Zinner member	Khuff Formation (part)	?	?
	Wordian		Kas Formation A				
		OSPZ5		Clastics Ga'ara Formation (subsurface)		Khuff transition section Upper Gharif Member (part)	Sardhai Formation

684

685

686

687

688

689

690

691

692

693

694

695

696 Figure6

