

A Practical Guide to Analysing the Force-Time Curve of Isometric Tasks in Excel

AUTHORS:

Shyam Chavda – MSc, PGCHE, CSCS, ASCC, CES¹, Anthony N Turner – PhD, CSCS*D¹,
Paul Comfort – PhD, CSCS, ASCC², G Gregory Haff – PhD^{2,3}, Steve Williams – MSc,
PGCHE¹, Chris Bishop – MSc, PGCHE, ASCC¹ and Jason P Lake – PhD⁴.

INSTITUTION:

1. London Sports Institute, Middlesex University, Allianz Park, Greenlands Lane, NW4 1RL
2. Directorate of Sport, Exercise and Physiotherapy, University of Salford, Salford, Greater Manchester, UK
3. Centre for Exercise and Sports Science Research, Edith Cowan University, Joondalup, Australia.
4. Chichester Institute of Sport, University of Chichester, College Lane, Chichester, PO19 6PE.

CORRESPONDENCE:

Email: s.chavda@mdx.ac.uk

Tel No: (+44)20 8411 2854

Address: As per above

Shyam Chavda is a strength and conditioning coach and technical tutor at the London Sport Institute, Middlesex University. He is also the performance scientist for British Weightlifting and head coach at Middlesex University weightlifting club.

Anthony N Turner is the director of postgraduate programmes in sport at the London Sport Institute, Middlesex University.

Paul Comfort is a reader in Strength and Conditioning and the program leader for the Masters degree in Strength and Conditioning at the University of Salford.

G. Gregory Haff is a Professor of Strength and Conditioning and course coordinator of the MSc in Exercise Science (Strength and Conditioning) at Edith Cowan University. He is also an Honorary Professor in the Directorate of Sport, Exercise and Physiotherapy at the University of Salford.

Steve Williams is a lecturer in biomechanics at the London Sport Institute, Middlesex University.

Chris Bishop is a strength and conditioning coach at the London Sport Institute, Middlesex University, where he is also the programme leader for the MSc in Strength and Conditioning.

Jason Lake is a Reader in Sport and Exercise Biomechanics and program leader of the MSc in Strength and Conditioning in the Chichester Institute of Sport at the University of Chichester

A Practical Guide to Analysing the Force-Time Curve of Isometric Tasks in Excel

Abstract

Understanding force generating capabilities of athletes is an important facet of strength diagnostics. The utilisation of isometric tasks such as the isometric squat and isometric mid-thigh pull are therefore popular methods used to gain a deeper understanding as to what strength characteristics have changed over a given period. This article aims to provide information on how to understand and analyse the force time curve of isometric tasks in Microsoft Excel, thus providing practitioners an inexpensive and accessible alternative to readily available software on the market.

Keywords: force; Rate of force Development; Isometric Strength; excel

Introduction

Lower-body neuromuscular force production characteristics have previously been determined utilising isometric tasks such as the isometric mid-thigh pull (IMTP) (34, 36, 42-44, 49, 51), isometric squat (IS) (3, 12, 36, 39, 47), isometric leg press (7, 55), and isometric dynamometry (38, 41, 46). Relative to multi-joint isometric assessments, such as the IS and IMTP, isolating a single joint requires specialist equipment such as a dynamometer which in most cases will not be financially viable or practical. Although testing multi-joint isometrics requires a force plate and a specialised isometric rack, a variety of testing systems are available ranging from customised laboratory set ups (Figure 1a), to more portable and affordable set ups (Figure 1b). Previous literature has shown conflicting evidence regarding single joint isometrics and their

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

24 correlations to dynamic performance (15, 35, 38, 40, 53), suggesting that they do not simulate
25 body positions representative of dynamic tasks such as jumping (2, 54), and do not best
26 represent the transfer of forces through the kinetic chain and are therefore considered a poor
27 indicator of athletic performance (36). Conversely, the IS and IMTP are used more frequently
28 (12, 23, 26, 28, 36, 47, 49, 51), potentially due to affordability (relative to a dynamometer), set
29 up efficiency, and greater replication of multi-joint positions demonstrated within sporting
30 actions. Although inconsistencies in methodologies and analysis of the force time curve is
31 evident among the literature (14), variables assessed during multi-joint isometric assessments
32 still demonstrate stronger relationships to dynamic tasks in comparison to single-joint isometric
33 assessments (23, 24, 33, 34, 45, 47, 52), with less conflicting evidence reported (36). Therefore,
34 it is unsurprising that multi-joint isometric tests such as the IS and IMTP are a more popular
35 choice amongst practitioners and will therefore be the focus of this article.

36
37 **INSERT FIGURE 1a and b HERE NEXT TO EACHOTHER – LAB (a) and PORTABLE
38 (b) ISO RIG**

39
40 The IS and IMTP data have been measured using crane scales (50), single -axial load cells (28)
41 and more commonly force plates, that are positioned underneath a fixed bar or within a custom
42 testing rack (Figure 1 a and b). The effort an athlete applies to the bar by pushing (IS) or pulling
43 (IMTP) against it is transmitted through the ground and the vertical ground reaction force
44 (vGRF) is recorded. Important methodological factors that can influence the quality of this
45 vGRF data are set up and sampling frequency (5, 14, 19, 31). Additionally, subsequent analysis
46 of the force-time data can be affected by the way in which the start of the pull / push is
47 determined (17). Since not all institution or teams have access to software that can

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48 automatically provide variables of interest, the aim of this article is to provide readers with
49 methodological guidelines for IS and IMTP data collection and a practical guide on how to
50 develop a spreadsheet to analyse the force-time curve to obtain appropriate variables of interest.

51

52 **Variables for Consideration**

53 *Peak Force*

54 Typically, the IS and IMTP are used to identify an athlete's ability to generate maximal force
55 during a given task, this is termed peak force (PF). This variable has been shown to be produced
56 at between 200 and 400 ms of test performance (23, 29) and represents the largest value on the
57 force-time curve (Figure 2). Researchers have found strong relationships between PF and
58 dynamic tasks, such as one repetition maximum (1RM) back squats ($r = 0.79 - 0.97$) (3, 34,
59 51), weightlifting performance ($r = 0.80$) (4, 26), vertical jump height ($r = 0.72 - 0.95$) (29, 34,
60 49), change of direction ($r = -0.657$) and sprint (5, 10, 15 and 20 meters) performance ($r = -$
61 $0.62 - 0.69$) (49). It has been posited that this is because the ability to generate large maximal
62 forces (both absolute and relative) underpins one's capacity to accelerate sports-related mass,
63 typically referring to the athlete during jumping, sprinting, and changing direction (45, 47, 52).
64 Since PF is a highly reliable variable in both IMTP (CV = 1.7% -3.7%, ICC = 0.97- 0.99) (16,
65 18, 26) and IS (CV = 0.9%, ICC = 0.97) (6), and strongly related to performance, this would
66 suggest that the ability to produce high forces is desirable for athletic performance. However,
67 to provide practitioners with valuable information that isoinertial tests cannot, insights into
68 what specific strength quality has improved to contribute to a specific outcome (e.g. jumping
69 higher or throwing an implement further) (3) would typically be related to time. Since, the
70 execution of dynamic athletic tasks such as jumping are generally constrained by time (i.e. the
71 athlete cannot afford to execute the task over more time or would need to reduce the time of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

72 execution), PF alone may not provide enough information about the adaptation of strength
73 qualities. This may be why force at specific time points and the rate of force development
74 (RFD) (12, 34, 38, 39, 42- 44, 46, 47, 49, 51) are useful to include into IS and IMTP force-time
75 data analysis to help maximise insight into an athlete's performance capacity.

76

77 **INSERT FIGURE 2 HERE – FTC IMAGE**

78

79 *Time related force values*

80 Force at specific time points is another metric often reported in the literature (17, 18, 28, 51).
81 This metric simply reports the force value at a given time point and can provide insight about
82 the change in force at time points of interest following training interventions. That is to say that
83 if an athlete is able to produce 800 N of force at 100 ms one would hope for an upward shift
84 following a training intervention focused on speed and power so that they may then achieve
85 950 N at the same time point. Time related force values have demonstrated good reliability
86 (CV = 2.3%-2.7% and 6.2-8.0%, ICC = 0.95-1.00 and 0.921-0.968) (18, 26). Force at 0 -90, -
87 100, -150, -200 and -250 ms have also been shown to have moderate to strong correlations with
88 back squat 1RM (r = 0.757-0.816), (28), CMJ height (r = 0.346), RSImod (r = 0.416-0.426)
89 and 1 RM power clean performance (r = 0.569-0.659) (18). Given its relationship to athletic
90 tasks, and its reliability, force at specific time points may provide a useful insight into the
91 athlete's physical capacity. Following an intervention an increase in this value at given time
92 point (i.e 200 ms) would infer that RFD and impulse (discussed below) would have also
93 increased relative to the onset (i.e baseline) (Figure 2). Since the calculation of RFD amplifies

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

94 signal noise using force at specific time points would introduce less error and simplify data
95 analysis.

96

97 **RFD**

98 RFD describes the gradient (steepness) of the force-time curve and is underpinned by
99 differentiation of force. This is achieved by dividing the change in force (ΔF [where $\Delta =$
100 change]) by the change in time (Δt) (26). The RFD is typically calculated across specific epochs
101 associated with athletic movements as shown in Figure 2. Average RFD (aRFD) is calculated
102 by dividing PF by time to PF. However, this metric has low reliability (CV <15%, ICC = 0.74)
103 (26), potentially due to PF being achieved at different time points between trials, day and
104 individuals. Therefore, to address this and to provide practitioners with a clearer picture of
105 force generating capabilities during times relevant to sporting tasks, utilising pre-determined
106 epochs is a preferred method of analysis (12, 16-19, 26, 28, 47, 51). RFD can be calculated
107 from the initiation of the start of the task (0 ms) to a specific time point (e.g., 0-100 ms) of
108 which the force at 100 ms is then divided by 0.100 (100 ms) (1). In order to define the steepest
109 part of the curve researchers have also utilised peak RFD (pRFD) where epochs of equal length
110 sequentially measure RFD using consecutive time periods up the slope (i.e., 50-100, 100-150
111 ms, etc). Haff et al (26) demonstrated that pRFD is reliable when a 20 ms moving average was
112 used as opposed to 50, 30, 10, 5 and 2 ms, all of which displayed CV's greater than 15% and
113 low ICC's.

114
115 The relationship between measures of RFD and athletic performance have previously
116 demonstrated moderate to strong relationships (3, 49) in tasks such as hang clean 1RM ($r =$

117 0.668- 0.701), vertical jump height ($r = 0.556-0.570$) and 1RM squat, both full and partial ($r =$
118 0.423-0.554), with small and non-significant relationships also having been reported within
119 front squat 1RM ($r = 0.119 - 0.466$), pro and lane agility ($r = -0.179- -0.378$) and
120 countermovement and squat jump (CMJ and SJ, respectively) height ($r = -0.04 - 0.13$), (34,
121 44, 49). This disparity may be attributed to a multitude of factors such as; small epochs being
122 too sensitive and therefore unreliable (31), different onset thresholds and different sampling
123 frequencies utilised in the research. Dos'Santos et al. (16) concluded that to obtain reliable
124 RFD and time-specific force values data could be sampled as low as 500 Hz, however, in order
125 to capture all relevant signals, the authors suggest a capture frequency of 1,000 Hz. Also due
126 to inherent noise experienced from the force plate an onset threshold of pre-pull quiet standing
127 mean + 5 standard deviations (5SD) sufficiently accounts for both signal and human 'noise'
128 (17). However, in fixed laboratory set ups (figure 1a) noise may likely be less and therefore a
129 lower standard deviation may suffice. Since using either an automated method (e.g. + 5
130 standard deviations) or manual inspection to determine the onset will have an impact on time-
131 related force variables more research is needed to determine the best onset method based on
132 the set-up in which the data is being collected.

133

134 Early phase RFD (<50 ms) may be unreliable due to electro-mechanical delay (EMD). This is
135 the time between the onset of muscle activity and the onset of mechanical output. This is best
136 described by Folland et al (21) who utilised a dynamometer to measure a series of voluntary
137 and electrically evoked isometric contractions of the quadriceps. Their findings reported high
138 interindividual variability (CV = 48%) in RFD during 0-50 ms during isometric knee extension,
139 with electrically evoked contractions showing over half this variability in the twitch (CV =
140 22%) and octet (eight pulses at 300 Hz) (CV = 19%) conditions. They conclude that this
141 variability is caused by neural factors, supported by the high interindividual variability (CV =

142 38%) presented in the activation of the quadriceps, as identified through electromyography
143 (EMG). Since EMD is <13 ms for voluntary contractions (48), the advised sampling frequency
144 of 1,000 Hz by the authors makes it less likely to miss any signal relevant to the isometric task.
145 It should also be noted that as a virtue of differentiating the force-time data signal noise is
146 amplified, therefore potentially accounting for the variability often reported in RFD measures.

147

148 *Impulse*

149 Impulse (area under the curve) is of high importance in sporting tasks since it explains the
150 change in momentum (mass × velocity) as a consequence of the product of net force (force less
151 body [and additional external] weight) and how long it is applied for. During isometric tasks,
152 impulse can be calculated using the same approach as that mentioned for RFD, multiplying
153 instead of dividing force by time:

$$154 \underline{F}\Delta t = m\Delta v$$

155 Where F is force, t is time, m is mass, and v is velocity. Since velocity is 0 throughout isometric
156 tasks and should therefore not be presented, the amount of force an athlete can apply and the
157 time over which they can apply it, can provide insight into the athlete's velocity capacity to
158 enable change in their momentum during non-isometric athletic tasks. Therefore, at a constant
159 body mass, a greater impulse at a given epoch would suggest an increase in the athlete's
160 capacity to generate greater movement velocity. Since impulse can be achieved through a
161 variation of high force and less time as well as low force and more time, it is important to
162 understand that during sporting tasks the athlete will rely on the amount of force they can apply
163 with respect to the time they have to apply it. For example, during top end sprinting it is
164 undesirable for the athlete to generate greater vertical impulse through longer ground contact,

165 applying more force in the typically shorter available time is desirable. In lesser time
166 constrained movements, such as throwing an implement (i.e., a shot putt), the athlete who
167 applies the greatest net force over a longer time period would be able to throw the implement
168 further since they have a longer time to apply force to the shot. Relationships between impulse
169 and non-time constrained athletic tasks have been found to have moderate correlations. For
170 example, IS impulse at 250 ms, and at 90-120 degrees knee flexion, both displayed strong to
171 moderate correlation with 1RM squat ($r = 0.7, 0.58$) (3). This is unsurprising because back
172 squat 1RM movement velocity is low (due to the inverse load-velocity relationship). This
173 means that the athlete is therefore required to produce the necessary force over any time period
174 to complete rep (bar and body mass displacement).

175

176 Literature on impulse collected from isolated joint isometrics have reported similar CV and
177 ICC values to that of RFD, with early phase force production (0- 50 ms) shown to be the most
178 variable (CV = 16.6% and 18.7%, ICC= 0.80 and 0.77), although this improves at 0-100 ms
179 (CV= 6.8% and 9.8%, ICC= 0.90 and 0.82) and 100-150 ms (CV= 10.5% and 8.4% , ICC=
180 0.62 and 0.77) for RFD and impulse respectively (10). Guppy et al (22) had also reported that
181 during IMTP, measures of impulse across a range of epochs were more reliable (CV = 4.3 –
182 8.70%, ICC = 0.89 – 0.97) than RFD when assessing different hip and knee angles. It should
183 therefore be noted that as a virtue of differentiating the force-time curve RFD can be more
184 susceptible to increased error and is magnified when using smaller epochs (31). Therefore, the
185 reader may wish to check the reliability of RFD and impulse within their population to help
186 determine which is more reliable, noting that impulse is indicative of sporting actions.

187 **Test Set Up**

188 **Isometric Mid-Thigh Pull**

189 For users to collect PF, RFD and impulse, the correct set up of the IMTP and IS must be
190 considered to ensure testing consistency. The IMTP is a position with high similarities to that
191 of the power position in the clean, whereby the largest vGRF is produced when the torso is in
192 an upright position (20, 23, 26). There has been a variety of set up methods used in the literature
193 for the IMTP ranging from a set position at the mid-point of the knee and hip joint (14, 51),
194 and individualised knee and hip angles based on the clean power position (start of the second
195 pull) (22, 23, 26, 29, 36, 43). Research from Comfort et al (13) compared varying hip and knee
196 angles commonly used in previous research (125° and 145°, and 120°,130°,140° and 150°,
197 respectively) to a self-selected position, where the bar was situated at the mid-thigh. They
198 reported that there were no significant differences between kinetic variables at the differing
199 angles for 100, 200 and 300 ms, but it is worth noting that the self-selected posture, based on
200 the start of the second pull, resulted in joint angles in line with Haff et al (23-25). Within- and
201 between-session reliability was also reported to be highly reliable for all kinetic measures (ICC
202 = 0.849 – 0.993), with impulse at 130° knee flexion and 125° hip flexion the least reliable at
203 all time points (ICC = 0.731-0.739). This may suggest that the hip and knee angle can be self-
204 selected if the athlete is instructed to adopt the posture required for the start of the second pull,
205 and then recorded for reproducibility by the tester. More recent research from Dos' Santos and
206 colleagues (19) investigated the effects of hip angle on PF and RFD. They concluded that to
207 optimise kinetic output and reduce pre-tension during the weighing period (period prior to
208 onset) a hip and knee angle of 145° should be adopted. This is further supported by the findings
209 of Guppy et al (22) who presented greater means across PF, force at specific time points and
210 impulse at specific epochs during traditional IMTP set up (145° at knee and hip) compared to
211 3 alternative positions (145° and 120°, 120° and 125° and 120° and 145° for the knee and hip
212 angles, respectively.). They also reported low to moderate CV's (4 – 11.1%) and high ICC's
213 (0.86 – 0.98) suggesting that utilising a knee and hip position of ~145° may not only optimise

214 kinetic performance, but also provide highly reliable data. Standardising a knee and hip angle
215 utilising a goniometer is best practice because it enables reproducibility of the test within the
216 same individual, which is critical if utilising the IMTP for monitoring purposes. The angle
217 ranges at the knee and hip are presented in Figure 3, and are typical of that previously presented
218 in the literature (14, 22, 23, 26, 29, 36, 43).

219
220 **INSERT FIGURE 3 HERE – IMTP POSITION IMAGE**

221
222 It should also be noted that to enable the athlete to generate maximum force during the IMTP
223 prior research suggests utilising weightlifting straps and where possible athletic tape to strap
224 the hands of the athlete to the bar to prevent the grip being compromised, which may limit the
225 force output (23, 26, 29).

226 227 **Isometric Squat**

228 Different knee and hip angles have been used for the isometric squat, ranging from 90 - 140°
229 of knee flexion and 110 - 140° of hip flexion (9). Research from Marchetti et al. (32) has shown
230 that 90° of knee flexion produces the highest overall muscle activation in the quadriceps,
231 hamstrings, and glutes when compared to 20° and 140°. The limitation of this study, however,
232 was that they only compared muscle activity and not ground reaction force. Therefore, it would
233 be unjust to assume that a higher muscle activity at 90° would produce higher vGRF as the
234 heightened muscle activity may also be due to stabilisation at such depth. Since the IMTP
235 replicates the power position of the clean during which the greatest amounts of force (relative
236 to other positions) have been shown to be produced, it would seem logical to utilise a knee

237 angle $\sim 140^\circ$ for the IS (Figure 4). This has been examined by Brady et al (8) who compared
238 the reliability of specific kinetic variables during IMTP and IS using the same knee and hip
239 angle (knee angle $136 \pm 3^\circ$ and hip angle $137 \pm 2^\circ$). Their results suggested that absolute peak
240 force, relative peak force, allometrically scaled peak force, RFD at 0 – 200 and 250 ms and
241 impulse between 0 – 300 ms were deemed reliable (CV < 10%, ICC > 0.8).

242
243 **INSERT FIGURE 4 HERE – IS POSITION IMAGE **

244
245 It is important that when athletes set up for these tasks that they take a low-level of pre-tension
246 (48), conversely the athlete should not take so much tension that they are actively pushing or
247 pulling against the bar. Since all of the of force acquisition software that the authors are aware
248 of display force-time data as it is recorded and/or at trial completion, the authors suggest that
249 the reader visually inspect the force-time curve and discard any readings that do not appear to
250 have a flat baseline or display an obvious countermovement before force application (Figure 5
251 a and b, respectively). Alternatively, the reader could use a more robust post analysis method
252 of inclusion and exclusion criteria of trials which is later discussed in the “*Exclusion of Trials*”
253 section. Trials with clear countermovement’s should not be used because this will lead to
254 difficulties in determining trial initiation thus compromising the accuracy of time related data.
255 This issue can be avoided by providing clear and concise verbal instruction, instructing the
256 athlete to “get into position” and to “focus on pushing the ground as hard and as fast as
257 possible” (27). Providing these kinds of instructions has been shown to produce significantly
258 greater peak force ($p < 0.001$, ES = 0.33, CI 95%) than internal focus of attention (27).

260 **INSERT FIGURE 5a and b HERE ONTOP OF EACHOTHER – FORCE TRACE

261 EXAMPLE **

262

263 An important factor to consider with the IMTP and IS that is often overlooked is the type of
264 rig (where rig refers to the way the bar is positioned) and bar that is used. Where possible, users
265 are recommended to use a purpose-built isometric rig with minimal to no movement and a cold
266 rolled steel bar. This is to minimise the amount of force lost via bar flexion or rig movement.
267 Not doing this may potentially provide misleading outputs, particularly with respect to RFD
268 and other time related variables. It becomes clear that obtaining force-time data during
269 isometric tasks requires a consistent and reliable method to be used from test set-up to data
270 extraction (14) (refer to instruction table below). Since there is such a large discrepancy in the
271 research on position and reliability, the authors suggest readers use table 1 as a guideline to
272 best fit their practice and to then test the reliability of the chosen isometric task within their
273 own settings.

274

275 **INSERT TABLE 1 HERE – INSTRUCTIONS TO COLLECT IS AND IMTP DATA **

276

277 **Analyzing Data**

278 ***Template Set-up***

279 Firstly, a template should be created so that the raw data can be copied into and the variables
280 of interest calculated. All equations presented in appendix 1 to 3 will relate to the row and
281 column numbers presented in Figure 6, however, readers can adjust the layout as they see fit.

282

283 **INSERT FIGURE 6 HERE – DATA TEMPLATE IMAGE **

284

285 ***Baseline Measure***

286 In order to best collect a quiet baseline, the authors suggest calibration of the force plate without
287 the athlete on it. This will zero the force plate and lessen the likelihood of noise from
288 perturbations generated by the athlete. Secondly, since not all acquisition software's will give
289 a live reading it is pivotal that the athlete is strapped to the bar and is in position in order to
290 obtain a good, stable baseline in which the 5SD is calculated. From this point the acquisition
291 of data can begin and the baseline would likely be very low if 1) they are familiar with the test
292 and 2) have stayed still once in position. Once the data has been reliably collected and the data
293 extracted, the raw force-time data can be pasted into cells A2 and B2. This is important so that
294 subsequent variables can be calculated in column C, labelled "Net Force (N)". If key variables
295 are then derived from the raw force-time data athletes with greater mass will be favoured.
296 Unlike other common athletic tasks collected on a force plate (i.e., countermovement jump),
297 there is no period where body weight can be calculated from the force-time data. The tension
298 taken on the bar will create a vertical force in addition to the athlete's bodyweight. Therefore,
299 an average period of approximately 1-2 s should be calculated from the flattest part of the force-
300 time data before initiation (Figure 5a) (equation b, appendix 1). This now needs to be subtracted
301 from the absolute force in column B, to provide net force in column C.

302

303 ***Defining the start***

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

304 There has been a considerable lack of consistency with the way that the start of isometric tasks
305 has been identified in force-time, with previous literature utilising absolute values (28) and
306 manual detection (45). More recently a comparison of reliability on PF and RFD measures
307 between manual and algorithm-based detection was conducted by Carroll et al. (11). Results
308 showed near perfect reliability for both methods, but manual detection yielded greater CV's
309 for RFD measures, both within (CV = 16.25 - 20.59%) and between testers (CV = 32.27 -
310 42.17%) with greater reliability prevalent in the larger epochs (200ms vs 50ms). Thus, proving
311 problematic when using manual based analysis for detection of significant change. Carroll et
312 al. (11) concluded that utilising an automated start point for isometric testing may provide
313 superior between tester reliability when compared to manual suggesting that different
314 examiners can be used when collecting data, thus potentially saving time and resources.
315 However, it should be noted that the automated start points presented by Carroll et al (11) is
316 not defined within the paper and that further research is needed to determine if this method is
317 more reliable than other methods of detection. With this in mind the authors suggest utilising
318 a five-standard deviation threshold relative to the baseline noise (i.e., when the athlete has taken
319 pre- tension), for multiple reasons, 1) it accounts for any noise or perturbations in force during
320 the quiet standing period, 2) it provides greater certainty that the onset of contraction identifies
321 a true change in force (16), and 3) it requires less time than manually analysing data which may
322 not be conducive to the reader's work flow. Before defining this point, it is suggested that the
323 reader creates a "scatter with smooth lines" graph of the absolute data to determine where a
324 quiet period exists before the initiation of force application. This would be represented by a
325 relatively constant force for a period of no less than one second. From this the reader can hover
326 the cursor over the graph and obtain row values relating to a "Baseline Start" and "Baseline
327 End" and insert these values in cells G4 and G5 respectively.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

328 Now the baseline is determined the starting threshold can be determined using equations c to e
329 (appendix 1). This will calculate and present the average force and 5 SD from this period,
330 summing them to provide the start threshold. This method accounts for both signal and human
331 noise and provides more certainty that the onset of contraction identifies a true change in force,
332 as previously explained. However, some fixed isometric racks may experience less noise,
333 therefore a smaller SD may be used at the user's discretion. Using the MATCH and INDEX
334 functions, the location (row) of this value can be found with its associated time point (equation
335 f and g, respectively. Appendix 1).

336

337 *Exclusion of Trials*

338 During the collection of isometric force-time data practitioners need to consider reducing
339 between trial variation by utilising exclusion criteria. Previous research (30) has utilised a PF
340 difference between trials of 250 N, of which a third trial would be taken should that value be
341 exceeded. This ensures the intent of the individual during the test remains consistent and can
342 also be an indication on whether they have warmed up sufficiently.

343 Since baseline + 5SD is determined as a meaningful change in force (16) for the onset of the
344 isometric task, the authors suggest that using a baseline – 5SD may provide an appropriate
345 method of determining whether a countermovement prior to the initiation of the pull is
346 meaningful enough to discard. Previously methods utilising arbitrary values (16) and manual
347 inspection (46) have been used to discard force-time curves displaying countermovement's.
348 Since there is no agreed method for the exclusion of trials, the authors rational is based on
349 previous literature by Owens et al (37) who looked at the onset of movement during a CMJ.
350 Owens suggested that when vGRF minus BW changed ± 5 SD, meaningful movement had
351 occurred that falls outside of the “noise” of the baseline. Although he went on further to suggest

1 352 that the initiation of the jump should in fact be taken – 30 ms from this identified point to ensure
2 353 velocity is zero, during an isometric task velocity should not to be considered. Therefore, as a
3
4 354 virtue of + 5 SD being meaningful change to determine the onset, - 5SD could be used to
5
6
7 355 determine a meaningful countermovement.
8
9

10 356 The – 5 SD exclusion method can be integrated easily into the sheet to show the user whether
11
12 357 the trial is to be kept or discarded. In column I10 to I14 insert the headings as outlined in figure
13
14
15 358 6. Insert the relevant calculations in J10- J14, this will 1) calculate the baseline – 5 SD (equation
16
17 359 h, appendix 2) and 2) look for the lowest value (equation I, appendix 2) between the end of the
18
19
20 360 baseline (user defined) and the start of the task (equation j and k, appendix 2). From this an IFS
21
22 361 function can be utilised to tell the user whether the minimum value (i.e. the countermovement)
23
24
25 362 is lower than that of the bassline – 5SD (equation l, appendix 2).
26
27

28 363 Alternatively, those with software’s displaying live force-time curves, a threshold baseline of
29
30
31 364 10% bodyweight can be used to ensure that the athlete is holding a stable position without too
32
33 365 much pretension or unloading, therefore potentially pertaining to a better onset, however,
34
35
36 366 further research is needed as to define an acceptable baseline.
37
38

39 367

42 368 *Calculations*

45 369 *Peak Force*

47
48
49 370 Because peak force is the greatest value achieved by the athlete during the task, the MAX
50
51 371 function can be used to find this value from the “Net Force (N)” data range (equation m,
52
53
54 372 appendix 3). Once found, the row it occurs at can be located using a MATCH function
55
56 373 (equation n, appendix 3), this can then be converted to a time in seconds by using the INDEX
57
58
59
60
61
62
63
64
65

374 function (equation o, appendix 3). Finally, the time between the onset and peak force (time to
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

375 peak force) can also be calculated using equations p (Appendix 3).

376

377 *RFD and force at specific time points*

378 Average RFD is the first RFD value that can be calculated by dividing the peak force by the
379 time to peak force (equation q, appendix 3) although this has found to be unreliable (26). To
380 calculate RFD from the onset to pre-determined time points, the reader first needs to determine
381 what time points they are interested in. For this example, we use 100 ms, but once the reader
382 understands how this is done the same methods can be applied to any time interval. In the RFD
383 cell (G23) type in the time frame of interest in seconds (i.e. 100ms is 0.1s). From here 100 ms
384 can be located relative to the onset (equation r, appendix 3), thus allowing the use of a MATCH
385 function to define the cell number (equation s, appendix 3). This can be used to then determine
386 force occurring at 100ms (equation t, appendix 3), which when subtracted from the onset force
387 and divided by the time point of interest, will provide RFD (equation u, appendix 3). For all
388 other time points of interest, readers can repeat this process, alternatively, the time point of
389 interest (cell G23) can be changed, which in turn will change the corresponding RFD values.

390 Peak RFD provides information on the steepest part of the curve using consecutive epochs up
391 the force-time curve (e.g. 100ms to 200ms). To calculate this variable, two-time points must
392 first be defined. If the reader has already set up pre-determined time points for RFD (and force
393 at specific time points) as discussed above, then extraction of peak RFD is simplified. For the
394 epochs of interest simply insert equation v (Appendix 3), where “Fz of above (N)” for each
395 time point is taken away from each other (e.g. Fz of above at 100 ms – Fz of above at 200 ms),
396 divided by the time interval of interest (e.g. 100 ms).

397 *Impulse*

398 Impulse can be calculated from any period of interest (e.g. 0 – 100 ms or 100-200 ms). To do
399 this, force specified at a specific time point can be multiplied by the time point of interest.
400 Alternatively, the trapezoidal rule of calculating the area under the curve can be added within
401 the epoch of interest. In order to achieve this column D must be set up using equation w
402 (Appendix 3) which in turn will calculate impulse for each time point. Then an INDEX function
403 can be used to sum the impulse between two-time points (equation x and y, appendix 3) which
404 were already defined during the RFD calculation, this can be repeated for any other time period.

405

406 **Graphical Representations**

407 Graphical representation of the isometric task can be useful for the reader to plot time points
408 of interest on to the force-time curve, which in turn can help athletes and coaches contextualise
409 the meaning of some of the variables. To do this, first select column C “Fz-BW (N)” and create
410 a “Scatter with smooth lines”. Points of interest, such as peak force and time related variables
411 (e.g. RFD, impulse) can be highlighted within the force-time data by right clicking on the
412 graph, selecting “Select data”, “Add” and adding in the name of the point of interest (e.g. peak
413 force) followed by the cell it is situated in under “Series X values” followed by the value in the
414 “Series Y values”. To make this visible within the force-time curve, right click and select
415 “Change chart type”, where the point of interest “chart type” can be changed to scatter. This
416 will need to be completed for all points of interest should users want them to be define on the
417 force-time curve and when complete will look similar to that depicted in Figure 7.

418

419 ****INSERT FIGURE 7 HERE – POI IMAGE****

420

1
2
3 421 **Summary**
4
5

6 422 When collecting IS or IMTP vGRF data it is clear that there are common variables of interest
7
8
9 423 as reported within the research (3, 17, 18, 26, 28, 34, 51). These include peak force, RFD (and
10
11 424 their associated measures), force at specific time points, and impulse. Generally, it is agreed
12
13
14 425 that a majority of the variables are reliable, with the exception of average RFD. The reliability
15
16 426 of RFD and their associated measures needs to be approached with caution since there is a large
17
18
19 427 disparity within the literature which suggests that earlier time points may not be reliable. The
20
21 428 efficacy of these variables will also be dependent on the onset threshold used, the sampling
22
23
24 429 frequency and the position the isometric task is conducted, as well as the type. The authors
25
26 430 suggest that the reader tests the reliability and sensitivity of extrapolated variables within their
27
28
29 431 own working environment and can do so by collecting multiple trials and checking the
30
31 432 variables CV. Utilising a consistent methodology and the suggestions presented in this article
32
33 433 will help readers ensure they are obtaining the best possible information within their practice.
34
35

36 434

37
38
39
40 435 **Figure 1.** Laboratory fixed isometric rack set up (a) and portable isometric rack set up (b).
41
42

43 436 **Figure 2.** Force Time curve highlighting typical variables of interest.
44
45

46
47 437 **Figure 3.** Isometric Mid-Thigh (IMTP) Pull Set up and force applied by the subject (red
48
49 438 arrows) and the corresponding vGRF (green arrows).
50
51

52 439 **Figure 4.** Isometric Squat (IS) set up and force applied by the subject (red arrows) and the
53
54 440 corresponding vGRF (green arrows)
55
56
57
58
59
60
61
62
63
64
65

441 **Figure 5.** Example of a raw force time curve displaying a steady baseline (a) and a baseline
442 with a countermovement (b), with force on the x axis and cell number on the y axis. Each trial
443 met a peak force of 1826 and 2023 N, respectively. In this instance both trials fall within the
444 250 N exclusion range.

445 **Figure 6.** Data extrapolation template.

446 **Figure 7.** Line graph depicting the points of interest as calculated within the spreadsheet.

447

448

449 ****INSERT APPENDIX TABLES 1-3 AT END****

450 **References**

- 451 1. Aagaard P, Simonsen EB, Andersen JL, Magnusson P, and Dyhre-Poulsen P. Increased
452 rate of force development and neural drive of human skeletal muscle following
453 resistance training. *J Appl Physiol* 93: 1318-1326, 2002.
- 454
455 2. Baker D, Wilson G, and Carlyon B. Generality versus specificity: A comparison of
456 dynamic and isometric measures of strength and speed-strength. *Eur J Appl Physiol*
457 *Occup Physiol* 68: 350-355, 1994.
- 458
459 3. Bazyler CD, Beckham GK, and Sato K. The use of the isometric squat as a measure of
460 strength and explosiveness. *J Strength Cond Res* 29: 1386-1392, 2015.

461

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- 462 4. Beckham G, Mizuguchi S, Carter C, Sato K, Ramsey M, Lamont H, Hornsby G, Haff
463 G, and Stone M. Relationships of isometric mid-thigh pull variables to weightlifting
464 performance. *J Sports Med Phys Fitness* 53: 573-581, 2013.
 - 465
 - 466 5. Beckham GK, Sato K, Mizuguchi S, Haff GG, and Stone MH. Effect of body position
467 on force production during the isometric mid-thigh pull. *J Strength Cond Res.* 32: 48-
468 56: 2018.
 - 469
 - 470 6. Blazevich AJ, Gill N, and Newton RU. Reliability and validity of two isometric squat
471 tests. *J Strength Cond Res* 16: 298-304, 2002.
 - 472
 - 473 7. Bogdanis GC, Tsoukos A, Kaloheri O, Terzis G, Veligekas P, and Brown LE.
474 Comparison between Unilateral and Bilateral Plyometric Training on Single and
475 Double Leg Jumping Performance and Strength. *J Strength Cond Res* 33: 633-640,
476 2019.
 - 477
 - 478 8. Brady CJ, Harrison AJ, Flanagan EP, Haff GG, and Comyns TM. A comparison of the
479 isometric mid-thigh pull and isometric squat: intraday reliability, usefulness and the
480 magnitude of difference between tests. *Int J Sports Physiol Perform* 28: 1-25, 2017.
 - 481
 - 482 9. Brady CJ, Harrison AJ, Comyns TM. A review of the reliability of biomechanical
483 variables produced during the isometric mid-thigh pull and isometric squat and the
484 reporting of normative data. *Sports Biomech* 21: 1-25, 2018.

- 486 10. Buckthorpe MW, Hannah R, Pain MTG, Folland JP. Reliability of neuromuscular
1
2 487 measurements during explosive isometric contractions, with special reference to
3
4 488 electromyography normalisation techniques. *Muscle and Nerve* 46: 566-576, 2012.
5
6
7 489
8
9
10 490 11. Carroll KM, Wagle JP, Sato K, DeWeese BH, Mizuguchi S, and Stone MH. Reliability
11
12 491 of a commercially available and algorithm-based kinetic analysis software compared to
13
14 492 manual-based software. *Sports Biomech* 18:1-9, 2019.
15
16
17 493
18
19 494 12. Comfort P, Graham-Smith P, Matthews MJ, and Bamber C. Strength and power
20
21 495 characteristics in English elite rugby league players. *J Strength Cond Res* 25: 1374-
22
23 1384, 2011.
24 496
25
26 497
27
28
29 498 13. Comfort P, Jones PA, McMahon JJ, and Newton R. Effect of knee and trunk angle on
30
31 499 kinetic variables during the isometric midthigh pull: Test-retest reliability. *Int J Sports*
32
33 *Physiol Perform* 10: 58-63, 2015.
34 500
35
36 501
37
38
39 502 14. Comfort P, Dos'Santos T, Beckham G K, Stone M H, Guppy S N, and Haff GG.
40
41 503 Standardisation and methodological considerations for the isometric mid-thigh pull.
42
43 504 *Strength Cond J*, 2018. [e-pub ahead of print].
44
45
46 505
47
48
49 506 15. De Ruiter CJ, Van Leeuwen D, Heijblom A, Bobbert MF, and De Haan A. Fast
50
51 507 unilateral isometric knee extension torque development and bilateral jump height. *Med*
52
53 508 *Sci Sports Exerc* 38: 1843-1852, 2006.
54
55
56 509
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

510 16. Dos'Santos T, Jones, P A, Kelly J, McMahon J J, Comfort P, and Thomas C. Effect of
511 sampling frequency on isometric midhigh-pull kinetics. *Int J Sports Physiol Perform*
512 11: 255-260, 2016.
513

514 17. Dos'Santos T, Jones, P A, Comfort P, and Thomas C. Effect of Different Onset
515 Thresholds on Isometric Mid-Thigh Pull Force-Time Variables. *J Strength Cond Res*
516 31: 3463-3473, 2017a.
517
518

519 18. Dos'Santos T, Thomas C, Comfort P, McMahon J J, and Jones, P A. Relationship
520 between isometric force-time characteristics and dynamic performance. *Sports* 5: 1 -
521 12, 2017b.
522

523 19. Dos'Santos T, Thomas C, Jones, P A, McMahon J J, and Comfort P. The effect of hip
524 joint angle on isometric mid-thigh pull kinetics. *J Strength Cond Res* 31: 2748-2757,
525 2017c.
526

527 20. Enoka RM. The pull in Olympic weightlifting. *Med Sci Sports* 11: 131-137, 1979.
528

529 21. Folland JP, Buckthorpe MW, and Hannah R. Human capacity for explosive force
530 production: Neural and contractile determinants. *Scand J Med Sci Sports* 24: 894-906,
531 2014.
532

- 533 22. Guppy SN, Brady CJ, Kotani Y, Stone MH, Medic N, and Haff GG. The effect of
1 altering body posture and barbell position on the between-session reliability of force-
2 534 time curve characteristics in the isometric mid-thigh pull. *Sports* 6: 162-177, 2018.
3
4 535
5
6 536
7
8
9 537 23. Haff GG, Stone MH, O'Bryant HS, Harman E, Dinan C, Johnson R, and Han K-H.
10 Force-time dependent characteristics of dynamic and isometric muscle actions. *J*
11 538 *Strength Cond Res* 11: 269-272, 1997.
12
13 539
14
15 540
16
17 541 24. Haff GG, Carlock JM, Hartman MJ, Kilgore JL, Kawamori N, Jackson JR, Morris RT,
18 Sands WA, and Stone MH. Force-time curve characteristics of dynamic and isometric
19 542 muscle actions of elite women Olympic weightlifters. *J Strength Cond Res* 19: 741-
20 543 748, 2005.
21
22 544
23
24 545
25
26 546 25. Haff GG, Jackson JR, Kawamori N, Carlock JM, Hartman MJ, Kilgore JL, Morris RT,
27 Ramsey MW, Sands WA, and Stone MH. Force-time curve characteristics and
28 547 hormonal alterations during an eleven-week training period in elite women
29 548 weightlifters. *J Strength Cond Res* 22: 433-446, 2008.
30
31 549
32 550
33
34 551 26. Haff GG, Ruben RP, Lider J, Twine C, and Cormie P. A comparison of methods for
35 determining the rate of force development during isometric mid-thigh clean pulls. *J*
36 552 *Strength Cond Res* 29: 386-395, 2015.
37
38 553
39
40 554
41
42 555 27. Halperin I, Williams KJ, Martin DT, and Chapman DW. The effects of attentional
43 focusing instructions on force production during the isometric midthigh pull. *J Strength*
44 556 *Cond Res* 30: 919-923, 2016.
45
46 557
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

558

1
2 559 28. James LP, Roberts LA, Haff GG, Kelly VG, and Beckman EM. Validity and reliability
3
4 of a portable isometric mid-thigh clean pull. *J Strength Cond Res* 31: 1378-1386, 2017.
5 560

6
7 561
8
9 562 29. Kawamori N, Rossi SJ, Justice BD, Haff EE, Pistilli EE, O'Bryant HS, Stone MH, and
10
11 Haff GG. Peak force and rate of force development during isometric and dynamic mid-
12 563
13 thigh clean pulls performed at various intensities. *J Strength Cond Res* 20: 483-491,
14 564
15 2006.
16
17 565

18
19 566
20
21 567 30. Kraska JM, Ramsey MW, Haff GG, Fethke N, Sands WA, Stone ME and Stone MH.
22
23 Relationship between strength characteristics and unweighted and weighted vertical
24 568
25 jump height. *Int J Sports Physiol Perform* 4: 461-473, 2009.
26
27 569

28
29 570
30
31 571
32
33 572 31. Maffiuletti N A, Aagaard P, Blazevich A J, Folland J, Tillin N, and Duchateau J. Rate
34
35 of force development: physiological and methodological considerations. *Eur J Appl*
36 573
37 *Physiol* 116: 1091-1116, 2016.
38
39 574

40
41 575
42
43 576 32. Marchetti PH, da Silva JJ, Schoenfeld BJ, Nardi PSM, Pecoraro SL, Greve JMD, and
44
45 Hartigan E. Muscle activation differs between three different knee joint-angle positions
46 577
47 during maximal isometric back squat exercise. *J Sports Med* 2016: 1-6,2016.
48
49 578

50
51 579
52
53 580 33. McGuigan MR, and Winchester JB. The relationship between isometric and dynamic
54
55 strength in college football players. *J Sports Sci Med* 7: 101-105, 2008.
56 581

57
58 582
59
60
61
62
63
64
65

- 583 34. McGuigan MR, Newton MJ, Winchester JB, and Nelson AG. Relationship between
1 isometric and dynamic strength in recreationally trained men. *J Strength Cond Res* 24
2 584 : 2570-2573, 2010.
3
4 585
5
6 586
- 7 587 35. McKinlay BJ, Wallace PJ, Dotan R, Long D, Craig T, Gabriel DA, and Falk B.
8
9 Isometric and dynamic strength and neuromuscular attributes as predictors of vertical
10 588 jump performance in 11-13-year-old male athletes. *Appl Physiol Nutr Metab* 42: 924-
11
12 589 930, 2017.
13
14 590
15
16 591
- 17 592 36. Nuzzo JL, McBride JM, Cormie P, and McCaulley GO. Relationship between
18
19 countermovement jump performance and multijoint isometric and dynamic tests of
20 593 strength. *J Strength Cond Res* 22: 699-707, 2008.
21
22 594
23
24 595
- 25 596 37. Owens NJ, Watkins J, Kilduff LP, Bevan HR, and Bennett MA. Development of a
26
27 criterion method to determine peak mechanical power output in a countermovement
28 597 jump. *J Strength Cond Res* 28: 1552-1558, 2014.
29
30 598
31
32 599
- 33 600 38. Pääsuke M, Ereline J, and Gapeyeva H. Knee extension strength and vertical jumping
34
35 performance in Nordic combined athletes. *J Sports Med Phys Fit* 41: 354-361, 2001.
36 601
37
38 602
- 39 603 39. Rahimi A, Viale F, Dalleau G, and Lacour J-R. Force/velocity and power/velocity
40
41 relationship in squat exercises. *Eur Appl Physiol* 84: 227 – 232, 2001.
42 604
43
44 605
- 45 606 40. Requena B, Gonzalez-Badillo JJ, Villareal ESS, Ereline J, Garcia I, Gapeyeva H, and
46
47 Pääsuke M. Function performance, maximal strength and power characteristics in
48 607
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

608 isometric and dynamic actions of lower extremities in soccer players. *J Strength Cond*
1
2 609 *Res* 23: 1391-1401, 2009.
3
4 610
5
6
7 611
8
9
10 612 41. Sahaly R, Vandewalle H, and Monod DH. Maximal voluntary force and rate of force
11
12 613 development in humans – importance of instruction. *Eur J Appl Physiol* 85: 345-350,
13
14 614 2001.
15
16
17 615
18
19 616 42. Sheppard JM, Chapman D, and Taylor K-L. An evaluation of a strength qualities
20
21 617 assessment method for the lower body. *J Australian Strength Cond* 19: 4-10, 2011.
22
23
24 618
25
26 619 43. Stone MH, Sands WA, Carlock J, Callan S, Dickie D, Daigle K, Cotton J, Smith SL,
27
28
29 620 and Hartman M. The importance of isometric maximum strength and peak rate-of-force
30
31 621 development in sprint cycling. *J Strength Cond Res* 18: 878-884, 2004.
32
33
34 622
35
36 623 44. Thomas C, Jones PA, Rothwell J, Chiang CY, and Comfort P. An investigation into the
37
38
39 624 relationship between maximum isometric strength and vertical jump performance. *J*
40
41 625 *Strength Cond Res* 29: 2176-2185, 2015.
42
43
44 626
45
46 627 45. Thomas C, Dos'Santos T, Comfort P, and Jones PA. Relationship between isometric
47
48
49 628 strength, sprint and change of direction speed in male academy cricketers. *J Trainology*,
50
51 629 5: 18-23, 2016.
52
53 630
54
55
56
57
58
59
60
61
62
63
64
65

- 631 46. Tillin NA, Jimenez-Reyes P, Pain MTG and Folland JP. Neuromuscular performance
1
2 632 of explosive power athletes versus untrained individuals. *Med Sci Sports Exerc* 42: 781-
3
4 633 790, 2010.
5
6
7 634
- 8
9 635 47. Tillin NA, Pain MTG, and Folland JP. Explosive force production during isometric
10
11 636 squats correlates with athletic performance in rugby union players. *J Sport Sci* 31: 66-
12
13 637 76, 2013.
14
15
16 638
- 17
18 639 48. Tillin NA, Pain MTG, and Folland JP. Identification of contraction onset during
19
20 640 explosive contractions. Response to Thompson et al. “Consistency of rapid muscle
21
22 641 force characteristics: influence of muscle contraction onset detection methodology” [J
23
24 642 Electromyogr Kinesiol 2012;22(6):893–900]. *J Electromyogr Kinesiol* 23: 991-994,
25
26 643 2013b.
27
28
29 644
- 30
31 645 49. Townsend JR, Bender D, Vantrease W, Hudy J, Huet K, Williamson C, Bechke E,
32
33 646 Serfini P, and Mangine GT. Isometric mid-thigh pull performance is associated with
34
35 647 athletic performance and sprinting kinetics in division 1 men and women’s basketball
36
37 648 players. *J Strength Cond Res*, 2017. [e-pub ahead of print].
38
39
40
41 649
- 42
43 650 50. Urquhart M, Bishop C, and Turner AN. Validation of a crane scale for the assessment
44
45 651 of portable isometric mid-thigh pulls. *J Australian Strength Cond* 26: 28-33, 2018.
46
47
48
49 652
- 50
51 653 51. Wang R, Hoffman JR, Tanigawa S, Miramonti AA, La Monica MB, Beyer KS, Church
52
53 654 DD, Fukuda DH, and Stout JR. Isometric mid-thigh pull correlates with strength, sprint
54
55
56
57
58
59
60
61
62
63
64
65

- 655 and agility performance in collegiate rugby union players. *J Strength Cond Res* 30:
1
2 656 3051-3056, 2016.
3
4
5 657
6
7 658 52. West DJ, Owen NJ, Jones MR, Bracken M, Cook CJ, Cunningham DJ, Shearer DA,
8
9 Finn CV, Newton RU, Crewther BT, and Kilduff LP. Relationships between force–time
10 659 characteristics of the isometric midhigh pull and dynamic performance in professional
11 660 rugby league players. *J Strength Cond Res* 25: 3070–3075, 2011.
12
13
14 661
15
16
17 662
18
19 663 53. Wilson GJ, Lyttle AD, Ostrowski KJ, and Murphy AJ. Assessing dynamic
20
21 performance: A Comparison of rate of force development tests. *J Strength Cond Res* 9:
22 664 176-181, 1995.
23
24 665
25
26
27 666
28
29 667 54. Wilson GJ, and Murphy AJ. The efficacy of isokinetic, isometric and vertical jump tests
30
31 in exercise science. *Aust J Sci Med Sport* 27:20-24, 1995.
32 668
33
34 669
35
36 670 55. Zaras ND, Stasinaki AN, Methenitis SK, Krase AA, Karampatsos GP, Georgiadis GV,
37
38 Spengos KM, and Terzis GD. Rate of Force Development, Muscle Architecture, and
39 671 Performance in Young Competitive Track and Field Throwers. *J Strength Cond Res*
40
41 672 30: 81-92, 2016.
42
43 673
44
45
46 674
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1. *Instructions to collect IS and IMTP data.*

Isometric Mid-Thigh Pull	Isometric Squat
1) Zero the force plate while the athlete is not stood on it.	
2) Set the bar up so the athlete's knee and hip angles are between 130-145 and 140-145°, respectively (as shown in Figure 3). ** Ensure the athletes grip is optimised by using straps and athletic tape.	2) Set the bar up so the athlete's knee and hip angles are between 90-120 and 120-150°, respectively (as shown in Figure 4).
3) Once the athlete has taken their respective position they should be instructed to remain as still as possible without actively pulling or pushing on the bar. 4) Instruct the athlete that following a countdown from 3 they must “focus on pushing the ground as hard and as fast as possible” and continue doing so until told to stop. 5) Acquire a trial length of ~5 seconds which would include a ~2 second quiet stance and ~3-4 second pull. 6) Provide consistent encouragement throughout trials to ensure maximum effort. 7) Analyse the force-time curve to ensure there is no increase in force or a large countermovement before the pull (refer, to figure 5 a and b).	

Appendix 1 – Excel equations relating to defining the onset of the pull.

Cell Name	Equation	Excel calculation
a. Time Point	=1/Sample Frequency	=1/\$G\$1
b. Baseline weight (N)	= AVERAGE(INDEX(Fz array, Baseline start row):INDEX(Fz array,Baseline end row))	=AVERAGE(INDEX(B:B,G4):INDEX(B:B,G5))
c. F at start (N)	=AVERAGE(INDEX(Fz-BW array, Baseline start row):INDEX(Fz-BW array,Baseline end row))	=AVERAGE(INDEX(C:C,G4):INDEX(C:C,G5))
d. F SD (N)	=STDEV(INDEX(Fz-BW array, Baseline start row):INDEX(Fz-BW array, Baseline end row))	=STDEV(INDEX(C:C,G4):INDEX(C:C,G5))
e. Start of Pull + 5SD (N)	=F at start + (5*Fz SD)	=G10+(5*G11)
f. Start of Pull Cell	=MATCH(Start of Pull + 5SD, Fz-BW array)	=MATCH(G12,C:C)
g. Start of Pull Time (s)	=INDEX(Time array, Start of Pull Cell)	=INDEX(A:A,G13)

*Dollar sign (\$) holds the cell column or row prior to it.

Appendix 2. Excel equations to identify whether the trial should be excluded based on minus five standard deviations of the baseline.

Cell Name	Equation	Excel calculation
h. Baseline – 5SD	=F at Start-(5*F SD (N))	= G10-(5*G11)
i. Min. CMv	=MIN(INDEX(Fz-BL (N) ,Baseline End):INDEX(Fz-BL (N) , Start of Pull Cell))	=MIN(INDEX(C:C,G5):INDEX(C:C,G13))
j. Cells from BL End	=MATCH(Min CMv ,INDEX(Fz-BL (N),Baseline End):INDEX(Fz-BL (N), Start of Pull Cell),-1)	=MATCH(J11,INDEX(C:C,G5):INDEX(C:C,G13),-1)
k. Actual Cell	=Baseline End + Cells from BL End	=G5+J12
l. Outcome	=IFS(Min CMv>Baseline-5SD,"Include", Min CMv < Baseline-5SD,"Exclude")	=IFS(J11>J10,"Include",J11<J10,"Exclude")

Appendix 3 – Excel equations to extract peak force and time related variables.

Cell Name	Equation	Excel calculation
m. Fz Max Value (N)	=MAX(Fz-BW array)	=MAX(C:C)
n. Fz Max Cell	=MATCH(Fz Max Value,Fz-BW array, 0)	=MATCH(G16,C:C,0)
o. Fz Max Time (s)	=INDEX(Time array, Fz Max Cell)	=INDEX(A:A,G17)
p. Time to Fz Max (s)	=Fz Max Time – Start of Pull Time	=G18-G14
RFD, Force at specific time points and Impulse.	All RFD, force at specific time points and impulse measures, with the exception of aRFD can be repeated to extract information on the required epoch's . Alternatively, users can manually change the time value to obtain epochs.	
q. Average RFD (N·s)	=Fz Max Value / Time of Fz Max	=G16/G19
r. Time at above (s)	= Start of Pull Time + RFD @	=\$G\$14+G23
s. Cell of above	=MATCH(Time at above, Time array,1)	=MATCH(G24,A:A,1)

t. Fz of above (N)	=INDEX(Fz-BW array, Cell of above)	=INDEX(C:C,G25)
u. RFD of above (N·s)	=(Fz of above - Start of Pull + 5SD)/RFD @	=(G\$26-G\$12)/G23
v. RFD 0.1 -0.2 s (Ns)	=(Fz of above at RFD 0.1 -Fz of above at RFD 0.2)/Time intervals	=(G33-G26)/G\$44
w. Impulse (Column D)	=(AVERAGE(Fz-BL:Fz-BL))/(1*sample frequency)	=(AVERAGE(C2:C3))/(1*G\$1)
x. Impulse of above (N·s)	=SUM(INDEX(Impulse array, Start of Pull Cell):INDEX(Impulse array, Cell of above))	=SUM(INDEX(D:D,G13):INDEX(D:D,G25))
y. Impulse 0.1-0.2 (Ns)	=SUM(INDEX(Impulse array, Cell of above at RFD 0.1):INDEX(Impulse array, Cell of above at RFD 0.2))	=SUM(INDEX(D:D,G25):INDEX(D:D,G32))

*Dollar sign (\$) holds the cell column or row prior to it.

Figure 1a

[Click here to access/download;Figure;Fig 1a-Lab.jpg](#)








Figure 1b


[Click here to access/download;Figure;Fig 1b-Portable.jpg](#)














Figure 6

[Click here to access/download;Figure;Fig 6- Excel template.jpg](#)

